Title: Integrating patterns of thermal tolerance and phenotypic plasticity with population genetics to improve understanding of vulnerability to warming in a widespread copepod

Running Title: Factors affecting vulnerability to warming

Authors:

Matthew C. Sasaki & Hans G. Dam

University of Connecticut, Department of Marine Sciences, Groton, CT, USA

Corresponding author: matthew.sasaki@uconn.edu

Abstract:

Differences in population vulnerability to warming are defined by spatial patterns in thermal adaptation. These patterns may be driven by natural selection over spatial environmental gradients, but can also be shaped by gene flow, especially in marine taxa with high dispersal potential. Understanding and predicting organismal responses to warming requires disentangling the opposing effects of selection and gene flow. We begin by documenting genetic divergence of thermal tolerance and developmental phenotypic plasticity. Ten populations of the widespread copepod *Acartia tonsa* were collected from sites across a large thermal gradient, ranging from the Florida Keys to Northern New Brunswick, Canada (spanning over 20 degrees latitude). Thermal performance curves from common garden experiments revealed local adaptation at the sampling range extremes, with thermal tolerance increasing at low latitudes and decreasing at high latitudes. The opposite pattern was observed in phenotypic plasticity, which was strongest at high latitudes. Over a large portion of the sampled range, however, we observed a remarkable lack of differentiation of thermal performance curves. To examine whether this lack of divergence is the result of selection for a generalist performance curve or constraint by gene flow, we analyzed cytochrome oxidase I mtDNA sequences, which revealed abundant genetic

diversity and widely-distributed haplotypes. Strong divergence in thermal performance within genetic clades, however, suggests that the pace of thermal adaptation can be relatively rapid. The combined insight from the laboratory physiological experiments and genetic data indicate that gene flow constrains differentiation of thermal performance curves. This balance between gene flow and selection has implications for patterns of vulnerability to warming. Taking both genetic differentiation and phenotypic plasticity into account, our results suggest that local adaptation does not increase vulnerability to warming, and that low latitude populations in general may be more vulnerable to predicted temperature change over the next century.

Keywords:

Climate change, Climate Variability Hypothesis, Copepod, Gene flow, Local adaptation, Macrophysiology, Phenotypic plasticity, Plankton, Thermal performance, Rapid adaptation

Introduction:

Temperature affects processes at every level of biological organization (Angilletta, 2009; Hochachka & Somero, 2002). The rapid warming of the world's oceans (Cheng *et al.*, 2019a; Cheng *et al.*, 2019b; Roemmich *et al.*, 2015; Saba *et al.*, 2016; Wijffels *et al.*, 2016) presents a significant threat to contemporary marine biodiversity (Bryndum-Buchholz *et al.*, 2019; Parmesan, 2006). In addition to the increase in mean ocean temperature, significant increases in the magnitude and frequency of acute events like heat waves have been predicted (Lorenzo & Mantua, 2016; Meehl, 2004; Perkins *et al.*, 2012). These acute events have significant consequences for organisms, and therefore cannot be ignored in predictions of biotic response to climate change (Campbell-Staton *et al.*, 2017; Leicht *et al.*, 2017; Stoks *et al.*, 2017; Sydeman *et al.*, 2013; Ummenhofer & Meehl, 2017; Wernberg *et al.*, 2016). Vulnerability to these climatic changes is established by pre-existing spatial patterns in thermal adaptation. Characterizing

patterns of thermal adaptation and determining their underlying causes is, therefore, directly related to our ability to predict vulnerability and responses of the biota to climate change (Moran *et al.*, 2016; Sorte *et al.*, 2011).

Macrophysiology, the study of variation in physiological traits across space and time (Chown *et al.*, 2004), often yields evidence for adaptation across environmental gradients. Latitudinal thermal gradients, for example, are well-known drivers of local adaptation of thermal tolerance (Addo-Bediako *et al.*, 2000; Castañeda *et al.*, 2015; Gaitán-Espitia *et al.*, 2017; Pereira *et al.*, 2017; Yampolsky *et al.*, 2013). These patterns in adaptation across large spatial scales are often attributed to selection acting on a set of populations, as gene flow is assumed to be relatively low over large distances. This important assumption may not always hold for marine taxa which often have high dispersal potentials (Bowen *et al.*, 2016; Carlton *et al.*, 2017; Cowen *et al.*, 2006; Cowen & Sponaugle, 2009; Gélín *et al.*, 2017; Kinlan & Gaines, 2003; Sexton & Norris, 2008). However, dispersal dynamics in the ocean can be complex (McManus & Woodson, 2012), preventing easy generalization or prediction of connectivity. Instead, genetic markers are often required to estimate levels of connectivity between populations (Palumbi, 2003).

Adaptive genetic differentiation and phenotypic plasticity are two of the main mechanisms used by organisms to cope with variation in the thermal environment. (Angilletta, 2009; Dam, 2013; Magozzi & Calosi, 2014; Somero, 2010; Sparks *et al.*, 2017). Adaptive genetic differentiation is well-known to produce significant variation in phenotypes (Hochachka & Somero, 2002). Phenotypic plasticity, the capacity of a single genotype to produce multiple phenotypes in response to different environmental conditions, can also have large effects (Ayrinhac *et al.*, 2004; Chown *et al.*, 2004, West-Eberhard 2003). Several types of phenotypic plasticity, including acclimation (Stillman, 2003), hardening (Sørensen *et al.*, 2001), and developmental

phenotypic plasticity (Pereira *et al.*, 2017), have all been shown to have strong effects on organismal thermal tolerance. Both mechanisms are likely to play important roles in determining organismal responses to climate change (Hoffmann & Sgro, 2011; Reusch, 2013). Importantly, plasticity acts within generations, and might therefore provide a mechanism for rapid response to environmental variability (Chevin *et al.*, 2010; Chown *et al.*, 2007; Merilä & Hendry, 2014; Seebacher & Grigaltchik, 2014). Plasticity may also prevent the loss of cryptic genetic diversity by shielding genotypes from selection (Friedrich & Meyer, 2016; Pfennig *et al.*, 2010; Schlichting, 2004). This is in stark contrast to selection on standing genetic diversity, which may result in strong demographic bottlenecks and the loss of genetic diversity (Corbett-Detig *et al.*, 2015; Hoffmann & Sgrò, 2011; Kellermann *et al.*, 2009). Studies that examine both mechanisms across large spatial scales are needed.

Characterizing the spatial patterning of phenotypic plasticity and adaptive genetic differentiation is important for predictions of population vulnerability to warming. The Climate Variability Hypothesis (CVH) (Janzen, 1967; Stevens, 1989) posits that thermal tolerance should correspond to the mean temperature experienced by a population whereas phenotypic plasticity should evolve in response to variability in the thermal environment. This hypothesis has accumulated support over time, especially in terrestrial and freshwater systems (Deutsch *et al.*, 2008; Sunday *et al.*, 2010), but still lacks robust experimental validation in the marine realm. It has also been proposed that patterns in the evolution of phenotypic plasticity may be the result of a trade-off between thermal tolerance and the strength of phenotypic plasticity (Stillman, 2003), where higher thermal tolerance evolves at the expense of the capability to modify the phenotype via plasticity.

Patterns in adaptation can, however, also be strongly influenced by gene flow (Lenormand, 2002). The “Gene Flow vs. Selection” issue has been at the heart of evolutionary ecology for decades (Blanquert *et al.* 2013; Slatkin, 1985; 1987). Successful gene flow between populations can strongly impede local adaptation (Garant *et al.*, 2007; Hendry & Taylor, 2004; Lenormand, 2002; Moore *et al.*, 2007; Nosil & Crespi, 2004), and phenotypic divergence is often correlated with the degree of isolation (Mayr, 1963). However, low levels of gene flow might also promote local adaptation by increasing the genetic diversity contained within a population (Garant *et al.*, 2007; Tallmon *et al.*, 2004). The potential for interaction between selection and gene flow makes integrated approaches to studying evolutionary physiology critical for robust characterization of spatial patterns in adaptation. Taking both selection and gene flow into account may be needed to explain observed patterns of adaptation (Dionne *et al.*, 2008; Moore & Hendry 2005).

Spatial patterns in both adaptation and predicted warming interact to produce what is likely to be spatially heterogeneous vulnerability to warming. Understanding which populations are more vulnerable to warming, and why, is critical for effective management and conservation of diversity. Previous work has suggested that warm-adapted, low latitude species or populations are more vulnerable to climate change as they already experiencing temperatures near their thermal maxima (Comte & Olden, 2017; Tewksbury *et al.*, 2008; Vinagre *et al.*, 2016).

However, this is not a universal observation, and populations from mid- to high-latitudes have also been predicted to be more vulnerable (Bennett *et al.*, 2015; Calosi *et al.*, 2008; Fusi *et al.*, 2015). Additionally, these predictions are often based on measurements of thermal tolerance.

This is insufficient, however, as phenotypic plasticity may also play a large role in determining vulnerability to climate change (Burggren, 2018; Chown *et al.*, 2007; Magozzi & Calosi, 2015; Sparks *et al.*, 2017). Examining spatial patterns in both thermal tolerance and the strength of phenotypic plasticity may provide more robust estimates of vulnerability.

Copepods are the most abundant metazoans in the ocean (Humes, 1994; Turner, 2004). They play an important role in transferring energy from primary producers to secondary consumers and are therefore tightly linked to global biogeochemical systems (Menden-Deuer & Kiørboe, 2016), marine trophic webs (Turner, 2004), and commercial fisheries (Castonguay *et al.*, 2008; Dam and Baumann, 2017; Friedland *et al.*, 2012). Studies of thermal adaptation in copepods have a long history (Bradley, 1978; Lonsdale & Levinton, 1985), but local adaptation has largely been ignored, especially in pelagic species (but see Smolina *et al.*, 2016). The calanoid copepod *Acartia tonsa* often dominates coastal and estuarine environments. With a large latitudinal distribution (Turner, 1981) and a life-history amenable to laboratory culturing, this is an ideal model species for studying spatial patterns in adaptation. Previous studies of *Acartia* species have observed local adaptation to several different factors, including exposure to toxic dinoflagellates (Colin & Dam, 2002; 2004), salinity (Plough *et al.*, 2018), and pH (Aguilera *et al.*, 2016). Limited evidence also exists that local adaptation to temperature occurs across large geographic scales (González, 1974; Sasaki *et al.*, 2019). This past work suggests that low latitude populations may be more vulnerable to warming (Sasaki *et al.*, 2019), but conclusions are limited by the lack of spatial coverage and the number of populations used. Additionally, no population genetic information was included. Population genetic studies have shown genetic structuring in this taxon to be complex, with abundant cryptic diversity (Caudill & Bucklin, 2004; Chen & Hare, 2011). The effects of this genetic diversity on patterns of adaptation has largely been ignored (but see Plough *et al.* 2018). Here we examine both large- and fine-scale spatial patterns in the effects of genetic differentiation and phenotypic plasticity on the thermal performance curves (TPCs) of *Acartia tonsa*. We observe clear latitudinal variation in both thermal tolerance and the magnitude of the plastic response, but also find a remarkable lack of divergence over a large portion of the sampled range. Paired with insights from the analysis of

cytochrome oxidase I (COI) mtDNA sequence data, we suggest that this lack of divergence is due to constraint by gene flow rather than selection for a generalist performance curve. These spatial patterns in adaptation will likely affect vulnerability to warming, with southern populations being the most vulnerable.

Methods:

Sampling and Culture Maintenance

Copepods were collected from ten sites spanning 21 degrees of latitude, ranging from the Florida Keys to the Northumberland Strait in Canada (Fig. 1; Table 1). Sites selected cover a range of thermal environments, with mean monthly temperatures ranging from 9.05 – 25.1°C, and average monthly temperature ranges varying between 1.38 – 12.88°C. At each site, samples were collected in surface plankton tows using a 250-µm mesh plankton net with a solid cod end. Water column depth was less than 10 m at all sites. Salinity and surface water temperature were recorded at the time of collection. Within 3 hours of collection, mature *Acartia tonsa* individuals were visually identified using a dissection microscope and sorted into 0.6-µm filtered sea water (FSW), with salinity and temperature adjusted to match collection conditions. Each culture began with more than 1000 mature females and abundant males to ensure fertilization. Additional individuals were preserved in 95% molecular grade ethanol for later genetic analysis. All samples and cultures were then transported by car back to the University of Connecticut, Avery Point campus. Copepods were transported in temperature-controlled containers with particular care to maintain temperature and salinity near collection conditions. Aquarium bubblers were used to keep containers well oxygenated. Copepods were fed with a mixture of a green flagellate (*Tetraselmis* sp.) and a small diatom (*Thalassiosira weissflogii*) during transport. In the laboratory, live cultures were gradually brought to 18°C and 30 practical salinity units (psu) and

then maintained for several generations under constant conditions to minimize the effects of previous environmental acclimation. Aquarium bubblers were used to ensure cultures were well oxygenated. The water of each culture was changed weekly. During this period, cultures were fed *ad libitum* a mixture of a green flagellate (*Tetraselmis* sp.), a small diatom (*Thalassiosira weissflogii*), and a cryptomonad (*Rhodomonas salina*). Phytoplankton were cultured semi-continuously in F/2 medium (without silica for *Tetraselmis* and *Rhodomonas*) with a 12 hr:12 hr light:dark cycle at 18°C. Genetic samples were kept at -80°C until extraction.

Figure 1: Map of sampling locations in the North Atlantic and Gulf of Mexico. The inset is a closer view of the southern sampling sites, to better show the spatial arrangement. Sampled site names are abbreviated: Shediac Bay (SB), St. Andrew (SA), Maine (ME), Connecticut (CT), New Smyrna (NS), St. Petersburg (SP), Manatee River (MR), Punta Gorda (PG), Punta Rasa (PR), and Key Largo (KL). Additional site details can be found in Table 1.

Table 1: Collection site details for each of the ten populations recovered. Site names are provided, along with the two-letter abbreviation used. Water body refers to the general area where each site is located. The latitude of each site is provided, along with the salinity of collection. High salinity refers to sites with a salinity greater than 28 psu, while the low salinity sites both had a salinity of 5 psu. Climatological temperature data for each of the sites are also included. Raw temperature data was acquired from the MODIS-aqua sea surface temperature database and summarized to monthly intervals using CDO (Schulzweida *et al.*, 2009).

Population	Abbr.	Water Body	Latitude	Salinity	Monthly Mean Temp. (°C)	Monthly Maximum Temp. (°C)	Monthly Minimum Temp. (°C)	Monthly Temperature Range (°C)	Monthly Temperature Variance (°C)
Key Largo	KL	Florida - Atlantic	25.2838	High	25.10	29.56	23.73	2.87	1.91
Punta Rasa	PR	Florida - Gulf Coast	26.4835	High	24.79	29.42	19.35	10.47	8.17
Punta Gorda	PG	Florida - Gulf Coast	26.9403	Low	24.69	28.80	20.04	9.05	7.06
Manatee River	MR	Florida - Gulf Coast	27.5056	Low	20.77	21.84	19.87	1.99	1.79
St. Petersburg	SP	Florida - Gulf Coast	27.5073	High	24.02	29.50	17.66	1.38	9.59
New Smyrna	NS	Florida - Atlantic	29.0374	High	22.84	26.71	18.60	8.45	6.35
Connecticut	CT	Long Island Sound	41.3205	High	12.52	18.69	6.55	12.88	17.41
Machias	MA	Gulf of Maine	44.6150	High	9.05	11.87	6.41	5.81	5.77
St. Andrew	SA	Bay of Fundy	45.1000	High	9.37	13.06	5.75	7.72	9.81
Shediac Bay	SB	Northumberland Strait	46.272	High	12.38	15.76	8.31	12.06	19.66

Heat Stresses and Performance Curve Estimation

After two generations, a split-brood common garden experiment was used to examine the effects of genetic differentiation and developmental phenotypic plasticity on the thermal performance curve (TPC) of *Acartia tonsa*. For each population, eggs were collected and randomly split between two developmental temperatures, 18°C and 22°C. All other conditions were kept the same between treatments. Upon reaching maturity, females were collected from both developmental conditions and exposed to a 24-hour acute heat stress. Healthy individuals were placed into 1.5 mL of FSW in a 2.0 mL microfuge tube, which was then partially capped to allow for gas exchange with the atmosphere. Only one female was placed into each tube. After all tubes were filled, they were left to rest for one hour at their respective developmental temperatures. Tubes were then placed into dry baths and exposed to a single temperature, ranging from 18°C to 38°C. The number of females per temperature varied, with fewer individuals exposed to the lowest and highest temperatures (where survivorship was expected to be least variable) and more individuals at intermediate temperatures. At least six females were used per temperature. Each female experienced only one temperature throughout the duration of the assay and was used for only one assay. After 24 hours, tubes were collected, and survivorship was determined visually using a dissection scope. Copepods were marked as alive if there was an active response to external stimuli or visible gut passage movement. Evaporation (and therefore fluctuations in salinity) was negligible during the heat stress.

All statistical analyses were performed using the software package R (R Core Team, 2016). Logistic regressions were used to estimate TPCs for the two developmental treatments for all ten populations. A three-way ANOVA was used to examine the effects of genetic differentiation and phenotypic plasticity on the TPCs (survival as a function of stress temperature, population, and

developmental temperature). An effect of genetic differentiation would be indicated by a significant population term, while an effect of developmental phenotypic plasticity would be indicated by a significant developmental temperature term. Population differences in the strength of phenotypic plasticity would be indicated by a significant population x developmental temperature term (a heterogeneity of slopes test).

Each curve was also summarized by estimation of LD₅₀ values (the temperature at which 50% survivorship would be observed). This is a common metric for thermal tolerance. We define the difference in LD₅₀ values between the two developmental treatments (ΔLD_{50}) as the strength of developmental phenotypic plasticity. Standard error for ΔLD_{50} values were calculated as $\sqrt{(\text{SE}_{18\text{C}}^2 + \text{SE}_{22\text{C}}^2)}$, where SE_{18C} and SE_{22C} are the standard error estimates for LD₅₀ from the 18°C and 22°C developmental temperature groups, respectively. The effect of plasticity was also estimated by calculating the average increase at all survivorship levels from 10% - 90%, which we call $\Delta\text{LD}_{\text{average}}$. This takes into account any potential population-specific non-linearities in the effect of phenotypic plasticity across the TPC. We also estimated LD₁₀ (temperature of 10% survival) and ΔLD_{10} values as estimates of thermal limits.

Climatologies and Correlations

Daily satellite sea-surface temperature (SST) measurements for the years 2000 - 2017 were retrieved for each geographic location from the MODIS-Aqua database (<https://oceancolor.gsfc.nasa.gov/data/aqua/>). Because *A. tonsa* has generation times on the order of weeks to a month, this data was summarized as average monthly climatological parameters (mean, maximum, and minimum temperatures as well as temperature range and variance; Table 1) using the command line Climate Data Operators (CDO; Schulzweida *et al.*, 2009).

The CVH postulates that thermal tolerance (LD_{50}) should respond to some mean representation of the thermal environment, while the strength of the phenotypic plasticity (ΔLD_{50}) should reflect the variation in the thermal environment. To test these predictions, we regressed our metrics of thermal adaptation against the environmental parameters of interest ($LD_{50} \sim$ developmental temperature + monthly mean + monthly maximum + monthly minimum; $\Delta LD_{50} \sim$ monthly temperature variance + monthly temperature range). A linear regression was also used to examine the relationship between thermal tolerance and the strength of plasticity. Trade-offs between thermal tolerance and the strength of phenotypic plastic (Stillman, 2003) would be suggested by a significant negative relationship between LD_{50} and ΔLD_{50} . Correlations between the environmental parameters and latitude, as well as with each other were also examined.

Vulnerability to Warming

The statistical model used to estimate TPCs from the common garden experiment can also be used to predict what would happen if parameters like developmental temperature and stress temperature change. We use this to estimate vulnerability to warming across the ten populations examined. For each population, we estimated the TPC with the mean monthly temperature as the developmental temperature, using the *predict.glm* function in R (*stats* package). Survivorship at the mean monthly maximum temperature was then taken from this modelled TPC. The magnitude of predicted warming at each site over the next century was visually estimated from a high resolution model of warming in the North Atlantic (Saba *et al.* 2016). Future survivorship was estimated in a similar manner using these future temperature values (Supp. Table 1). Vulnerability was estimated as the difference between current and future survivorship values; a positive difference indicates an increase in survivorship (less vulnerable to warming) while a negative value indicates a decrease in survivorship (more vulnerable to warming).

DNA Extraction, Amplification, and Sequencing

Mitochondrial cytochrome-oxidase I (COI) sequence data was generated for individuals from all ten populations (n = 17 - 34 per population). DNA was extracted using a Qiagen Blood and Tissue kit following the manufacturer's instructions. Extracted DNA was eluted in 50 µl of elution buffer (25 µl twice) and stored at -20°C. COI sequences were amplified by polymerase chain reaction (PCR) using mtCOI primers LCO1490 (forward: GGTCAACAAATCATAAAGATATTGG) and HCO2198 (reverse: TAAACTTCAGGGTGACCAAAAAATCA) (Folmer *et al.*, 1994). All PCR reactions were performed in 24 µl volumes with 13 µl ExTaq HS polymerase (Takara Bio Inc.), 1 µL each forward and reverse primers, 5 µl genomic DNA, and 5 µl ultrapure molecular grade water. The optimized PCR protocol began with an initial denaturation of 94°C for 3 minutes followed by 35 cycles of denaturation at 94°C for 45 seconds, annealing at 48°C for 45 seconds, and extension at 72°C for 45 seconds. The protocol ended with a final extension at 72°C for 7 minutes. Amplification success and product length were confirmed visually using a 1.2% agarose gel post-stained with GelRed (Biotium Inc.). Successful amplification products were then purified using an ExoSAP-IT PCR clean-up kit (ThermoFisher Scientific) following manufacturer's instructions before being sent to Eurofins Genomics for forward and reverse strand sequencing.

Sequence Analysis

Consensus sequences were generated for each individual using forward and reverse strands in UniPro (Okonechnikov *et al.*, 2012). Sequences were aligned using Clustal-W (Thompson *et al.*, 1994) and then visually checked. Species identity of each sequence was verified by BLAST search in NCBI's GenBank database (Sayers *et al.*, 2018). Population genetic summary statistics (nucleotide diversity (π), haplotype diversity (Hd), and average number of nucleotide differences

between haplotypes) were calculated using DNaSP v6 (Librado & Rozas, 2009). A haplotype network using an infinite site model was computed using the R package *Pegas* (Paradis, 2010). MigrateN was used to estimate mutation-scaled population size and migration rate values (Beerli, 2006; Beerli & Felsenstein, 2001). The transition-transversion ratio was set to 20. A uniform migration rate prior was set with a minimum of 0 and a maximum of 2000, while the population size prior was set with a minimum of 0.001 and a maximum of 0.1. Analyses entailed a single long MCMC chain with 4 concurrent replicates and a static heating scheme. 10,000,000 steps were recorded with a burn-in of 500,000. All other settings used default values. The number of migrants per generation (N_M) was then calculated as $N_{M(pop1 \rightarrow pop2)} = \Theta_{pop1} * M_{pop1 \rightarrow pop2}$. A linear regression between the number of migrants per generation and the pairwise population difference in LD_{50} was used to investigate constraint of thermal adaptation by gene flow.

Results:

Differentiation of Thermal Performance Curves

A total of 6144 females were used to estimate the thermal performance curves (Fig. 2). The ANOVA results for the TPCs yield a significant effect of both population and developmental temperature (both $p < 2.2 \times 10^{-16}$; Table 2), indicating an influence of both genetic differentiation and developmental phenotypic plasticity, respectively. The significant interaction term between stress temperature and population ($p < 2.2 \times 10^{-16}$) indicates that genetic differentiation results in changes to the shape of the performance curve, while the significant interaction term between population and developmental temperature ($p = 1.45 \times 10^{-7}$) indicates that the strength of phenotypic plasticity differs between populations.

297 Table 2: Results of an ANOVA on the logistic regression of survivorship against stress
 298 temperature, population, and developmental temperature. All terms are statistically significant (*p*
 299 values << 0.0002)

	Df	Deviance	Resid. Df	Resid. Dev	Pr(>Chi)
Stress Temp	1	3024.26	6140	5158.1	< 2.2e-16
Population	9	351.38	6131	4806.8	< 2.2e-16
Dev Temp	1	266.91	6130	4539.8	< 2.2e-16
Stress Temp * Pop	9	102.14	6121	4437.7	< 2.2e-16
Stress Temp * Dev Temp	1	23.95	6120	4413.8	9.90E-07
Pop * Dev Temp	9	49.31	6111	4364.4	1.45E-07
Stress Temp * Pop * Dev Temp	9	32.45	6102	4332	0.0001665

300

301 These results are clearly evident in the TPCs (Fig. 2). Shediad Bay, the northernmost population,
 302 has a TPC shifted towards cooler temperatures while the southernmost population, Key Largo,
 303 has a TPC shifted towards warmer temperatures. Two groups of intermediate TPCs are also
 304 observed, one containing the populations from Manatee River, Punta Gorda, and Punta Rasa,
 305 with the other comprised of all remaining populations. This pattern can also be seen in the LD₅₀
 306 values of the 18°C developmental temperature group (Fig. 3). Differences are reduced in the
 307 22°C developmental temperature group.

308

309 Figure 2: Thermal survivorship curves for each of the ten populations (different colors) and
 310 developmental temperatures (solid vs. dashed lines). Survivorship was measured using
 311 individuals from a split-brood, common garden experiment and a 24-hour acute heat shock.
 312 Survivorship was recorded as binary data (1 = survived, 0 = died), with curves estimated by
 313 logistic regression.

314

315 Figure 3: Reaction norms of thermal tolerance (LD_{50}) for the ten populations, shown in different
316 colors. LD_{50} was calculated as the temperature at which 50% survivorship was observed in the
317 thermal performance curve. Error bars show standard errors. The slope of the individual norms
318 represents the strength of developmental phenotypic plasticity (ΔLD_{50}).

319 Thermal tolerance increased with warmer developmental temperature in all populations,
320 suggesting a ubiquitous effect of developmental phenotypic plasticity. Shediad Bay, the
321 northernmost population had the largest strength of phenotypic plasticity while Manatee River, a
322 population from the Gulf Coast of Florida, had the smallest. The various crossed reaction norms
323 also indicate variation in the strength of plasticity. The effects of developmental phenotypic

plasticity were stronger for thermal tolerance than thermal limits ($\Delta LD_{50} > \Delta LD_{10}$; Supp. Fig. 1). There are no large differences between ΔLD_{50} and $\Delta LD_{average}$ (Supp. Fig. 2), so we will focus only on ΔLD_{50} for the sake of uniformity with the LD_{50} metric of thermal tolerance.

Lack of Differentiation in Thermal Performance Curves

Interestingly, we also observed a striking lack of differentiation in the TPCs of populations from a large portion of the sampling range, indicated by the boxes in Fig. 4. Thermal tolerance values for the St. Petersburg (SP), New Smyrna (NS), Connecticut (CT), Maine (ME), and Saint Andrew (SA) populations, spanning over 20 degrees latitude and originating from drastically different thermal environments, were remarkably similar. This lack of differentiation is also seen in the strength of developmental phenotypic plasticity.

Environmental Correlations

Despite the lack of differentiation between some populations, LD_{50} and ΔLD_{50} were both significantly correlated with latitude ($p = 0.0017$ and 0.008 respectively; Table 3; Fig. 4). ANOVA results show LD_{50} to be significantly correlated with both mean monthly temperature ($p = 3.7 \times 10^{-4}$) and mean monthly minimum temperature ($p = 0.0012$; Table 4; Fig. 5). ΔLD_{50} is significantly correlated with mean monthly temperature variance ($p = 0.015$) but not mean monthly temperature range ($p = 0.58$; Table 4; Fig. 5). Both mean monthly temperature and mean monthly temperature variance were significantly correlated with latitude ($p = 1.81 \times 10^{-6}$ and 0.045 respectively; Supp. Fig. 3, 4) but not with each other ($p = 0.14$; Supp. Fig. 5). There is a significant correlation between LD_{50} and ΔLD_{50} ($p = 0.0023$; Fig. 6).

Table 3: ANOVA results for the correlation between thermal tolerance and the strength of phenotypic plasticity with latitude.

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
LD₅₀					
Dev. Temp.	1	45.923	45.923	23.3632	0.0001834
Latitude	1	27.812	27.812	14.1493	0.0017056
Dev. Temp. x Latitude	1	5.4	5.4	2.7474	0.1168863
Residuals	16	31.45	1.966		
ΔLD₅₀					
Latitude	1	11.1685	11.1685	12.196	0.008171
Residuals	8	7.3261	0.9158		

Table 4: ANOVA results for the correlation between thermal tolerance and the strength of phenotypic plasticity with environmental parameters.

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
LD₅₀					
Dev. Temp.	1	45.923	45.923	37.1471	2.05E-05
Monthly Mean	1	25.722	25.722	20.8062	0.0003746
Monthly Maximum	1	0.693	0.693	0.5603	0.4657218
Monthly Minimum	1	19.704	19.704	15.9388	0.0011775
Residuals	15	18.544	1.236		
ΔLD₅₀					
Monthly Variance	1	10.8028	10.8028	10.3074	0.01485
Monthly Range	1	0.3555	0.3555	0.3392	0.57859
Residuals	7	7.3364	1.0481

350

351 Figure 4: Latitudinal patterns in thermal adaptation. A) Thermal tolerance (LD₅₀) is plotted
 352 against latitude of origin. Filled circles represent thermal tolerance values from the 18°C
 353 developmental temperature while unfilled circles are values from the 22°C developmental
 354 temperature. B) Developmental phenotypic plasticity (Δ LD₅₀) is plotted against latitude of
 355 origin. In both panels, error bars represent standard error. Boxes highlight the latitudinal region
 356 with limited divergence of thermal performance curves.

357

358 Figure 5: Correlation between thermal performance metrics and environmental parameters, as
 359 predicted by the Climate Variability Hypothesis. A) Thermal tolerance (LD₅₀) is correlated with
 360 mean monthly temperature. Filled circles represent thermal tolerance values from the 18°C
 361 developmental temperature while empty circles are values from the 22°C developmental
 362 temperature. B) Phenotypic plasticity (ΔLD₅₀) is correlated with mean monthly temperature
 363 variance. In both panels, error bars represent standard error.

364

365 Figure 6: Linear correlation between thermal tolerance (LD_{50}) and the strength of developmental
366 phenotypic plasticity (ΔLD_{50}) for the ten populations. Error bars represent standard error.

367 Vulnerability to Warming

368 Change in survivorship at mean monthly maximum temperatures ranged from -16% to +9.8%
369 (Supp. Table 1). There was a general trend in vulnerability across latitudes (Fig. 7). Locally
370 adapted populations from low latitudes either saw almost no change (KL and MR) or a reduction
371 in survivorship (PG and PR). The locally adapted population from high latitudes (SB) saw an
372 increase in survivorship. Generally, populations without differentiation of TPCs saw a decrease
373 in survivorship at low latitudes (NS and SP) but an increase in survivorship at high latitudes (CT
374 and ME).

375

376 Figure 7: Vulnerability to predicted warming over the next century. Vulnerability is estimated as
 377 the change in survivorship at the mean monthly maximum temperature if the developmental
 378 temperature is assumed to be the mean monthly temperature. Positive changes represent an
 379 increase in survivorship while negative changes represent a decrease in survivorship. Estimates
 380 of the magnitude of warming and the temperature values used for estimating change in
 381 survivorship for each population can be found in Supp. Table 1.

382 Genetic Diversity and Gene Flow

383 A total of 228 COI sequences (aligned length: 562 base pairs) were used for the genetic analyses.
 384 COI sequences revealed high levels of genetic diversity (Table 5). Large variation between
 385 populations was observed in haplotype diversity (H_d ; 0.189 - 0.941), nucleotide diversity (π ;
 386 0.0029 - 0.079), and the average number of nucleotide differences between the haplotypes (1.47 -
 387 40.46).

Table 5: Population genetic summary statistics, as calculated by DNAsP.

Population	Individuals	Haplotypes	Haplotype Diversity	Nucleotide Diversity	Average Number of Nucleotide Differences
KL	31	2	0.67527	0.00526	2.69
PR	21	11	0.85714	0.03549	18.17
PG	19	1	0.66082	0.01597	8.17
MR	34	5	0.94118	0.07903	40.46
SP	20	8	0.84211	0.0768	39.32
NS	17	6	0.83824	0.00287	1.47
CT	22	8	0.77922	0.0565	26.71
ME	20	11	0.18947	0.02072	10.61
SA	21	18	0.74286	0.03633	18.60
SB	23	3	0.74704	0.01583	8.11

In all, 73 unique haplotypes were identified, which segregated into four major clades (Fig. 8). Each clade was represented in populations from across a large geographic area, and most populations contained individuals from multiple clades. Additionally, we observed several instances of shared haplotypes between geographically distant populations (highlighted in Fig. 8). As exceptions to this, the Key Largo, Punta Gorda, Manatee River, and Shediack Bay populations, each of which were characterized by differentiated thermal performance curves, were all largely represented by a single dominant haplotype each, which was generally not recovered from other populations. The dominant single haplotype is, however, shared between Punta Gorda and Manatee River. Estimates of the number of migrants from MigrateN varied greatly between populations, ranging between 2.6 to 107.7 per generation. There was a negative correlation between the number of migrants exchanged and the pairwise population difference in LD₅₀ values ($p = 0.045$; Supp. Fig. 6). However, this correlation explained only a very small proportion of the variance ($R^2 = 0.07$).

403
404 Figure 8: Mitochondrial COI haplotype network for *Acartia tonsa* sequences recovered from
405 each of the ten populations. Individual circles represent unique haplotypes, with the size of each
406 proportional to the haplotype's frequency. Connecting bar lengths are proportional to the number
407 of base pair differences between haplotypes. Circles are colored based on the sequence
408 population of origin. Shared haplotypes are labelled with the sampling sites they were recovered
409 from. Four distinct clades are observed and labelled A – D.

Discussion:

Thermal tolerance and the strength of phenotypic plasticity both vary strongly among populations of *Acartia tonsa* from across the North Atlantic and Gulf of Mexico. This variation conforms to the predictions of the CVH (Janzen, 1967; Stevens, 1989), but may also be explained by Stillman's hypothesis regarding the trade-off between thermal tolerance and the strength of phenotypic plasticity (Stillman 2003). We also observed a surprising lack of differentiation of thermal performance across a large portion of the sampled range. Mitochondrial COI sequences suggest that this lack of differentiation is due to constraint by gene flow, rather than selection for a generalist performance curve. The patterns we observed in both thermal tolerance and the strength of phenotypic plasticity may result in regional differences in vulnerability to warming, with low latitude populations being the most vulnerable.

Patterns in Thermal Adaptation

Key Largo (KL), the southernmost population, had the highest thermal tolerance of the ten populations examined, followed by three populations from the Gulf of Mexico. The northernmost population, Shediack Bay (SB), had the lowest thermal tolerance but the largest strength of phenotypic plasticity. Manatee River (MR), one of the Gulf of Mexico populations with higher thermal tolerance, had the smallest strength of phenotypic plasticity. The CVH (Janzen, 1967; Stevens, 1989) predicts higher thermal tolerance in warmer environments and increased phenotypic plasticity in more variable environments. The results of our study are consistent with these predictions. We observed significant correlations between I) thermal tolerance and mean monthly temperature and II) the strength of phenotypic plasticity and mean monthly temperature variance. The observed patterns, however, may also be explained by a trade-off between thermal tolerance and phenotypic plasticity (Stillman, 2003). We observed a

significant negative relationship between thermal tolerance and the strength of phenotypic plasticity, but no significant correlation between mean monthly temperature and mean monthly temperature variance. Together, these observations might indicate that the trade-off proposed by Stillman (2003) has a strong influence on shaping patterns of adaptation in this taxon, as there is no covariance between environmental parameters that would drive the observed correlation between LD_{50} and ΔLD_{50} . The few studies that have estimated the strength of selection on phenotypic plasticity generally indicate relatively weak selection (Arnold *et al.*, 2019). It is not implausible, therefore, that observed spatial patterns in phenotypic plasticity might instead be driven by patterns of selection for increased thermal tolerance. However, support for this hypothesis appears to be limited and species specific (Gunderson & Stillman, 2015), warranting investigation in additional species.

The negative relationship between thermal tolerance and the strength of phenotypic plasticity might reflect a role of genetic accommodation or “plasticity first” evolutionary change (Scheiner *et al.*, 2017; Kelly, 2019; Levis & Pfennig, 2016; Levis *et al.*, 2017; Pfennig *et al.*, 2010; Pigliucci *et al.*, 2006; Price *et al.*, 2003; West-Eberhard, 2005; 2003), wherein phenotypic modification by plasticity becomes canalized or fixed by genetic mechanisms, resulting in the loss of capacity for plastic change. The Manatee River - St. Petersburg (MR - SP) comparison best illustrates this in our study. These sites are in close proximity to each other (a feature discussed further below) but exhibit strong differences in both thermal tolerance and the strength of phenotypic plasticity. St. Petersburg had a lower thermal tolerance in the 18°C developmental temperature group, but larger strengths of phenotypic plasticity than Manatee River. Increasing developmental temperature reduces the difference in thermal tolerance between the two populations. If we assume that these two populations share an ancestral phenotype similar to the contemporary St. Petersburg population, the elevated thermal tolerance but reduced plasticity in

Manatee River may reflect the fixation of changes originally induced by phenotypic plasticity in the Manatee River population. However, this is the only population pair where this is observed (the Punta Gorda (PG), Punta Rasa (PR), and Key Largo (KL) populations all had comparable or higher thermal tolerance values but larger strengths of phenotypic plasticity than the Manatee River population), suggesting that there may be several alternative evolutionary mechanisms at play across larger spatial scales.

Spatial Scales of Adaptation - Gene flow vs. Selection

Local adaptation within the range of dispersal, so called microgeographic adaptation (Richardson *et al.*, 2014), has garnered increased attention. “Microgeographic” may be a misleading term in marine systems though, as dispersal kernels can encompass hundreds of kilometers (Cowen & Sponaugle, 2009; Kinlan & Gaines, 2003; Kinlan *et al.*, 2005). The lack of differentiation we observed over a large geographic area, ranging from the Gulf of Mexico to the Bay of Fundy, is in agreement with the expectation of high levels of gene flow in pelagic copepods and other planktonic taxa. The wide distribution of genetic clades, several instances of shared haplotypes between distant populations, and the negative correlation between number of migrants and the pairwise difference in LD₅₀ suggests that gene flow can be strong enough to constrain the adaptive divergence of thermal performance curves in this taxon. Direct dispersal across these large distances within the short generation times of *A. tonsa* is unlikely. Instead, dispersal may follow a stepping-stone model (Kimura & Weiss, 1964), as is common in other coastal and estuarine taxa (Hellberg, 1995; Burrige *et al.*, 2004; Williams *et al.*, 2008; Ragionieri *et al.*, 2010; Crandall *et al.*, 2012).

In stark contrast to this pattern of long-distance connectivity, we also observed significant differentiation of TPCs over spatial scales of less than ten kilometers, suggestive of

microgeographic adaptation. Copepods from the Manatee River (MR) and St. Petersburg (SP) sites were collected from either side of a strong salinity gradient, with the salinity at the St. Petersburg collection site approaching full oceanic levels (32 psu) and the Manatee River site strongly influenced by riverine input (5 psu). Adaptation to salinity has been shown to correspond with reproductive isolation in other populations of *Acartia tonsa* (Plough *et al.*, 2018). In this case, dispersal and gene flow may be strongly limited by this salinity gradient, allowing for the local adaptation of the thermal performance curve in the Manatee River population, while the St. Petersburg population remains under the constraining influence of gene flow from other environments. This is supported by the population genetic data; the Manatee River individuals are almost entirely represented by a single haplotype in Clade B, while St. Petersburg individuals are represented by several different haplotypes in clades A, C, and D. No haplotypes are shared between these two sites, despite their distinct geographic proximity. It should also be noted that the major haplotype found at Manatee River represents the only haplotype found at Punta Gorda (PG), the other low salinity site included in this study. However, unlike in the Manatee River - St. Petersburg comparison, the Punta Gorda (PG) and the adjacent Punta Rasa (PR) populations share a similar TPC, despite being genetically distinct. The apparent local adaptation of both populations suggests that factors other than salinity may also contribute to isolation and the reduction of gene flow between Punta Rasa and other sites.

Interestingly, differentiation of TPCs occurs predominantly within clades rather than between clades, suggesting a rapid rate of thermal adaptation relative to the differentiation of the COI marker region. Each clade contains several distinct thermal phenotypes. Clade A for example contains populations representing three of the distinct groups of TPCs. This appears to be true for both temperature and salinity conditions, as clade B is recovered from warm and fresh sites like Manatee River (MR) and Punta Gorda (PG) as well as cold and high salinity sites like St.

Andrew (SA). The generally large capacity for plasticity observed in *Acartia tonsa* may promote this wide distribution of clades across environmental conditions. Phenotypic plasticity may allow migrants to survive in a wider range of environments, thus increasing gene flow and constraining local adaptation (Crispo, 2008; Thibert-Plante & Hendry, 2011).

Of course, as is the case for all single-gene markers studies, our results must be interpreted with caution. Single-gene markers are vulnerable to incomplete lineage sorting (Nichols, 2001), and may obscure important patterns in genetic structuring. Genome-scale data would provide many more markers for population genomic assessment of structure and gene flow, but for the purposes of this study COI sequences provide robust evidence for where, across the examined range, gene flow may be potentially restricting adaptive divergence of performance curves.

Vulnerability to Climate Change

Because copepods are key components in marine ecosystems and biogeochemical cycles determining their susceptibility to warming is essential for predicting the fate of the oceanic biota in light of climate change. An increasingly large body of literature has recognized the important role phenotypic plasticity may play in determining organismal responses to rapid climate change (Somero, 2010). Much of this literature has focused on acclimation or hardening, but our results show clearly that developmental phenotypic plasticity also deserves increased scrutiny as a factor affecting vulnerability (Burggren, 2018). In most of the populations examined, we observed an almost 1:1 relationship between the increase in thermal tolerance compared with the increase in developmental temperature. Strong phenotypic plasticity like this may reduce vulnerability to rapid climate change by providing a mechanism for correspondingly rapid phenotypic change. However, developmental phenotypic plasticity had a weaker effect on thermal limits than thermal tolerance ($\Delta LD_{10} < \Delta LD_{50}$; Supp. Fig. 1). The apparent constraint of upper thermal limits

suggests that accommodation of future warming by plasticity alone may increase vulnerability to extreme temperature events like heatwaves (Meehl, 2004; Perkins *et al.*, 2012), as the difference between environmental temperature and thermal limits decreases.

In addition to strong phenotypic plasticity, the high levels of genetic diversity, a large potential for gene flow, and the apparently rapid rate of thermal adaptation (when gene flow is weak) would also suggest reduced vulnerability to warming. Selection on standing genetic variation may provide a rapid response to change (Pantel *et al.*, 2015; Barrett & Schluter, 2008; Torda *et al.*, 2017). Plasticity may also play a role in increasing or maintaining migration between populations, indirectly supporting evolutionary rescue (Crispo, 2008). Paired with the high levels of standing genetic diversity observed both within and between populations, increasing migration success may reduce vulnerability. This makes understanding oceanographic effects of climate change an important prerequisite for predicting biotic responses, as changes in ocean current patterns may strongly affect the potential for gene flow between populations. Reductions in gene flow may promote local adaptation, thus reducing vulnerability, while an increase in gene flow could strongly increase vulnerability to climate change if existing local adaptation is eroded by gene swamping (Lenormand, 2002).

Vulnerability will also be affected by the pre-existing spatial patterns in adaptation. Using the current and predicted temperatures at each site, we observed large variation in the potential change in survivorship at maximum temperatures. In the southern range, warming generally had little effect on survivorship at the maximum temperature in locally adapted populations, likely due to accommodation by developmental phenotypic plasticity. While the response was mixed, non-differentiated populations (those constrained by gene flow) may be strongly affected by warming; the St. Petersburg (SP) population saw the largest decrease in survivorship. Both non-

differentiated and locally adapted populations from higher latitudes generally saw increased survivorship. Stronger predicted warming in this region may drive a larger increase in thermal tolerance due to developmental phenotypic plasticity, especially in the northernmost population Shediak Bay (SB), which had the largest strength of phenotypic plasticity. In general, local adaptation to increased temperature does not appear to increase vulnerability to warming, contrary to what has been previously suggested for warm-adapted tropical species (Somero, 2010; Tewksbury *et al.*, 2008; Nguyen *et al.*, 2011). However, this is likely highly regionally specific; the second largest decrease in survivorship was predicted for one of the locally adapted populations, Punta Rasa (PR), from the Gulf of Mexico.

Our analysis incorporates population differences in both thermal tolerance and the strength of phenotypic plasticity to provide a more robust estimate of vulnerability. Previous work also indicated that warm-adapted low latitude populations may be more vulnerable to warming, as they already experience temperatures near their thermal limits (Sasaki *et al.*, 2019). This study refines that prediction with the inclusion of more populations from a wider range of thermal environments, and the integration of population genetic data. However, our analysis cannot account for other important factors that will also likely play a large role in determining vulnerability to climate change. Factors such as changing food quality and quantity (Gregg *et al.*, 2003; Van der Waal *et al.*, 2010; Paul *et al.*, 2015; Hixon & Arts, 2016; Dutkiewicz *et al.*, 2019), changes in predation pressure (Broitman *et al.*, 2009; Rall *et al.*, 2009; De Block *et al.*, 2013; Allan *et al.*, 2015), phenological mismatches between copepods, their prey, and their predators (Edwards & Richardson, 2004; Søreide *et al.*, 2010; Brown *et al.*, 2016), changes in behavior (Kearney *et al.*, 2009; Marshall *et al.*, 2013; Nagelkerken & Munday, 2015), and changing direction or magnitude of gene flow between populations all might strongly shape population vulnerability to climate change. Our analysis also assumes that populations are not undergoing

range shifts or further evolutionary adaptation. Despite these limitations, our results provide a representative baseline estimate of vulnerability incorporating several different adaptive mechanisms.

The major conclusions of this study are possible only through the integration of physiological experiments and molecular ecology. On their own, the thermal performance curves cannot differentiate the potential explanations for the lack of divergence observed across large distances, selection for a generalist performance curve or gene flow over evolutionary timescales. The population genetic insights alone are not enough to infer vulnerability. We demonstrate that tight integration between different fields can provide a more comprehensive understanding of the factors determining vulnerability to climate change.

Acknowledgements:

We thank Dr. Ann Bucklin for helpful comments on the analysis of the molecular data, and Dr. Kendra Daly for assistance sampling in St. Petersburg. Research was supported by grants NSF-OCE 1559180 and CT Sea Grant R/LR-25, a Research Council grant from the University of Connecticut, and graduate research fellowships from the Department of Marine Sciences, University of Connecticut, USA.

References:

Addo-Bediako, A., Chown, S., & Gaston, K. (2000). Thermal tolerance, climatic variability and latitude. *Proceedings of the Royal Society B: Biological Sciences*, 267(1445), 739-745.
<https://doi.org/10.1098/rspb.2000.1065>

594 Aguilera, V. M., Vargas, C. A., Lardies, M. A., & Poupin, M. J. (2016). Adaptive variability to
595 low-pH river discharges in *Acartia tonsa* and stress responses to high pCO₂ conditions. *Marine*
596 *Ecology*, 37(1), 215–226. <https://doi.org/10.1111/maec.12282>

597 Allan, B. J., Domenici, P., Munday, P. L., & McCormick, M. I. (2015). Feeling the heat: the
598 effect of acute temperature changes on predator–prey interactions in coral reef fish. *Conservation*
599 *Physiology*, 3(1), cov011. <https://doi.org/10.1093/conphys/cov011>

600 Angilletta, M. J. (2009). Thermal adaptation: a theoretical and empirical analysis. New York,
601 USA: *Oxford University Press Inc.* DOI:10.1093/acprof:oso/9780198570875.001.1

602 Arnold, P. A., Nicotra, A. B., & Kruuk, L. E. (2019). Sparse evidence for selection on
603 phenotypic plasticity in response to temperature. *Philosophical Transactions of the Royal Society*
604 *B*, 374(1768), 20180185. <https://doi.org/10.1098/rstb.2018.0185>

605 Ayrinhac, A., Debat, V., Gibert, P., & Kister, A. (2004). Cold adaptation in geographical
606 populations of *Drosophila melanogaster*: phenotypic plasticity is more important than genetic
607 variability. *Functional Ecology*, 18(5), 700–706. [https://doi.org/10.1111/j.0269-](https://doi.org/10.1111/j.0269-8463.2004.00904.x)
608 [8463.2004.00904.x](https://doi.org/10.1111/j.0269-8463.2004.00904.x)

609 Barrett, R., & Schluter, D. (2008). Adaptation from standing genetic variation. *Trends in Ecology*
610 *& Evolution*, 23(1), 38–44. <https://doi.org/10.1016/j.tree.2007.09.008>

611 Beerli, P. (2006). Comparison of Bayesian and maximum-likelihood inference of population
612 genetic parameters. *Bioinformatics*, 22(3), 341–345.
613 <https://doi.org/10.1093/bioinformatics/bti803>

614 Beerli, P., & Felsenstein, J. (2001). Maximum likelihood estimation of a migration matrix and
615 effective population sizes in n subpopulations by using a coalescent approach. *Proceedings of the*
616 *National Academy of Sciences*, 98(8), 4563-4568. <https://doi.org/10.1073/pnas.081068098>

617 Bennett, S., Wernberg, T., Joy, B. A., De Bettignies, T., & Campbell, A. H. (2015). Central and
618 rear-edge populations can be equally vulnerable to warming. *Nature Communications*, 6(10280),
619 1-7. <https://doi.org/10.1038/ncomms10280>

620 Blanquart, F., Koltz, O., Nuismer, S. L., & Gandon, S. (2013). A practical guide to measuring
621 local adaptation. *Ecology letters*, 16(9), 1195-1205. <https://doi.org/10.1111/ele.12150>

622 Bowen, B. W., Gaither, M. R., DiBattista, J. D., Iacchei, M., Andrews, K. R., Grant, S. W.,
623 Toonen, R. J., & Briggs, J. C. (2016). Comparative phylogeography of the ocean planet.
624 *Proceedings of the National Academy of Sciences*, 113(29), 7962-7969.
625 <https://doi.org/10.1073/pnas.1602404113>

626 Bradley, B. P. (1978). Genetic and physiological adaptation of the copepod *Eurytemora affinis* to
627 seasonal temperatures. *Genetics*, 90(1), 193-205.

628 Broitman, B. R., Szathmary, P. L., Mislán, K. A. S., Blanchette, C. A., & Helmuth, B. (2009).
629 Predator-prey interactions under climate change: the importance of habitat vs body
630 temperature. *Oikos*, 118(2), 219-224. <https://doi.org/10.1111/j.1600-0706.2008.17075.x>

631 Brown, C. J., Fulton, E. A., Hobday, A. J., Matear, R. J., Possingham, H. P., Bulman, C.,
632 Griffiths, S. P., Lozano-Montes, H., Martin, J. M., Metcalf, S., Okey, T. A., Watson, R., &
633 Richardson, A. J. (2010). Effects of climate-driven primary production change on marine food

- 634 webs: implications for fisheries and conservation. *Global Change Biology*, 16(4), 1194-1212.
635 <https://doi.org/10.1111/j.1365-2486.2009.02046.x>
- 636 Brown, C. J., O'Connor, M. I., Poloczanska, E. S., Schoeman, D. S., Buckley, L. B., Burrows, M.
637 T., Duarte, C. M., Halpern, B. S., Pandolfi, J. M., Parmesan, C., & Richardson, A. J. (2016).
638 Ecological and methodological drivers of species' distribution and phenology responses to
639 climate change. *Global change biology*, 22(4), 1548-1560. <https://doi.org/10.1111/gcb.13184>
- 640 Bryndum-Buchholz, A., Tittensor, D. P., Blanchard, J. L., Cheung, W. W., Coll, M., Galbraith,
641 E. D., Jennings, S., Galbraith, E. D., & Lotze, H. K. (2019). Twenty-first-century climate change
642 impacts on marine animal biomass and ecosystem structure across ocean basins. *Global Change*
643 *Biology*, 25(2), 459–472. <https://doi.org/10.1111/gcb.14512>
- 644 Burggren, W. (2018). Developmental phenotypic plasticity helps bridge stochastic weather
645 events associated with climate change. *Journal of Experimental Biology*, 221(9), jeb161984.
646 <https://doi.org/10.1242/jeb.161984>
- 647 Burridge, C. P., Hurt, A. C., Farrington, L. W., Coutin, P. C., & Austin, C. M. (2004). Stepping
648 stone gene flow in an estuarine-dwelling sparid from south-east Australia. *Journal of Fish*
649 *Biology*, 64(4), 805-819. <https://doi.org/10.1111/j.1095-8649.2004.0347.x>
- 650 Calosi, P., Bilton, D. T., & Spicer, J. I. (2007). Thermal tolerance, acclimatory capacity and
651 vulnerability to global climate change. *Biology letters*, 4(1), 99-102.
652 <https://doi.org/10.1098/rsbl.2007.0408>

653 Campbell-Staton, S. C., Cheviron, Z. A., Rochette, N., Catchen, J., Losos, J. B., & Edwards, S.
654 V. (2017). Winter storms drive rapid phenotypic, regulatory, and genomic shifts in the green
655 anole lizard. *Science*, 357(6350), 495–498. <https://doi.org/10.1126/science.aam5512>

656 Carlton, J. T., Chapman, J. W., Geller, J. B., Miller, J. A., Carlton, D. A., McCuller, M. I.,
657 Treneman, N. C., Steves, B. P., & Ruiz, G. M. (2017). Tsunami-driven rafting: Transoceanic
658 species dispersal and implications for marine biogeography. *Science*, 357(6358), 1402–1406.
659 <https://doi.org/10.1126/science.aao1498>

660 Castañeda, L. E., Rezende, E. L., & Santos, M. (2015). Heat tolerance in *Drosophila subobscura*
661 along a latitudinal gradient: Contrasting patterns between plastic and genetic responses.
662 *Evolution*, 69(10), 2721–2734. <https://doi.org/10.1111/evo.12757>

663 Castonguay, M., Plourde, S., Robert, D., Runge, J. A., & Fortier, L. (2008). Copepod production
664 drives recruitment in a marine fish. *Canadian Journal of Fisheries and Aquatic Sciences*, 65(8),
665 1528–1531. <https://doi.org/10.1139/f08-126>

666 Caudill, C. C., & Bucklin, A. (2004). Molecular Phylogeography and Evolutionary History of the
667 Estuarine Copepod, *Acartia tonsa*, on the Northwest Atlantic Coast. *Hydrobiologia*, 511(1–3),
668 91–102. <https://doi.org/10.1023/b:hydr.0000014032.05680.9d>

669 Chen, G., & Hare, M. P. (2011). Cryptic diversity and comparative phylogeography of the
670 estuarine copepod *Acartia tonsa* on the US Atlantic coast. *Molecular Ecology*, 20(11), 2425–
671 2441. <https://doi.org/10.1111/j.1365-294x.2011.05079.x>

672 Cheng, L., Abraham, J., Hausfather, Z., & Trenberth, K. E. (2019a). How fast are the oceans
673 warming? *Science*, 363(6423), 128–129. <https://doi.org/10.1126/science.aav7619>

674 Cheng, L., Zhu, J., Abraham, J., Trenberth, K. E., Fasullo, J. T., Zhang, B., Yu, F., Wan, L.
675 Chen, X., & Song, X. (2019b). 2018 Continues Record Global Ocean Warming. *Advances in*
676 *Atmospheric Sciences*, 36(3), 249-252. <https://doi.org/10.1007/s00376-019-8276-x>

677 Chevin, L.-M., Lande, R., & Mace, G. M. (2010). Adaptation, Plasticity, and Extinction in a
678 Changing Environment: Towards a Predictive Theory. *PLoS Biology*, 8(4), e1000357 8.
679 <https://doi.org/10.1371/journal.pbio.1000357>

680 Chown, S., Gaston, K., & Robinson, D. (2004). Macrophysiology: large-scale patterns in
681 physiological traits and their ecological implications. *Functional Ecology*. 18(2), 159-167.
682 <https://doi.org/10.1111/j.0269-8463.2004.00825.x>

683 Chown, S. L., Slabber, S., McGeoch, M. A., Janion, C., & Leinaas, H. (2007). Phenotypic
684 plasticity mediates climate change responses among invasive and indigenous arthropods.
685 *Proceedings of the Royal Society B: Biological Sciences*, 274(1625), 2531–2537.
686 <https://doi.org/10.1098/rspb.2007.0772>

687 Colin, S. P., & Dam, H. (2002a). Latitudinal differentiation in the effects of the toxic
688 dinoflagellate *Alexandrium* spp. on the feeding and reproduction of populations of the copepod
689 *Acartia hudsonica*. *Harmful Algae*, 1(1), 113 125. [https://doi.org/10.1016/s1568-9883\(02\)00007-](https://doi.org/10.1016/s1568-9883(02)00007-0)
690 0

691 Colin, S. P., & Dam, H. (2002b). Testing for toxic effects of prey on zooplankton using sole
692 versus mixed diets. *Limnology and Oceanography*. 46(5), 1430-1437.
693 <https://doi.org/10.4319/lo.2002.47.5.1430>

694 Comte, L., & Olden, J. D. (2017). Climatic vulnerability of the world's freshwater and marine
695 fishes. *Nature Climate Change*, 7(10), 1-6. <https://doi.org/10.1038/nclimate3382>

696 Corbett-Detig, R. B., Hartl, D. L., & Sackton, T. B. (2015). Natural selection constrains neutral
697 diversity across a wide range of species. *PLoS Biology*, 13(4), e1002112 25.
698 <https://doi.org/10.1371/journal.pbio.1002112>

699 Cowen, R. K., Paris, C., & Srinivasan, A. (2006). Scaling of connectivity in marine populations.
700 *Science*, 311(5760), 522-527. <https://doi.org/10.1126/science.1122039>

701 Cowen, R. K., & Sponaugle, S. (2009). Larval dispersal and marine population connectivity.
702 *Annual Review of Marine Science*, 1(1), 443-466.
703 <https://doi.org/10.1146/annurev.marine.010908.163757>

704 Crandall, E. D., Treml, E. A., & Barber, P. H. (2012). Coalescent and biophysical models of
705 stepping-stone gene flow in neritid snails. *Molecular Ecology*, 21(22), 5579-5598.
706 <https://doi.org/10.1111/mec.12031>

707 Crispo, E. (2008). Modifying effects of phenotypic plasticity on interactions among natural
708 selection, adaptation and gene flow. *Journal of Evolutionary Biology*, 21(6), 1460-1469.
709 <https://doi.org/10.1111/j.1420-9101.2008.01592.x>

710 Dam, H. G. (2013). Evolutionary adaptation of marine zooplankton to global change. *Annual*
711 *Review of Marine Science*, 5(1), 349-370. [https://doi.org/10.1146/annurev-marine-121211-](https://doi.org/10.1146/annurev-marine-121211-172229)
712 [172229](https://doi.org/10.1146/annurev-marine-121211-172229)

713 Dam, H. G., & Baumann, H. (2017). Climate change, zooplankton and fisheries. *Climate Change*
714 *Impacts on Fisheries and Aquaculture: A Global Analysis*, 2, 851-874.

715 De Block, M., Pauwels, K., Van Den Broeck, M., De Meester, L., & Stoks, R. (2013). Local
716 genetic adaptation generates latitude-specific effects of warming on predator–prey
717 interactions. *Global Change Biology*, 19(3), 689-696. <https://doi.org/10.1111/gcb.12089>

718 Deutsch, C. A., Tewksbury, J. J., Huey, R. B., Sheldon, K. S., Ghalambor, C. K., Haak, D. C., &
719 Martin, P. R. (2008). Impacts of climate warming on terrestrial ectotherms across latitude.
720 *Proceedings of the National Academy of Sciences of the United States of America*, 105(18),
721 6668-6672. <https://doi.org/10.1073/pnas.0709472105>

722 Dionne, M., Caron, F., Dodson, J. J., & Bernatchez, L. (2008). Landscape genetics and
723 hierarchical genetic structure in Atlantic salmon: the interaction of gene flow and local
724 adaptation. *Molecular Ecology*, 17(10), 2382-2396. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-294x.2008.03771.x)
725 [294x.2008.03771.x](https://doi.org/10.1111/j.1365-294x.2008.03771.x)

726 Dutkiewicz, S., Hickman, A. E., Jahn, O., Henson, S., Beaulieu, C., & Monier, E. (2019). Ocean
727 colour signature of climate change. *Nature communications*, 10(1), 578.
728 <https://doi.org/10.1038/s41467-019-08457-x>

729 Edwards, M., & Richardson, A. J. (2004). Impact of climate change on marine pelagic
730 phenology and trophic mismatch. *Nature*, 430(7002), 881. <https://doi.org/10.1038/nature02808>

731 Folmer, O., Hoeh, W., Black, M., & Vrijenhoek, R. (1994). Conserved primers for PCR
732 amplification of mitochondrial DNA from different invertebrate phyla. *Molecular Marine*
733 *Biology and Biotechnology*, 3, 294-299.

734 Friedland, K. D., Stock, C., Drinkwater, K. F., Link, J. S., Leaf, R. T., Shank, B. V., Rose, J. M.,
735 Pilskaln, C. H., & Fogarty, M. J. (2012). Pathways between primary production and fisheries

736 yields of large marine ecosystems. *PLoS ONE*, 7(1), e28945 11.

737 <https://doi.org/10.1371/journal.pone.0028945>

738 Friedrich, S., & Meyer, A. (2016). How plasticity, genetic assimilation and cryptic genetic
739 variation may contribute to adaptive radiations. *Molecular Ecology*, 26(1), 330-350.

740 <https://doi.org/10.1111/mec.13880>

741 Fusi, M., Giomi, F., Babbini, S., Daffonchio, D., McQuaid, C. D., Porri, F., & Cannicci, S.
742 (2015). Thermal specialization across large geographical scales predicts the resilience of
743 mangrove crab populations to global warming. *Oikos*, 124(6), 784-795.

744 <https://doi.org/10.1111/oik.01757>

745 Gaitán-Espitia, J., Bacigalupe, L. D., Opitz, T., Lagos, N. A., Osore, S., & Lardies, M. A.
746 (2017). Exploring physiological plasticity and local thermal adaptation in an intertidal crab along
747 a latitudinal cline. *Journal of Thermal Biology*, 68, 14-20.

748 <https://doi.org/10.1016/j.jtherbio.2017.02.011>

749 Garant, D., Forde, S. E., & Hendry, A. P. (2007). The multifarious effects of dispersal and gene
750 flow on contemporary adaptation. *Functional Ecology*, 21(3), 434-443.

751 <https://doi.org/10.1111/j.1365-2435.2006.01228.x>

752 Gélín, P., Fauvelot, C., Mehn, V., Bureau, S., Rouzé, H., & Magalon, H. (2017). Superclone
753 expansion, long-distance clonal dispersal and local genetic structuring in the coral *Pocillopora*
754 *damicornis* Type β in Reunion Island, South Western Indian Ocean. *PLoS ONE*, 12(1), e0169692
755 19. <https://doi.org/10.1371/journal.pone.0169692>

756 González, J. (1974). Critical thermal maxima and upper lethal temperatures for the calanoid
757 copepods *Acartia tonsa* and *A. clausi*. *Marine Biology*, 27(3), 219-223.
758 <https://doi.org/10.1007/BF00391947>

759 Gregg, W. W., Conkright, M. E., Ginoux, P., O'Reilly, J. E., & Casey, N. W. (2003). Ocean
760 primary production and climate: Global decadal changes. *Geophysical Research Letters*, 30(15).
761 <https://doi.org/10.1029/2003GL016889>

762 Gunderson, A., & Stillman, J. (2015). Plasticity in thermal tolerance has limited potential to
763 buffer ectotherms from global warming. *Proceedings of the Royal Society B: Biological*
764 *Sciences*, 282(1808), 20150401. <https://doi.org/10.1098/rspb.2015.0401>

765 Hellberg, M. E. (1995). Stepping-stone gene flow in the solitary coral *Balanophyllia elegans*:
766 equilibrium and nonequilibrium at different spatial scales. *Marine Biology*, 123(3), 573-581.
767 <https://doi.org/10.1007/BF00349236>

768 Hendry, A. P., & Taylor, E. B. (2004). How much of the variation in adaptive divergence can be
769 explained by gene flow? An evaluation using lake-stream stickleback pairs. *Evolution*, 58(10),
770 2319-2331. <https://doi.org/10.1111/j.0014-3820.2004.tb01606.x>

771 Hixson, S. M., & Arts, M. T. (2016). Climate warming is predicted to reduce omega-3, long-
772 chain, polyunsaturated fatty acid production in phytoplankton. *Global Change Biology*, 22(8),
773 2744-2755. <https://doi.org/10.1111/gcb.13295>

774 Hochachka, P. W., & Somero, G. N. (2002). Biochemical adaptation: mechanism and process in
775 physiological evolution. New York, USA: Oxford University Press

776 Hoffmann, A. A., & Sgrò, C. (2011). Climate change and evolutionary adaptation. *Nature*,
777 470(7335), 479-485. <https://doi.org/10.1038/nature09670>

778 Humes A.G. (1994). How many copepods? In: Ferrari F.D., Bradley B.P. (eds) Ecology and
779 Morphology of Copepods. Developments in Hydrobiology, vol 102. Springer, Dordrecht

780 Janzen, D. (1967). Why mountain passes are higher in the tropics. *The American Naturalist*.
781 101(919), 233-249. <https://doi.org/10.1086/282487>

782 Kearney, M., Shine, R., & Porter, W. P. (2009). The potential for behavioral thermoregulation to
783 buffer “cold-blooded” animals against climate warming. *Proceedings of the National Academy of*
784 *Sciences*, 106(10), 3835-3840. <https://doi.org/10.1073/pnas.0808913106>

785 Kellermann, V., van Heerwaarden, B., Sgrò, C. M., & Hoffmann, A. A. (2009). Fundamental
786 evolutionary limits in ecological traits drive *Drosophila* species distributions. *Science*,
787 325(5945), 1244–1246. <https://doi.org/10.1126/science.1175443>

788 Kelly, M. (2019). Adaptation to climate change through genetic accommodation and assimilation
789 of plastic phenotypes. *Philosophical Transactions of the Royal Society B*, 374(1768), 20180176.
790 <https://doi.org/10.1098/rstb.2018.0176>

791 Kimura, M., & Weiss, G. (1964). The stepping stone model of population structure and the
792 decrease of genetic correlation with distance. *Genetics*, 49(4), 561-576.

793 Kinlan, B. P., & Gaines, S. D. (2003). Propagule dispersal in marine and terrestrial
794 environments: a community perspective. *Ecology*, 84(8), 2007-2020. [https://doi.org/10.1890/01-](https://doi.org/10.1890/01-0622)
795 0622

796 Kinlan, B. P., Gaines, S. D., & Lester, S. E. (2005). Propagule dispersal and the scales of marine
797 community process. *Diversity and Distributions*, 11(2), 139–148. <https://doi.org/10.1111/j.1366->
798 9516.2005.00158.x

799 Leicht, K., Seppälä, K., & Seppälä, O. (2017). Potential for adaptation to climate change: family-
800 level variation in fitness-related traits and their responses to heat waves in a snail population.
801 *BMC Evolutionary Biology*, 17(1). <https://doi.org/10.1186/s12862-017-0988-x>

802 Lenormand, T. (2002). Gene flow and the limits to natural selection. *Trends in Ecology &*
803 *Evolution*, 17(4), 183–189. [https://doi.org/10.1016/s0169-5347\(02\)02497-7](https://doi.org/10.1016/s0169-5347(02)02497-7)

804 Levis, N. A., & Pfennig, D. W. (2016). Evaluating ‘plasticity-first’ evolution in nature: key
805 criteria and empirical approaches. *Trends in Ecology & Evolution*, 31(7), 563-574.
806 <https://doi.org/10.1016/j.tree.2016.03.012>

807 Levis, N., Serrato-Capuchina, A., & Pfennig, D. (2017). Genetic accommodation in the wild:
808 evolution of gene expression plasticity during character displacement. *Journal of Evolutionary*
809 *Biology*, 30(9), 1712–1723. <https://doi.org/10.1111/jeb.13133>

810 Librado, P., & Rozas, J. (2009). DnaSP v5: a software for comprehensive analysis of DNA
811 polymorphism data. *Bioinformatics*, 25(11), 1451-1452.
812 <https://doi.org/10.1093/bioinformatics/btp187>

813 Lonsdale, D. J., & Levinton, J. S. (1985). Latitudinal differentiation in copepod growth: an
814 adaptation to temperature. *Ecology*, 66(5), 1397–1403. <https://doi.org/10.2307/1938002>

815 Lorenzo, E., & Mantua, N. (2016). Multi-year persistence of the 2014/15 North Pacific marine
816 heatwave. *Nature Climate Change*, 6(11), 1042–1047. <https://doi.org/10.1038/nclimate3082>

- 817 Magozzi, S., & Calosi, P. (2014). Integrating metabolic performance, thermal tolerance, and
818 plasticity enables for more accurate predictions on species vulnerability to acute and chronic
819 effects of global warming. *Global Change Biology*, 21(1), 181-194.
820 <https://doi.org/10.1111/gcb.12695>
- 821 Marshall, D. J., Baharuddin, N., & McQuaid, C. D. (2013). Behaviour moderates climate
822 warming vulnerability in high-rocky-shore snails: interactions of habitat use, energy
823 consumption and environmental temperature. *Marine biology*, 160(9), 2525-2530.
824 <https://doi.org/10.1007/s00227-013-2245-1>
- 825 Mayr, E. (1963). Animal species and evolution. Cambridge, USA: *Harvard University Press*.
- 826 McManus, M., & Woodson, B. C. (2012). Plankton distribution and ocean dispersal. *Journal of*
827 *Experimental Biology*, 215(6), 1008–1016. <https://doi.org/10.1242/jeb.059014>
- 828 Meehl, G. (2004). More intense, more frequent, and longer lasting heat waves in the 21st
829 century. *Science*, 305(5686), 994-997. <https://doi.org/10.1126/science.1098704>
- 830 Menden-Deuer, S., & Kiørboe, T. (2016). Small bugs with a big impact: linking plankton
831 ecology with ecosystem processes. *Journal of Plankton Research*, 38(4), 1036–1043.
832 <https://doi.org/10.1093/plankt/fbw049>
- 833 Merilä, J., & Hendry, A. P. (2014). Climate change, adaptation, and phenotypic plasticity: the
834 problem and the evidence. *Evolutionary Applications*, 7(1), 1-14.
835 <https://doi.org/10.1111/eva.12137>

836 Moore, J.-S., Gow, J. L., Taylor, E. B., & Hendry, A. P. (2007). Quantifying the constraining
837 influence of gene flow on adaptive divergence in the lake-stream threespine stickleback system.
838 *Evolution*, 61(8), 2015–2026. <https://doi.org/10.1111/j.1558-5646.2007.00168.x>

839 Moore, J.-S., & Hendry, A. P. (2005). Both selection and gene flow are necessary to explain
840 adaptive divergence: evidence from clinal variation in stream stickleback. *Evolutionary Ecology*
841 *Research*, 7, 871–886.

842 Moran, E. V., Hartig, F., & Bell, D. M. (2016). Intraspecific trait variation across scales:
843 implications for understanding global change responses. *Global Change Biology*, 22(1), 137–
844 150. <https://doi.org/10.1111/gcb.13000>

845 Nagelkerken, I., & Munday, P. L. (2016). Animal behaviour shapes the ecological effects of
846 ocean acidification and warming: moving from individual to community-level responses. *Global*
847 *Change Biology*, 22(3), 974–989. <https://doi.org/10.1111/gcb.13167>

848 Nguyen, K. D. T., Morley, S. A., Lai, C. H., Clark, M. S., Tan, K. S., Bates, A. E., & Peck, L. S.
849 (2011). Upper temperature limits of tropical marine ectotherms: global warming
850 implications. *PLoS One*, 6(12), e29340. <https://doi.org/10.1371/journal.pone.0029340>

851 Nichols, R. (2001). Gene trees and species trees are not the same. *Trends in Ecology &*
852 *Evolution*, 16(7), 358–364. [https://doi.org/10.1016/s0169-5347\(01\)02203-0](https://doi.org/10.1016/s0169-5347(01)02203-0)

853 Nosil, P., & Crespi, B. (2004). Does gene flow constrain adaptive divergence or vice versa? A
854 test using ecomorphology and sexual isolation in *Timema cristinae* walking-sticks. *Evolution*,
855 58(1), 102–112. <https://doi.org/10.1111/j.0014-3820.2004.tb01577.x>

856 Okonechnikov, K., Golosova, O., & Fursov, M. (2012). Unipro UGENE: a unified
857 bioinformatics toolkit. *Bioinformatics*, 28(8), 1166–1167.
858 <https://doi.org/10.1093/bioinformatics/bts091>

859 Palumbi, S. (2003). Population genetics, demographic connectivity, and the design of marine
860 reserves. *Ecological Applications*, 13(1), S146–S158. [https://doi.org/10.1890/1051-0761\(2003\)013\[0146:PGDCAT\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2003)013[0146:PGDCAT]2.0.CO;2)

862 Pantel, J. H., Duvivier, C., & Meester, L. (2015). Rapid local adaptation mediates zooplankton
863 community assembly in experimental mesocosms. *Ecology Letters*, 18(10), 992–1000.
864 <https://doi.org/10.1111/ele.12480>

865 Paradis, E. (2010). pegas: an R package for population genetics with an integrated–modular
866 approach. *Bioinformatics*, 26(3), 419–420. <https://doi.org/10.1093/bioinformatics/btp696>

867 Parmesan, C. (2006). Ecological and evolutionary responses to recent climate change. *Annual
868 Review of Ecology, Evolution, and Systematics*, 37(1), 637–669.
869 <https://doi.org/10.1146/annurev.ecolsys.37.091305.110100>

870 Paul, C., Matthiessen, B., & Sommer, U. (2015). Warming, but not enhanced CO₂ concentration,
871 quantitatively and qualitatively affects phytoplankton biomass. *Marine Ecology Progress
872 Series*, 528, 39–51. <https://doi.org/10.3354/meps11264>

873 Pereira, R. J., Sasaki, M. C., & Burton, R. S. (2017). Adaptation to a latitudinal thermal gradient
874 within a widespread copepod species: the contributions of genetic divergence and phenotypic
875 plasticity. *Proceedings of the Royal Society B*, 284(1853), 20170236.
876 <https://doi.org/10.1098/rspb.2017.0236>

877 Perkins, S., Alexander, L., & Nairn, J. (2012). Increasing frequency, intensity and duration of
878 observed global heatwaves and warm spells. *Geophysical Research Letters*, 39(20), L20714.
879 <https://doi.org/10.1029/2012gl053361>

880 Pfennig, D. W., Wund, M. A., Snell-Rood, E. C., Cruickshank, T., Schlichting, C. D., &
881 Moczek, A. P. (2010). Phenotypic plasticity's impacts on diversification and speciation. *Trends*
882 *in Ecology & Evolution*, 25(8), 459-467. <https://doi.org/10.1016/j.tree.2010.05.006>

883 Pigliucci, M., Murren, C. J., & Schlichting, C. D. (2006). Phenotypic plasticity and evolution by
884 genetic assimilation. *Journal of Experimental Biology*, 209(12), 2362–2367.
885 <https://doi.org/10.1242/jeb.02070>

886 Plough, L., Fitzgerald, C., Plummer, A., & Pierson, J. (2018). Reproductive isolation and
887 morphological divergence between cryptic lineages of the copepod *Acartia tonsa* in Chesapeake
888 Bay. *Marine Ecology Progress Series*, 597, 99–113. <https://doi.org/10.3354/meps12569>

889 Price, T. D., Qvarnström, A., & Irwin, D. E. (2003). The role of phenotypic plasticity in driving
890 genetic evolution. *Proceedings of the Royal Society of London. Series B: Biological Sciences*,
891 270(1523), 1433–1440. <https://doi.org/10.1098/rspb.2003.2372>

892 R Core Team. 2016 R: A language and environment for statistical computing. R Foundation for
893 Statistical Computing, Vienna, Austria. URL: <https://www.R-project.org/>

894 Ragionieri, L., Cannicci, S., Schubart, C. D., & Fratini, S. (2010). Gene flow and demographic
895 history of the mangrove crab *Neosarmatium meinerti*: A case study from the western Indian
896 Ocean. *Estuarine, Coastal and Shelf Science*, 86(2), 179-188.
897 <https://doi.org/10.1016/j.ecss.2009.11.002>

898 Rall, B. C., Vucic-Pestic, O., Ehnes, R. B., Emmerson, M., & Brose, U. (2010). Temperature,
899 predator–prey interaction strength and population stability. *Global Change Biology*, 16(8), 2145-
900 2157. <https://doi.org/10.1111/j.1365-2486.2009.02124.x>

901 Reusch, T. B. (2014). Climate change in the oceans: evolutionary versus phenotypically plastic
902 responses of marine animals and plants. *Evolutionary applications*, 7(1), 104-122.
903 <https://doi.org/10.1111/eva.12109>

904 Richardson, J. L., Urban, M. C., Bolnick, D. I., & Skelly, D. K. (2014). Microgeographic
905 adaptation and the spatial scale of evolution. *Trends in Ecology & Evolution*, 29(3), 165-176.
906 <https://doi.org/10.1016/j.tree.2014.01.002>

907 Roemmich, D., Church, J., Gilson, J., Monselesan, D., Sutton, P., & Wijffels, S. (2015).
908 Unabated planetary warming and its ocean structure since 2006. *Nature Climate Change*, 5(3),
909 240-245. <https://doi.org/10.1038/nclimate2513>

910 Saba, V. S., Griffies, S. M., Anderson, W. G., Winton, M., Alexander, M. A., Delworth, T. L.,
911 Hare, J. A., Harrison, M. J., Rosati, A., Vecchi, G. A., & Zhang, R. (2016). Enhanced warming
912 of the Northwest Atlantic Ocean under climate change. *Journal of Geophysical Research:*
913 *Oceans*, 121(1), 118–132. <https://doi.org/10.1002/2015jc011346>

914 Sasaki, M. C., Hedberg, S., Richardson, K., & Dam, H. (2019). Complex interactions between
915 local adaptation, phenotypic plasticity and sex affect vulnerability to warming in a widespread
916 marine copepod. *Royal Society Open Science*, 6(3), 182115 9.
917 <https://doi.org/10.1098/rsos.182115>

918 Sayers, E. W., Cavanaugh, M., Clark, K., Ostell, J., Pruitt, K. D., & Karsch-Mizrachi, I. (2018).
919 GenBank. *Nucleic Acids Research*, 47(D1), 94-99. <https://doi.org/10.1093/nar/gky989>

920 Scheiner, S., Barfield, M., & Holt, R. D. (2017). The genetics of phenotypic plasticity. XV.
921 Genetic assimilation, the Baldwin effect, and evolutionary rescue. *Ecology and Evolution*,
922 19(21), 8788-8803. <https://doi.org/10.1002/ece3.3429>

923 Schlichting, C. D. (2004). The role of phenotypic plasticity in diversification. In: T. J. DeWitt &
924 S. M. Scheiner (Eds.), *Phenotypic Plasticity: Functional and Conceptual Approaches* (pp. 191-
925 200). New York, USA: Oxford University Press.

926 Schulzweida, U., Kornblueh, L., & Quast, R. (2009). Climate Data Operators. User's Guide.

927 Seebacher, F., & Grigaltchik, V. S. (2014). Embryonic developmental temperatures modulate
928 thermal acclimation of performance curves in tadpoles of the frog *Limnodynastes peronii*. *PLoS*
929 *ONE*, 9(9), e106492. <https://doi.org/10.1371/journal.pone.0106492>

930 Sexton, P. F., & Norris, R. D. (2008). Dispersal and biogeography of marine plankton: Long-
931 distance dispersal of the foraminifer *Truncorotalia truncatulinoides*. *Geology*, 36(11), 899-902.
932 <https://doi.org/10.1130/g25232a.1>

933 Slatkin, M. (1985). Rare alleles as indicators of gene flow. *Evolution*, 39(1), 53-65.
934 <https://doi.org/10.1111/j.1558-5646.1985.tb04079.x>

935 Slatkin, M. (1987). Gene flow and the geographic structure of natural populations. *Science*,
936 236(4803), 787-792. <https://doi.org/10.1126/science.3576198>

- 937 Smolina, I., Harmer, R., Lindeque, P., & Hoarau, G. (2016). Reduced up-regulation of gene
938 expression in response to elevated temperatures in the mid-Atlantic population of *Calanus*
939 *finmarchicus*. *Journal of Experimental Marine Biology and Ecology*, 485, 88–93.
940 <https://doi.org/10.1016/j.jembe.2016.09.003>
- 941 Somero, G. N. (2010). The physiology of climate change: how potentials for acclimatization and
942 genetic adaptation will determine “winners” and “losers”. *The Journal of Experimental Biology*,
943 213(6), 912-920. <https://doi.org/10.1242/jeb.037473>
- 944 Søreide, J. E., Leu, E. V. A., Berge, J., Graeve, M., & Falk-Petersen, S. T. I. G. (2010). Timing
945 of blooms, algal food quality and *Calanus glacialis* reproduction and growth in a changing
946 Arctic. *Global change biology*, 16(11), 3154-3163. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2486.2010.02175.x)
947 2486.2010.02175.x
- 948 Sørensen, J. G., Dahlgaard, J., & Loeschcke, V. (2001). Genetic variation in thermal tolerance
949 among natural populations of *Drosophila buzzatii*: down regulation of Hsp70 expression and
950 variation in heat stress resistance traits. *Functional Ecology*, 15(3), 289-296.
951 <https://doi.org/10.1046/j.1365-2435.2001.00525.x>
- 952 Sorte, C. J., Jones, S. J., & Miller, L. P. (2011). Geographic variation in temperature tolerance as
953 an indicator of potential population responses to climate change. *Journal of Experimental*
954 *Marine Biology and Ecology*, 400(1), 209-217. <https://doi.org/10.1016/j.jembe.2011.02.009>
- 955 Sparks, M. M., Westley, P. A., Falke, J. A., & Quinn, T. P. (2017). Thermal adaptation and
956 phenotypic plasticity in a warming world: Insights from common garden experiments on Alaskan
957 sockeye salmon. *Global Change Biology*, 23(12), 5203–5217. <https://doi.org/10.1111/gcb.13782>

958 Stevens, G. (1989). The latitudinal gradient in geographical range: how so many species coexist
959 in the tropics. *American Naturalist*, 133(2), 240-256. <https://doi.org/10.1086/284913>

960 Stillman, J. H. (2003). Acclimation capacity underlies susceptibility to climate change. *Science*,
961 301(5629), 65–65. <https://doi.org/10.1126/science.1083073>

962 Stoks, R., Verheyen, J., Dievel, M., & Tüzün, N. (2017). Daily temperature variation and
963 extreme high temperatures drive performance and biotic interactions in a warming world.
964 *Current Opinion in Insect Science*, 23, 35–42. <https://doi.org/10.1016/j.cois.2017.06.008>

965 Sunday, J., Bates, A., & Dulvy, N. (2010). Global analysis of thermal tolerance and latitude in
966 ectotherms. *Proceedings of the Royal Society B: Biological Sciences*, 278(1713), 1823–1830.
967 <https://doi.org/10.1098/rspb.2010.1295>

968 Sydemann, W. J., Santora, J. A., Thompson, S., Marinovic, B., & Lorenzo, E. (2013). Increasing
969 variance in North Pacific climate relates to unprecedented ecosystem variability off California.
970 *Global Change Biology*, 19(6), 1662-1675. <https://doi.org/10.1111/gcb.12165>

971 Tallmon, D. A., Luikart, G., & Waples, R. S. (2004). The alluring simplicity and complex reality
972 of genetic rescue. *Trends in Ecology & Evolution*, 19(9), 489–496.
973 <https://doi.org/10.1016/j.tree.2004.07.003>

974 Tewksbury, J. J., Huey, R. B., & Deutsch, C. A. (2008). Putting the heat on tropical
975 animals. *Science*, 320(5881), 1296-1297. <https://doi.org/10.1126/science.1159328>

976 Thibert-Plante, X., & Hendry, A. (2011). The consequences of phenotypic plasticity for
977 ecological speciation. *Journal of Evolutionary Biology*, 24(2), 326–342.
978 <https://doi.org/10.1111/j.1420-9101.2010.02169.x>

979 Thompson, J., Higgins, D., & Gibson, T. (1994). CLUSTAL W: improving the sensitivity of
980 progressive multiple sequence alignment through sequence weighting. *Nucleic Acids Research*,
981 22(22), 4673-4680. <https://doi.org/10.1093/nar/22.22.4673>

982 Torda, G., Donelson, J., Aranda, M., Barshis, D. J., Bay, L., Berumen, M. L., Bourne, D. G.,
983 Cantin, N., Foret, S., Matz, M., Miller, D. J., Moya, A., Putnam, H. M., Ravasi, T., van Oppen,
984 M. J. H., Vega Thurber, R., Vidal-Dupiol, J., Voolstra, C. R., Watson, S.-A., Whitelaw, E.,
985 Willis, B. L., & Munday, P. L. (2017). Rapid adaptive responses to climate change in corals.
986 *Nature Climate Change*, 7(9), 627-636. <https://doi.org/10.1038/nclimate3374>

987 Turner, J. (1981). Latitudinal patterns of calanoid and cyclopoid copepod diversity in estuarine
988 waters of eastern North America. *Journal of Biogeography*, 8(5), 369.
989 <https://www.jstor.org/stable/2844757>

990 Turner, J. (2004). The importance of small planktonic copepods and their roles in pelagic marine
991 food webs. *Zoological Studies*, 43(2), 255-266.

992 Ummenhofer, C. C., & Meehl, G. A. (2017). Extreme weather and climate events with ecological
993 relevance: a review. *Philosophical Transactions of the Royal Society B*, 372(1723), 20160135.
994 <https://doi.org/10.1098/rstb.2016.0135>

995 Van de Waal, D. B., Verschoor, A. M., Verspagen, J. M., van Donk, E., & Huisman, J. (2010).
996 Climate-driven changes in the ecological stoichiometry of aquatic ecosystems. *Frontiers in*
997 *Ecology and the Environment*, 8(3), 145-152. <https://doi.org/10.1890/080178>

998 Vinagre, C., Leal, I., Mendonca, V., Madeira, D., Narciso, L., Diniz, M. S., & Flores, A. A.
999 (2016). Vulnerability to climate warming and acclimation capacity of tropical and temperate

1000 coastal organisms. *Ecological indicators*, 62, 317-327.

1001 <https://doi.org/10.1016/j.ecolind.2015.11.010>

1002 Wernberg, T., Bennett, S., Babcock, R. C., de Bettignies, T., Cure, K., Depczynski, M., Dufois,
1003 F., Fromont, J., Fulton, C. J., Hovey, R. K., Harvey, E. S., Holmes, T. H., Kendrick, G. A.,
1004 Radford, B., Santana-Garcon, J., Saunders, B. J., Smale, D. A., Thomsen, M. S., Tuckett, C. A.,
1005 Tuya, F., Vanderklift, M. A., & Wilson, S. (2016). Climate-driven regime shift of a temperate
1006 marine ecosystem. *Science*, 353(6295), 169–172. <https://doi.org/10.1126/science.aad8745>

1007 West-Eberhard, M. J. (2005). Phenotypic accommodation: adaptive innovation due to
1008 developmental plasticity. *Journal of Experimental Zoology. Part B, Molecular and*
1009 *Developmental Evolution*, 304(6), 610–618. <https://doi.org/10.1002/jez.b.21071>

1010 West-Eberhard, M. J. (2003). Developmental plasticity and evolution. *Oxford Publishers*.

1011 Wijffels, S., Roemmich, D., Monselesan, D., Church, J., & Gilson, J. (2016). Ocean temperatures
1012 chronicle the ongoing warming of Earth. *Nature Publishing Group*, 6(2), 116-118.
1013 <https://doi.org/10.1038/nclimate2924>

1014 Williams, D. A., Brown, S. D., & Crawford, D. L. (2008). Contemporary and historical
1015 influences on the genetic structure of the estuarine-dependent Gulf killifish *Fundulus*
1016 *grandis*. *Marine Ecology Progress Series*, 373, 111-121. <https://doi.org/10.3354/meps07742>

1017 Yampolsky, L., Schaer, T., & Ebert, D. (2013). Adaptive phenotypic plasticity and local
1018 adaptation for temperature tolerance in freshwater zooplankton. *Proceedings of the Royal Society*
1019 *B: Biological Sciences*, 281(1776), 20132744. <https://doi.org/10.1098/rspb.2013.2744>

1020

Supplementary Table

Supp. Table 1: Temperature data used for predictions of vulnerability to warming. Current temperature data was acquired from the MODIS-aqua SST database. Warming was estimated from a high-resolution model of warming in the North Atlantic (Saba *et al.* 2016). Change in survivorship at maximum temperatures was estimated using TPCs based on a developmental temperature equal to the mean temperature.

Pop	Latitude	Current Mean Temp (°C)	Current Max Temp (°C)	Future Mean Temp (°C)	Future Max Temp (°C)	Estimated Temp Increase (°C)	Change in Survivorship
KL	25.283891	25.1	29.56	26.6	31.06	1.5	-0.0023
PR	26.483536	24.79	29.42	26.29	30.92	1.5	-0.15
PG	26.940398	24.69	28.8	26.19	30.3	1.5	-0.032
MR	27.505606	20.77	21.84	22.27	23.34	1.5	-0.0049
SP	27.507322	24.02	29.5	25.52	31	1.5	-0.16
NS	29.037478	22.84	26.71	24.34	28.21	1.5	-0.011
CT	41.320591	12.52	18.69	14.52	20.69	2	0.065
ME	44.615086	9.05	11.87	12.55	15.37	3.5	0.098
SA	45.100059	9.37	13.06	12.87	16.56	3.5	3.50E-04
SB	46.27291	12.38	15.76	13.88	17.26	1.5	0.054

Supplementary Figures

Supp. Fig. 1: Change in thermal limits vs. change in thermal tolerance by developmental phenotypic plasticity. The solid line represents a 1:1 relationship. Points that fall below the line represent a larger increase in thermal tolerance than thermal limits.

1038

1039 Supp. Fig. 2: Average increase in survivorship at every dosage level. No major differences are
 1040 observed between this metric and ΔLD_{50} values, which are represented by black crosses.

1041

1042 Supp. Fig. 3: Correlation between monthly mean temperature and latitude. A significant

1043 correlation is observed.

1044

1045 Supp. Fig. 4: Correlation between monthly mean temperature variance and latitude. A significant
1046 correlation is observed.

1047

1048 Supp. Fig. 5: Correlation between the two explanatory environmental variables (mean monthly
 1049 temperature and mean monthly temperature variance). No significant relationship between the
 1050 variables is observed.

1051

1052 Supp. Fig. 6 – Correlation between the estimated number of migrants exchanged between two
 1053 populations and the corresponding pairwise difference in thermal tolerance values between those
 1054 populations.