Integrated analysis of the miRNA-mRNA network associated with LMP1 gene in nasopharyngeal carcinoma

Short title: A co-analysis of EBV-associated genes in NPC

Autho list: Yang Yang^a, Wen Liu^a, Yan Zhang^{a,b}, Shuo Wu^a, Bing Luo^{a*}

Affiliation:

^a: Department of Pathogenic Biology, Qingdao University Medical College, 38 Dengzhou Road, Qingdao, 266071, China

^b: Department of Clinical Laboratory, Central Hospital of Zibo, 54 Gongqingtan Road, Zibo, 255036, China

* Corresponding author: Bing Luo

Affiliation: Department of Pathogenic Biology, Qingdao University Medical College, 38 Dengzhou Road, Qingdao, 266071, China. Tel: 86-532-8299108.

Abstract: Epstein-Barr virus oncogenic latent membrane protein 1 (LMP1) has been known to play important roles in nasopharyngeal carcinoma (NPC). LMP1 gene also induced a variety of microRNAs (miRNAs) which bear pivotal roles in regulation of mRNAs expression. However, little was known about the global change of mRNAs and miRNAs induced by LMP1 gene in NPC. In our study, one NPC tissue microarray profile and two LMP1-associated microarray expression profiles data were downloaded from the Gene Expression Omnibus database. A protein-protein interaction network was constructed by using bioinformatics platform Gene-Cloud of Biotechnology Information (NCBI). 78 differentially expressed miRNAs and 3322 differentially expressed genes were identified in order to generate a macroscopic network between miRNAs and mRNAs associated with LMP1 gene. In addition, two significant models were generated to illustrate the expression tendency. Our study provided a way to reveal the interaction between miRNAs and mRNAs in LMP1 axis, bringing insights into the pathogenesis of NPC.

Keywords: latent membrane protein 1; Epstein-Barr virus; micro RNA; nasopharyngeal carcinoma

33 Introduction

34 Nasopharyngeal carcinoma is a primary cancer arising from the nasopharynx. The disease
35 commonly distributes in certain regions of East Asia and Africa, especially in Guangdong prov-
36 ince of China [1]. It is generally considered that genetic susceptibility, Epstein-Barr virus (EBV)
37 infection and environmental factors were involved in the pathogenesis of NPC. Linkage analysis
38 showed that several susceptibility loci were related to the etiology of NPC in China [2,3]. Con-
39 sumption of salt-preserved fish may also play an important role in NPC [4].

40 EBV is an enveloped, double-stranded human DNA herpesvirus. It infects more than 90% of
41 the global adult people being for life and in most carriers the infection is usually asymptomatic.
42 Nevertheless, EBV has oncogenic properties for several malignancies including Hodgkin's disease,
43 Burkitt's lymphoma, as well as undifferentiated NPC [5]. The consistent detection of virus in NPC
44 cases and the continuous expression of virus genes indicate that EBV is crucial for the malignant
45 growth. The circulating cell free EBV DNA is well-recognized biomarker in undifferentiated NPC
46 [6]. EBV latently infects NPC cells by expressing multiple virus genes including EBV nuclear
47 antigens 1 (EBNA1), the latent membrane proteins (LMP1) , 2A, and 2B and viral noncoding
48 RNAs. Among them, LMP1 oncoprotein may play a critical role in the initiation and progression
49 of NPC as well as the invasion and metastasis [7-9]. LMP1 is a 66kDa integral membrane protein
50 consisting of a cytoplasmic amino-terminus, six transmembrane domains and a large cytoplasmic
51 C-terminal tail comprising three regions: transformation effector site (TES)/C-terminal activating
52 regions (CTAR) 1, 2 and 3. CTAR1, CTAR2 are responsible for recruiting cellular signaling mol-
53 ecules of the tumor necrosis factor receptor associated factor (TRAF) family and TRAF-associated
54 death domain protein (TRADD) and the activation of the nuclear factor kap-
55 pa-light-chain-enhancer of activated B cells (NF- κ B) and c-Jun N-terminal protein kinase (JNK)
56 /DNA-binding activity of activator protein-1 (AP-1) signaling [10,11].

57 It is well recognized that the many small non-coding microRNAs (miRNAs) play pivotal role
58 in regulation the homeostasis of mRNAs expression in physiological and pathological process [12].
59 MiRNAs can bind to the specific regions in the target mRNAs, thus leading to the degradation or
60 repression of the mRNAs [13]. As a multifunctional gene, LMP1 was proved to modulate many
61 kinds of miRNAs in B cells and epithelial cells [14-16]. Most of these studies have focused on the
62 interaction of single miRNA with the carcinoma cells, thus a landscape of interaction between
63 miRNAs and mRNAs was needed to illustrate the global change induced by LMP1 in NPC.

64 In the present study, we investigated three expression profiles involving LMP1-associated
65 miRNAs and mRNAs data and NPC tissue samples from the Gene Expression Omnibus (GEO)
66 database (<http://www.ncbi.nlm.nih.gov/geo/>). We analyzed the differentially expressed genes
67 (DEGs) and miRNAs (DE-miRNAs) and performed the pathway analysis. The protein-protein
68 interaction (PPI) network and the mRNA-miRNA interaction network were constructed to obtain
69 the key genes and miRNAs in NPC.

71 Materials and Methods

72 Data Collection

73 Gene expression profiles were downloaded as raw signals from GEO. The dataset GSE12452
74 comprised 31 NPC tissues and 10 normal nasopharyngeal tissues. The dataset GSE29297 involved
75 a series of LMP1 TES2-stimulated HEK-293 cell line samples at different time points. The dataset

GSE26596 valuated the miRNAs expression in NPC cell line TW03 transfected with LMP1. TW03 was an EBV-negative cell line derived from a lymphoepitheliomatous undifferentiated carcinoma in Taiwan province [17]. The sequences of the LMP1 gene encoded by different EBV strains has been shown to have a degree of variation in NPC. The LMP1 vector used in dataset GSE29596 was as described [18]. Vector encoding LMP1 TES2 (amino acids 351-386) was used in GSE29297. Both of the two vector covered part of C-terminal activating region.

Differential Expression Analysis

An online platform Gene-Cloud of Biotechnology Information (GCBI) was used to interpret, normalized, log2 scaled the datasets GSE12452 and GSE29297. This platform integrated with biology, computer science, medicine, informatics, mathematics and other disciplines. Microarray probe signals with absolute value of fold change (FC)>1.2, *P*-value<0.05, and false discovery rate (FDR) <0.05 were considered to be statistically differential [19,20]. Co-expression networks were constructed based upon contribution degrees and models of Series Test of Cluster (STC) were established by random permutation scheme (<http://college.gcbi.com.cn/helpme>). GSE26596 was analysed via the R-programming language-based dataset analysis tool GEO2R (<http://www.ncbi.nlm.nih.gov/geo/geo2r/>). This interactive tool was allowed to screen different expression miRNAs between TW03 cell line and TW03 cell line transfected with LMP1 in dataset GSE26596 [21].

GO and pathway analysis

The functional annotation of the DEGs and DE-miRNAs were performed by GCBI platform. Gene Ontology (GO) (<http://www.geneontology.org>) classification was used to analysis the DEGs function including biological processes, molecular function, and cellular component [22]. The Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis was utilized to explore the pathway of DEGs.

Gene regulation network and miRNA-mRNA network

Micro RNA target genes were predicted mainly based on the bioinformatic platform miRWalk 2.0 [23]. The platform integrates information from 12 miRNA-target databases, including prediction datasets and validated information: MiRWalk, MicroT4, miRanda, miRBridge, miRDB, miRMap, miRNAMap, PICTAR2, PITA, RNA22, RNAhybrid, Targetscan. The 3' untranslated regions for the target genes were used as the primary base-pairing regions of miRNAs [24]. The miRNA-gene pairs that were common in at least five databases and met with *P*<0.05 were regard as reliable. The miRNA target mRNAs were then compared with the genes in STC models and the overlapped genes were filtered for further analysis. The LMP1 associated miRNA-mRNA interactomes was further visualized using the Cytoscape 3.5.1 platform [25]. In addition, DEGs of GSE29297 and GSE12452 were used to construct a protein-protein interaction networks based on GCBI platform. The details of the network construction algorithm were described on GCBI platform.

Results

Identification of DEGs and DE-miRNAs associated with LMP1

In the present study, the mRNA profiling dataset GSE29297 was analyzed on GCBI platform. Samples in the dataset were divided into five groups according to the TES2 stimulation period. 3109 probes for 2787 DEGs were identified after background correction and normalization (*P* value<0.05, *Q* value<0.05, Figure 1). 41 samples (including 31 NPC samples and 10 normal tissues) in the dataset GSE12452 were divided into two groups using GCBI tools. 2566 probes for

120 1815 genes were up-regulated and 2021 probes for 1507 genes were down-regulated. The criteria
121 were set to adjust P value <0.05 and $FC>1.5$ (Figure 2). 372 DEGs were up-regulated and 245
122 DEGs were down-regulated in both datasets GSE29297 and GSE12452. For dataset GSE26596,
123 72 miRNAs were up-regulated, and 6 miRNAs were down-regulated (adjust P value <0.05 ,
124 $FC>1.5$).

125 **Models of Series Test of Cluster**

126 The expression tendency of gene clusters of dataset GSE29297 was calculated on GCBI
127 platform. Among the 20 expression STC models constructed, 15 models were estimated as statis-
128 tical significant (all P values <0.05 , Figure 3). Notably, a descending expression model STC32
129 including 1358 probes for 1070 genes and an ascending model STC49 including 1422 genes were
130 identified as significantly enriched, which may be affected by the LMP1 associated up-regulated
131 miRNAs.

132 **Prediction of miRNA-target genes**

133 The potentially targeted mRNAs were predicted using the integrated bioinformatic tool
134 miRWalk 2.0. Among the 78 DE-miRNAs, 11 miRNAs showed up-regulated expression with at
135 least 20 fold change, including hsa-miR-134, hsa-miR-7, hsa-miR-432 and miR-515 family such
136 as hsa-miR-516-3p, hsa-miR-520b, hsa-miR-520d, hsa-miR-520e, hsa-miR-523. While only 6
137 miRNAs down-regulated mildly, including the hsa-miR-15a, hsa-miR-107, mmu-miR-140*,
138 hsa-miR-149, hsa-miR-194, hsa-miR-103. The target genes of several miRNAs were later ob-
139 served with the STC models.

140 **Significant functions and pathway enrichment analysis.**

141 Gene ontology and KEGG pathway enrichment analysis were performed on the DEGs to re-
142 veal their biological significance in NPC on GCBI platform.

143 The results showed that the DEGs in dataset GSE29297 were significantly involved in the
144 molecular functions of protein binding, zinc ion binding and DNA binding and sequence-specific
145 DNA binding transcription factor activity. In terms of biological processes, the DEGs were mainly
146 enriched in the process of positive and negative regulation of transcription. Mitogen-activated
147 protein kinase (MAPK) signaling pathway, phosphatidylinositol-3-kinase-Serine/threoninekinase
148 (PI3K-AKT) signaling pathway and pathways of human T-cell leukemia virus type 1 (HTLV-I)
149 and hepatitis B were predominately significant in the KEGG pathway analysis.

150 GO enrichment analysis revealed that the DEGs of dataset GSE12452 were significantly en-
151 riched in protein and ATP binding, DNA and RNA binding as well as mitotic cell cycle. KEGG
152 pathway analysis showed that the DEGs were significantly enriched in the metabolic pathway, the
153 PI3K signaling pathway and pathways associated with cell cycle and RNA transport.

154 **Network Analysis**

155 The protein-protein coexpression network showed that a number of genes presented higher
156 node degree including exportin 1(XPO1), eukaryotic translation initiation factor 3(EIF3E), tran-
157 scription elongation regulator 1(TCERG1), anaphase promoting complex subunit 5 (ANAPC5),
158 cytochrome c, somatic (CYCS) (Figure 4). The pathway relation network showed that MAPK
159 signaling pathway, pathway of cell cycle and apoptosis and p53 signaling pathway were signifi-
160 cant-enriched downstream pathways, while the pathway in cancer regulated significantly upstream
161 (Figure 5).

162 The time-associated DEGs in STC32 and STC49 models and the up-regulated miRNAs iden-
163 tified in GSE26596 were used to generate a LMP1-specific miRNA-RNA interacting network. Top

164 5 miRNAs were chosen as core miRNAs to built the network (Figure 6).

165 Discussion

166 NPC featured an EBV type $\square$ latency which was characterized by the expression of LMP1
167 [7]. High level miRNAs such as the BamHI-A rightward transcripts (BARTs), with the combina-
168 tion of the human noncoding RNAs induced by the virus, may affect the process of the
169 oncogenesis. However, a variety of ncRNAs may hinder our understanding about the LMP1 axis.
170 This integrated study aimed to seek a clue about the interaction between LMP1 and its induced
171 miRNAs in the universe of the signal network.

172 A co-expression analysis of the datasets GSE12452 and GSE29297 displayed a 36-hub genes
173 network. Among these hub genes, the major export receptors of mRNA (XPO1) gene was pre-
174 dicted to co-express with 22 genes in NPC cell and the expression level was 1.8 fold up-regulated
175 ($P < 0.01$). The gene can be regulated by hsa-miR-155 and hsa-miR-373*, which were upregulated
176 in GSE26596. This dataset comprised the expression profile of EBV-negative NPC cell line TW03
177 and the LMP1-transfected TW03 cell line. We proposed that the up-regulated expression levels of
178 miRNAs act as a response to attenuate the up regulated mRNAs such as the XPO1.

179 An overlapped analysis of the datasets GSE12452 and GSE29297 revealed that the DEGs
180 were significantly involved in the PI3K-AKT signaling pathway. Studies have shown that LMP1
181 can activate the PI3K-AKT pathway and evade tumor suppressor responses in many ways. LMP1
182 over-expression led to upregulated level of mi-155-Ubiquitin1 axis to activate PI3K-AKT pathway
183 in NPC cells, which promoted the radioresistance of NPC [26]. In diffuse large B-cell lymphoma
184 the PI3K-AKT signaling pathway was significantly activated by the overexpression of miR-155 in
185 DHL16 cells, while in OCL-Ly3 cells knockdown of miR-155 attenuate the AKT activity [27].
186 Correspondingly, in GSE26596, miRNA-155 was 12.73 fold upregulated (adjust $P < 0.01$), which
187 indicating that the PI3K-AKT signaling pathway may possibly be regulated by LMP1 via
188 miR-155 in NPC.

189 Analysis showed that most genes in GSE29297 were enriched in two STC expression models
190 in LMP1 TES2-stimulated HEK-293 cell line, named STC32 and STC49. The ascendant expres-
191 sion model STC49 indicated that 1787 probes for 1422 genes were up-regulated in pace with the
192 TES2 stimulation. Pathway analysis showed that these genes were primarily enriched in the pro-
193 cesses such as ubiquitin mediated proteolysis and transforming growth factor- β signaling pathway.
194 It has been proved that LMP1 can regulate ubiquitination by the activation of Interferon regulatory
195 factor 7 [28]. While in the down-regulated gene model STC32, the genes were enriched in the
196 process of mesenchymal to epithelial transition and cell communication. Several studies have
197 shown that LMP1 can promote the progress of epithelial to mesenchymal transition, thus contrib-
198 utes to the metastatic nature of NPC [29,30].

199 Integrate analysis revealed a miR-515 family including hsa-miR-520e, hsa-miR-523,
200 hsa-miR-520b, hsa-miR-516-3p and hsa-miR-520d was significantly up-regulated. The targets of
201 miR-515 family target genes were predicted to cover large number of genes in the down-regulated
202 model STC32. However, as shown in figure.4, more putative interactions were in the ascending
203 model STC49, hinting that the up-regulation of miRNAs may be a restrain means or a remedial
204 measure to counteract the up-regulation of particular mRNAs. For example, analysis of miRNA
205 hsa-miR-520e targeted genes showed that the down-regulated genes were primarily enriched in
206 glypican pathway, while the up-regulated genes were mainly enriched in vascular endothelial

growth factor signaling pathway, hinting that hsa-miR-520e may play a negative role in balancing the abnormal increasement of specific miRNAs. Another study showed that miR-520 acted as a tumor-suppressive factor by direct targeting of transcription factor P65 and thus inhibit the NF- κ B signaling to reduce the expression of the pro-inflammatory cytokines [31].

Recently, it has been reported that hsa-miR-134 played crucial roles in abundant and complicated pathways, including KRAS signal pathway, Notch pathway and EGFR pathway [32,33]. In dataset GSE2927, both up-regulated genes such as SMAD6, MYCN, PRLR, CYTH3, BMP3, PDE5A, and down-regulated genes such as BCL2, TGFB2, FOXP2, PKD2, TAF4B, and TSN1 were predicted to be targets of miR-134. As many studies showed, miR-134 not only functions as a tumor repressor, but also acts as a cancer promoter. High expression of miR-134 contributes to head and neck carcinogenesis by targeting the WW Domain-Containing Oxidoreductase (WWOX) gene [34]. However, overexpression of miR-134 can inhibit the cell cycle progression of human human ovarian cancer stem cells and decrease the tumorigenicity in nude mice [35]. These studies hinting that a dysregulation of miR-134 may participate in the development of NPC.

Nevertheless, our study had its points which were not rigorous enough, for example, the dataset GSE29297 used was limited in only one cite TES2, the DEGs and the DE-miRNAs involved in our study might associate with the sample sorts, the pathological stage and the condition of cell lines. In addition, more experiments were necessary to validate our results.

Conclusion

In conclusion, by taking advantage of bioinformatic tools and GEO profiles, a protein-protein network and a miRNA-mRNA interaction network were constructed. Our result revealed another layer of gene regulation network in the LMP1-associated gene expression axis, which would provide a better understanding of the interaction of mRNAs and miRNAs. The interaction relationship may open a way to explore the potential use of miRNA in NPC.

Acknowledgments

This research was supported by the National Natural Science Foundation of China (NSFC 81571995) and the Natural Science Foundation of Shandong Province (ZR2015HM069). The authors appreciated Linlin Wang for providing an account of GCBI tools.

Authors' contributions

BL and QS conceived the project and designed experiments. WL, YZ, SW and YY participated in performance of the experiments and the drafting of the manuscript. All authors read and approved the final manuscript.

Ethics, consent and permissions

Competing interests The authors declared that they have no competing interests.

Ethical approval The study was approved by Ethics Committee of Qingdao University Medical College.

Consent to publish All the datasets used in the study were obtained from GEO public database, which were equipped with informed consent.

245 Reference

- 246 [1] MA Wei K-R, Zheng R-S, Zhang S-W, Liang Z-H, Li Z-M, Chen W-Q et al. Nasopharyngeal carcinoma incidence and mortality in China 2013. *Chin J Cancer*. 2017;36:90.
- 247
- 248 [2] Feng, B.J., Huang, W., Shugart, Y.Y., et al. Genome-wide scan for familial nasopharyngeal carcinoma reveals evidence of linkage to chromosome 4. *Nat Genet*. 2002; 31(4): 395.
- 249
- 250 DOI:10.1038/ng932.
- 251 [3] Xiong, W., Zeng, Z.Y., Xia, J.H., Xia, K., Shen, S.R., Li, X.L., Hu, D.X., Tan, C., Xiang, J.J., Zhou, J., Deng, H., Fan, S.Q., Li, W.F., Wang, R., Zhou, M., Zhu, S.G., Lü, H.B., Qian, J., Zhang, B.C., Wang, J.R., Ma, J., Xiao, B.Y., Huang, H., Zhang, Q.H., Zhou, Y.H., Luo, X.M., Zhou, H.D., Yang, Y.X., Dai, H.P., Feng, G.Y., Pan, Q., Wu, L.Q., He, L., Li, G.Y., et al. A susceptibility locus at chromosome 3p21 linked to familial nasopharyngeal carcinoma. *Cancer Res*.2004;64(6):1972-4. DOI:10.1158/0008-5472.CAN-03-3253.
- 252
- 253
- 254
- 255
- 256
- 257 [4] Ren, Z.F., Liu, W.S., Qin, H.D., Xu, Y.F., Yu, D.D., Feng, Q.S., Chen, L.Z., Shu, X.O., Zeng, Y.X., Jia, W.H., et al. Effect of family history of cancers and environmental factors on risk of nasopharyngeal carcinoma in Guangdong, China. *Cancer Epidemiol*. 2010;34(4):419-24. DOI: 10.1016/j.canep.2010.04.011.
- 258
- 259
- 260
- 261 [5] Taylor, G.S., Long, H.M., Brooks, J.M., Rickinson, A.B., Hislop, A.D., et al. The immunology of Epstein-Barr virus-induced disease. *Annu Rev Immunol*. 2015;33:787-821.
- 262
- 263 DOI:10.1146/annurev-immunol-032414-112326.
- 264 [6] Lo, Y.M., Chan, L.Y., Lo, K.W., Leung, S.F., Zhang, J., Chan, A.T., Lee, J.C., Hjelm, N.M., Johnson, P.J., Huang, D.P., et al. Quantitative analysis of cell-free Epstein-Barr virus DNA in plasma of patients with nasopharyngeal carcinoma. *Cancer Res*.,1999;59(6):1188-91.
- 265
- 266
- 267 [7] Yoshizaki, T., Kondo, S., Wakisaka, N., Muro, S., Endo, K., Sugimoto, H., Nakanishi, S., Tsuji, A., Ito, M., et al. Pathogenic role of Epstein-Barr virus latent membrane protein-1 in the development of nasopharyngeal carcinoma. *Cancer Lett*. 2013;337(1):1-7.
- 268
- 269
- 270 DOI:10.1016/j.canlet.2013.05.018.
- 271 [8] Horikawa, T., Yoshizaki, T., Sheen T.S., Lee, S.Y., Furukawa, M., et al. Association of latent membrane protein 1 and matrix metalloproteinase 9 with metastasis in nasopharyngeal carcinoma. *Cancer*. 2000;89(4):715-23.
- 272
- 273
- 274 [9] Lee, D.C., Chua, D.T., Wei, W.I., Sham, J.S., Lau, A.S., et al. Induction of matrix metalloproteinases by Epstein-Barr virus latent membrane protein 1 isolated from nasopharyngeal carcinoma. *Biomed Pharmacother*. 2007;61(9):520-6. DOI:10.1016/j.biopha.2007.08.007.
- 275
- 276
- 277 [10] Ersing, I., Bernhardt, K., Gewurz, B.E., et al. NF-κB and IRF7 Pathway Activation by Epstein-Barr Virus Latent Membrane Protein 1. *Viruses*. 2013;5(6):1587-606.
- 278
- 279 DOI:10.3390/v5061587.
- 280 [11] Devergne, O., Hatzivassiliou, E., Izumi, K.M., Kaye, K.M., Kleijnen, M.F., Kieff, E., Mosialos, G., et al. Association of TRAF1, TRAF2, and TRAF3 with an Epstein-Barr virus LMP1 domain important for B-lymphocyte transformation: role in NF-kappaB activation. *Mol Cell Biol*. 1996;16(12):7098-108.
- 281
- 282
- 283
- 284 [12] Hammond S M. An overview of microRNAs. *Adv Drug Deliv Rev*. 2015;87:3-14. DOI: 10.1016/j.addr.2015.05.001.
- 285
- 286 [13] Hammond, S. M. Dicing and slicing: the core machinery of the RNA interference pathway. *FEBS Lett*. 2005;579(26):5822-9. DOI:10.1016/j.febslet.2005.08.079.
- 287

- 288 [14] Li, G., Wu, Z., Peng, Y., Liu, X., Lu, J., Wang, L., Pan, Q., He, M.L., Li, X.P., et al.
289 MicroRNA-10b induced by Epstein-Barr virus-encoded latent membrane protein-1 promotes the
290 metastasis of human nasopharyngeal carcinoma cells. *Cancer Lett.*2010;299(1):29-36.
291 DOI:10.1016/j.canlet.2010.07.021.
- 292 [15] Du, Z.M., Hu, L.F., Wang, H.Y., Yan, L.X., Zeng, Y.X., Shao, J.Y., Ernberg, I., et al.
293 Upregulation of MiR-155 in nasopharyngeal carcinoma is partly driven by LMP1 and LMP2A and
294 downregulates a negative prognostic marker JMJD1A. *Plos One.* 2011;6(4):e19137.
295 DOI:10.1371/journal.pone.0019137.
- 296 [16] Xia, H., Ng, S.S., Jiang, S., Cheung, W.K., Sze, J., Bian, X.W., Kung, H.F., Lin, M.C., et
297 al. miR-200a-mediated downregulation of ZEB2 and CTNNB1 differentially inhibits nasopha-
298 ryngeal carcinoma cell growth, migration and invasion. *Biochem Biophys Res Commun.*
299 2010;391(1):535-41.
- 300 [17] Teramoto N, Maeda A, Kobayashi K, et al. Epstein-Barr virus infection to Epstein-Barr
301 virus-negative nasopharyngeal carcinoma cell line TW03 enhances its tumorigenicity. *Laboratory*
302 *investigation*, 2000;80(3):303-312.
- 303 [18].Hu LF, Zabarovsky ER, Chen F, Cao SL, Ernberg I, Klein G, Winberg G, et al. Isolation
304 and sequencing of the Epstein-Barr virus BNLF-1 gene (LMP1) from a Chinese nasopharyngeal
305 carcinoma. *J Gen Virol.* 1991;72(Pt10):2399–2409.
- 306 [19] Tusher, V.G., Tibshirani, R., Chu, G., et al. Significance analysis of microarrays applied
307 to the ionizing radiation response. *Proc Natl Acad Sci U S A.* 2011;98(9):5116-21.
308 DOI:10.1073/pnas.091062498.
- 309 [20] Reiner, A., Yekutieli, D., Benjamini, Y., et al. Identifying differentially expressed genes
310 using false discovery rate controlling procedures. *Bioinformatics.* 2003,19(3):368-75.
- 311 [21] Barrett, T., Wilhite, S.E., Ledoux, P., Evangelista, C., Kim, I.F., Tomashevsky, M., Mar-
312 shall, K.A., Phillippy, K.H., Sherman, P.M., Holko, M., Yefanov, A., Lee, H., Zhang, N., Robert-
313 son, C.L., Serova, N., Davis, S., Soboleva, A., et al. NCBI GEO: archive for functional genomics
314 data sets--update. *Nucleic Acids Res.* 2013;41 (Database issue), D991-5.
315 DOI:10.1093/nar/gks1193.
- 316 [22]Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P.,
317 Dolinski, K., Dwight, S.S., Eppig, J.T., Harris, M.A., Hill, D.P., Issel-Tarver, L., Kasarskis, A.,
318 Lewis, S., Matese, J.C., Richardson, J.E., Ringwald, M., Rubin, G.M., Sherlock, G., et al. Gene
319 ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet.*
320 2000;25(1):25-9. DOI:10.1038/75556.
- 321 [23] Dweep, Harsh., N., Gretz., et al. miRWalk2.0: a comprehensive atlas of mi-
322 croRNA-target interactions. *Nat Methods.*2015;12(8):697. DOI:10.1038/nmeth.3485.
- 323 [24] Rehmsmeier, M., Steffen, P., Hochsmann, M., Giegerich, R., et al. Fast and effective
324 prediction of microRNA/target duplexes. *RNA.* 2004;10(10):1507-17.
- 325 [25] Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N.,
326 Schwikowski, B., Ideker, T., et al. Cytoscape: a software environment for integrated models of
327 biomolecular interaction networks. *Genome Res.* 2003;13(11):2498-504.
328 DOI:10.1101/gr.1239303.
- 329 [26] Yang, F., Liu, Q., Hu, C.M., et al. Epstein-Barr virus-encoded LMP1 increases miR-155
330 expression, which promotes radioresistance of nasopharyngeal carcinoma via suppressing
331 UBQLN1. *Eur Rev Med Pharmacol Sci.* 2015;19(23):4507-15.

[27] Huang, X., Shen, Y., Liu, M., Bi, C., Jiang, C., Iqbal, J., McKeithan, T.W., Chan, W.C., Ding, S.J., Fu, K., et al. Quantitative proteomics reveals that miR-155 regulates the PI3K-AKT pathway in diffuse large B-cell lymphoma. *Am J Pathol*.2012;181(1):26-33. DOI:10.1016/j.ajpath.2012.03.013.

[28] Ning, S., Pagano, J.S., Barber, G.N., et al. IRF7: activation, regulation, modification and function. *Genes Immun*.2011;12(6):399-414. DOI:10.1038/gene.2011.21.

[29] Wasil, L.R., Shair, K.H., et al. Epstein-Barr virus LMP1 induces focal adhesions and epithelial cell migration through effects on integrin- α 5 and N-cadherin. *Oncogenesis*.2015;4:e171. DOI:10.1038/oncsis.2015.31.

[30] Sides, M.D., Klingsberg, R.C., Shan, B., Gordon, K.A., Nguyen, H.T., Lin, Z., Takahashi, T., Flemington, E.K., Lasky, J.A., et al. The Epstein-Barr virus latent membrane protein 1 and transforming growth factor- β 1 synergistically induce epithelial-mesenchymal transition in lung epithelial cells. *Am J Respir Cell Mol Biol*. 2011;44(6):852-62. DOI:10.1165/rcmb.2009-0232OC.

[31] Keklikoglou, I., Koerner, C., Schmidt, C., Zhang, J.D., Heckmann, D., Shavinskaya, A., Allgayer, H., Gückel, B., Fehm, T., Schneeweiss, A., Sahin, O., Wiemann, S., Tschulena, U., et al. MicroRNA-520/373 family functions as a tumor suppressor in estrogen receptor negative breast cancer by targeting NF- κ B and TGF- β signaling pathways. *Oncogene*.2012;31(37):4150-63. DOI:10.1038/onc.2011.571.

[32] Liu, Y., Zhang, M., Qian, J., Bao, M., Meng, X., Zhang, S., Zhang, L., Zhao, R., Li, S., Cao, Q., Li, P., Ju, X., Lu, Q., Li, J., Shao, P., Qin, C., Yin, C., et al. miR-134 functions as a tumor suppressor in cell proliferation and epithelial-to-mesenchymal Transition by targeting KRAS in renal cell carcinoma cells. *DNA Cell Biol*. 2015;34(6):429-36.

[33] Gao, Y., Liu, T., Huang, Y., et al. MicroRNA-134 suppresses endometrial cancer stem cells by targeting POGLUT1 and Notch pathway proteins. *FEBS Lett*. 2015;589(2):207-14. DOI:10.1016/j.febslet.2014.12.002.

[34] Ekizoglu, S., Bulut, P., Karaman, E., Kilic, E., Buyru, N., et al. Epigenetic and genetic alterations affect the WWOX gene in head and neck squamous cell carcinoma. *PLoS One*.2015; 10(1):e0115353. DOI:10.1371/journal.pone.0115353.

[35] Chang, C., Liu, T., Huang, Y., Qin, W., Yang, H., Chen, J., et al. MicroRNA-134-3p is a novel potential inhibitor of human ovarian cancer stem cells by targeting RAB27A. *Gene*. 2017;605:99-107. DOI:10.1016/j.gene.2016.12.030.

364 **Figure legends**

365

366 Figure 1. Heat map showing the differential expression pattern of genes in dataset
367 GSE29297. The x-axis represents samples and the y-axis represents the genes. The
368 bar on the top indicates the log₂-scaled expression level, up-regulated genes were
369 presented in red, while down-regulated in green.

370

371 Figure 2. Heat map showing the differential expression pattern of genes in dataset
372 GSE12452. The x-axis represents samples and the y-axis represents the genes. The
373 bar on the top indicates the log₂-scaled expression level, up-regulated genes were
374 presented in red, while down-regulated in green.

375

376 Figure 3. STC models. The profile represent the series test of cluster (STC), profile
377 number represent the STC models. P-value: adjusted P-value represents the signifi-
378 cant levels of the genes in a certain model compared with theoretical genes. The dia-
379 gram was present in red lines as the P-value of the model<0.05, otherwise in blue
380 lines. Numbers in horizontal ordinate were relative periods of the LMP1-TES2 gene
381 stimulation on the HEK293 cell line. The ordinate numbers represent relative grades
382 of expression levels.

383

384 Figure 4. Gene co-expression network of datasets GSE29297 and GSE12452. The size
385 of the node represents the degree of the gene. The positive correlation coefficient was
386 presented in red line, the negative was in blue.

387

388 Figure 5. Pathway relation network of datasets GSE29297 and GSE12452. The arrow
389 in the network directed to the downstream pathways. The size of the node represents
390 the degree of pathway.

391

392 Figure 6. Interaction network between top 5 up-regulated miRNAs in GSE26596 with
393 expression models STC32 and STC49. Up-regulated genes were filled in Red dia-
394 mond, and down-regulated genes green ellipse. The miRNAs were filled in blue dia-
395 mond. These genes were regarded as targets of the miRNAs in predicted and validate
396 databases as described.

397

Profile 32,p-value:0

Profile 49,p-value:0

Profile 52,p-value:3.336e-44

Profile 26,p-value:5.176e-40

Profile 55,p-value:1.879e-32

Profile 50,p-value:5.802e-26

Profile 58,p-value:4.726e-22

Profile 23,p-value:5.999e-20

Profile 31,p-value:3.832e-14

Profile 29,p-value:2.732e-11

Profile 80,p-value:4.872e-09

Profile 53,p-value:0.0006728

Profile 57,p-value:0.001209

Profile 79,p-value:0.008018

Profile 77,p-value:0.02458

Profile 2,p-value:0.05255

Profile 4,p-value:0.0919

Profile 30,p-value:0.449

Profile 68,p-value:0.5316

Profile 36,p-value:0.6101

Figure 4