

1 **No association of mitochondrial DNA levels in trophoctodermal cells with the developmental**
2 **competence of the blastocyst and pregnancy outcomes**

3 Ritu G¹, Geetha Veerasigamani², Mohammed C. Ashraf², Sankalp Singh², Saniya Laheri³,
4 Deepak Modi³

5 ¹Department of Medical Genetic, Craft Hospital and research Center, Chandappura, Kodungallur,
6 Pin - 680664, Thrissur District, Kerala. India.

7 ²Department of Reproductive Medicine, Craft Hospital and Research Center, Chandappura,
8 Kodungallur, Pin- 680664, Thrissur District, Kerala. India.

9 ³Molecular and Cellular Biology Laboratory, National Institute for Research in Reproductive
10 Health (ICMR), J. M. Street, Parel, Mumbai - 400012, India.

11 Correspondence address. E mail: raorit@gmail.com

12 Running title: Mitochondrial DNA levels in outcomes of assisted reproduction.

13 Key words: Mitochondrial DNA/ Next Generation Sequencing/ blastocyst/ implantation/ ploidy/
14 maternal age/morphology

15

16 **Abstract**

17 **Study question**

18 Can mitochondrial DNA (mtDNA) levels in trophoctodermal cells of the blastocyst predict the
19 blastocyst quality, ploidy status, implantation rate and clinical outcomes?

20 **Summary answer**

21 mtDNA levels in trophoctodermal cells of the blastocyst do not associate with the blastocyst
22 quality, ploidy status, implantation potential and clinical outcomes, but can differentiate between
23 aneuploid and euploid blastocysts.

24 **What we already know**

25 mtDNA levels in the trophoctodermal cells have been suggested to be associated with blastocyst
26 morphology, ploidy and implantation rates, and has been proposed as biomarker to assess
27 blastocyst quality and predict clinical outcomes. However, discrepancies exist if mtDNA levels
28 could serve as a marker for the same.

29 **Study design and duration**

30 Retrospective analysis of mtDNA levels in trophoctodermal cells obtained from blastocysts
31 undergoing preimplantation genetic testing for aneuploidy (PGT-A) at Craft Hospital & Research
32 Center, Kerala from January 2016- July 2017.

33 **Participants/materials and methods**

34 Study included data from 287 blastocyst from (61) couples who underwent PGT-A using next
35 generation sequencing (NGS). Levels of mtDNA in trophoctodermal cells of the blastocyst were

36 estimated by the NGS. Comparison of mtDNA levels with maternal age, blastocyst morphology,
37 ploidy status, implantation rates, miscarriage rates and live birth rate was done.

38 **Main results**

39 The levels of mtDNA in the trophoctoderm of the blastocyst did not correlate with maternal age.
40 There was no significant difference in the mtDNA levels between grade 1 and grade 2 blastocyst.
41 Euploid blastocyst had significantly lower amounts of mtDNA levels in trophoctodermal cells of
42 the blastocyst were compared to aneuploid blastocyst. No significant differences were seen
43 between mtDNA levels and implanting and non-implanting blastocysts or those resulted into
44 miscarriage or live birth.

45 **Limitations**

46 The study is limited by a small sample size and hence type II error cannot be ruled out.

47 **Wider Implications**

48 The study does not support the potential use of mtDNA levels in the trophoctodermal cells as
49 biomarker for blastocyst quality and predicting clinical outcomes needs.

50 **Study funding/competing interest(s)**

51 There is no external funding for the study. There is no conflict of interest.

52 **Introduction**

53 Assisted reproduction (ART) has witnessed significant progress in the last three decades and has
54 benefitted many infertile couples. Despite, the advancement in ART, the take-home baby rates are
55 still low (Sadeghi et al., 2012). Various factors which contribute to success rate of ART include
56 maternal and paternal age, gamete quality, endometrial receptivity and most importantly embryo
57 quality (Colaco and Sakkas, 2018; Miao et al., 2009; Oron et al., 2014; Revel, 2012). The current
58 embryo selection methods rely on assessment of embryo morphology with a subjective grading
59 criteria or a real time monitoring of the embryonic development and assessment of multiple
60 quantitative endpoints. However, none of these have been of great aid in improving pregnancy
61 rates (Bromer and Seli, 2008; Capalbo et.al., 2014; Minasia et.al., 2016). The advent of single cell
62 genetic analysis, has made it apparent that a large number of embryos developed *in vitro* are
63 chromosomally aneuploid (Demko et.al., 2016). This has prompted the evaluation of chromosomal
64 complement of the embryos by preimplantation genetic testing for aneuploidy (PGT-A). While the
65 introduction of PGT-A has led to reduction in miscarriage, the improvements in live birth rate
66 following PGT-A remains relatively low and as many as 50% of embryos diagnosed as “euploid”
67 by PGT-A do not implant (Capalbo et al., 2015; Chang et.al., 2016). Thus, genomic aneuploidy
68 may not be a sole factor responsible for failure of implantation and hence there is a need to identify
69 embryo selection markers for improving pregnancy rates.

70 Recently, levels of mitochondrial DNA (mtDNA) has emerged as possible marker for embryo
71 selection (Cecchino et al., 2019, Humaidan et al., 2018, Kim & Seli, 2019). Fragouli and group
72 were the first to report that high levels of mtDNA in embryos derived from older mothers or
73 embryos with aneuploid and embryos with higher levels of mtDNA rarely implant (Fragouli et.al.,
74 2015). These initial findings were subsequently confirmed in a large blinded retrospective study

75 by same group. Large cohort studies have suggested, the quantity of mtDNA as a new biomarker
76 to assist embryo selection (Fragouli et.al., 2017; Ravichandran et.al., 2017). The above studies
77 were confirmed by two more groups independently (de Los Santos et al., 2018; Diez-Juan et al.,
78 2015). However, three additional studies have failed to show any such correlation. Treff and group
79 reported no association between mtDNA levels in implanted and non-implanted embryos (Treff
80 et. al., 2017). In another study the authors argued that the differences in mtDNA content could
81 arise out of differences in genomic DNA content, and hence a normalization factor was used,
82 which led to the conclusion that the levels of mtDNA are near identical between blastocysts
83 stratified by ploidy, maternal age, or implantation potential (Victor et.al., 2017). Another study by
84 Qui and group failed to establish any correlation between mtDNA, embryo quality and clinical
85 outcomes (Qui et al., 2018). Thus, the levels of mtDNA in predicting the embryo quality,
86 implantation potential of embryos and pregnancy outcomes remains controversial.

87 In the current study, we present our experience with use of trophectodermal mtDNA levels to
88 determine blastocyst quality, implantation potential of blastocyst and clinical outcomes that have
89 undergone PGT-A in a clinical setup. Our results reveal no significant association of
90 trophectodermal mtDNA levels with maternal age, implantation potential and clinical outcomes,
91 however mtDNA levels in trophectodermal cells of the blastocyst can differentiate between
92 euploid and aneuploid blastocyst.

93 **Materials and Methods**

94 This was a retrospective study of next generation sequencing (NGS) workflow data from patients
95 who underwent PGT-A between January 2016 to July 2017 at Craft hospital and Research Center,
96 Kerala, India. The study was approved by the ethical review board of Craft Hospital & Research

97 Center, Kodungallur, Kerala (ethics no: 002/21/3/2019).

98 **Study group**

99 At the Craft Hospital and Research Center we routinely offer PGT-A to patients with recurrent
100 implantation failure, recurrent pregnancy loss and advanced maternal age (above 35yrs). The
101 patient characteristics are described in Table 1. A total of 61 couples agreed to undergo PGT-A by
102 NGS from January 2016 to July 2017. Written informed consent was obtained from
103 patients/couples prior to the PGT-A.

104 **Ovarian stimulation, Intracytoplasmic sperm injection (ICSI) and embryo culture**

105 Controlled ovarian stimulation (COS) was done with antagonist protocol using gonadotropins with
106 dosage between 150 – 300IU depending on age and body mass index. Oocytes were aspirated
107 under local anesthesia after 36 h of agonist trigger. Denudation was done after 1h of oocyte
108 retrieval. ICSI was performed as described previously (Velde et al., 1998). After ICSI, fertilization
109 check was done next day, followed by day 3 embryo quality check. Good quality day 3 cleavage
110 stage embryos (embryos with 6-8 cells, equal size of blastomere and cytoplasmic fragmentation
111 less than 10%) were continued to grow till blastocyst stage. Zygotes were cultured in VITROMED
112 culture medium for 5-6 days. Blastocysts quality was graded as described previously (Gardner et
113 al., 2000; Sen et al., 2013). Patients that required oocyte accumulation, the oocytes were frozen
114 within 30 mins after denudation and ICSI done later.

115 **Embryo/Blastocyst Biopsy and vitrification/freezing down (of blastocyst)**

116 Trophectoderm biopsy was done on day 5 embryos (blastocyst stage) as described previously
117 (Capalbo et al., 2015). Briefly laser assisted hatching was performed using LYKOS Laser

118 (Hamilton Thorne; MA, USA) on day 3 embryo. On day 5, herniating blastocyst were selected,
119 and 5-10 trophectoderm cells were removed by suction followed by laser pulsation. The
120 trophectoderm cells were collected in phosphate buffer saline (PBS) and stored at -80°C until
121 further processing. Vitrification method was used to freeze down the blastocyst. The blastocysts
122 were equilibrated in equilibration solution for 12 to 15 mins, followed by transfer to vitrification
123 solution. This was finally transferred to cryolock containing liquid nitrogen.

124 **Whole genome amplification and Next generation sequencing for trophectodermal cells**

125 Whole genome amplification (WGA) on each biopsy was performed using the Rubicon PicoPLEX
126 WGA kit (Agilent, CA, USA) as per manufacture's recommendations. Following WGA, next
127 generation sequencing (NGS) of the trophectodermal biopsies was carried out (Well et al., 2014).
128 For constructing WGA library, Ion Xpress Plus fragment library kit, and Ion Xpress barcode
129 adapters 1–32 kit were performed as per manufacturer's instructions (Thermo Fisher Scientific).
130 150 ng of WGA DNA was fragmented to generate 280 base pair fragments using Ion Shear Plus
131 reagent for 4 mins. Purification of fragmented DNA was done with Agencourt AMPure XP reagent
132 beads (Beckman Coulter, CA, USA), followed by barcoded adaptor ligation, nick repair and
133 purification as per manufacturer's instructions. E-Gel Size Select (Thermo Fisher Scientific)
134 agarose gel was used to select a peak size of 280 base pairs, followed by amplification of DNA
135 with 10 cycles of polymerase chain reaction (PCR) using Platinum PCR SuperMix High Fidelity
136 (Thermo Fisher Scientific). Individual libraries were diluted to 100 pM. On Ion 520 Chip, a pool
137 of 24 samples were loaded. Ion Sphere particles containing amplified DNA were prepared with
138 Ion PI Template OT2 200 Kit v3 (Thermo Fisher Scientific). Template-positive Ion Sphere
139 particles were enriched with the Ion OneTouch ES (Thermo Fisher Scientific). They were later
140 sequenced with Ion 520 Chip and Ion PI Sequencing 200 Kit v3 on the Ion S5 instrument (Thermo

141 Fisher Scientific). Approximately 3 million reads were obtained for each barcoded sample.

142 **Analysis of mtDNA levels**

143 The ploidy status and mtDNA levels were analyzed for all the blastocysts undergoing PGT-A with

144 NGS was assessed with Ion Reporter Cloud based software 5.3 (Thermo Fisher Scientific, MA,
145 USA). The mitochondrial DNA levels were calculated:

146 by Ion Reporter Software.

147 **Embryo transfer**

148 Total of 287 blastocysts were investigated for mtDNA levels in trophectodermal cells, out of which

149 68 euploid embryos were selected who underwent frozen embryo transfer. The endometrium was

150 prepared for transplantation using hormone replacement protocol. Estradiol valerate was

151 administered from day 2 of cycle, in a dose dependent manner. Serial ultrasound monitoring was

152 done to check endometrium thickness (10mm), after which oral and vaginal progesterone were

153 administered and frozen embryo transfer was done. The vitrified blastocyst in the cryolock were

154 directly placed in transfer solution (TS), followed by washing with sucrose solution at 37°C. Serum

155 beta human chorionic gonadotropin levels were measured using HCG STAT Elecsys assay on

156 Cobas E601 Immunology Analyzer (Roche, Basel, Switzerland) after 2 weeks of embryo transfer

157 to detect biochemical pregnancy. Clinical pregnancy was determined by transvaginal ultrasound

158 was done at 6 weeks to see an intrauterine gestation sac.

159 **Statistical analysis**

160 Linear regression analysis was done to study the correlation between maternal age at the time of

161 oocyte retrieval and mtDNA from trophectodermal cell of blastocyst. Correlation between mtDNA

162 level and blastocyst morphology, ploidy status and pregnancy outcomes were carried out. One-
163 way ANOVA using Tukey's all column comparison test was performed using GraphPad Prism,
164 version 5. $p < 0.05$ was accepted as statistically significant.

165 **Results**




Figure 1

Correlation of mitochondrial DNA (mtDNA) levels in trophoctodermal with maternal age at the time of oocyte retrieval. Values on X axis are maternal age in years. Y axis is mtDNA levels in the trophoctodermal cells. Each dot represents data form one patient. The data is derived from 61 independent samples. $p < 0.05$ was accepted as statistically significant.

166

167 **No correlation between mtDNA levels in trophoctodermal cells of the blastocyst and**
168 **maternal age at the time of oocyte retrieval**

169 To investigate the correlation between mtDNA levels in the trophoctodermal cells of blastocyst
170 and maternal age at the time of oocyte retrieval, mtDNA levels were analyzed from
171 trophoctodermal cells of blastocyst generated/collected from 61 women (between the range of 24-
172 46 with an average age of 32.6 ± 4.13). Linear regression analysis did not show any correlation
173 between mtDNA levels in the trophoctodermal cells of blastocyst and maternal age at the time of
174 oocyte retrieval (Fig.1).




Figure 2

Association of levels of mitochondrial DNA (mtDNA) in trophoctodermal cells with blastocyst morphology and ploidy status. (A) Comparison of mtDNA levels in Grade 1 and Grade 2 blastocysts. (B) Comparison of mtDNA levels in euploid and aneuploid blastocysts (C) Comparison of mtDNA levels in euploid, monosomic, trisomic and polysomic blastocyst. In all the graphs, values on Y axis is mtDNA levels estimated by next generation sequencing. The numbers (n) of blastocysts are given in each case. Each dot represents data form one patient. $p < 0.05$ was accepted as statistically significant.

175

176 **mtDNA levels in the trophoctodermal cells of the do not correlate with blastocyst morphology**

177 The predominant criteria for selection of blastocyst is morphological grading of the blastocyst. In
178 order to evaluate if there was a correlation between the mtDNA levels in the trophoctodermal cells
179 and grade 1 (n= 131) and grade 2 (n= 151) blastocyst were analyzed. PGT-A was not performed
180 on grade 3 blastocyst and thus mtDNA levels are not calculated. The results revealed no difference
181 in the mtDNA levels between grade 1 and grade 2 blastocysts (Fig.2A).

182 **mtDNA levels in trophoctodermal cells is higher in aneuploid blastocysts**

183 To study the correlation between mtDNA levels and ploidy status, the blastocysts were classified
184 into euploid and aneuploid. Out of 287 blastocysts analyzed, 170 were euploid and 117 were
185 aneuploid. Among 117 aneuploid blastocyst, 41 were monosomic, 31 were trisomic and 45 were
186 polysomic (aneuploidy of more than 1 chromosome), mtDNA levels were significantly high in

187 aneuploid blastocysts compared to euploid blastocyst ($p < 0.05$, Fig. 2B). The mtDNA levels were
188 significantly higher only in polysomic blastocysts as compared to euploid and monosomic ($p <$
189 0.05 , Fig 2C)




Figure 3

Association of mitochondrial DNA (mtDNA) levels in trophoctodermal cells implanted euploid embryos, and clinical outcomes. Values on Y axis is mtDNA levels. The numbers (n) of blastocysts are given in each case. Each dot represents data form one. $p < 0.05$ was accepted as statistically significant

190

191 **Levels of mtDNA in trophoctodermal cells of blastocyst do not correlate with embryo**
192 **implantation rate and clinical outcomes**

193 Implantation rate is defined as the number of gestational sacs seen at 6weeks of gestation in
194 ultrasound divided by the total number of embryos transferred. All the couples had atleast one
195 euploid embryo for transfer. Of these 4 couples had double embryo transfer. The remaining 57
196 couples had single embryo transfer. Out of the 68 blastocyst, 46 blastocysts implanted, whereas
197 22 blastocysts failed to implant. The overall implantation rate was 67.6%. There was no statistical

198 difference in levels of mtDNA of implanted blastocysts vs non- implanted. The couples who were
199 positive for clinical pregnancy were followed up until end of term. There was no statistical
200 significance difference in mtDNA levels and pregnancy that resulted in miscarriage or live birth
201 (Fig 3).

202 **Discussion**

203 In the present study, we show the mtDNA levels in trophectodermal cells were high in aneuploid
204 blastocysts; however, these mtDNA levels in trophectodermal cells could not predict blastocyst
205 morphology, implantation potential and clinical outcomes.

206 The ultimate goal for researchers in the field of assisted reproduction is to devise strategies that
207 can improve implantation rates and live birth. Sophisticated morphological grading of blastocyst
208 and pre-implantation genetic screening have helped identify euploid embryos. Even with the
209 elimination of aneuploid embryos, one-quarter of euploid embryos fail to implant, suggesting
210 additional factors may be involved (Seli, 2016; Well, 2017). In recent years, mtDNA levels has
211 gained significant importance in determination of embryo quality and pregnancy outcomes
212 (Cecchino et al., 2019, Humaidan et al., 2018, Kim & Seli, 2019). Mitochondria is involved in
213 various critical cellular processes such as energy generation in the form of ATP, homeostasis,
214 amino acid synthesis and apoptosis. Elevated levels of mtDNA indicate dysfunctional
215 mitochondrial machinery or compensatory mechanism to fulfill the requirements of the ever-
216 growing embryo due to defective organelle (Leese 2002). It is well established that maternal age
217 has detrimental effect on pregnancy outcomes (Miao et al., 2009). Earlier studies on mtDNA levels
218 were low in cumulus cells or the oocytes from in the trophectodermal cells women with advanced
219 age (Boucret et al., 2015; Chan et al., 2005; Onigo et al., 2016). Fragouli and group reported high

220 mtDNA levels in embryos from women with advanced maternal age at the time of oocyte retrieval
221 (Fragouli et.al., 2015; 2017; Ravichandran et al., 2017). However, in this study we did not observe
222 any correlation between maternal age (24-46 years) at the time of oocyte retrieval and mtDNA
223 levels in trophoctodermal cells of blastocyst. Similar to our findings, other studies also failed to
224 report any correlation between maternal age with mtDNA levels in the blastocysts (de Los Santos
225 et al., 2018; Diez-Juan et al., 2015; Klimczak et al., 2018; Victor et al., 2017). Together the results
226 indicate, maternal age does not contribute to alterations in mtDNA levels in the blastocysts.

227 A good quality embryo is vital for the success of ART. Gardner and Schoolcraft grading system is
228 widely used to assess the morphology of the embryo (Gardner et al., 2000). The quiet embryo
229 hypothesis by Leese suggest that good quality embryo have low metabolism, whereas the embryo
230 under stress tend to be more metabolically active (Leese 2002). On the basis of this hypothesis,
231 various studies have analyzed the levels of mtDNA and its association of the developmental
232 competence. Studies have reported high levels of mtDNA in poor quality embryos (de Los Santos
233 et al., 2018; Diez-Juan et al., 2015; Klimczak et al., 2018; Murakoshi et al., 2013). Interestingly,
234 in our study, we did not find any correlation between morphological grading of blastocysts and
235 mtDNA levels in the trophoctodermal cells. In concordance with our study, Qui et al., also failed
236 to establish any such correlation (Qui et al., 2018). Thus, association between mtDNA levels in
237 trophoctodermal cells and blastocyst quality is debatable. We next tested the levels of mtDNA in
238 trophoctodermal cells and its association with ploidy status of the blastocyst. We observed higher
239 levels of mtDNA in trophoctodermal cells of aneuploid blastocysts as compared to euploid
240 blastocysts. Interestingly, the high level of mtDNA in trophoctodermal cells was restricted to
241 polysomic blastocysts. Our results are consistent with earlier study, wherein polysomic embryos
242 showed elevated levels of mtDNA (Fragouli et.al., 2015). Interestingly, in another study, embryos

243 with monosomies showed higher mtDNA levels as compared to embryos trisomies (de Los Santos
244 et al., 2018). Whereas, some studies fail to show any association between chromosomal status of
245 the blastocyst and mtDNA levels (Qui et al., 2018; Victor et.al., 2017). The association of mtDNA
246 levels and the ploidy status of the blastocyst require further investigation.

247 Transfer of morphologically and developmentally competent blastocyst to the uterus does not
248 always guarantee pregnancy. Fragouli and group showed euploid embryos with higher mtDNA
249 levels failed to implant, whereas embryos with low mtDNA implanted successfully (Fragouli et
250 al., 2015; 2017). Therefore, we checked if alterations in levels of mtDNA in trophoctodermal cells
251 of euploid blastocyst could impact its implantation potential. However, our data failed to establish
252 any correlation between levels of mtDNA in trophoctodermal cells and implantation potential of
253 blastocyst. This is in concordance to earlier data where, no correlation such was observed (Qui et
254 al., 2018; Victor et al., 2017). Finally, we asked if the mtDNA levels could predict outcomes of
255 implanted blastocysts. The results revealed the levels of mtDNA in the blastocyst had no
256 correlation with the pregnancy outcomes like miscarriage or live birth. Our study is in concordance
257 to earlier studies where Qui et al., also failed to establish any correlation between mtDNA levels
258 in blastocyst and clinical outcomes (Qui et al., 2018). Taken together, the use of mtDNA levels in
259 the blastocyst as biomarker to predict implantation outcomes and pregnancy remains inconclusive.

260 From above it is evident that there are differences in reports from different studies. What could be
261 the possible reason for the discrepancies should be addressed. In a recent study, embryos from
262 women with higher body mass index (BMI) showed higher mtDNA copy number; and high levels
263 of maternal serum progesterone inversely correlated with mtDNA levels (de Los Santos et al.,
264 2018). It could be possible that there could be differences in BMI or serum progesterone levels in
265 women in different studies. However, in our study, the baseline BMI and serum progesterone

266 levels did not significantly differ between the different groups and had no effect of mtDNA levels
267 (data not shown). It is possible that technical differences in quantification of mtDNA levels,
268 sample to sample variability, quantifying number of cells during biopsy, sample storage method
269 ethics are some reasons for differences in results (Fragouli et al., 2013; Humaidan et al., 2018;
270 Well, 2017). There is a need of defined selection criteria and standardized procedure to study if
271 there is correlation between mtDNA and ART outcomes.

272 In summary, the present study fails to support the notion that trophoctodermal mtDNA level can
273 predict the embryo quality, implantation ability of the blastocysts or pregnancy outcomes. The
274 limiting factor of our study could be sample size, which fails to account for type II error. Larger
275 study size along with standardized protocol for mtDNA evaluation, quantification and culture
276 methods is the need of hour.

277 **Acknowledgement**

278 SL is grateful to Council of Scientific & Industrial Research (CSIR), India for Senior Research
279 Fellowship.

280 **Author's role**

281 G.V., R.N., M.A., and S.S., the principal investigators take primary responsibility of the paper.
282 G.V., R.N.,M.A., did the clinical work, counselled the patients and collected the samples. RN
283 performed the experiments analyzed the data. D.M., and S.L., contributed in data analysis, data
284 interpretation. All the authors contributed in manuscript preparation.

285 **Funding**

286 No specific funding was obtained for this study.

287 **Conflict of interest**

288 The authors do not have conflict of interest.

289 **References**

- 290 Boucrot L, Chao De La Barca JM, Moriniere C, Desquiret V, Ferre-L'Hotellier V, Descamps P,
291 Marcaillou C, Reynier P, Procaccio V, May-Panloup P. Relationship between diminished ovarian
292 reserve and mitochondrial biogenesis in cumulus cells. *Hum Rep.* 2015; **30** :1653-1664.
- 293 Bromer, JG, Seli, E. Assessment of embryo viability in assisted reproductive technology:
294 shortcomings of current approaches and the emerging role of metabolomics. *Curr Opin Obstet*
295 *Gynecol* 2008; **20** :234-241.
- 296 Capalbo A, Rienzi L, Cimadomo D, Maggiulli R, Elliott T, Wright G, Nagy ZP, Ubaldi FM.
297 Correlation between standard blastocyst morphology, euploidy and implantation: an
298 observational study in two centers involving 956 screened blastocysts. *Hum Reprod* 2014; **29**
299 :1173–1181.
- 300 Capalbo, A, Filippo MU, Danilo C, Roberta M, Cristina P, Ludovica D, Federica S, Laura B,
301 Roberta V, Laura R. Consistent and reproducible outcomes of blastocyst biopsy and aneuploidy
302 screening across different biopsy practitioners: a multicentre study involving 2586 embryo
303 biopsies. *Hum Reprod* 2015; **31** :199–208.
- 304 Cecchino, Gustavo N., and Juan A. Garcia-Velasco. "Mitochondrial DNA copy number as a
305 predictor of embryo viability." *Fert Steri* 2019; **111** :205-211.
- 306 Chan CC, Liu VW, Lau EY, Yeung WS, Ng EH, Ho PC. Mitochondrial DNA content and 4977
307 bp deletion in unfertilized oocytes. *Mol Hum Reprod.* 2005; **11** :843–846.
- 308 Chang J, Boulet SL, Jeng G, Flowers L, Kissin DM. Outcomes of in vitro fertilization with
309 preimplantation genetic diagnosis: an analysis of the United States Assisted Reproductive
310 Technology Surveillance Data, 2011–2012. *Fertil Steril* 2016; **105** :394-400.

- 311 Colaco S, Sakkas D. Paternal factors contributing to embryo quality. *J Assist Reproduct Genet*
312 2018; **25**:19531-19616.
- 313 de los Santos, Maria J, Antonio DJ, Amparo M, Amparo M, Marcos Mr, Carmen R, Antonio P.
314 "Variables associated with mitochondrial copy number in human blastocysts: what can we learn
315 from trophectoderm biopsies?." *Fert Steril* 2018; **109** :110-117.
- 316 Demko ZP, Simon AL, McCoy RC, Petrov DA, Rabinowitz M. Effects of maternal age on
317 euploidy rates in a large cohort of embryos analyzed with 24-chromosome single-nucleotide
318 polymorphism–based preimplantation genetic screening. *Fertil Steril* 2016; **105** :1307–1313.
- 319 Diez-Juan A, Rubio C, Marin C, Martinez S, Al-Asmar N, Riboldi M, Díaz-Gimeno P, Valbuena
320 D, Simón C. Mitochondrial DNA content as a viability score in human euploid embryos: less is
321 better. *Fertil Steril* 2015; **104** :534–541.
- 322 Fragouli E, Alfarawati S, Spath K, Jaroudi S, Sarasa J, Enciso M, Wells D. The origin and impact
323 of embryonic aneuploidy. *Hum Genet* 2013; **132** :1001–1013.
- 324 Fragouli E, McCaffrey C, Ravichandran K, Spath K, Grifo JA, Munné S, Wells D. Clinical
325 implications of mitochondrial DNA quantification on pregnancy outcomes: a blinded prospective
326 non-selection study. *Human Reprod* 2017; **32**: 2340–2347.
- 327 Fragouli E, Spath K, Alfarawati S, Kaper F, Craig A, Michel CE, Kokocinski F, Cohen J, Munne
328 S, Wells D. Altered levels of mitochondrial DNA are associated with female age, aneuploidy, and
329 provide an independent measure of embryonic implantation potential. *PLoS Genet* 2015; **3**
330 :e1005241.
- 331 Gardner DK, Lane M, Stevens J, Schlenker T, Schoolcraft WB. Blastocyst score affects
332 implantation and pregnancy outcome: towards a single blastocyst transfer, *Fertil Steril* 2000; **73**
333 :1155-1158.

- 334 Humaidan, Peter, Stine Gry Kristensen, and Kevin Coetzee. "Mitochondrial DNA, a new
335 biomarker of embryonic implantation potential: fact or fiction?" *Fertil Steril* 2018; **109** :61-62.
- 336 Kim J, Seli E. Mitochondria as a Biomarker for IVF Outcome. *Reproduct* 2019; **157** : 235–242.
- 337 Klimczak, Amber M., Lucia E. Pacheco, Kelsey E. Lewis, Niloofar Massahi, Jon P. Richards,
338 William G. Kearns, Antonio F. Saad, and John R. Crochet. "Embryonal mitochondrial DNA:
339 relationship to embryo quality and transfer outcomes." *J Assist Reproduct Genet* 2018; **35** :871-
340 877.
- 341 Leese, Henry J. "Quiet please, do not disturb: a hypothesis of embryo metabolism and
342 viability." *BioEssays* 2002; **24** :845-849.
- 343 Miao L, Kikuchi K, Sun Y, Schatten H. Oocyte aging: cellular and molecular changes,
344 developmental potential and reversal possibility. *Hum Reprod Update*. 2009; **15** :573-585.
- 345 Minasi MG, Colasante A, Riccio T, Ruberti A, Casciani V, Scarselli F, Spinella F, Fiorentino F,
346 Varricchio MT, Greco E. Correlation between aneuploidy, standard morphology evaluation and
347 morphokinetic development in 1730 biopsied blastocysts: a consecutive case series study. *Hum*
348 *Reprod* 2016; **31** :2245–2254.
- 349 Murakoshi Y, Sueoka K, Takahashi K, Sato S, Sakura T, Tajima H, Yoshimura Y, Embryo
350 developmental capability and pregnancy outcome are related to the mitochondrial DNA copy
351 number and ooplasmic volume. *J Assist Reproduct Genet* 2013; **30** :1367-1375.
- 352 Ogino M, Tsubamoto H, Sakata K, Oohama N, Hayakawa H, Kojima T, Shigeta M, Shibahara H.
353 Mitochondrial DNA copy number in cumulus cells is a strong predictor of obtaining good-quality
354 embryos after IVF. *J Assist Reproduct Genet* 2016; **33** :367-371.
- 355 Oron G, Son WY, Buckett W, Tulandi T, Holzner H. The association between embryo quality
356 and perinatal outcome of singletons born after single embryo transfers: a pilot study. *Hum Reprod*.

357 2014; **29** :1444–1451.

358 Qiu Y, Chen S, Chen D, Liu P, Xia J, Yang L, Song Z, Shi Q, Xie L, Zhu Z, Du Y. The
359 mitochondrial DNA content can not predict the embryo viability. *bioRxiv*. 2018; **1** :445940.

360 Ravichandran K, McCaffrey C, Grifo J, Morales A, Perloe M, Munne S, Wells D, Fragouli E.
361 Mitochondrial DNA quantification as a tool for embryo viability assessment: retrospective
362 analysis of data from single euploid blastocyst transfers. *Hum Reprod* 2017; **32** :1282–1292.

363 Revel, A. Defective endometrial receptivity. *Fertil Steril* 2012; **97** :1028-1032.

364 Sen S, Dixit A, Thakur C, Gokral J, Hinduja I, Zaveri K, Thangaraj K, Modi D. Association of
365 progesterone receptor gene polymorphism with male infertility and clinical outcome of ICSI.
366 *Journal of assisted reproduction and genetics*. 2013; **30** :1133-1139.

367 Seli E. Mitochondrial DNA as a biomarker for in-vitro fertilization outcome.
368 *Curr Opin Obstet Gynecol*, 2016; **28** :158-163.

369 Treff NR, Zhan Y, Tao X, Olcha M, Han M, Rajchel J, Morrison L, Morin SJ, Scott Jr RT. Levels
370 of trophoctoderm mitochondrial DNA do not predict the reproductive potential of sibling
371 embryos. *Hum Reprod*. 2017; **32** :954-962.

372 Van de Velde H, De Vos A, Joris H, Nagy ZP, Van Steirteghem AC. Effect of timing of oocyte
373 denudation and micro-injection on survival, fertilization and embryo quality after
374 intracytoplasmic sperm injection. *Hum Reprod*. 1998; **13** :3160-3164.

375 Victor AR, Brake AJ, Tyndall JC, Griffin DK, Zouves CG, Barnes FL, Viotti M. Accurate
376 quantitation of mitochondrial DNA reveals uniform levels in human blastocysts irrespective of
377 ploidy, age, or implantation potential. *Fertil Steril* 2017; **107** :34-42.

378 Wells D, Kaur K, Grifo J, Glassner M, Taylor JC, Fragouli E, Munne S. Clinical utilisation of a
379 rapid low-pass whole genome sequencing technique for the diagnosis of aneuploidy in human

- 380 embryos prior to implantation. *Journal of medical genetics*. 2014; **51**:553-62.
- 381 Wells D. Mitochondrial DNA quantity as a biomarker for blastocyst implantation potential. *Fertil*
- 382 *Steril*. 2017; **108** :742-747.