Title

Role of Triose Phosphate Utilization in photosynthetic response of rice to variable carbon dioxide levels and plant source-sink relations

Running Title

Role of TPU in photosynthesis under source-sink imbalance

Authors

Denis Fabre^{1,2}, Xinyou Yin³, Michael Dingkuhn^{1,2}, Anne Clément-Vidal^{1,2}, Sandrine Roques^{1,2}, Lauriane Rouan^{1,2}, Armelle Soutiras^{1,2}, Delphine Luquet^{1,2}

Contact information

1: CIRAD, UMR AGAP, F-34398 Montpellier, France.

2: Univ Montpellier, CIRAD, INRA, Montpellier SupAgro, Montpellier, France

3: Centre for Crop Systems Analysis, Department of Plant Sciences, Wageningen University & Research, Wageningen, Netherlands

Email address of each authors:

denis.fabre@cirad.fr

xinyou.yin@wur.nl

michael.dingkuhn@cirad.fr

anne.clement-vidal@cirad.fr

sandrine.roques@cirad.fr

lauriane.rouan@cirad.fr

armelle.soutiras@cirad.fr

delphine.luquet@cirad.fr

Corresponding author

Denis Fabre

CIRAD, UMR AGAP, F-34398 Montpellier, France

denis.fabre@cirad.fr

Phone: +33 4 67 61 71 40

Number of tables: 2

Number of figures: 5

Word count: 6270

Number of supplementary data: 5 (4 tables, 1 figure)

Highlight

This study provide new insights in the effect of C source-sink relationships on rice photosynthesis. TPU should be considered in photosynthesis studies under severe source-sink imbalance at elevated CO₂.

Abstract

This study aimed to understand the physiological bases of rice photosynthesis response to C source-sink imbalances, with focus on dynamics of the photosynthetic parameter TPU (Triose Phosphate Utilization). A dedicated experiment was replicated twice on IR64 indica rice cultivar in controlled environments. Plants were grown under the current ambient CO₂ concentration until heading, thereafter, two CO₂ treatments (400 and 800 $\mu\text{mol mol}^{-1}$) were compared in the presence and absence of a panicle pruning treatment modifying the C sink. At two weeks after heading, photosynthetic parameters derived from CO₂ response curves, and nonstructural carbohydrate content of flag leaf and internodes were measured 3-4 times of day. Spikelet number per panicle and flag leaf area on the main culm were recorded. Net C assimilation and TPU decreased progressively after midday in panicle-pruned plants, especially under 800 $\mu\text{mol mol}^{-1}$. This TPU reduction was explained by sucrose accumulation in the flag leaf resulting from the sink limitation. It is suggested that TPU is involved in rice photosynthesis regulation under elevated CO₂ conditions, and that sink limitation effects should be considered in crop models.

Keyword index

Rice; CO₂ enrichment; triose phosphate utilization; source-sink; photosynthesis; sink feedback; sucrose; climate change.

Introduction

Increasing world population and negative effects of global climate change on agricultural production require increased and more climate-resilient crop yields (Ainsworth, 2008; Ort et al., 2015; von Caemmerer et al., 2012). Rice (*Oryza sativa* L.) is the staple food for almost half of the population on Earth (GRiSP, 2013). To meet rice demand in 2050, its production has to increase by 2.4% annually until 2050 (Mohanty et al., 2013; Ray et al., 2013). This must be achieved in the context of climate change that is expected to have mostly negative effects on crop yields (Porter et al., 2014). But air CO₂ elevation (e-CO₂), expected to reach 600 to 700 $\mu\text{mol mol}^{-1}$ in 2050 (IPCC 2016), will affect C₃ crops like rice positively if efficiently used by photosynthesis. To achieve the production goal, leaf photosynthesis is a key leverage for improving crops (Evans, 2013; Lawson et al., 2012; Long et al., 2015; Ort et al., 2015), including rice (Makino, 2011; Yoshida et al., 2008; Yoshida and Horie, 2009).

A key requirement for achieving high crop productivity is to optimize carbon source-sink balance in the plants. E-CO₂ can perturb plant carbon (C) source-sink balance as it can increase C source more than the sink (White et al., 2016), leading to leaf carbohydrate accumulation that may down-regulate photosynthesis (Burnett et al., 2016; Paul and Foyer, 2001; Shimono et al., 2010; White et al., 2016). Source-sink interactions have been intensively studied during the last two decades (Chang et al., 2017). Whether and when plant growth and production is limited by C source (chiefly, photosynthesis) or sink (demand for organ growth) is still a key research question for agronomists, plant physiologists, biochemists and crop modelers (Burnett et al., 2016).

For agronomists, this question is particularly relevant during the grain filling period (Tang et al., 2017; Wei et al., 2018; Yang and Zhang, 2010; Zhang et al., 2017). Some studies used pruning treatments to manipulate the C source (leaf) and/or sink (grains) (Cock and Yoshida, 1973; Hasegawa et al., 2013, 2016; Jing et al., 2016; Nakano et al., 1995, 2017; Shimono et al., 2010; Shinano et al., 2006), and some of these were conducted with an e-CO₂ treatment. Conflicting results were reported regarding the response of photosynthesis to e-CO₂ but all studies agreed that plants with larger sink capacity benefitted more from e-CO₂.

Several physiological studies dealt with the role of non-structural carbohydrate (NSC) in C source-sink relationships under abiotic constraints such as drought (e.g. Dingkuhn et al., 2007). Experimental manipulations of plant C source and/or sink strength demonstrated that

photosynthetic rate depends on C sink strength (Ainsworth and Bush, 2011; Lemoine et al., 2013; Osorio et al., 2014). Accumulation of NSC commonly occurs in leaves of plants grown under e-CO₂ but down-regulation of photosynthesis is not always observed (Leakey et al., 2009; Wang et al., 2015), suggesting that feedbacks on photosynthesis are complex. Part of this complexity might be explained by partitioning of leaf NSC between sucrose and starch, controlled by day length (Mengin et al., 2017; Pokhilko and Ebenhoh, 2015; Sharkey, 2015; Sulpice et al., 2014), other environmental variables such as water deficit (Luquet et al., 2008), or time of day (Bläsing et al., 2005; Gibon et al., 2006). Plants lacking in sink capacity show reduced phloem loading. Rice is particularly effective in its capacity to export NSC from source leaves, suggesting that its photosynthetic response to e-CO₂ should be efficient (Makino and Mae, 1999), but little is known on the mechanisms.

Biochemical studies on C source-sink relationships have focused on two key parameters: the utilisation of triose phosphate produced in the Calvin cycle for sucrose and starch synthesis, and ribulose biphosphate (RuBP) regeneration by inorganic phosphate (Pi) recycling, which is related to sugar turnover (Leegood and Furbank, 1986; Paul and Foyer, 2001; Paul and Pellny, 2003; Sharkey, 1985). Analysis of leaf photosynthesis classically considers three limiting steps according to a biochemical photosynthesis model (the FvCB model hereafter) described by Farquhar et al. (1980), later extended by Sharkey (1985), involving the key parameters: i) Rubisco activity (V_{cmax}), ii) photosynthetic electron transport rate (J_{max}) determining the ability to regenerate RuBP substrate for Rubisco, and iii) Triose Phosphate Utilization (TPU) driving the synthesis of sucrose from sugar precursors in the Calvin-Benson cycle. Thereby, TPU acts as a short-term sink that commits carbon to end-products and is closely linked to triose phosphate conversion into sucrose or starch. High sink capacity accelerates the utilization of triose phosphate for sugar synthesis and export *via* phloem. It accelerates Pi recycling and RuBP regeneration in the Calvin cycle (Gibson et al., 2011; Kant et al., 2012; Kaschuk et al., 2009; Paul and Foyer, 2001; Paul and Pellny, 2003). TPU limitation occurs primarily at high CO₂ or sink-limited situations (Leegood and Furbank, 1986; Sharkey, 1985).

The FvCB model is commonly used as a module in crop models (Wu et al., 2016). Currently, photosynthesis is thought to be limited mainly by either V_{cmax} or J_{max} , whereas TPU has received less attention and is mostly ignored by crop models (Long and Bernacchi, 2003; von Caemmerer, 2000) because its regulation is largely unknown (Yang et al., 2016). However, TPU as a link between sugar production (source) and consumption (sink) may become

functionally important for crop models when addressing future climatic scenarios and e-CO₂ (Busch and Sage, 2016), particularly in sink-limiting situations (Asseng et al., 2017; Lombardozzi et al., 2017).

As the relations between source and sink activities at the crop, plant and process levels are complex, and there is a need to integrate the different levels. For this purpose, the present study aims to explore the role of TPU in the regulation of photosynthesis in response to C source-sink relationships. A dedicated experiment was designed to observe photosynthetic parameters, the dynamics of C source-sink ratio at plant or leaf level, and NSC partitioning between soluble sugars and starch. Results are expected to provide insights on whether TPU influences photosynthetic rate in current and future climatic scenarios, and should thus be considered in crop modelling.

Material and methods

Plant material and growth conditions

Seeds of high yielding *indica* rice cultivar from the Philippines, IR64, were germinated on wet filter paper and transplanted to 4L pots filled with EGO 140 substrate (17%N-10%P-14%K, pH = 5). Basal fertilizer was applied using Basacot 6 M (Compo Expert) at 2 g l⁻¹, 11%N-9%P-19%K +2%Mg. A second application was performed (topdressing) just before the heading stage to avoid post-floral nitrogen deficiency. Experiment was undertaken twice in the same growth chambers, in November 2016 (Exp1) and February 2017 (Exp2), using the same environmental conditions.

For each experiment, 60 plants were grown and divided between two identical growth chambers (microclima MC1750E, Snijders, Netherlands) at CIRAD, Montpellier, France. The two chambers were maintained at 12-h photoperiod, with day/night temperatures of 29/22°C, air humidity of 65/80% and daytime radiation of 1200 μmol photons m⁻² s⁻¹ photosynthetically active radiation (PAR) at plant tops. The 30 pots per chamber were rotated regularly to compensate for heterogeneity. They were arranged at 35-cm plant spacing in a completely randomized design with five replicates (potted plants). Pots were irrigated to maintain soil moisture at field capacity level.

At heading stage (80 days after transplanting), all panicles of half of the plants in each growth chamber were excised (pruning treatment PR, first experimental factor). Non-PR plants were called controls. The second factor was CO₂ treatment: In chamber 1, CO₂ level was set at 400 $\mu\text{mol mol}^{-1}$ during the whole experiment (ambient treatment); in chamber 2, CO₂ level was maintained at 400 $\mu\text{mol mol}^{-1}$ until the onset of heading, then switched to 800 $\mu\text{mol mol}^{-1}$ (e-CO₂ treatment) for 15 days, the main period of grain filling (Cho et al., 1988). At the end of the e-CO₂ period, physiological and biochemical measurements were performed. The combination of PR and CO₂ treatments at grain filling stage was chosen to achieve maximal C source-sink differences and to avoid the appearance of new sinks (panicles) during differential treatments.

For Exp1 in each growth chamber, photosynthesis, biochemical and biomass measurements (see details below) were carried out at three times of day: morning, midday, and afternoon, at +1h, +6h, and +9h after dawn, respectively, on 5 consecutive days. Measurements were done on a total of 60 plants (2 PR treatments x 3 times of day of sampling x 2 e-CO₂ levels x 5 biological replications).

For Exp2, in each growth chamber and for each treatment, photosynthesis, biochemical and biomass measurements carried out at midday, afternoon and evening; at +6h, +9h and +11h after dawn, respectively. As for Exp1, these were done for 5 consecutive days, resulting in a total of 60 measured plants.

Leaf photosynthesis measurement

Leaf photosynthesis parameters were measured on the flag leaf on the main culm 2 weeks after heading, using two portable photosynthesis systems (GFS-3100, Walz, Germany) identically calibrated and used to measure simultaneously plants at each CO₂ level. The measurements were made *in situ* using saturating PPFD light (1500 $\mu\text{mol m}^{-2} \text{s}^{-1}$ of PAR), controlled leaf temperature at 29°C, relative humidity in the cuvette set at 65%, and constant air flow rate through the cuvette of 800 ml min^{-1} . We used a large exchange area cuvette of 8 cm^2 to limit border effects known to affect photosynthesis measurement at high [CO₂] (Long and Bernacchi, 2003). Net photosynthesis CO₂ response curves (A/C_i) were obtained over a range of external CO₂ levels in the following order: 400, 300, 200, 100, 50, 400, 600, 800, 1000, 1200, 1400, 1600, 1800 and 2000 $\mu\text{mol mol}^{-1}$. At each step, gas exchange variables were recorded upon

reaching steady-state (7-8 min per step, coefficient of variation <1%). In subsequent analysis, net photosynthesis (A), stomatal conductance (g_s) and intercellular CO₂ concentration (C_i) were determined as the value measured at the 400 $\mu\text{mol mol}^{-1}$ CO₂ step of the curve. Chlorophyll fluorescence was measured for each CO₂ step simultaneously using Walz PAM-fluorimeter 3055FL, integrated into the photosynthesis equipment. The steady-state fluorescence yield (F_s) was measured after registering the gas-exchange parameters. A saturating light pulse (8000 $\mu\text{mol m}^{-2} \text{s}^{-1}$ during 0.8 s) was applied to achieve the light-adapted maximum fluorescence (F_m'). The operating PSII photochemical efficiency (ϕPSII) was determined as $\phi\text{PSII} = (F_m' - F_s)/F_m'$.

To fit the FvCB model of C₃ photosynthesis to experimental data, we used non-linear fitting procedure developed by Sharkey (2016), version 2, using the Rubisco kinetic parameters determined by temperature response functions according to Bernacchi (2002). The three main photosynthesis limitations, maximum carboxylation rate (V_{cmax}), electron transport rate (J_{max}), and triose phosphate utilisation (TPU), were estimated simultaneously, along with mesophyll conductance (g_m), by minimizing the sum of squares of the residuals. Independent measurements of day-time respiration (R_d) were made on some plants using the procedure of Yin et al. (2011), and an average value of R_d was used as a constant in the fitting procedure to avoid over-parameterization. Fluorescence measurements of ϕPSII were used to study the rate-limiting process for each level on the CO₂ curve, particularly to study the transition to TPU limitation as ϕPSII declines at high C_i (Sharkey 2016). To allow treatment comparisons, all parameters were scaled to a constant temperature of 25°C. In total, 120 CO₂ response curves were analyzed.

Sugar content analysis

Immediately after A/C_i curve measurements, the same leaf was sampled to measure non-structural carbohydrate content (NSC: starch, sucrose, glucose, fructose). Segments of the corresponding culm (top internode below the peduncle and bottom-most elongated internode) were also analyzed. Prior to grinding by ball grinder (Mixer mill MM 200, Retsch, Germany), the samples were frozen in liquid nitrogen. Sugars were extracted 3x from 20 mg samples with 1 mL of 80% ethanol for 30 min at 75°C, then centrifuged 10 min at 9500 g (Mikro 200, Hettich centrifuge). Soluble sugars (sucrose, glucose, and fructose) were contained in the supernatant and starch in the sediment. Supernatant was filtered in the presence of polyvinyl

polypyrrolidone and activated carbon to eliminate pigments and polyphenols. After evaporation of solute with Speedvac (RC 1022 and RCT 90, Jouan SA, Saint Herblain, France), soluble sugars were quantified by high performance ionic chromatography (HPIC, standard Dionex) with pulsed amperometric detection (HPAE-PAD). The sediment was solubilized with 0.02 N NaOH at 90°C for 1.5 h then hydrolyzed with α -amylglucosidase at 50°C, pH 4.2 for 1.5 h. Starch was quantified as described in Boehringer (Pomeranz and Meloan, 1994) with 5 μ L of hexokinase (glucose-6-phosphate dehydrogenase), followed by photometry of NADPH at 340 nm (spectrophotometer UV/VIS V-530, Jasco Corporation, Tokyo, Japan).

Leaf Nitrogen Content and Mass per Area

On each plant, segments of the leaf used for measuring CO₂ curve was used for determining the nitrogen content in % dw (Nm; mg N g⁻¹ dw of leaf blade) and specific leaf area (SLA; cm² g⁻¹). Nitrogen content per leaf area (Na; g N m⁻²) was obtained as Nm divided by SLA. The area of each sample was measured with a leaf area meter (Li-3100 Li-Cor) then oven-dried until constant weight (48 h at 70°C). Total nitrogen (N) was analyzed based by Dumas combustion method using a LECO TruMac Nitrogen analyzer, and potassium content (K) was measured in addition in Exp2 using an ICP-OES spectrometer 700 Series (Agilent Technologies). A relative indicator of chlorophyll content, SPAD, was also measured on the same leaf using a SPAD-502 (Minolta, Ltd., Japan).

Plant growth and biomass measurements

After sampling for biochemical analyses, all the aerial parts of plants were collected. Leaf blade, sheath, culm and panicle dw per plant (DM) were measured after drying samples at 70°C during 48 h (adding *a posteriori* the DM of organ segments sampled previously). Tillers and panicles were counted and total plant green leaf area measured, using a leaf area meter (Li-3100 Li-Cor, Lincoln, NE, USA). A proxy for the source-sink ratio was estimated at the time of photosynthesis measurements as the main-culm flag leaf area to fertile spikelet number ratio.

Statistical analysis

A three-way analysis of variance (ANOVA) of pruning treatment (PR), CO₂, sampling time and interaction effects on each measured parameter was performed for each experiment combined using the PROC MIXED method of the SAS package (SAS Institute Inc., NC, USA, version 9.04). A multiple comparison of means and Tukey's test ($\alpha=0,05$) was then performed.

Carbohydrate variables were log-transformed to stabilize variance. An analysis of covariance was performed to study the relationship between TPU, sugar contents, CO₂ and pruning treatments, using the PROC GLIMMIX method of the SAS package. Blocking effects (time of the day) were considered as random effects (Piepho et al., 2003). No experiment effect was observed on parameters measured at the same time in both experiments (illustrated by box plots in Fig. S1 for *A*, TPU and flag leaf sucrose content only).

Results

Photosynthetic parameter responses to C source-sink imbalance

Under ambient [CO₂], leaf photosynthesis (*A*) was significantly reduced by PR treatment ($P < 0.001$, Table S1), mainly in the afternoon during which *A* declined for all treatments. This was supported by a significant interaction observed between pruning treatment and time of day of measurement ($P < 0.001$, Table S1). This result was amplified under elevated [CO₂], with a reduction of *A* by 50% in the evening for PR compared to control plants (Fig. 1A).

No significant effects of the experimental factors were observed on leaf chlorophyll content (SPAD). Stomatal conductance (g_s) decreased along the day at both CO₂ concentrations, with significant effects of time of day ($P < 0.001$, Table S1). g_s was significantly decreased by PR treatment ($P < 0.001$) which interacted with [CO₂] ($P < 0.05$) without any significant variation of intercellular CO₂ concentration (C_i) (Table 1). Decrease of g_s was significant only in the afternoon under ambient [CO₂] but already from midday onwards under elevated [CO₂].

Before estimating g_m and other derived parameters from measured A/C_i curves, we assessed the shape of A/C_i response curves (replicate means) for each treatment along the day (Fig. 2A). Within high C_i levels, *A* responded little to a change in C_i . It even declined with increasing C_i for the PR-treated plants (Fig. 2A), suggesting an inhibition of *A* by TPU limitation. It is difficult to rely on only A/C_i curves to determine the transition from RuBP-regeneration limitation to TPU limitation since they usually occur together under high [CO₂] (Bernacchi et al., 2013; Long and Bernacchi, 2003). We used chlorophyll fluorescence-based data on the operating efficiency of PSII electron flow ($\phi PSII$) measured concomitantly with *A* to detect the C_i above which TPU limited *A*. When this is the case, $\phi PSII$ declines (Sharkey, 2016). The decline of $\phi PSII$ was observed above a C_i of 825 $\mu\text{mol mol}^{-1}$ in the evening, in plants exposed

to ambient $[\text{CO}_2]$ and pruning treatment (Fig. 2B, left). It occurred at C_i above 742 and 363 $\mu\text{mol mol}^{-1}$ under elevated CO_2 condition in the afternoon and in the evening, respectively (Fig. 2B, right). As these C_i thresholds were mostly higher than the C_i at A measurement (Fig. 2B), TPU did not limit A under the experimental conditions. However, under the most severe sink limitation (PR, 800 $\mu\text{mol mol}^{-1}$ $[\text{CO}_2]$, evening) TPU was close to limiting levels.

We applied the procedure of Sharkey (2016) to fit each A/C_i curve to derive estimates of g_m and biochemical photosynthetic parameters. All values of g_m were high, $> 10 \mu\text{mol m}^{-2} \text{s}^{-1} \text{Pa}^{-1}$ (Table 1), values known not to limit photosynthesis. Therefore, our estimates of g_m did not explain differences in A among treatments or times of day.

A significant decrease was observed for V_{cmax} in response to PR treatment (for both experiments: $P < 0.001$, Table S1). However, this reduction depended on CO_2 treatment (interaction PR $\times$ CO_2 at $P < 0.05$, Table S1) and was significant only in the afternoon and evening under elevated CO_2 (Table 1). Mean reduction in PR was 29% compared to control in the afternoon under elevated CO_2 (Fig. 1C). Regarding J_{max} , a significant effect of PR treatment was observed ($P < 0.001$, Table S1), despite no significant numerical decrease of J_{max} in PR treatment compared to control as shown in Table 1 and Fig. 1D. A time-of-day effect was also observed ($P < 0.05$).

Although A/C_i curves for control plants grown at 400 $\mu\text{mol mol}^{-1}$ $[\text{CO}_2]$ did not show TPU limitation (Fig. 2), TPU was significantly reduced by PR treatment ($P < 0.001$, Table S1). A decline of TPU after noon was observed in both $[\text{CO}_2]$ treatments (Fig. 1B), resulting in a highly significant time-of-day effect ($P < 0.001$). This decrease was particularly strong under e- CO_2 when combined with PR treatment, which led to significant interaction effects (PR $\times$ CO_2 , $P < 0.05$). In this latter situation, significant differences between control and PR plants were observed in the afternoon. The decrease of TPU caused by PR in the evening was 40% under elevated $[\text{CO}_2]$ and 13% under ambient $[\text{CO}_2]$ (Fig. 1B).

Nonstructural carbohydrate response to C source-sink imbalance

Leaf sucrose concentration in PR plants was significantly higher than in control plants in the afternoon ($P < 0.001$ for both PR and time-of-day effects, Table S2). No interaction effects between these factors were observed. The $[\text{CO}_2]$ effect on leaf sucrose was smaller ($P < 0.05$). Hexose concentration in the flag leaf was not affected by any of the experimental factors.

PR reduced sucrose concentration in the basal internode on the main culm ($P < 0.001$, Tables 2 and S2), without significant variations along the day. Similar results were observed for hexose concentration in the lower internode, but at about 35-fold lower concentrations than sucrose (Table 2). As soluble sugar content (hexose and sucrose) was similar in basal and upper internodes, results are presented only for basal internodes.

PR increased starch concentration in both top and bottom internodes on the main culm ($P < 0.001$; Tables 2 and S2), whereby no interaction between CO_2 and PR treatments was observed. No time-of-day effect was observed for starch concentration in internodes.

No PR and CO_2 effects were observed on leaf starch concentration, but there was a significant time-of-day effect ($P < 0.001$, Tables 2 and S2), causing a continuous increase of leaf starch concentration along the day (Table 2).

Plant growth response to C source-sink imbalance

PR significantly increased culm dry matter (by 50-60%, $P < 0.001$, Table S3 and Table S4) and sheath dry matter (by 12-20%, $P < 0.001$). No $[\text{CO}_2]$ effect and no interaction between factors were observed (Table S3). Panicle dry weight sampled two weeks after heading in the control plants was 280% higher under elevated $[\text{CO}_2]$ compared to ambient $[\text{CO}_2]$ ($P < 0.001$, Table S3 and Table S4), suggesting a strong stimulation of CO_2 enrichment on grain filling. By contrast, none of the factors affected plant total leaf dry matter, tiller number, panicle number and the SLA of the flag leaves used for photosynthesis measurement (Table S3). The same was true for nitrogen and potassium contents of the flag leaf, except for a significant reduction ($P < 0.05$) of nitrogen content under elevated $[\text{CO}_2]$, particularly on control plants (Table S3 and S4). Dry matter of plant roots was also measured at the end of the experiments. No significant effect of experimental factors were observed (data not presented).

Correlations between photosynthetic and biochemical parameters

A positive linear correlation was observed between A and TPU ($R^2 = 0.64$, $P < 0.001$) across all combinations of $[\text{CO}_2]$ and pruning treatments, and across all times of day (Fig. 3). The strongest treatment-specific correlation was observed when PR treatment was combined with high $[\text{CO}_2]$, i.e. for the treatment combination causing the highest C source-sink ratio ($R^2 = 0.72$,

P<0.001). The corresponding correlation between A and V_{cmax} was also significant but weaker ($R^2=0.60$; data not presented).

Analysis of covariance was performed to study the relationship between nonstructural carbohydrate and TPU variations. Flag leaf sucrose concentration was by far the most predictive factor of TPU variation ($P<0.001$) (Table S5). This was supported by the negative, linear correlations ($R^2=0.66$ for controls, $R^2=0.40$ for PR) observed between flag leaf sucrose concentration and TPU (Fig. 4). The two linear correlations showed a similar slope (-5.4 for control and -6.1 for pruned) but with lower TPU value at the intercepts in the case of pruned plants. An effect of starch concentration in the lower internodes was also observed ($P=0.01$) but it was smaller than that of leaf sucrose.

Finally, a negative correlation was found between TPU and plant C source-sink ratio measured two weeks after heading (R^2 of 0.45, $P<0.01$; Fig. 5), defined as the ratio of flag leaf area over fertile spikelet number of the corresponding panicle (measured only for Exp2).

Discussion

The physiology and biochemistry of leaf photosynthesis of major crops such as rice are well studied. So are the relations between sources, sinks and the formation of grain yield at the plant or crop scale. These processes are necessarily inter-dependent but little is known on the feedbacks causing interaction. We hypothesized that (1) source-sink imbalances are locally expressed as variations of TPU in the leaf, and (2) TPU would limit photosynthetic rate when C_i exceeds a critical level. To test the hypothesis, we manipulated the source with CO_2 enrichment and the sink with panicle pruning. The results confirmed hypothesis (1). However, in our experiments C_i did not exceed critical levels causing TPU limitation for A (Hyp.2), although it came close to that level in the afternoon under combined pruning and e- CO_2 . The strong reductions of A , accompanied by local accumulation of assimilates, confirmed the presence of feedback inhibition of photosynthesis under sink limitation.

Photosynthesis down-regulation under C sink limitation

Elevated $[\text{CO}_2]$ enhances plant C source capacity in C_3 plants and potentially, if plant sinks are insufficiently plastic, the C source-sink ratio. Our study showed that photosynthesis decreased along the day. The extent of the decrease depended on C sink limitation induced by sink pruning and/or source stimulation with CO_2 . Declining photosynthesis along the day was previously

reported under non-modified C source-sink balance (Ishihara and Saitoh, 1987; Koyama and Takemoto, 2014; Yang et al., 2008). Our observations on control plants under ambient $[\text{CO}_2]$ confirmed this trend, whereby enhanced source and pruned sinks further amplified it. Reductions in A attained 50% for both factors combined at the end of the day despite constant light resources. Sink limitation effects of this magnitude have not been noticed for rice previously. A similar effect, however, has been reported for wheat (King et al., 1967) after removing ears under ambient $[\text{CO}_2]$.

In the present study, CO_2 enrichment was applied during two weeks following heading. A small but significant reduction of N content per leaf area was observed under high $[\text{CO}_2]$, particularly in control plants. As indicated by leaf N concentration per unit dry mass, which was 36 mg g^{-1} for the elevated CO_2 treatment, N was above the empirical observation to affect growth, reported to be 28 mg g^{-1} (Seneweera et al., 2005: study made at $700 \mu\text{mol mol}^{-1} \text{CO}_2$). Leaf N concentration can decrease under CO_2 enrichment due to dilution, causing reduced photosynthesis (Ainsworth and Long, 2005; Leakey et al., 2009; Nakano et al., 1995; Yin 2013). This was avoided in this study by limiting CO_2 treatment to two weeks, at a stage when leaves were not expanding anymore. It was also reported that under elevated CO_2 and sink limitation, a high leaf N content could alleviate a photosynthetic down-regulation during the day (Makino et al., 1997; Seneweera et al., 2002). In our experiments, however, a diurnal decline of photosynthesis happened in all treatments despite ample N resources.

We also investigated leaf potassium concentration in Exp2 because K deficiency potentially affects photosynthesis through stomatal responses *via* osmoregulation in guard cells (Jin et al., 2011; Wang et al., 2013; Weng et al., 2007), and also assimilate transport in phloem (Gerardeaux et al., 2010). No potassium deficiency was observed that could explain the observed variations in photosynthesis.

A reduction of stomatal conductance was observed under sink pruning treatment, as reported for many plants, e.g., citrus (Nebauer et al., 2011; Urban, 2004) and coffee (DaMatta et al., 2008). Compared to control plants, panicle-pruned plants showed a smaller increase of photosynthetic rate in response to e- CO_2 , whereby pruning always reduced stomatal conductance. However, there was no significant difference in C_i between control and pruned plants when measured at a given atmospheric $[\text{CO}_2]$. Pruning thus decreased A at an unchanged C_i level, indicating that the photosynthetic capacity of the leaf was affected. Shimono et al.

(2010) also reported that rice plants with pruned panicles under ambient and elevated CO₂ had unaltered C_i levels.

No CO₂ effect was observed on stomatal conductance, in contrast to reports by Ainsworth (2008) and Yoshimoto et al. (2005). In our study, a supplemental dose of N fertilizer was applied just before heading stage. This might have maintained high stomatal conductance as previously shown in rice (Shimoda, 2012; Shimoda and Maruyama, 2014). Similar to another study evaluating short-term CO₂ enrichment effects on mesophyll conductance (Tazoe et al., 2009), no PR and CO₂ effect was observed on mesophyll conductance g_m, a parameter known to be sensitive to environment and estimation method (Flexas et al., 2008; Pons et al., 2009; Singaas et al., 2004; Sun et al., 2014). In our experimental conditions involving a 14-day CO₂ enrichment, g_m was very high and thus did not limit photosynthesis. Rice generally has high g_m as compared with other species (van der Putten et al., 2018), probably because rice leaves have high chloroplasts coverage on the mesophyll cell periphery (Busch et al., 2013). Therefore, g_m was not responsible for the decline in *A* observed under sink limitation. In addition, no difference was observed for SLA, a morphological trait that can affect genotypic photosynthetic capacity (Dingkuhn et al., 1998).

The down-regulation of photosynthesis observed under elevated CO₂ in the afternoon under C source-sink imbalance is a phenomenon that escapes observation if photosynthesis time courses are studied day-to-day and not within the day. This phenomenon is not captured in measurements at daily intervals (e.g., Makino and Mae, 1999), commonly done in the morning or at noon (Pérez-Harguindeguy et al., 2013). Our results suggest that it is crucial to capture diurnal changes of photosynthesis when studying source-sink effects on photosynthetic rate, or when estimating cumulative photosynthesis for a day.

TPU effect on photosynthesis under sink limitation

Generally, there are transitions from one limiting factor to another, although they may be masked by a concomitant decline of several factors through feedbacks. Along *A*/C_i curves, *A* is limited by Rubisco activity (characterized by *V*_{max}) at low C_i, by RuBP regeneration (characterized by *J*_{max}) at higher C_i, and potentially by TPU at even higher C_i. A TPU limitation is characterized by a lack of sensitivity of *A* to, or by a slight decline of *A* with, increases of CO₂ partial pressure (Sharkey, 1985). TPU limitation can be further ascertained by a decline of φPSII with increasing C_i (Sharkey, 2016). When TPU limitation occurs, photosynthesis is

affected by shortage of Pi (Paul and Foyer, 2001; Sharkey and Vanderveer, 1989) needed for ATP synthesis. In the absence of a strong C sink, TPU can be rate-limited by sucrose synthesis, thereby decreasing Pi recycling rate (Paul and Pellny, 2003). This limited regeneration is reflected in the rate at which the intermediate products of CO₂ fixation (triose-phosphate) are converted to starch and sucrose and accumulated locally.

The patterns of diurnal decline of TPU generally mirrored those of *A* (Fig. 1), resulting in a highly significant correlation between the two variables (Fig. 3). This correlation in itself, however, is not proof of a rate limitation by TPU. The *A/C_i* curves in Fig. 2A suggest that with increasing sink limitation (combinations of factors e-CO₂, pruning, time of day), *A* tended to plateau, or even decline, at high *C_i* values. Chlorophyll fluorescence-based quantum yield efficiency, measured concurrently with gas exchange under exposure to 14 levels of [CO₂], indicated the critical *C_i* above which TPU limited *A* for each treatment (Fig. 2B). In most of the situations studied, the critical *C_i* incurring TPU limitation was higher than the observed *C_i*. In one particular situation, however (e-CO₂, pruning, evening), the critical *C_i* was 363 μmol mol⁻¹ and thus, similar to the observed *C_i* value. Consequently, although TPU decreased most strongly among the biochemical photosynthetic parameters under sink limitation, it probably did not limit *A* except, possibly, for the treatment causing the strongest source-sink imbalance.

Further studies should determine if TPU can limit *A* in a climate change context with elevated ambient CO₂, and/or under lower temperatures to which TPU is very sensitive (Busch and Sage, 2016; Cen and Sage, 2005), and for plants having lesser phenotypic plasticity and assimilate transport capacity than rice. TPU limitations have been reported under low temperature (Sage and Kubien, 2007; Sage and Sharkey, 1987) or high CO₂ environments (Cen and Sage, 2005; Sharkey et al., 1986). In such cases, mutual adjustment of *V_{max}* and TPU is observed, as these parameters decrease concurrently and in strict stoichiometry (McClain and Sharkey, 2018). Indeed, we observed a strong decrease of *V_{max}* at high [CO₂] in the pruned plants (Fig. 1) concurrently with TPU, suggesting a co-adjustment. Changes in *V_{max}* may have contributed to the observed decrease in photosynthesis under e-CO₂ (Makino et al., 2000; Shimono et al., 2010), possibly due to a loss of Rubisco (Long et al., 2004). It was also shown that TPU limitation activates energy-dependent quenching (*q_E*), resulting in a deactivation of Rubisco (Sage et al., 1989; Sharkey et al., 1986). To enable photosynthesis, the carbon reduction cycle needs to regenerate RuBP, consuming ATP and NADPH produced through photosynthetic electron transport in the chloroplast. This process can be evaluated by *J_{max}* parameter (RuBP

regeneration) which can also limit photosynthesis. Although we observed a significant decrease in $J_{\max}$ during the day in response to sink pruning, variations were too small to explain the observed variation of A . The two parameters correlating most with A were $V_{\max}$ and TPU, suggesting a tight coordination between TPU and Rubisco capacity.

Our findings suggest that TPU and A of rice generally decline in the afternoon, and particularly when sink is restricted. This may potentially cause overestimation of whole-day photosynthesis in crop models that do not consider rate limitations to assimilate export from the leaf (Lombardozzi et al., 2017, 2018). TPU is situated at the interface between production and consumption (or removal) of photosynthates. Thus, the mechanisms controlling this parameter can only be understood in a whole-plant context including assimilate transport and partitioning among sinkd (Yang et al., 2016).

Sugar partitioning effect on photosynthesis and TPU regulation

Similar to our results, Morita et al., (2016), Shimono et al. (2010), Thompson et al. (2017) and Zhu et al. (2016) found that leaf sucrose concentration increased more than hexose concentrations under e-CO₂. No increase in starch concentration in the flag leaf was observed under those conditions (Shimono et al., 2010). This can be explained by the large capacity of rice to accumulate carbohydrates in culm. Moreover, leaf starch concentration in the flag leaf (100 to 300 $\mu\text{g cm}^{-2}$) was below empirical critical values (600 $\mu\text{g cm}^{-2}$) reported to affect photosynthesis in rice (Weng and Chen, 1991), suggesting that leaf starch accumulation did not affect A .

We observed an increase in starch concentration in culm internodes in panicle-pruned plants, probably because internodes acted as alternative sinks for the panicle. Culm sucrose remobilization decreased under pruning and possibly explained the increase in starch. It may have acted as a physiological signal regulating photosynthesis as reported for sugarcane (McCormick et al., 2009; Wang et al., 2018).

Flag leaf sucrose concentration was identified as the main nonstructural carbohydrate affected by pruning treatment and [CO₂], showing a continuous increase along the day. Photosynthesis is inhibited by leaf carbohydrate accumulation (Goldschmidt and Huber, 1992; Paul and Pellny, 2003). In this study, a negative linear relation was observed between TPU and leaf sucrose content. A theory of TPU control by Pi availability, mediated by sugar production, was

proposed by Paul and Pellny (2003). According to this theory, TPU can limit photosynthetic rate through a reduced export carbon from the Calvin-Benson cycle, which in turn is related to the rate at which sugar phosphates are dephosphorylated and end-products are produced. Thus, the production and export of sucrose is essential for sustaining photosynthesis. We suggest that in our study, leaf sucrose was exported to plant sinks such as the panicle in control plants, preventing excessive build-up in the leaf. For panicle-pruned plants, sucrose could not be exported sufficiently during the afternoon. Some export occurred to the top internode, where starch concentration increased, but sucrose concentration increased in the flag leaf due to the smaller sink (Huber and Huber, 1992; Paul and Foyer, 2001). In this case, Sucrose Phosphate Synthase (SPS) feedback inhibition occurs because of an increase in the phosphorylation state of the enzyme (Huber et al., 1989). It has been shown that SPS is a substrate for SNF-1 related protein kinases, modulating SPS activity when sucrose accumulates (Sugden et al., 1999). Inhibition of sucrose synthesis may reduce export of TP from the chloroplast, causing a drop in Pi in the cytosol, leading to decrease in TPU (Paul and Pellny, 2003).

While the negative correlation between TPU and flag-leaf sucrose concentration showed a similar slope for the panicle-pruned and control plants, TPU was lower in the pruning treatment at any given sucrose concentration (reduced intercept, Fig. 4). Thus, the TPU vs. [sucrose] relationship was not the same for control and panicle-pruning treatments, and [sucrose] alone could not explain the TPU decline under C source-sink imbalance. Possibly, additional feedbacks on TPU occurred, e.g. via phloem sucrose concentration. Sucrose concentration in the leaf phloem depends on the rate of sucrose loading at the source and unloading at the sink end (Chiou and Bush, 1998; Li et al., 2003). Panicle pruning probably led to high sucrose concentrations in the leaf phloem as sucrose transport is mainly operated by phloem in rice (Regmi et al., 2016). Photo-assimilates would build up in the mesophyll (Chiou and Bush, 1998) and decrease TPU as previously described.

The ratio of flag-leaf area over the fertile spikelet number of the corresponding panicle provides a rough proxy for local C source-sink ratio during grain filling. It correlated negatively with TPU (Fig. 5), suggesting that morphology-based phenotypic plasticity causing variation in C source-sink ratio can affect TPU. A recent study also reported the effect of sink strength on sucrose partitioning that may be used to increase grain yield in rice (Morey et al., 2018). More research is needed to understand how whole-plant source-sink interactions affect crops' ability to utilize rising CO₂ levels.

This study provided new insights into the effect of C source-sink relationships on rice photosynthesis and in particular, its parameter TPU. A significant down-regulation of photosynthesis (up to 50%) was demonstrated during the 2nd half of the day in response to sink limitation. TPU strongly decreased along with *A* and it was negatively correlated with flag-leaf sucrose concentration, suggesting sugar feedback inhibition of *A*. It is suggested that photosynthesis measurements performed in the morning, as commonly practiced, may not reliably represent the plant's diurnal photosynthetic performance, particularly under CO₂ enrichment or sink limitation.

Although TPU decline mirrored the decline of *A* under sink limitation, its rate-limiting effect on *A* could not be confirmed, except possibly at the end of the day for the combination of e-CO₂ and panicle pruning. Only under these specific conditions, the observed *C_i* was similar to the critical *C_i* above which quantum yield efficiency decreased. TPU may thus play an important role in photosynthesis regulation only under extreme source-sink imbalance that may occur in plants that poorly adjust sinks and assimilate transport to increased assimilation potential.

Based on these results, it will be interesting to explore the photosynthetic responses of genotypes differing in source-sink ratio and the adaptive plasticity of sinks in CO₂-enriched environments.

Acknowledgments

This work was financially supported by CIRAD, French Agricultural Research Centre for International Development; <http://www.cirad.fr>). We thank Audrey Dardou, Paul Pruvost, Christian Chaine, Remy Michel for their valuable assistance in technical support, and the biochemical phenotyping platform of the AGAP research unit at CIRAD.

Supplementary data

Fig. S1: Boxplot on various parameters for the two experiments

Table S1: Summary of 3-way analysis of variance of physiological flag leaf variables

Table S2: Summary of 3-way analysis of variance of non-structural carbohydrate concentrations

Table S3: Summary of 3-way analysis of variance of plant biomass and morphological variables

Table S4: Plant growth characteristics

Table S5: Type III test of fixed effects for co-variance analysis for TPU with carbohydrates and treatment

References

Ainsworth EA. 2008. Rice production in a changing climate: a meta-analysis of responses to elevated carbon dioxide and elevated ozone concentration. *Global Change Biology* **14**, 1642–1650.

Ainsworth EA, Bush DR. 2011. Carbohydrate Export from the Leaf: A Highly Regulated Process and Target to Enhance Photosynthesis and Productivity. *PLANT PHYSIOLOGY* **155**, 64–69.

Ainsworth EA, Long SP. 2005. What have we learned from 15 years of free-air CO₂ enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO₂. *New Phytologist* **165**, 351–372.

Asseng S, Kassie BT, Labra MH, Amador C, Calderini DF. 2017. Simulating the impact of source-sink manipulations in wheat. *Field Crops Research* **202**, 47–56.

Bernacchi CJ. 2002. Temperature Response of Mesophyll Conductance. Implications for the Determination of Rubisco Enzyme Kinetics and for Limitations to Photosynthesis in Vivo. *PLANT PHYSIOLOGY* **130**, 1992–1998.

Bernacchi CJ, Bagley JE, Serbin SP, Ruiz-Vera UM, Rosenthal DM, Vanloocke A. 2013. Modelling C₃ photosynthesis from the chloroplast to the ecosystem. *Plant, Cell & Environment* **36**, 1641–1657.

Bläsing OE, Gibon Y, Günther M, Höhne M, Morcuende R, Osuna D, Thimm O, Usadel B, Scheible W-R, Stitt M. 2005. Sugars and Circadian Regulation Make Major Contributions to the Global Regulation of Diurnal Gene Expression in Arabidopsis. *The Plant Cell* **17**, 3257–3281.

Burnett AC, Rogers A, Rees M, Osborne CP. 2016. Carbon source-sink limitations differ between two species with contrasting growth strategies: Source-sink limitations vary with growth strategy. *Plant, Cell & Environment* **39**, 2460–2472.

Busch FA, Sage RF. 2016. The sensitivity of photosynthesis to O₂ and CO₂ concentration identifies strong Rubisco control above the thermal optimum. *New Phytologist*.

Busch FA, Sage TL, Cousins AB, Sage RF. 2013. C₃ plants enhance rates of photosynthesis by reassimilating photorespired and respired CO₂. *Plant, Cell & Environment* **36**, 200–212.

- von Caemmerer S.** 2000. *Biochemical Models of Leaf Photosynthesis*.
- von Caemmerer S, Quick WP, Furbank RT.** 2012. The Development of C4 Rice: Current Progress and Future Challenges. *Science* **336**, 1671–1672.
- Cen Y-P, Sage RF.** 2005. The Regulation of Rubisco Activity in Response to Variation in Temperature and Atmospheric CO2 Partial Pressure in Sweet Potato. *Plant Physiology* **139**, 979–990.
- Chang T-G, Zhu X-G, Raines C.** 2017. Source-sink interaction: a century old concept under the light of modern molecular systems biology. *Journal of Experimental Botany* **68**, 4417–4431.
- Chiou TJ, Bush DR.** 1998. Sucrose is a signal molecule in assimilate partitioning. *Proceedings of the National Academy of Sciences of the United States of America* **95**, 4784–4788.
- Cho D, Jong S, Son S, Park YK.** 1988. Studies on the duration and rate of grain filling in rice (*Oryza sativa* L.). II. Difference between the parts of a panicle. *Kor J Crop Sci*, 5–11.
- Cock JH, Yoshida S.** 1973. Changing sink and source relations in rice (*Oryza sativa* L.) using carbon dioxide enrichment in the field. *Soil Science and Plant Nutrition* **19**, 229–234.
- DaMatta FM, Cunha RL, Antunes WC, Martins SCV, Araujo WL, Fernie AR, Moraes GABK.** 2008. In field-grown coffee trees source-sink manipulation alters photosynthetic rates, independently of carbon metabolism, via alterations in stomatal function. *The New Phytologist* **178**, 348–357.
- Dingkuhn M, Jones MP, Johnson DE, Sow A.** 1998. Growth and yield potential of *Oryza sativa* and *O. glaberrima* upland rice cultivars and their interspecific progenies. *Field Crops Research* **57**, 57–69.
- Dingkuhn M, Luquet D, Clément-Vidal A, Tambour L, Kim HK, Song YH.** 2007. Is plant growth driven by sink regulation? Implications for crop models, phenotyping approaches and ideotypes. *Frontis* **21**, 155–168.
- Evans JR.** 2013. Improving Photosynthesis. *PLANT PHYSIOLOGY* **162**, 1780–1793.
- Farquhar G v, von Caemmerer S von, Berry JA.** 1980. A biochemical model of photosynthetic CO2 assimilation in leaves of C3 species. *Planta* **149**, 78–90.
- Flexas J, Ribas-Carbó M, Diaz-Espejo A, Galmés J, Medrano H.** 2008. Mesophyll conductance to CO2: current knowledge and future prospects. *Plant, Cell & Environment* **31**, 602–621.
- Gerardeaux E, Jordan-Meille L, Constantin J, Pellerin S, Dingkuhn M.** 2010. Changes in plant morphology and dry matter partitioning caused by potassium deficiency in *Gossypium*

hirsutum (L.). Environmental and Experimental Botany **67**, 451–459.

Gibon Y, Usadel B, Blaesing OE, Kamlage B, Hoehne M, Trethewey R, Stitt M. 2006. Integration of metabolite with transcript and enzyme activity profiling during diurnal cycles in Arabidopsis rosettes. Genome Biology **7**, R76.

Gibson K, Park J-S, Nagai Y, et al. 2011. Exploiting leaf starch synthesis as a transient sink to elevate photosynthesis, plant productivity and yields. Plant Science **181**, 275–281.

Goldschmidt EE, Huber SC. 1992. Regulation of Photosynthesis by End-Product Accumulation in Leaves of Plants Storing Starch, Sucrose, and Hexose Sugars. Plant Physiology **99**, 1443–1448.

Hasegawa T, Sakai H, Tokida T, et al. 2013. Rice cultivar responses to elevated CO₂ at two free-air CO₂ enrichment (FACE) sites in Japan. Functional Plant Biology **40**, 148.

Hasegawa T, Sakai H, Tokida T, Usui Y, Yoshimoto M, Fukuoka M, Nakamura H, Shimono H, Okada M. 2016. Rice Free-Air Carbon Dioxide Enrichment Studies to Improve Assessment of Climate Change Effects on Rice Agriculture. Improving Modeling Tools to Assess Climate Change Effects on Crop Response *advancesinagric*, 45–68.

Huber SC, Huber JL. 1992. Role of Sucrose-Phosphate Synthase in Sucrose Metabolism in Leaves. Plant Physiology **99**, 1275–1278.

Huber JL, Huber SC, Nielsen TH. 1989. Protein phosphorylation as a mechanism for regulation of spinach leaf sucrose-phosphate synthase activity. Archives of Biochemistry and Biophysics **270**, 681–690.

Ishihara K, Saitoh K. 1987. Diurnal Courses of Photosynthesis, Transpiration, and Diffusive Conductance in the Single-leaf of the Rice Plants Grown in the Paddy Field under Submerged Condition. Japanese Journal of Crop Science **56**, 8–17.

Jin SH, Huang JQ, Li XQ, Zheng BS, Wu JS, Wang ZJ, Liu GH, Chen M. 2011. Effects of potassium supply on limitations of photosynthesis by mesophyll diffusion conductance in *Carya cathayensis*. Tree Physiology **31**, 1142–1151.

Jing L, Wu Y, Zhuang S, Wang Y, Zhu J, Wang Y, Yang L. 2016. Effects of CO₂ enrichment and spikelet removal on rice quality under open-air field conditions. Journal of Integrative Agriculture **15**, 2012–2022.

Kant S, Seneweera S, Rodin J, Materne M, Burch D, Rothstein SJ, Spangenberg G. 2012. Improving yield potential in crops under elevated CO₂: Integrating the photosynthetic and nitrogen utilization efficiencies. Frontiers in Plant Science **3**.

Kaschuk G, Kuyper TW, Leffelaar PA, Hungria M, Giller KE. 2009. Are the rates of photosynthesis stimulated by the carbon sink strength of rhizobial and arbuscular mycorrhizal

symbioses? Soil Biology and Biochemistry **41**, 1233–1244.

King RW, Wardlaw IF, Evans LT. 1967. Effect of assimilate utilization on photosynthetic rate in wheat. *Planta* **77**, 261–276.

Koyama K, Takemoto S. 2014. Morning reduction of photosynthetic capacity before midday depression. *Scientific Reports* **4**.

Lawson T, Kramer DM, Raines CA. 2012. Improving yield by exploiting mechanisms underlying natural variation of photosynthesis. *Current Opinion in Biotechnology* **23**, 215–220.

Leakey ADB, Ainsworth EA, Bernacchi CJ, Rogers A, Long SP, Ort DR. 2009. Elevated CO₂ effects on plant carbon, nitrogen, and water relations: six important lessons from FACE. *Journal of Experimental Botany* **60**, 2859–2876.

Leegood RC, Furbank RT. 1986. Stimulation of photosynthesis by 2% oxygen at low temperatures is restored by phosphate. *Planta* **168**, 84–93.

Lemoine R, Camera SL, Atanassova R, et al. 2013. Source-to-sink transport of sugar and regulation by environmental factors. *Frontiers in Plant Science* **4**.

LI C, WEISS D, GOLDSCHMIDT EE. 2003. Girdling Affects Carbohydrate-related Gene Expression in Leaves, Bark and Roots of Alternate-bearing Citrus Trees. *Annals of Botany* **92**, 137–143.

Lombardozzi D, Smith NG, Cheng SJ, Bonan GB, Dukes JS, Fisher R, Sharkey TD, Rogers A. 2017. The impact of including export limitation in photosynthetic models from leaf to global scales. *AGU Fall Meeting Abstracts* **21**.

Lombardozzi DL, Smith NG, Cheng SJ, Dukes JS, Sharkey TD, Alistair Rogers, Fisher R, Bonan GB. 2018. Triose phosphate limitation in photosynthesis models reduces leaf photosynthesis and global terrestrial carbon storage. *Environmental Research Letters* **13**, 074025.

Long SP, Ainsworth EA, Rogers A, Ort DR. 2004. Rising atmospheric carbon dioxide: plants FACE the Future*. *Annu. Rev. Plant Biol.* **55**, 591–628.

Long SP, Bernacchi CJ. 2003. Gas exchange measurements, what can they tell us about the underlying limitations to photosynthesis? Procedures and sources of error. *Journal of Experimental Botany* **54**, 2393–2401.

Long SP, Marshall-Colon A, Zhu X-G. 2015. Meeting the Global Food Demand of the Future by Engineering Crop Photosynthesis and Yield Potential. *Cell* **161**, 56–66.

Luquet D, Clément-Vidal A, Fabre D, This D, Sonderegger N, Dingkuhn M. 2008. Orchestration of transpiration, growth and carbohydrate dynamics in rice during a dry-down

cycle. *Functional Plant Biology* **35**, 689–704.

Makino A. 2011. Photosynthesis, Grain Yield, and Nitrogen Utilization in Rice and Wheat. *Plant Physiology* **155**, 125–129.

Makino A, Harada M, Sato T, Nakano H, Mae T. 1997. Growth and N allocation in rice plants under CO₂ enrichment. *Plant Physiology* **115**, 199–203.

Makino A, Mae T. 1999. Photosynthesis and plant growth at elevated levels of CO₂. *Plant and Cell Physiology* **40**, 999–1006.

Makino A, Nakano H, Mae T, Shimada T, Yamamoto N. 2000. Photosynthesis, plant growth and N allocation in transgenic rice plants with decreased Rubisco under CO₂ enrichment. *Journal of Experimental Botany* **51**, 383–389.

McClain AM, Sharkey TD. 2018. Triose phosphate utilization and beyond: from photosynthesis to end-product synthesis. *bioRxiv*, 434928.

McCormick AJ, Watt DA, Cramer MD. 2009. Supply and demand: sink regulation of sugar accumulation in sugarcane. *Journal of Experimental Botany* **60**, 357–364.

Mengin V, Pyl E-T, Alexandre Moraes T, Sulpice R, Krohn N, Encke B, Stitt M. 2017. Photosynthate partitioning to starch in *Arabidopsis thaliana* is insensitive to light intensity but sensitive to photoperiod due to a restriction on growth in the light in short photoperiods. *Plant, Cell & Environment* **40**, 2608–2627.

Mohanty S, Wassmann R, Nelson A, Moya P, Jagadish SVK. 2013. Rice and climate change: significance for food security and vulnerability. International Rice Research Institute.

Morey SR, Hirose T, Hashida Y, Miyao A, Hirochika H, Ohsugi R, Yamagishi J, Aoki N. 2018. Genetic Evidence for the Role of a Rice Vacuolar Invertase as a Molecular Sink Strength Determinant. *Rice* **11**.

Morita R, Inoue K, Ikeda K, Hatanaka T, Misoo S, Fukayama H. 2016. Starch Content in Leaf Sheath Controlled by CO₂-Responsive CCT Protein is a Potential Determinant of Photosynthetic Capacity in Rice. *Plant and Cell Physiology* **57**, 2334–2341.

Nakano H, Makino A, Mae T. 1995. Effects of panicle removal on the photosynthetic characteristics of the flag leaf of rice plants during the ripening stage. *Plant and cell physiology* **36**, 653–659.

Nakano H, Yoshinaga S, Takai T, et al. 2017. Quantitative trait loci for large sink capacity enhance rice grain yield under free-air CO₂ enrichment conditions. *Scientific Reports* **7**.

Nebauer SG, Renau-Morata B, Guardiola JL, Molina R-V. 2011. Photosynthesis down-regulation precedes carbohydrate accumulation under sink limitation in Citrus. *Tree Physiology* **31**, 169–177.

Ort DR, Merchant SS, Alric J, et al. 2015. Redesigning photosynthesis to sustainably meet global food and bioenergy demand. *Proceedings of the National Academy of Sciences of the United States of America* **112**, 8529–8536.

Osorio S, Ruan Y-L, Fernie AR. 2014. An update on source-to-sink carbon partitioning in tomato. *Frontiers in Plant Science* **5**.

Paul MJ, Foyer CH. 2001. Sink regulation of photosynthesis. *Journal of Experimental Botany* **52**, 1383–1400.

Paul MJ, Pellny TK. 2003. Carbon metabolite feedback regulation of leaf photosynthesis and development. *Journal of Experimental Botany* **54**, 539–547.

Pérez-Harguindeguy N, Díaz S, Garnier E, et al. 2013. New handbook for standardised measurement of plant functional traits worldwide. *Australian Journal of Botany* **61**, 167.

Piepho HP, Buchse A, Emrich K. 2003. A Hitchhiker’s Guide to Mixed Models for Randomized Experiments. *Journal of Agronomy and Crop Science* **189**, 310–322.

Pokhilko A, Ebenhoh O. 2015. Mathematical modelling of diurnal regulation of carbohydrate allocation by osmo-related processes in plants. *Journal of The Royal Society Interface* **12**, 20141357–20141357.

Pomeranz Y, Meloan CE. 1994. Enzymatic Methods. In: Pomeranz Y, In: Meloan CE, eds. *Food Analysis: Theory and Practice*. Boston, MA: Springer US, 506–531.

Pons TL, Flexas J, von Caemmerer S, Evans JR, Genty B, Ribas-Carbo M, Brugnoli E. 2009. Estimating mesophyll conductance to CO₂: methodology, potential errors, and recommendations. *Journal of Experimental Botany* **60**, 2217–2234.

Porter J, Xie L, Challinor AJ, Cochrane K, Howden S, Iqbal M, Lobell D, Travasso M. 2014. *Food security and food production systems*.

van der Putten PEL, Yin X, Struik PC. 2018. Calibration matters: On the procedure of using the chlorophyll fluorescence method to estimate mesophyll conductance. *Journal of Plant Physiology* **220**, 167–172.

Ray DK, Mueller ND, West PC, Foley JA. 2013. Yield Trends Are Insufficient to Double Global Crop Production by 2050. *PLOS ONE* **8**, e66428.

Regmi KC, Zhang S, Gaxiola RA. 2016. Apoplasmic loading in the rice phloem supported by the presence of sucrose synthase and plasma membrane-localized proton pyrophosphatase. *Annals of Botany* **117**, 257–268.

Sage RF, Kubien DS. 2007. The temperature response of C₃ and C₄ photosynthesis. *Plant, Cell & Environment* **30**, 1086–1106.

Sage RF, Sharkey TD. 1987. The Effect of Temperature on the Occurrence of O₂ and

CO₂ Insensitive Photosynthesis in Field Grown Plants. *Plant Physiology* **84**, 658–664.

Sage RF, Sharkey TD, Seemann JR. 1989. Acclimation of photosynthesis to elevated CO₂ in five C₃ species. *Plant Physiology* **89**, 590–596.

Seneweera SP, Conroy JP, Ishimaru K, Ghannoum O, Okada M, Lieffering M, Kim HY, Kobayashi K. 2002. Changes in source-sink relations during development influence photosynthetic acclimation of rice to free air CO₂ enrichment (FACE). *Functional plant biology* **29**, 945–953.

Seneweera S, Makino A, Mae T, Basra AS. 2005. Response of Rice to *p* (CO₂) Enrichment: The Relationship Between Photosynthesis and Nitrogen Metabolism. *Journal of Crop Improvement* **13**, 31–53.

Sharkey TD. 1985. Photosynthesis in intact leaves of C₃ plants: Physics, physiology and rate limitations. *The Botanical Review* **51**, 53–105.

Sharkey TD. 2015. Understanding carbon partitioning and its role in determining plant growth: Understanding carbon partitioning. *Plant, Cell & Environment* **38**, 1963–1964.

Sharkey TD. 2016. What gas exchange data can tell us about photosynthesis: Gas exchange data and photosynthesis. *Plant, Cell & Environment* **39**, 1161–1163.

Sharkey TD, Stitt M, Heineke D, Gerhardt R, Raschke K, Heldt HW. 1986. Limitation of Photosynthesis by Carbon Metabolism : II. O₂-Insensitive CO₂ Uptake Results from Limitation Of Triose Phosphate Utilization. *Plant Physiology* **81**, 1123–1129.

Sharkey TD, Vanderveer PJ. 1989. Stromal phosphate concentration is low during feedback limited photosynthesis. *Plant Physiology* **91**, 679–684.

Shimoda S. 2012. Contribution of leaf nitrogen to photosynthetic gas exchange in contrasting rice (*Oryza sativa* L.) cultivars during the grain-filling period. *Photosynthetica* **50**, 387–394.

Shimoda S, Maruyama A. 2014. Rice varietal differences in responses of stomatal gas exchange to supplemental nitrogen application. *Photosynthetica* **52**, 397–403.

Shimono H, Suzuki K, Aoki K, Hasegawa T, Okada M. 2010. Effect of panicle removal on photosynthetic acclimation under elevated CO₂ in rice. *Photosynthetica* **48**, 530–536.

Shinano T, Osawa M, Soejima H, Osaki M. 2006. Effect of panicle removal on cytokinin level in the xylem and nitrogen uptake activity of rice. *Soil Science & Plant Nutrition* **52**, 331–340.

Singsaas EL, Ort DR, Delucia EH. 2004. Elevated CO₂ effects on mesophyll conductance and its consequences for interpreting photosynthetic physiology. *Plant, Cell and Environment* **27**, 41–50.

Sugden C, Donaghy PG, Halford NG, Hardie DG. 1999. Two SNF1-Related Protein

Kinases from Spinach Leaf Phosphorylate and Inactivate 3-Hydroxy-3-Methylglutaryl-Coenzyme A Reductase, Nitrate Reductase, and Sucrose Phosphate Synthase in Vitro. *Plant Physiology* **120**, 257–274.

Sulpice R, Flis A, Ivakov AA, Apelt F, Krohn N, Encke B, Abel C, Feil R, Lunn JE, Stitt M. 2014. Arabidopsis Coordinates the Diurnal Regulation of Carbon Allocation and Growth across a Wide Range of Photoperiods. *Molecular Plant* **7**, 137–155.

Sun Y, Gu L, Dickinson RE, et al. 2014. Asymmetrical effects of mesophyll conductance on fundamental photosynthetic parameters and their relationships estimated from leaf gas exchange measurements: Asymmetrical mesophyll conductance effects. *Plant, Cell & Environment* **37**, 978–994.

Tang L, Gao H, Yoshihiro H, Koki H, Tetsuya N, Liu T, Tatsuhiko S, Xu Z. 2017. Erect panicle super rice varieties enhance yield by harvest index advantages in high nitrogen and density conditions. *Journal of Integrative Agriculture* **16**, 1467–1473.

Tazoe Y, von Caemmerer S, Badger MR, Evans JR. 2009. Light and CO₂ do not affect the mesophyll conductance to CO₂ diffusion in wheat leaves. *Journal of Experimental Botany* **60**, 2291–2301.

Thompson M, Gamage D, Hirotsu N, Martin A, Seneweera S. 2017. Effects of Elevated Carbon Dioxide on Photosynthesis and Carbon Partitioning: A Perspective on Root Sugar Sensing and Hormonal Crosstalk. *Frontiers in Physiology* **8**.

Urban L. 2004. Effect of fruit load and girdling on leaf photosynthesis in *Mangifera indica* L. *Journal of Experimental Botany* **55**, 2075–2085.

Wang J, Wang C, Chen N, Xiong Z, Wolfe D, Zou J. 2015. Response of rice production to elevated [CO₂] and its interaction with rising temperature or nitrogen supply: a meta-analysis. *Climatic Change* **130**, 529–543.

Wang J, Zhao T, Yang B, Zhang S. 2018. Sucrose Metabolism and Regulation in Sugarcane. *Journal of Plant Physiology & Pathology* **2017**.

Wang M, Zheng Q, Shen Q, Guo S. 2013. The Critical Role of Potassium in Plant Stress Response. *International Journal of Molecular Sciences* **14**, 7370–7390.

Wei H, Meng T, Li X, Dai Q, Zhang H, Yin X. 2018. Sink-source relationship during rice grain filling is associated with grain nitrogen concentration. *Field Crops Research* **215**, 23–38.

Weng J-H, Chen C-Y. 1991. Effect of Accumulated Nonstructural Carbohydrates on Photosynthesis of Rice Leaves. *Japanese Journal of Crop Science* **60**, 320–321.

Weng X-Y, Zheng C-J, Xu H-X, Sun J-Y. 2007. Characteristics of photosynthesis and functions of the water–water cycle in rice (*Oryza sativa*) leaves in response to potassium

deficiency. *Physiologia Plantarum* **131**, 614–621.

White AC, Rogers A, Rees M, Osborne CP. 2016. How can we make plants grow faster? A source–sink perspective on growth rate. *Journal of Experimental Botany* **67**, 31–45.

Wu A, Song Y, van Oosterom EJ, Hammer GL. 2016. Connecting Biochemical Photosynthesis Models with Crop Models to Support Crop Improvement. *Frontiers in Plant Science* **7**.

Yang F, Liang Z, Wang Z, Chen Y. 2008. Relationship Between Diurnal Changes of Net Photosynthetic Rate and Influencing Factors in Rice under Saline Sodic Stress. *Rice Science* **15**, 119–124.

Yang JT, Preiser AL, Li Z, Weise SE, Sharkey TD. 2016. Triose phosphate use limitation of photosynthesis: short-term and long-term effects. *Planta* **243**, 687–698.

Yang J, Zhang J. 2010. Crop management techniques to enhance harvest index in rice. *Journal of Experimental Botany* **61**, 3177–3189.

Yin X, Sun Z, Struik PC, Gu J. 2011. Evaluating a new method to estimate the rate of leaf respiration in the light by analysis of combined gas exchange and chlorophyll fluorescence measurements. *Journal of Experimental Botany* **62**, 3489–3499.

Yoshida H, Horie T. 2009. A process model for explaining genotypic and environmental variation in growth and yield of rice based on measured plant N accumulation. *Field Crops Research* **113**, 227–237.

Yoshida H, Horie T, Shiraiwa T. 2008. A model for explaining genotypic and environmental variation in vegetative biomass growth in rice based on observed LAI and leaf nitrogen content. *Field Crops Research* **108**, 222–230.

Yoshimoto M, Oue H, Kobayashi K. 2005. Energy balance and water use efficiency of rice canopies under free-air CO₂ enrichment. *Agricultural and Forest Meteorology* **133**, 226–246.

Zhang S, He X, Zhao J, et al. 2017. Identification and validation of a novel major QTL for harvest index in rice (*Oryza sativa* L.). *Rice (New York, N.Y.)* **10**, 44.

Zhu C, Xu X, Wang D, Zhu J, Liu G, Seneweera S. 2016. Elevated atmospheric [CO₂] stimulates sugar accumulation and cellulose degradation rates of rice straw. *GCB Bioenergy* **8**, 579–587.

Table 1: Photosynthesis characteristics (A : for Exp1 and B : for Exp2) measured two weeks after heading on the flag leaf on the main culm of IR64 plants grown under two [CO₂] levels, with panicle pruned at heading (PR) or not (Control). Average values $\pm$ standard errors (n=5) are presented. For each column within a [CO₂] level, values followed by different letters differ significantly (P<0.05)

A

[CO ₂] 400 $\mu\text{mol mol}^{-1}$	Sampling	Treatment	A ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	g_s ($\text{mmol m}^{-2} \text{s}^{-1}$)	C_i ($\mu\text{mol mol}^{-1}$)	V_{cmax} ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	J_{max} ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	TPU ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	g_m ($\mu\text{mol m}^{-2} \text{s}^{-1} \text{Pa}^{-1}$)	SPAD
	Control									
	Morning		23,03 $\pm$ 1,18 a	598,15 $\pm$ 21,21 a	321,82 $\pm$ 9,67 a	93,80 $\pm$ 8,66 a	142,80 $\pm$ 10,98 a	10,74 $\pm$ 0,60 a	12,65 $\pm$ 0,57 a	45,66 $\pm$ 0,76 a
	Midday		21,84 $\pm$ 0,96 a	572,50 $\pm$ 28,37 a	309,75 $\pm$ 2,35 ab	96,50 $\pm$ 3,13 a	136,00 $\pm$ 3,55 a	10,05 $\pm$ 0,26 ab	13,10 $\pm$ 0,62 a	45,01 $\pm$ 0,74 a
	Afternoon		21,48 $\pm$ 1,54 a	525,32 $\pm$ 59,35 a	316,57 $\pm$ 6,15 ab	89,60 $\pm$ 7,78 a	126,20 $\pm$ 6,75 a	9,70 $\pm$ 0,37 ab	11,93 $\pm$ 0,95 a	44,40 $\pm$ 0,54 a
	PR									
	Morning		22,68 $\pm$ 0,74 a	541,06 $\pm$ 45,72 a	311,16 $\pm$ 6,33 ab	90,20 $\pm$ 4,55 a	132,00 $\pm$ 6,24 a	9,32 $\pm$ 0,36 ab	12,48 $\pm$ 0,16 a	47,00 $\pm$ 1,87 a
	Midday		19,38 $\pm$ 1,89 a	393,47 $\pm$ 91,60 ab	286,36 $\pm$ 15,43 ab	88,40 $\pm$ 4,85 a	123,20 $\pm$ 1,98 a	8,76 $\pm$ 0,49 ab	12,58 $\pm$ 0,72 a	46,86 $\pm$ 1,17 a
	Afternoon		18,38 $\pm$ 1,00 a	281,24 $\pm$ 33,01 b	277,85 $\pm$ 8,39 b	90,60 $\pm$ 6,13 a	125,80 $\pm$ 7,23 a	8,38 $\pm$ 0,57 b	12,38 $\pm$ 0,56 a	47,28 $\pm$ 0,92 a
	Control									
	Morning		24,28 $\pm$ 0,47 a	709,38 $\pm$ 27,47 a	326,58 $\pm$ 4,73 a	95,50 $\pm$ 3,51 a	144,75 $\pm$ 5,92 a	10,85 $\pm$ 0,41 a	13,21 $\pm$ 0,73 a	45,25 $\pm$ 0,80 a
	Midday		22,48 $\pm$ 01,09 a	662,15 $\pm$ 33,46 ab	327,55 $\pm$ 3,01 a	88,80 $\pm$ 4,24 a	134,60 $\pm$ 8,35 a	10,10 $\pm$ 0,58 a	13,01 $\pm$ 0,64 a	46,14 $\pm$ 0,27 a
	Afternoon		21,76 $\pm$ 1,41 a	576,29 $\pm$ 84,55 ab	317,58 $\pm$ 13,10 ab	94,00 $\pm$ 5,54 a	134,40 $\pm$ 5,01 a	9,94 $\pm$ 0,27 a	12,20 $\pm$ 1,06 a	47,04 $\pm$ 1,06 a
	PR									
	Morning		21,36 $\pm$ 1,29 a	576,28 $\pm$ 85,35 ab	312,40 $\pm$ 8,89 ab	85,50 $\pm$ 3,97 ab	136,25 $\pm$ 2,86 a	9,33 $\pm$ 0,20 ab	12,44 $\pm$ 0,66 a	46,16 $\pm$ 0,65 a
	Midday		19,98 $\pm$ 1,22 a	431,54 $\pm$ 55,71 b	301,39 $\pm$ 6,68 ab	85,20 $\pm$ 1,59 ab	128,40 $\pm$ 4,83 a	9,16 $\pm$ 0,35 ab	12,21 $\pm$ 0,53 a	47,05 $\pm$ 0,75 a
	Afternoon		10,94 $\pm$ 0,92 b	168,36 $\pm$ 9,15 c	282,15 $\pm$ 7,11 b	64,80 $\pm$ 9,51 b	116,40 $\pm$ 10,06 a	7,88 $\pm$ 0,39 b	12,50 $\pm$ 0,44 a	44,58 $\pm$ 1,70 a

B

[CO ₂] 400 $\mu\text{mol mol}^{-1}$	Sampling	Treatment	A ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	g_s ($\text{mmol m}^{-2} \text{s}^{-1}$)	C_i ($\mu\text{mol mol}^{-1}$)	V_{cmax} ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	J_{max} ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	TPU ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	g_m ($\mu\text{mol m}^{-2} \text{s}^{-1} \text{Pa}^{-1}$)	SPAD
	Control									
	Midday		21,44 $\pm$ 0,94 a	465,18 $\pm$ 38,50 a	291,70 $\pm$ 6,82 a	97,40 $\pm$ 3,50 a	144,60 $\pm$ 2,97 a	10,08 $\pm$ 0,30 a	12,12 $\pm$ 0,78 a	44,98 $\pm$ 1,24 a
	Afternoon		20,73 $\pm$ 0,46 ab	427,13 $\pm$ 37,20 ab	280,29 $\pm$ 6,75 ab	91,60 $\pm$ 7,59 a	141,20 $\pm$ 4,98 a	9,96 $\pm$ 0,16 a	12,93 $\pm$ 0,51 a	44,30 $\pm$ 1,53 a
	Evening		20,49 $\pm$ 0,52 ab	375,91 $\pm$ 5,72 ab	278,72 $\pm$ 9,05 bc	95,2 $\pm$ 2,95 a	131,00 $\pm$ 5,21 a	9,54 $\pm$ 0,21 ab	11,97 $\pm$ 0,89 a	44,06 $\pm$ 0,67 a
	PR									
	Midday		20,70 $\pm$ 0,56 ab	395,71 $\pm$ 36,66 ab	277,95 $\pm$ 5,77 ab	100,40 $\pm$ 5,31 a	133,60 $\pm$ 4,50 a	9,14 $\pm$ 0,27 abc	11,87 $\pm$ 0,85 a	45,48 $\pm$ 0,85 a
	Afternoon		19,32 $\pm$ 0,52 ab	296,42 $\pm$ 32,02 bc	251,56 $\pm$ 5,49 bc	96,80 $\pm$ 1,98 a	130,80 $\pm$ 4,21 a	8,74 $\pm$ 0,24 bc	11,30 $\pm$ 0,65 a	44,30 $\pm$ 1,49 a
	Evening		18,06 $\pm$ 0,55 b	261,15 $\pm$ 11,78 c	228,23 $\pm$ 12,32 c	87,00 $\pm$ 4,91 a	127,20 $\pm$ 3,72 a	8,30 $\pm$ 0,17 c	11,15 $\pm$ 0,72 a	45,09 $\pm$ 0,98 a
	Control									
	Midday		22,89 $\pm$ 0,51 a	560,16 $\pm$ 32,49 a	302,28 $\pm$ 6,24 a	93,80 $\pm$ 3,56 a	138,60 $\pm$ 5,79 a	10,38 $\pm$ 0,21 a	13,80 $\pm$ 0,59 a	47,80 $\pm$ 0,68 a
	Afternoon		22,26 $\pm$ 0,60 a	420,84 $\pm$ 39,11 b	289,06 $\pm$ 10,12 ab	91,80 $\pm$ 8,34 a	135,80 $\pm$ 5,97 a	10,06 $\pm$ 0,29 a	10,02 $\pm$ 0,50 c	46,76 $\pm$ 1,28 a
	Evening		21,35 $\pm$ 0,64 a	360,9 $\pm$ 10,46 bc	287,98 $\pm$ 13,67 ab	91,00 $\pm$ 2,56 a	129,60 $\pm$ 7,65 ab	9,70 $\pm$ 0,29 a	12,15 $\pm$ 0,51 abc	46,02 $\pm$ 0,56 a
	PR									
	Midday		20,63 $\pm$ 0,45 a	409,25 $\pm$ 13,58 b	279,86 $\pm$ 5,06 ab	88,60 $\pm$ 4,30 ab	128,80 $\pm$ 3,30 ab	9,42 $\pm$ 0,33 a	12,98 $\pm$ 0,72 ab	47,94 $\pm$ 1,69 a
	Afternoon		13,21 $\pm$ 0,37 b	275,55 $\pm$ 6,62 c	258,18 $\pm$ 2,12 b	67,60 $\pm$ 3,66 b	120,20 $\pm$ 4,27 ab	7,78 $\pm$ 0,07 b	10,62 $\pm$ 0,75 bc	47,50 $\pm$ 0,98 a
	Evening		10,85 $\pm$ 0,65 b	147,47 $\pm$ 6,35 d	256,23 $\pm$ 4,05 b	65,50 $\pm$ 5,47 b	107,00 $\pm$ 8,22 b	6,38 $\pm$ 0,35 c	10,10 $\pm$ 0,49 c	47,08 $\pm$ 0,73 a

Table 2: Nonstructural carbohydrate contents, raw data (A : for Exp1 and B : for Exp2) measured two weeks after heading in the flag leaf, the upper and lower internode on the main culm of IR64 plants grown under two air CO₂ levels, with panicle pruned at heading (PR) or not (Control). Average values $\pm$ standard errors (n=5) are presented. For each column within a [CO₂] level, values followed by different letters differ significantly (P<0.05).

A

[CO ₂] 400 $\mu\text{mol mol}^{-1}$	Sampling	Treatment	Sucrose flag leaf ($\mu\text{g cm}^{-2}$)	Sucrose lower internode (mg g^{-1} DM)	Hexose flag leaf ($\mu\text{g cm}^{-2}$)	Hexose lower internode (mg g^{-1} DM)	Starch flag leaf ($\mu\text{g cm}^{-2}$)	Starch upper internode (mg g^{-1} DM)	Starch lower internode (mg g^{-1} DM)
	Control								
	Morning		191,17 $\pm$ 23,57 b	150,67 $\pm$ 37,61 a	16,02 $\pm$ 4,95 a	11,65 $\pm$ 4,40 a	83,07 $\pm$ 35,01 a	134,76 $\pm$ 44,89 b	245,23 $\pm$ 60,21 b
	Midday		234,21 $\pm$ 45,89 ab	205,92 $\pm$ 15,09 a	28,76 $\pm$ 6,61 a	5,27 $\pm$ 0,79 ab	125,66 $\pm$ 19,73 a	187,78 $\pm$ 23,86 ab	275,73 $\pm$ 67,09 ab
	Afternoon		270,82 $\pm$ 24,87 ab	190,22 $\pm$ 38,72 a	25,75 $\pm$ 3,81 a	5,36 $\pm$ 1,01 ab	82,85 $\pm$ 12,73 a	163,16 $\pm$ 14,42 ab	233,97 $\pm$ 70,05 b
	PR								
	Morning		231,81 $\pm$ 36,16 b	103,53 $\pm$ 23,33 a	20,48 $\pm$ 6,40 a	2,19 $\pm$ 0,50 b	84,39 $\pm$ 31,10 a	311,88 $\pm$ 30,32 a	519,13 $\pm$ 22,72 a
	Midday		286,91 $\pm$ 28,58 ab	116,68 $\pm$ 7,17 a	26,50 $\pm$ 5,43 a	2,27 $\pm$ 0,29 b	53,02 $\pm$ 11,54 a	311,50 $\pm$ 38,84 a	424,59 $\pm$ 16,60 ab
	Afternoon		395,74 $\pm$ 34,50 a	116,83 $\pm$ 35,88 a	26,32 $\pm$ 4,18 a	2,93 $\pm$ 0,53 b	127,95 $\pm$ 35,12 a	323,19 $\pm$ 46,70 a	531,41 $\pm$ 44,68 a
[CO ₂] 800 $\mu\text{mol mol}^{-1}$	Sampling	Treatment	Sucrose flag leaf ($\mu\text{g cm}^{-2}$)	Sucrose lower internode (mg g^{-1} DM)	Hexose flag leaf ($\mu\text{g cm}^{-2}$)	Hexose lower internode (mg g^{-1} DM)	Starch flag leaf ($\mu\text{g cm}^{-2}$)	Starch upper internode (mg g^{-1} DM)	Starch lower internode (mg g^{-1} DM)
	Control								
	Morning		211,07 $\pm$ 17,99 d	162,40 $\pm$ 26,30 a	20,14 $\pm$ 4,95 a	5,39 $\pm$ 1,25 a	118,68 $\pm$ 57,70 a	466,75 $\pm$ 26,74 a	402,95 $\pm$ 87,42 a
	Midday		275,96 $\pm$ 20,25 cd	134,89 $\pm$ 32,73 a	35,35 $\pm$ 9,28 a	5,87 $\pm$ 2,80 a	140,17 $\pm$ 27,54 a	402,87 $\pm$ 83,40 a	361,65 $\pm$ 83,60 a
	Afternoon		374,44 $\pm$ 17,42 ab	109,29 $\pm$ 31,38 a	28,02 $\pm$ 2,32 a	3,73 $\pm$ 0,96 a	299,80 $\pm$ 69,52 a	403,31 $\pm$ 38,06 a	466,65 $\pm$ 55,21 a
	PR								
	Morning		289,59 $\pm$ 35,19 bcd	77,14 $\pm$ 36,31 a	28,59 $\pm$ 2,77 a	1,86 $\pm$ 0,77 a	231,67 $\pm$ 78,24 a	476,33 $\pm$ 71,84 a	580,20 $\pm$ 59,77 a
	Midday		361,39 $\pm$ 13,51 bc	81,31 $\pm$ 25,90 a	37,80 $\pm$ 5,49 a	1,82 $\pm$ 0,45 a	309,85 $\pm$ 85,62 a	357,17 $\pm$ 37,35 a	530,03 $\pm$ 40,33 a
	Afternoon		466,86 $\pm$ 24,82 a	80,86 $\pm$ 9,98 a	38,27 $\pm$ 1,63 a	2,17 $\pm$ 0,28 a	249,24 $\pm$ 69,02 a	511,71 $\pm$ 56,08 a	641,20 $\pm$ 26,05 a

B

[CO ₂] 400 $\mu\text{mol mol}^{-1}$	Sampling	Treatment	Sucrose flag leaf ($\mu\text{g cm}^{-2}$)	Sucrose lower internode (mg g^{-1} DM)	Hexose flag leaf ($\mu\text{g cm}^{-2}$)	Hexose lower internode (mg g^{-1} DM)	Starch flag leaf ($\mu\text{g cm}^{-2}$)	Starch upper internode (mg g^{-1} DM)	Starch lower internode (mg g^{-1} DM)
	Control								
	Midday		310,98 $\pm$ 55,11 b	161,37 $\pm$ 10,79 a	22,34 $\pm$ 5,64 a	12,99 $\pm$ 3,85 ab	44,20 $\pm$ 8,28 a	170,36 $\pm$ 47,47 ab	156,27 $\pm$ 41,56 a
	Afternoon		355,45 $\pm$ 28,85 b	180,03 $\pm$ 14,49 a	18,05 $\pm$ 1,10 a	14,71 $\pm$ 2,35 a	70,08 $\pm$ 10,64 a	93,63 $\pm$ 15,00 b	178,80 $\pm$ 35,69 a
	Evening		430,43 $\pm$ 59,40 ab	132,44 $\pm$ 12,12 a	21,55 $\pm$ 3,91 a	13,32 $\pm$ 0,83 b	117,66 $\pm$ 31,14 a	157,34 $\pm$ 55,66 ab	147,55 $\pm$ 65,72 a
	PR								
	Midday		311,76 $\pm$ 33,78 b	139,35 $\pm$ 13,67 a	17,17 $\pm$ 1,13 a	8,97 $\pm$ 2,03 ab	48,51 $\pm$ 8,99 a	170,59 $\pm$ 32,28 ab	362,25 $\pm$ 51,80 a
	Afternoon		490,53 $\pm$ 23,50 ab	141,84 $\pm$ 9,84 a	26,10 $\pm$ 1,42 a	8,00 $\pm$ 2,08 ab	94,55 $\pm$ 8,97 a	202,02 $\pm$ 19,43 ab	288,21 $\pm$ 68,60 a
	Evening		496,92 $\pm$ 37,84 a	134,44 $\pm$ 20,75 a	28,81 $\pm$ 3,65 a	5,72 $\pm$ 1,57 ab	128,38 $\pm$ 33,51 a	299,98 $\pm$ 53,68 a	294,26 $\pm$ 63,43 a
[CO ₂] 800 $\mu\text{mol mol}^{-1}$	Sampling	Treatment	Sucrose flag leaf ($\mu\text{g cm}^{-2}$)	Sucrose lower internode (mg g^{-1} DM)	Hexose flag leaf ($\mu\text{g cm}^{-2}$)	Hexose lower internode (mg g^{-1} DM)	Starch flag leaf ($\mu\text{g cm}^{-2}$)	Starch upper internode (mg g^{-1} DM)	Starch lower internode (mg g^{-1} DM)
	Control								
	Midday		267,33 $\pm$ 27,60 b	105,41 $\pm$ 3,38 a	20,17 $\pm$ 2,72 a	8,53 $\pm$ 4,35 a	53,16 $\pm$ 15,27 a	110,36 $\pm$ 16,20 b	184,22 $\pm$ 55,74 a
	Afternoon		352,10 $\pm$ 40,06 ab	93,10 $\pm$ 3,78 a	26,24 $\pm$ 4,07 a	9,40 $\pm$ 2,24 a	104,30 $\pm$ 17,14 a	54,03 $\pm$ 7,15 b	168,88 $\pm$ 60,77 a
	Evening		393,25 $\pm$ 18,79 a	106,22 $\pm$ 8,73 a	20 $\pm$ 2,31 a	3,72 $\pm$ 1,07 a	149,94 $\pm$ 23,65 a	80,99 $\pm$ 16,68 b	164,39 $\pm$ 52,57 a
	PR								
	Midday		382,65 $\pm$ 16,76 ab	97,01 $\pm$ 17,13 a	24,47 $\pm$ 3,23 a	4,23 $\pm$ 0,83 a	50,82 $\pm$ 11,22 a	282,52 $\pm$ 32,88 a	352,03 $\pm$ 49,52 a
	Afternoon		423,82 $\pm$ 33,98 a	85,91 $\pm$ 9,68 a	24,15 $\pm$ 4,96 a	2,35 $\pm$ 0,16 a	104,30 $\pm$ 32,82 a	240,88 $\pm$ 31,66 a	338,46 $\pm$ 37,24 a
	Evening		452,95 $\pm$ 12,27 a	96,18 $\pm$ 5,58 a	23,70 $\pm$ 3,51 a	3,84 $\pm$ 0,62 a	131,61 $\pm$ 39,77 a	267,47 $\pm$ 30,55 a	323,30 $\pm$ 37,70 a

Figure legends

Fig. 1: Effect of pruning and CO₂ treatment on photosynthetic parameters, (A) net assimilation rate A , (B) triose phosphate utilization TPU, (C) maximum carboxylation rate of Rubisco V_{cmax} , and (D) maximum rate of electron transport J_{max} . Measured at 2 weeks after heading on the flag leaf on the main culm of plants of IR64 rice genotype at two CO₂ levels (400 and 800 $\mu\text{mol mol}^{-1}$), in two growth chambers for experiments 1 and 2. Black symbol: Control (plants with panicles) and red symbol: PR (plants with panicle pruned). Measurements were carried out at morning (mor), midday (mid), afternoon (aft) and evening (eve) periods. Stars indicate significant difference at $p < 0.05$ (Tukey HSD test) among values. NS: not significant. Each point represents the mean of 5 values $\pm$ SE.

Fig. 2: Mean A/C_i curves for all treatment combination (A) and corresponding mean $\phi\text{PSII}/C_i$ curves (B) for experiments 1 and 2 (shown only for Pruning treatment which caused significant TPU decline). Dashed lines indicate mean C_i value (290 $\mu\text{mol mol}^{-1}$) for photosynthesis measurement at treatment CO₂ level (400 or 800 $\mu\text{mol mol}^{-1}$). Arrows in Fig. 2B indicates the C_i level at which TPU limitation may begin to occur. The CO₂ concentrations indicated in Figure headers are treatment conditions and not those administered when measuring CO₂ response, which comprised 14 different levels.

Fig. 3: Relationship between TPU and net photosynthesis (A) within experiments 1 and 2 and for each combination of CO₂ x panicle pruning treatment. With panicle and 400 $\mu\text{mol mol}^{-1}$ (black symbol), with panicle and 800 $\mu\text{mol mol}^{-1}$ (grey symbol), Panicles pruned and 400 $\mu\text{mol mol}^{-1}$ (red symbol), panicles pruned and 800 $\mu\text{mol mol}^{-1}$ (yellow symbol). Each point represents a single value.

Fig. 4: Relationship between TPU and Leaf sucrose content in experiments 1 and 2, separating control and panicle pruning treatment. With panicle at 400 $\mu\text{mol mol}^{-1}$ (black symbol), with panicle at 800 $\mu\text{mol mol}^{-1}$ (grey symbol), Panicles pruned at 400 $\mu\text{mol mol}^{-1}$ (red symbol), panicles pruned at 800 $\mu\text{mol mol}^{-1}$ (yellow symbol). Dashed lines represent linear regression for both pruning treatments. Each point is the average of 5 values and is presented with horizontal and vertical standard errors.

Fig. 5: Relationship between TPU and local source-sink ratio (defined as flag leaf area / spikelet number on the main culm) in experiments 2. Line represents linear regression.

Fig. 1:

Fig. 2:

Fig. 3:

Fig. 4:

Fig. 5: