Survey of intertidal ecosystem reveals a legacy of potentially toxic elements from industrial activity in the Skeena Estuary, British Columbia, Canada

Tom Sizmur^{1*}, Lily Campbell², Karina Dracott³, Megan Jones¹, Nelson J. O'Driscoll⁴ and Travis Gerwing^{2,5}

¹ Department of Geography and Environmental Science, University of Reading, Reading, UK

² Department of Biology, University of Victoria, Victoria, British Columbia, Canada.

³ North Coast Cetacean Research Initiative, Ocean Wise Conservation Association, Prince Rupert, British Columbia, Canada.

⁴ Department of Earth & Environmental Sciences, Acadia University, Wolfville, Nova Scotia, Canada.

⁵ Ecosystem Science and Management Program, University of Northern British Columbia, Prince George, British Columbia, Canada.

*Corresponding author

Email: t.sizmur@reading.ac.uk

Abstract

Relationships between concentrations of Potentially Toxic Elements (PTEs) in estuarine sediments and their impact benthic invertebrate communities are poorly understood. We sampled and analysed PTEs in sediments and benthic invertebrates from five sites surrounding the Skeena Estuary, including sites adjacent to an abandoned cannery and a decommissioned papermill. There was no indication that sediments of the salmon cannery are polluted, but acidic sediments adjacent to the papermill contained elevated concentrations of Cd, Cr, Hg and Pb. Benthic invertebrate community assemblages confirm that sediments have recovered from prior disturbances associated with discharge of papermill sludge. Oregon pill bugs (*Gnorimosphaeroma oregonensis*), observed at all five sites, feed on the fibers associated with the papermill discharge. Thus, *G. oregonensis* are useful biomonitors for quantifying the impact of the decommissioned papermill, and similar industrial development projects, on intertidal ecosystems along the north coast of British Columbia, Canada.

31 Introduction

32 Conflict between the economic benefits of industrial development and the potential impact of such
33 developments on the natural environment is most clearly expressed in remote communities, entirely
34 dependent on natural capital for survival [1-3]. Anthropogenic activities have impacted the
35 biogeochemical cycling of Potentially Toxic Elements (PTEs) in even the most remote ecosystems on
36 our planet [4-6]. Quantifying the disruption of natural biogeochemical cycles of these elements by
37 anthropogenic activities is difficult if the concentrations prior to industrial development are
38 unknown and may be redundant if investigations are only conducted after environmental damage
39 has occurred. Therefore, it is important to conduct baseline surveys to assess the bioavailability and
40 bioaccumulation of PTEs by organisms in remote locations to predict the likely impact of further
41 pollution on the ecosystem.

42

43 Intertidal estuarine sediments support ecologically and economically important ecosystems that
44 provide nursery habitats to several species (some that support important commercial fisheries), and
45 often host a diverse and abundant fauna [7-11]. Benthic invertebrates are often used as indicators of
46 environmental pollution since they live in sediments and are prey for many commercially important
47 fish species [12-14]. Benthic invertebrates inhabiting estuaries are inherently resistant to physical
48 and chemical change as they have adapted to living in a dynamic environment with wide spatial and
49 temporal ranges of chemical and physical properties, such as pH, redox potential, salinity, and
50 particle size [15-17]. Because the chemical and physical properties of estuarine sediment are
51 temporally and spatially dynamic, it is difficult to predict the impact of industrial development on
52 the fate and impact of PTEs [18]. We do not currently understand the relationship between
53 sediment properties, food web structure and PTE bioaccumulation by sediment-dwelling
54 invertebrates well enough to predict the impact of future emissions to the estuarine environment.
55 The objective of this study was to explore the relationship between sediment properties and the fate
56 of PTEs from historic industrial development in the Skeena Estuary and quantify the impact of PTEs
57 on the diversity of benthic invertebrates inhabiting intertidal sediments.

58

59 The Skeena is British Columbia's second largest river and provides important habitat for Pacific
60 salmon populations, upon which Canadian First Nations communities rely [2, 8, 19]. The estuary and
61 the intertidal areas towards the north of the Estuary along the coast of British Columbia are
62 important nursery habitats for juvenile salmon [8, 19]. Coastal areas to the north of the estuary

63 surrounding the small port cities of Prince Rupert and Port Edward have been extensively developed
64 [3-5, 20]. Industrial developments include an international port, a papermill, and several canneries.

65

66 Findings of previous surveys of the benthic invertebrates inhabiting the intertidal sediment in the
67 Skeena estuary reveal an infaunal community that is relatively undisturbed at the estuary scale, but
68 which still shows the scars of historic disturbance at finer-grained scales [10, 15, 21, 22]. Amphipods
69 are powerful indicator species [23-27] whose high densities throughout the Skeena estuary [10, 15]
70 suggests that current disturbances to intertidal areas are relatively limited. Similarly, we observe 40
71 intertidal species in this area, including multiple species at all trophic levels within the foodweb [10,
72 15], and such a complex community is often associated with non-disturbed habitats [4, 28-30].
73 Conversely, disturbed sites are often more easily invaded by invasive species [31, 32], and
74 Capitellidae (*Capitella capitata* species complex) polychaetas are often observed in disturbed
75 habitats, particularly areas that have been organically enriched [4, 27, 28, 33]. Capitellidae
76 polychaetas have a clumped distribution within the Skeena estuary and can be locally abundant at
77 the scale of a 1m² plot [10, 15]. Similarly, abundant but localized populations of the invasive
78 Cumacea *Nippoleucon hinumensis*, have been observed in the Skeena estuary [10, 15], and on other
79 historically impacted mudflats along BC's north coast [27]. While the universal presence and high
80 abundances of amphipods, coupled with the complexity of the intertidal community at the estuary
81 scale strongly suggest that these intertidal communities are relatively undisturbed, the locally
82 abundant populations of invasive species and Capitellidae polychaetas offer contradictory evidence.
83 These indicators of disturbance suggest that either current small-scale disturbances are occurring, or
84 this biological signal is a remnant of past disturbances.

85

86 Persistent localised biological signals of historic disturbances in mudflats that are currently relatively
87 undisturbed have been observed before on the north coast of BC [27]. To the best of our knowledge,
88 no current disturbances are occurring in the Skeena estuary that can explain the small-scale
89 indicators of disturbance observed in the biological community [10, 15, 22, 27]. Therefore, we
90 hypothesise that fine-grained biological indicators of disturbance in the Skeena estuary are a legacy
91 of past disturbances [10, 15]. It is likely that these past disturbances are at least partially related to
92 discharge from the papermill, which was released into the immediate near shore area (Porpoise
93 Bay), strongly depressing the invertebrate communities in this area during the 1970s [20, 34, 35]. As

94 such, we postulate that these intertidal areas have been passively recovering from this disturbance
95 for ~50 years.

96

97 Considerable developments have been proposed in the Skeena estuary, including oil and gas
98 pipelines, super-tanker routes, potash loading facilities, and a liquid natural gas (LNG) terminal. The
99 Skeena salmon run contributes an estimated \$110 million dollars annually to the local economy [36],
100 so pollution of the intertidal nursery habitat of the Skeena Estuary could have devastating
101 consequences on both the economy and ecosystem [37, 38]. It is therefore critical to predict the
102 impact that future developments will have on the bioaccumulation of PTEs by benthic invertebrates
103 inhabiting intertidal sediments of the Skeena Estuary. This work presents a survey of intertidal
104 sediments and invertebrates at five locations north of the Skeena estuary to examine baseline levels
105 of PTE concentrations at sites that may be recovering from previous disturbances and identify
106 organisms that could be used to biomonitor the impact of future industrial developments.

107

108 **Methods**

109 **Field sampling**

110 Sediment cores and benthic invertebrates were collected from five locations north of the Skeena
111 Estuary (Fig 1) during July 2017: Cassiar Cannery (CC), Inverness Passage (IP), Tyee Banks (TB),
112 Papermill Bay (PB), and Wolfe Cove (WC). Cassiar Cannery (N54° 10' 40.4, W130° 10' 40.4) is a
113 former salmon cannery that closed in 1983 and is now an ecotourism lodge. Inverness passage (N54°
114 10' 05.9, W130° 09' 40.4), a mudflat ~3 km closer to the Skeena mouth than Cassiar Cannery, has
115 intertidal habitat similar to Cassiar Cannery, but was never directly impacted by a cannery or other
116 anthropogenic development. Tyee Banks (N54° 11' 59.1, W129° 57' 36.7) is located 20km upstream
117 from the mouth of the Skeena River on a large intertidal mudflat. At some point in the past, this area
118 had a small-scale sawmill operating and accumulations of sawdust and woodchips are still present in
119 the upper intertidal sediment. Papermill Bay (N54° 13' 59.3, W130° 17' 07.5) is a small bay located
120 directly adjacent to a large decommissioned papermill. Finally, Wolfe Cove (N54° 14' 33.0, W130° 17'
121 34.5) is a mudflat located approximately 1 km from the papermill. The papermill was closed in 2001,
122 ceasing all operations and discharge [39-41].

123

Fig 1. Map showing sites adjacent to the Skeena Estuary, British Columbia, Canada where intertidal mudflats at Papermill Bay, Wolfe Cove, Cassiar Cannery, Inverness Passage, and Tye Banks were sampled during this project.

Sediments were sampled by pushing polycarbonate cores (5 cm diameter, 20 cm length) into the mud with a rubber mallet, digging out with a spade, and wrapping with cling film, before extruding and dividing into strata; 0-5 cm, 5-10 cm, 10-15 cm and 15-20cm. Sediment samples from each strata were dried at 40 °C for 48 hours and shipped to the UK for analysis. A total of 15 cores were collected from each site along five transects comprising three samples taken from the upper, mid and lower shore in a random stratified sampling design [42]. Sediments from the upper, mid and lower shore (but the same strata) were pooled and homogenized prior to analysis. At each location that a sediment core was collected, two more cores (7 cm diameter) were taken to a depth of at least 5 cm (well into the anoxic layer), transferred to a plastic bag in the field and then passed through a 250µm stainless steel sieve to retain sediment-dwelling benthic infauna. This infauna sampling strategy was supplemented by opportunistic digging in areas of the site in which infauna were clearly present (e.g. burrow openings or surface casting) to collect macrofauna. All invertebrates found in sufficient quantities for chemical analysis were identified to the species level [10, 15], pooled into a single sample for each site, rinsed in deionised water, frozen, shipped to the UK, and freeze-dried prior to analysis.

Laboratory analysis

The particle size distribution of sediments was determined using laser granulometry (Malvern Mastersizer 3000). Sub-samples were then ground to a fine powder using an agate ball mill and analysed for total organic carbon and nitrogen content using a Thermo Scientific Flash 2000 Organic Elemental Analyser. Details of quality control can be found in the supporting information file. Sediment pH was determined in a soil-water suspension after shaking with water for 15 min at a 1:10 w/v ratio based on BS7755-3.2 [43]. The total concentration of PTEs in sediments was determined by ICP-MS (Inductively Coupled Plasma Mass Spectrometry) analysis of 0.5 g sediment samples digested in reverse aqua regia (9 ml of nitric acid and 3 ml of hydrochloric acid) using a MARS 6 microwave digestion system, based on EPA [44]. After preliminary analysis of a large range of PTE concentrations, the elements selected for further investigation were Cadmium (Cd), Cobalt (Co), Chromium (Cr), Nickel (Ni), Lead (Pb) and Zinc (Zn). Details of quality control can be found in

the supporting information file. The total concentration of mercury in sediments was determined using thermal degradation – gold amalgamation atomic absorbance spectroscopy as outlined in EPA [45] using a Nippon MA-3000 analyser. Details of quality control can be found in the supporting information file. The bioavailability of PTEs to biota was estimated by extracting 2.5 g of sediments with 25 ml of 0.05M EDTA (Ethylenediaminetetraacetic acid) at 20°C for one hour, centrifuging, filtering and analysing PTEs (As, Cd, Co, Cr, Cu, Ni, Pb and Zn) in the extract using ICP-OES (Inductively Coupled Plasma Optical Emission Spectrometry). The total concentrations of PTEs in invertebrates were determined by ICP-MS analysis of 0.5 g of sample digested in slightly diluted nitric acid (2 ml of ultra-pure water and 8 ml of nitric acid) using a MARS 6 microwave digestion system.

Statistical analysis

The influence of site and sediment depth on sediment PTE concentrations and properties was quantified using analysis of variance (ANOVA) and permutational multivariate analysis of covariance (PERMANCOVA) [46, 47]. The relationship between PTE concentrations and sediment properties was first tested using PRIMER's RELATE function [48]. This function compares two resemblance matrices looking for any relationships. Relationships were further explored using principal component analyses (PCA) on the variance-covariance matrix of all sediment PTE and sediment property data. Relationships between the PTE concentrations observed in the sediments and in the collected invertebrates were examined in several configurations using RELATE and plotted using non-metric multidimensional (nMDS) scaling plots. A full description of the statistical analysis undertaken is provided in the supporting information file and outputs provided in S1, S2 and S3 Tables.

Results and Discussion

Inverness Passage, Wolfe Cove and Tyee Banks are suitable

reference sites

The PERMANCOVA analysis indicates that properties of sediments (pH, median particle diameter, C and N) are significantly influenced by both site and sediment depth, with site explaining more than 50% of the observed variance in the data (S1 Table). There is no significant ($p > 0.05$) difference in

the median sediment particle diameter between Inverness Passage and Cassiar Cannery, or between Wolfe Cove and Papermill Bay (Fig 2 and S3 Table). This observation supports our assumption that the sediment deposited in the sites potentially contaminated by industrial emissions (Cassiar Cannery and Papermill Bay) and their respective proximal reference sites (Inverness Passage and Wolfe Cove) have the same geogenic origin and are subject to a similar depositional environment. Thus, any differences in sediment chemical properties between potentially contaminated sites and their proximal reference sites we infer is due to anthropogenic influences.

Fig 2. Sediment properties; Sediment pH (A); Median sediment particle diameter (B); Percentage Organic Carbon content (C); and Percentage Total Nitrogen content (D) presented at four depths in sediments sampled from Cassiar Cannery, Inverness Passage, Papermill Bay, Wolfe Cove and Tye Banks intertidal mudflats.

Tye Banks has a significantly ($p < 0.05$) larger median particle size than Inverness Passage and Cassiar Cannery and a significantly ($p < 0.05$) smaller median particle size than Wolfe Cove and Papermill Bay (Fig 2 and S3 Table). The location of Tye Banks is closer to the mouth of the Skeena River and, therefore, only receives sediment from the Skeena, whereas the other four sites may also receive sediment from the Nass River, to the north of the Skeena Estuary [49]. However, because Tye Banks is much further from potential emission sources, the site can be used to determine whether proximal reference (Inverness Passage and Wolfe Cove) sites are contaminated with PTEs from the adjacent industrial activity.

Papermill Bay sediments are contaminated with Cd, Cr, Hg and Pb

The PERMANCOVA analysis (S2 Table) indicates significant differences in sediment total and EDTA extractable PTE concentrations between different sites, but not between different depths. We observed significantly ($p < 0.05$) greater concentrations of As, Cr, Hg and Pb in the Papermill Bay sediments than the proximal reference sediments at Wolfe Cove (Fig 3 and S3 Table). However, Co, Cu and Ni concentrations were significantly ($p < 0.05$) lower in the Papermill Bay sediments (Fig 3 and S3 Table). A Geoaccumulation index, calculated following [50], using Wolfe Cove as an uncontaminated reference site, indicates that Papermill Bay is 'unpolluted' with As, Cd, Co, Cr, Cu, Ni and Zn, but 'unpolluted to moderately polluted' with Pb and 'moderately polluted' with Hg (Table 1). When Tye Banks is used as the reference site, Papermill Bay is classified as 'unpolluted to

moderately polluted' with Cd, Cr and Hg. Thus, there is evidence to suggest that the sediments in Papermill Bay are contaminated with Cd, Cr, Hg and Pb, most likely emanating from the sludge [51] discharged by the decommissioned papermill on Watson Island (Fig 1).

Fig 3. Total concentrations of Potentially Toxic Elements (As, Cd, Co, Cr, Cu, Hg, Ni, Pb and Zn) with depth in sediments sampled from Cassiar Cannery, Inverness Passage, Papermill Bay, Wolfe Cove and Tyee Banks intertidal mudflats

Table 1. Geoaccumulation index (I_{geo})^a for potentially toxic elements in sediments from Cassiar Cannery (CC) and Papermill Bay (PB) using reference sites Tyee Banks (TB), Wolfe Cove (WC) and Inverness Passage (IP).

Site	Reference site	As	Cd	Co	Cr	Cu	Hg	Ni	Pb	Zn
CC	IP	-0.99	-0.71	-0.78	-0.57	-0.69	-0.69	-0.67	-0.58	-0.63
	TB	-0.48	-0.33	-0.44	-0.27	-0.01	-0.13	-0.26	0.08	-0.24
PB	WC	-0.10	-0.25	-1.08	-0.32	-1.14	1.61	-0.82	0.74	-0.60
	TB	-1.51	0.39	-0.90	0.35	-0.19	0.20	-1.01	-0.02	-0.45

^a Geoaccumulation index is calculated as $I_{geo} = \log_2 \frac{C_n}{1.5 \times B_n}$

when C_n = Concentration measured in sediment at site and B_n = Concentration measured in sediment at reference (background) site.

Sites are classified as unpolluted if $I_{geo} \leq 0$, unpolluted to moderately polluted if $0 < I_{geo} \leq 1$, moderately polluted if $1 < I_{geo} \leq 2$, moderate to strongly polluted if $2 < I_{geo} \leq 3$, strongly polluted if $3 < I_{geo} \leq 4$, strongly to extremely polluted if $4 < I_{geo} \leq 5$, and extremely polluted if $I_{geo} > 5$.

Discharges by papermills have previously been associated with pollution of the environment with PTEs, including Cr, Cd, Cu, Pb and Zn [52, 53]. This pollution associated with papermill operations can originate from a number of sources, such as metals in the wood entering the papermill, or atmospheric deposition of metals from the smoke stacks [52]. PTE-containing materials may also be used during the pulp making processes. For example, chromate bricks are used in the recovery furnace of kraft paper mills to reduce chemical attack by the spent liquors [54]. However, Cr concentrations observed in the sediments of Papermill Bay (Fig 3) are considerably lower than the 52.4 mg kg⁻¹ observed in a deposit of pulp covering the sediments of a Swiss lake [55], or

concentrations of up to 197 mg kg⁻¹ observed in a fibrous deposit in the Ångermanälven river estuary, on the east coast of Sweden (Apler et al 2019).

The carbon and nitrogen content of the sediments at Papermill Bay were significantly ($p < 0.05$) and considerably greater than all of the other sites (Fig 2 and S3 Table). It was evident from visual inspection of the sediments themselves that they contained a high proportion of organic fibers, presumably discharged into the bay from the papermill during the operational phase. The PTEs observed to be elevated in the Papermill Bay sediments (Cd, Cr, Hg and Pb) all bind strongly to organic matter in sediments [56-58] and so their presence could be due to (i) the discharge and deposition of PTE contaminated organic material, (ii) the discharge of organic material alongside the discharge of PTEs in contaminated effluent that later bind with the organic material, or (iii) the discharge of organic material which binds to naturally present PTEs in the sediment and prevents their export to the overlying water.

In contrast to the observations made in the Papermill Bay sediments, none of the PTEs analysed in the Cassiar Cannery sediments were found to be present in significantly ($p > 0.05$) greater concentrations than the Inverness Passage sediments (Fig 3 and S3 Table). In fact, Co and Ni concentrations were significantly lower in Cassiar Cannery sediments than at Inverness Passage. When Inverness Passage is used as a reference site, all the PTEs analysed can be classified by the Geoaccumulation index as 'unpolluted' (Table 1). This observation is a clear indicator that any impact resulting from either the former use of the Cassiar Cannery site (up until 1983) as a salmon cannery, or its current use as an ecotourism lodge, can no longer be observed in the concentrations of PTEs in the top 20 cm of sediment at this site. Any sediment contaminated by cannery waste at this site may have been buried by fresh sediment, rendering the PTEs inaccessible to benthic invertebrates.

Relationship between sediment properties and total and labile PTE concentrations

Alongside greater total concentrations of Cd, Cr, Hg and Pb in the Papermill Bay sediments, compared to Wolfe Cove, we observed significantly ($p < 0.05$) greater EDTA extractable concentrations of Cd, Co, Cr, Ni, Pb and Zn (Fig 4 and S3 Table), indicating a greater availability of

these elements to benthic invertebrates [59]. The significantly ($p < 0.05$) greater carbon content and the slightly finer texture of the Papermill Bay sediments, compared to Wolfe Cove (Fig 2 and S3 Table) should provide more sites for PTEs to bind to [58, 60, 61]. Thus, the greater concentrations of EDTA extractable PTEs is largely due to the Papermill Bay sediments being significantly ($p < 0.05$) more acidic than the Wolfe Cove sediments. Lower pH results in greater competition between PTEs and hydrogen ions for the binding sites on the sediment surface and leads to greater PTE lability [62, 63].

Fig 4. EDTA extractable (potentially bioavailable) Potentially Toxic Elements (Cd, Co, Cr, Cu, Ni, Pb and Zn) with depth in sediments sampled from Cassiar Cannery, Inverness Passage, Papermill Bay, Wolf Cove, and Tyee Banks intertidal mudflats

The RELATE revealed a statistically significant correlation between the concentration of potentially toxic elements and the sediment physicochemical properties resemblance matrices ($Rho: 0.46$; $p = 0.001$, Permutations = 9999). The relationship between total and labile PTE concentrations and sediment properties was further explored using PCA (Fig 5). The principal component scores plot (Fig 5B) reveals a clear separation between the Papermill Bay sediments and the proximal and remote references sites of Wolfe Cove and Tyee Banks, but no separation between Cassiar Cannery and Inverness Passage, re-enforcing the conclusion that the sediments of Cassiar Cannery show little evidence of anthropogenic contamination. Principal component 2 (Fig 5), separates the Inverness Passage and Cassiar Cannery sediments from the other three sites (Papermill Bay, Wolfe Cove, and Tyee Banks), which is attributed to a different geochemical matrix composition at Cassiar Cannery and Inverness Passage, as also observed by DelValls et al. [64]. Principal component 1 (Fig 5) reveals a separation with depth in the Papermill Bay sediments. Papermill Bay sediments have greater total and EDTA extractable concentrations of several PTEs, higher C and N, and lower pH, all of which also increase with depth in the Papermill Bay sediments. This observation of deeper layers of sediment with a lower pH and higher availability of PTEs indicates that contaminated sediment is overlain by less contaminated sediment, deposited since discharge of sludge ceased. Although most benthic invertebrates live in surficial sediments, it is important to consider the likelihood that deeper sediment will be exposed and mobilized either by development projects [65], or by bioturbation from deep burrowing invertebrates [66, 67].

Fig 5. Principal Components Analysis of sediment properties (pH median sediment particle diameter, Percentage Organic Carbon content and Percentage Total Nitrogen content) and total and EDTA extractable Potentially Toxic Elements (As, Cd, Co, Cr, Cu, Hg, Ni, Pb and Zn) in sediments sampled 0-5 cm, 5-10 cm, 10-15 cm and 15-20cm from Cassiar Cannery, Inverness Passage, Papermill Bay, Wolf Cove and Tyee Banks intertidal mudflats. (A) is the latent vectors for each variable plotted in the plane of principal component one and principal component two, (B) is the principal component scores of all the samples plotted in the plane of principal component one and principal component two. Error bars represent the standard deviations of 5 replicate samples taken from 5 different transects.

Sediments host a healthy invertebrate population with no relationship between PTE concentrations in sediments and invertebrates

Overall, 22 different taxa (listed in S4 Table, along with PTE concentrations) were observed in this study, with Wolf Cove and Cassiar Cannery the most biodiverse sites, with 16 and 14 species collected, respectively. These taxa represent a subset of the 40 intertidal species that are commonly observed in the Skeena Estuary [10, 15]. Findings of previous studies in the Skeena Estuary reveal an infaunal community predominantly dominated by cumaceans, polychaetes, oligochaetes, nematodes, copepods, amphipods, and bivalves [15, 68]. The 40 intertidal species observed in this area include multiple species at all trophic levels within the food web [10, 15], and such a complex community is often associated with non-disturbed habitats [4, 28]. This complexity was also observed in our study, as *Alitta brandti*, *Neries vexillosa*, *Glycinde picta*, ribbon worms and crabs can act as predators, while *Macoma balthica*, isopods, amphipods, and sessile polychaete worms are primary consumers [47, 69, 70]. We therefore provide evidence to support our previous research that indicates that the intertidal ecosystem has been passively recovering for ~50 years from past disturbances related to discharge from the papermill, which was released into the immediate near shore area (Porpoise Bay), strongly depressing the invertebrate communities in this area during the 1970s [20, 34, 35] and currently exhibit a community that is relatively healthy.

When all invertebrates collected at all five sites are considered, we found no relationship between PTE concentrations in invertebrates and either total (Rho: 0.27; $p = 0.32$, Permutations = 9999), EDTA extractable (Rho: 0.24; $p = 0.79$, Permutations = 9999), or both total and EDTA extractable (Rho: 0.78; $p = 0.08$, Permutations = 9999) PTEs in the sediment. This finding is confirmed by

inspecting a nMDS plot of invertebrate PTE loadings, which includes all organisms collected at all five sites (Fig 6A). Bivalves of the same species (e.g. *M. balthica*, *Mytilus edulis*, and *Mya arenaria*), collected at different sites, seem to cluster in multidimensional space. There is slightly more separation in the PTE concentrations measured in polychaete worms (e.g. *Abarenicola pacifica*, *N. vexillosa*, *Paranemertes peregrina*, *G. picta*, *A. brandti*, *Nephtys caeca*, *Neotrypaea californiensis*, and *Streblospio benedicti*) which are the cause of much of the dissimilarity in the dataset.

Fig 6. Non-metric multidimensional scaling plots (nMDS) of invertebrate PTE concentrations and the vector overlay (left hand side) of five intertidal mudflats along the north coast of British Columbia, Canada. CC: Cassiar Cannery. WC: Wolfe Cove. IP: Inverness Passage. PB: Papermill Bay. TB: Tye Banks with (A) the entire dataset considered or (B) only including the two benthic invertebrates found at all five sites; Baltic clams (*Macoma balthica*) and Oregon pill bugs (*Gnorimosphaeroma oregonensis*).

The suitability of *G. oregonensis* as biomonitors of sediment PTE bioavailability

The selection of one or more of benthic invertebrate as a biomonitor of PTE pollution needs to consider how cosmopolitan their distribution is and the pathway by which they are exposed to the PTE [71]. Mussels and clams acquire food from the water column by suspension feeding, and clams are also able to acquire food from surface sediments by way of their siphon. Isopods are more motile and acquire food by ingesting organic debris, usually on the surface of the sediment. Polychaete worms, as a class, have a more diverse range of feeding strategies, including suspension feeding with tentacles or a mucus net, deposit feeding selectively at the surface or at depth, and predation of other infauna [72, 73]. This large diversity of feeding strategies may have led to the greater range of PTE concentrations observed in Fig 6A, compared to the bivalves and crustacea. Furthermore, the soft tissues of polychaete worms may only represent exposure to PTEs in the recent past due to large temporal fluctuations [74-76]. The whole body (including soft tissues and shells) of bivalves may better represent a record of time-averaged bioavailability of PTEs in the water column over the lifetime of the organism, albeit not without difficulties in interpretation [77, 78].

There were only two benthic invertebrate species observed at all five sites; Baltic clams (*M. balthica*) and Oregon pill bugs (*G. oregonensis*). When PTE concentrations in *M. balthica*, and *G. oregonensis*

was compared to sediment concentrations, no relationship was observed with either total (Rho: 0.31; $p = 0.19$, Permutations = 9999), EDTA extractable (Rho: -0.0.; $p = 0.48$, Permutations = 9999), or both total and EDTA extractable (Rho: 0.21; $p = 0.28$, Permutations = 9999) sediment PTE concentrations. These findings contrast to numerous articles in the literature quantifying relationships between sediment and benthic invertebrate PTE concentrations [58, 71, 79, 80]. When we plot the PTE concentrations in *M. balthica*, and *G. oregonensis* in multidimensional space (Fig 6B), we reveal that the elemental profiles of *M. balthica*, and *G. oregonensis* contrast greatly. *M. balthica* collected at Papermill Bay contained lower concentrations of Cr, Co, Ni, Zn, Cd, As and Pb than those collected at the other four sites, but higher concentrations of Cu. In contrast, *G. oregonensis* collected at Papermill Bay have higher concentrations of Cr, Ni, Zn, and Pb than those collected at the other four sites.

Biota-Sediment Accumulation Factors (BSAFs) were calculated by dividing the concentrations of PTEs in the invertebrate tissues by the average concentration in the sediments from the site from which the invertebrates were collected (average across all transects and depths). Relationships between sediment properties and BSAFs reveal the importance of pH in explaining the difference in the bioaccumulation of PTEs by *M. balthica*, and *G. oregonensis*. For most PTEs there is a positive relationship between pH and the BSAF for *M. balthica* (Table 2), including a significant relationship with the Cr BSAF (Fig 7A). This is an unexpected finding that does not have an immediately obvious explanation. However, we observe a negative relationship between sediment pH and the BAF of most PTEs for *G. oregonensis* (Table 2), including Cr (Fig 7B). This relationship is intuitive since metal cations dissociate from mineral surfaces at lower pH levels [62, 63]. Because *G. oregonensis* feed on organic material on the surface of the sediments, the concentrations of PTEs associated with acidic sediments contaminated by papermill sludge at Papermill Bay, are more likely to be assimilated by *G. oregonensis* and become bioaccumulated in their tissues. A survey of intertidal areas adjacent to papermills on the British Columbia coastline (including the papermill on Watson island) identified *G. oregonensis* as tolerant of papermill impacted shorelines [81]. The greater abundance of isopods (including *G. oregonensis*) in close proximity to British Columbian papermills, also observed by Robin et al. [82], is attributed to the provision of pulp fibers from the mill as a food source creating a stressed environment which enables these isopods to thrive.

Fig 7. The relationship between Chromium (Cr) Biota-Sediment Accumulation Factor (BSAF) and sediment pH for (A) Baltic clams (*Macoma balthica*) and (B) Oregon pill bugs (*Gnorumosphaeroma oregonensis*) sampled at five intertidal mudflats along the north coast of British Columbia, Canada.

Table 2. Correlation coefficients for the relationship between sediment pH and Biota-Sediment Accumulation Factors of 8 potentially toxic elements in Baltic clams (*Macoma balthica*) and Oregon pill bugs (*Gnorimosphaeroma oregonensis*) at all five sampling locations in the Skeena Estuary.

Species	As	Cd	Co	Cr	Cu	Ni	Pb	Zn
<i>M Balthica</i>	0.4397	0.2564	0.9162	0.8255*	-0.8177	0.9461	0.5788	0.896
<i>G. oregonensis</i>	-0.1744	0.3758	-0.7362*	-0.8688*	-0.1895	-0.8117	-0.2083	-0.7732*

* = $p < 0.05$

Conclusions

We found evidence to indicate that the discharge of papermill sludge from the decommissioned papermill on Watson Island has changed the sediment geochemistry at Papermill Bay by reducing the pH, and increasing the total and EDTA extractable concentrations of Cd, Cr, and Pb. However, the benthic invertebrate community composition confirms that the population has recovered from previous disturbance. Oregon pill bugs (*G. oregonensis*) were one of only two benthic invertebrate species observed at all five sites we visited. *G. oregonensis* are a cosmopolitan species along the north coast of British Columbia and are tolerant of sites contaminated with papermill effluent because they use the fibers discharged as a food source. Thus, we conclude here that *G. oregonensis* make an excellent candidate biomonitor species to assess recovery from the environmental impact of the papermill on Watson Island and monitor the future impacts of similar industrial developments in the region.

Acknowledgements

Sean Allen is acknowledged for fieldwork assistance. Ilse Kamerling is acknowledged for assisting C and N analysis. Anne Dudley is acknowledged for assisting ICP-OES analysis. Andy Dodson is acknowledged for assisting ICP-MS analysis. Accommodation and access to the shoreline were graciously provided by Cassiar Cannery Ecotourism lodge.

References

1. Parlee BL. Avoiding the Resource Curse: Indigenous Communities and Canada's Oil Sands. *World Development*. 2015;74:425-36. doi: <http://dx.doi.org/10.1016/j.worlddev.2015.03.004>.
2. Gerwing TG, Cox K. Erosion of trust in government consultation will impede the creation of environmental policy. *Marine Policy*. 2017;83:126-7.
3. Waldichuk M, Bousfield EL. Amphipods in low-oxygen marine waters adjacent to a sulphite pulp mill. *Journal of the Fisheries Board of Canada*. 1962;19(6):1163-5.
4. Pearson TH, Rosenberg R. A comparative study of the effects on the marine environment of wastes from cellulose industries in Scotland and Sweden. *Ambio*. 1976;77-9.
5. Tuominen TM, Sekela MA. Dioxins and furans in sediment and fish from the vicinity of four inland pulp and/or paper mills and one petroleum refinery in British Columbia: Environment Canada. Conservation and Protection; 1992.
6. Yunker MB, Lachmuth CL, Cretney WJ, Fowler BR, Dangerfield N, White L, et al. Biota-sediment partitioning of aluminium smelter related PAHs and pulp mill related diterpenes by intertidal clams at Kitimat, British Columbia. *Marine Environmental Research*. 2011;72(3):105-26.
7. Kritzer JP, DeLucia M-B, Greene E, Shumway C, Topolski MF, Thomas-Blate J, et al. The Importance of Benthic Habitats for Coastal Fisheries. *BioScience*. 2016;66(4):274-84. doi: 10.1093/biosci/biw014.
8. Carr-Harris C, Gottesfeld AS, Moore JW. Juvenile Salmon usage of the Skeena River Estuary. *PloS One*. 2015;10(3):e0118988.
9. Schindler DE, Scheuerell MD, Moore JW, Gende SM, Francis TB, Palen WJ. Pacific salmon and the ecology of coastal ecosystems. *Frontiers in Ecology and the Environment*. 2003;1(1):31-7.
10. Gerwing TG, Gerwing AMA, Macdonald T, Cox K, Juanes F, Dudas SE. Intertidal soft-sediment community does not respond to disturbance as postulated by the intermediate disturbance hypothesis. *Journal of Sea Research*. 2017;129:22-8.
11. Gerwing TG, Allen Gerwing AM, Drolet D, Barbeau MA, Hamilton DJ. Spatiotemporal variation in biotic and abiotic features of eight intertidal mudflats in the Upper Bay of Fundy, Canada. *Northeastern Naturalist*. 2015;22(12):1-44.
12. Amoozadeh E, Malek M, Rashidinejad R, Nabavi S, Karbassi M, Ghayoumi R, et al. Marine organisms as heavy metal bioindicators in the Persian Gulf and the Gulf of Oman. *Environmental Science and Pollution Research*. 2014;21(3):2386-95.
13. Gómez Gesteira JL, Dauvin J-C. Amphipods are good bioindicators of the impact of oil spills on soft-bottom macrobenthic communities. *Marine Pollution Bulletin*. 2000;40(11):1017-27.

14. Gerwing TG, Hamilton DJ, Barbeau MA, Haralampides KA, Yamazaki G. Short-term response of a downstream marine system to the partial opening of a tidal-river causeway. *Estuaries and Coasts* 2017;40(3):717-25. doi: 10.1007/s12237-016-0173-2.
15. Gerwing TG, Allen Gerwing AM, Macdonald T, Cox K, Juanes F, Dudas SE. Assessing the relationship between community dispersion and disturbance in a soft-sediment ecosystem. *Marine Ecology*. 2018;39(4):e12505.
16. Scholz B, Liebezeit G. Microphytobenthic dynamics in a Wadden Sea intertidal flat—Part II: Seasonal and spatial variability of non-diatom community components in relation to abiotic parameters. *European Journal of Phycology*. 2012;47(2):120-37.
17. van Proosdij D, Milligan T, Bugden G, Butler K. A tale of two macro tidal estuaries: differential morphodynamic response of the intertidal zone to causeway construction. *Journal of Coastal Research*. 2009;56:772–6.
18. de Souza Machado AA, Spencer K, Kloas W, Toffolon M, Zarfl C. Metal fate and effects in estuaries: A review and conceptual model for better understanding of toxicity. *Science of The Total Environment*. 2016;541:268-81. doi: <http://dx.doi.org/10.1016/j.scitotenv.2015.09.045>.
19. Moore JW, Gordon J, Carr-Harris C, Gottesfeld AS, Wilson SM, Russell JH. Assessing estuaries as stopover habitats for juvenile Pacific salmon. *Marine Ecology Progress Series*. 2016;559:201-15.
20. Wilkes B, Dwernychuk LW. Environment studies in the marine receiving environment at the Skeena Cellulose pulp mill, Watson Island, BC. *Pulp & Paper Canada*. 1991;92:10.
21. Gerwing TG. Preliminary report of intertidal research along the north coast of British Columbia: Summer 2016. Report to the Kitsumkalum First Nations. 34 p. 2016.
22. Gerwing TG, Cox K, Gerwing AMA, Carr-Harris CN, Dudas SE, Juanes F. Depth to the apparent redox potential discontinuity (aRPD) as a parameter of interest in marine benthic habitat quality models. *International Journal of Sediment Research*. 2018;33(2):149-56.
23. Sánchez-Moyano JE, García-Gómez JC. The arthropod community, especially Crustacea, as a bioindicator in Algeciras Bay (Southern Spain) based on a spatial distribution. *Journal of Coastal research*. 1998;14(3):1119-33.
24. Thomas JD. Biological monitoring and tropical biodiversity in marine environments: A critique with recommendations, and comments on the use of amphipods as bioindicators. *Journal of Natural History*. 1993;27(4):795-806.
25. Conlan K. Amphipod crustaceans and environmental disturbance: a review. *Journal of Natural History*. 1994;28(3):519-54.
26. Picone M, Bergamin M, Novelli Alessandra A, Noventa S, Delaney E, Barbanti A, et al. Evaluation of *Corophium orientale* as bioindicator for Venice Lagoon: Sensitivity assessment and toxicity-score proposal. *Ecotoxicology and Environmental Safety*. 2008;70(1):174-84.

503 27. Gerwing TG, Campbell L, Allen Gerwing AM, Allen S, Cox K, Rogers M, et al. Potential
504 impacts of logging on intertidal infaunal communities within the Kitimat River estuary.
505 *Journal of Natural History*. 2018;52(43-44):2833-55.

506 28. Pearson TH, Rosenberg R. Macrobenthic succession in relation to organic enrichment and
507 pollution of the marine environment. *Oceanography and Marine Biology: An Annual*
508 *Review*. 1978;16:229-331.

509 29. Allen L. The application of biodiversity indicators to infer ecosystem health in
510 regenerating tropical forest: University of Glasgow; 2019.

511 30. Zhang W, Dulloo E, Kennedy G, Bailey A, Sandhu H, Nkonya E. Biodiversity and
512 Ecosystem Services. *Sustainable Food and Agriculture*: Elsevier; 2019. p. 137-52.

513 31. Smith MD, Knapp AK. Exotic plant species in a C 4-dominated grassland: invasibility,
514 disturbance, and community structure. *Oecologia*. 1999;120(4):605-12.

515 32. Burke MJW, Grime JP. An experimental study of plant community invasibility. *Ecology*.
516 1996;77(3):776-90.

517 33. Guerra-García JM, García-Gómez JC. Polychaete assemblages and sediment pollution in
518 a harbour with two opposing entrances. *Helgoland Marine Research*. 2004;58(3):183-91.

519 34. HC. Hatfield Consultants Ltd. Skeena environmental effects monitoring (EEM) pre-
520 design reference document. Prepared for Skeena Cellulose Inc. 1994.

521 35. Hoos LM. The Skeena River Estuary: status of environmental knowledge to 1975: Report
522 of the Estuary Working Group, Department of the Environment, Regional Board, Pacific
523 Region. Environment Canada, 1975 Contract No.: 3.

524 36. NIBR. Northwest Institute for Bioregional Research: Valuation of the wild Salmon
525 economy of the Skeena River watershed. 1-30 2006.

526 37. Hilborn R, Walters CJ. Differing goals of salmon management on the Skeena River.
527 *Journal of the Fisheries Board of Canada*. 1977;34(1):64-72.

528 38. Higgins RJ, Schouwenburg WJ. A biological assessment of fish utilization of the Skeena
529 River Estuary, with special reference to port development in Prince Rupert. Technical Report
530 1973-1. Department of Environment, Fisheries and Marine Services. Vancouver, BC, 1973.

531 39. Faggetter BA. Review of the Environmental and Socioeconomic
532 Impacts of Marine Pollution in the
533 North and Central Coast Regions of British
534 Columbia. 2008.

535 40. Yunker MB, Cretney WJ, Ikonomou MG. Assessment of chlorinated dibenzo-p-dioxin
536 and dibenzofuran trends in sediment and crab hepatopancreas from pulp mill and harbor sites
537 using multivariate- and index-based approaches. *Environmental Science & Technology*.
538 2002;36(9):1869-78. doi: 10.1021/es0112893. PubMed PMID: WOS:000175311900006.

539 41. Akenhead S. A review of the oceanography and marine ecology of Prince Rupert
540 Harbour a propos sewage outfalls. 1992.

541 42. Cox K, Black M, Filip N, Miller M, Mohns K, Mortimor J, et al. Comparison of
542 community assessment techniques and implications for diversity indices and species
543 accumulation curves. *Ecology and Evolution*. 2017;DOI: 10.1002/ece3.3580.

544 43. BS7755-3.2. Soil Quality. Part 3: Chemical Methods. Section 3.2: Determination of pH
545 British Standards Institution, London, UK. 1995.

546 44. EPA US. Method 3051A (SW-846): Microwave Assisted Acid Digestion of Sediments,
547 Sludges, and Oils, Revision 1. Washington, DC. 2007.

548 45. EPA US. Method 7473 (SW-846): Mercury in Solids and Solutions by Thermal
549 Decomposition, Amalgamation, and Atomic Absorption Spectrophotometry, Revision 0.
550 Washington, DC. 1998.

551 46. Anderson M, Gorley RN, Clarke RK. Permanova+ for Primer: Guide to software and
552 statistical methods. Plymouth, United Kingdom: PRIMER-E Ltd; 2008.

553 47. Gerwing TG, Drolet D, Hamilton DJ, Barbeau MA. Relative importance of biotic and
554 abiotic forces on the composition and dynamics of a soft-sediment intertidal community
555 PLoS One. 2016;11(1):11:e0147098.

556 48. Clarke KR, Gorley RN. PRIMER v7: user manual/tutorial 3rd ed. Plymouth, United
557 Kingdom: Primer-E Ltd; 2015.

558 49. McLaren P. The environmental implications of sediment transport in the waters of Prince
559 Rupert, British Columbia, Canada: A comparison between kinematic and dynamic
560 approaches. *Journal of Coastal Research*. 2016;32(3):465-82.

561 50. Muller G. Index of geoaccumulation in sediments of the Rhine River. *Geojournal*.
562 1969;2:108-18.

563 51. Monte MC, Fuente E, Blanco A, Negro C. Waste management from pulp and paper
564 production in the European Union. *Waste management*. 2009;29(1):293-308.

565 52. Hoffman E, Lyons J, Boxall J, Robertson C, Lake CB, Walker TR. Spatiotemporal
566 assessment (quarter century) of pulp mill metal (loid) contaminated sediment to inform
567 remediation decisions. *Environmental monitoring and assessment*. 2017;189(6):257.

568 53. Borah P, Singh P, Rangan L, Karak T, Mitra S. Mobility, bioavailability and ecological
569 risk assessment of cadmium and chromium in soils contaminated by paper mill wastes.
570 *Groundwater for Sustainable Development*. 2018;6:189-99.

571 54. Nriagu JO, Nieboer E. Chromium in the natural and human environments: John Wiley &
572 Sons; 1988.

573 55. Kienle C, Langer-Jaesrich M, Baumberger D, Hohmann D, Santiago S, Köhler H-R, et al.
574 Integrated toxicity evaluation of a pulp deposit using organisms of different trophic levels.
575 *Journal of soils and sediments*. 2013;13(9):1611-25.

- 576 56. Alvarez MB, Malla ME, Batistoni DA. Comparative assessment of two sequential
577 chemical extraction schemes for the fractionation of cadmium, chromium, lead and zinc in
578 surface coastal sediments. *Fresenius' journal of analytical chemistry*. 2001;369(1):81-90.
- 579 57. Chakraborty P, Babu PVR, Sarma VV. A study of lead and cadmium speciation in some
580 estuarine and coastal sediments. *Chemical Geology*. 2012;294:217-25.
- 581 58. Sizmur T, Canário J, Gerwing TG, Mallory ML, O'Driscoll NJ. Mercury and
582 methylmercury bioaccumulation by polychaete worms is governed by both feeding ecology
583 and mercury bioavailability in coastal mudflats. *Environmental Pollution*. 2013;176:18-25.
- 584 59. Weimin Y, Batley G, Ahsanullah M. The ability of sediment extractants to measure the
585 bioavailability of metals to three marine invertebrates. *Science of the Total Environment*.
586 1992;125:67-84.
- 587 60. Burton ED, Phillips IR, Hawker DW. Geochemical partitioning of copper, lead, and zinc
588 in benthic, estuarine sediment profiles. *Journal of environmental quality*. 2005;34(1):263-73.
- 589 61. Luoma SN. Processes affecting metal concentrations in estuarine and coastal marine
590 sediments. *Heavy metals in the marine environment: CRC Press*; 2017. p. 51-66.
- 591 62. Riba I, Delvalls TÁ, Forja JM, Gómez-Parra A. The influence of pH and salinity on the
592 toxicity of heavy metals in sediment to the estuarine clam *Ruditapes philippinarum*.
593 *Environmental toxicology and chemistry*. 2004;23(5):1100-7.
- 594 63. Zhang C, Yu Z-g, Zeng G-m, Jiang M, Yang Z-z, Cui F, et al. Effects of sediment
595 geochemical properties on heavy metal bioavailability. *Environment International*.
596 2014;73:270-81. doi: <https://doi.org/10.1016/j.envint.2014.08.010>.
- 597 64. DelValls TÁ, Forja J, González-Mazo E, Gómez-Parra A, Blasco J. Determining
598 contamination sources in marine sediments using multivariate analysis. *TrAC Trends in*
599 *Analytical Chemistry*. 1998;17(4):181-92.
- 600 65. Chapman PM, Anderson J. A decision-making framework for sediment contamination.
601 *Integrated environmental assessment and management*. 2005;1(3):163-73.
- 602 66. Ciutat A, Boudou A. Bioturbation effects on cadmium and zinc transfers from a
603 contaminated sediment and on metal bioavailability to benthic bivalves. *Environmental*
604 *toxicology and chemistry*. 2003;22(7):1574-81.
- 605 67. Sizmur T, Canário J, Edmonds S, Godfrey A, O'Driscoll NJ. The polychaete worm *Nereis*
606 *diversicolor* increases mercury lability and methylation in intertidal mudflats. *Environmental*
607 *Toxicology and Chemistry*. 2013;32(8):1888-95.
- 608 68. Gerwing TG, Allen Gerwing AM, Macdonald T, Cox K, Juanes F, Dudas SE. Intertidal
609 soft-sediment community does not respond to disturbance as postulated by the intermediate
610 disturbance hypothesis. *Journal of Sea Research*. 2017;129:22-8.
- 611 69. Light SF. The Light and Smith manual: intertidal invertebrates from central California to
612 Oregon. 4 ed. Cameron JT, editor: University of California Press, Berkely, CA, USA; 2007.

70. Gerwing TG, Kim JH, Hamilton DJ, Barbeau MA, Addison JA. Diet reconstruction using next-generation sequencing increases the known ecosystem usage by a shorebird. *The Auk*. 2016;133(2):168–77.
71. Rainbow PS. Biomonitoring of heavy metal availability in the marine environment. *Marine pollution bulletin*. 1995;31(4-12):183-92.
72. Fauchald K, Jumars PA. The diet of worms: a study of polychaete feeding guilds. *Oceanography and Marine Biology Annual Review*. 1979;17:193-284.
73. Pagliosa PR. Another diet of worms: the applicability of polychaete feeding guilds as a useful conceptual framework and biological variable. *Marine Ecology*. 2005;26:246-54.
74. Howard L, Brown B. Natural variations in tissue concentration of copper, zinc and iron in the polychaete *Nereis diversicolor*. *Marine biology*. 1983;78(1):87-97.
75. Saha M, Sarkar S, Bhattacharya B. Interspecific variation in heavy metal body concentrations in biota of Sunderban mangrove wetland, northeast India. *Environment International*. 2006;32(2):203-7.
76. Saiz-Salinas J, Francés-Zubillaga G. *Nereis diversicolor*: an unreliable biomonitor of metal contamination in the ‘Ría de Bilbao’ (Spain). *Marine Ecology*. 1997;18(2):113-25.
77. Schöne BR, Krause Jr RA. Retrospective environmental biomonitoring–Mussel Watch expanded. *Global and Planetary Change*. 2016;144:228-51.
78. Zuykov M, Pelletier E, Harper DAT. Bivalve mollusks in metal pollution studies: from bioaccumulation to biomonitoring. *Chemosphere*. 2013;93(2):201-8.
79. Caçador I, Costa J, Duarte B, Silva G, Medeiros J, Azeda C, et al. Macroinvertebrates and fishes as biomonitors of heavy metal concentration in the Seixal Bay (Tagus Estuary): which species perform better? *Ecological Indicators*. 2012;19:184-90.
80. Sarkar SK, Cabral H, Chatterjee M, Cardoso I, Bhattacharya AK, Satpathy KK, et al. Biomonitoring of heavy metals using the bivalve molluscs in Sunderban mangrove wetland, northeast coast of Bay of Bengal (India): possible risks to human health. *CLEAN–Soil, Air, Water*. 2008;36(2):187-94.
81. Bard S, M. A biological index to predict pulp mill pollution levels. *Water environment research*. 1998;70(1):108-22.
82. Robin J, Harger E, Nassichuk MD. Marine intertidal community responses to Kraft pulp mill effluent. *Water, Air, and Soil Pollution*. 1974;3(1):107-22.

Supporting information captions

The supporting information file contains additional text that elaborates on the quality control and statistical analysis undertaken in addition to the following tables:

649 **S1 Table. PERMANCOVA showing sediment properties (pH, median particle diameter, C and N)**
 650 **varied by site, depth, and transect.**
 651

652 **S2 Table. PERMANCOVA showing that sediment total and available (EDTA extractable) PTEs varied**
 653 **by site and transect.**
 654

655 **S3 Table. Analysis of Variance for sediment properties. F-statistics of a two-way ANOVA with ‘site’**
 656 **and ‘depth’ as the two factors. The last two columns indicate whether reference sites; Tyee Banks**
 657 **(TB), Wolfe Cove (WC) and Inverness Passage (IP) are significantly ($p < 0.05$) different from**
 658 **potentially contaminated sites (Cassiar Cannery and Papermill Bay).**
 659

660 **S4 Table. Concentrations (mg kg^{-1}) of Cr, Co, Ni, Cu, Zn, As, Cd and Pb in benthic invertebrates**
 661 **sampled from five intertidal mudflats along the north coast of British Columbia, Canada. CC:**
 662 **Cassiar Cannery. WC: Wolfe Cove. IP: Inverness Passage. PB: Papermill Bay. TB: Tyee Banks**
 663

■ Cassiar Cannery

□ Inverness Passage

▲ Papermill bay

△ Wolf Cove

● Tyee Banks

- Cassiar Cannery
- Inverness Passage
- ▲ Papermill bay
- △ Wolfe Cove
- Tyee Banks

- Cassiar Cannery
- Inverness Passage
- ▲ Papermill bay
- △ Wolf Cove
- Tyee Banks

A

B

2D Stress: 0

Site

- ▲ CC
- ▼ IP
- PB
- ◆ TB
- WC