1 **Loss of F-box Motif Encoding Gene SAF1 and RRM3 Together Leads to** 2 **Synthetic Growth Defect and Sensitivity to HU, MMS in *S.cerevisiae***

3 **Meenu Sharma¹ V. Verma¹, Narendra K Bairwa^{1#}**

4 ¹School of Biotechnology, Shri Mata Vaishno Devi University, Katra, Jammu & Kashmir,
5 India- 182320

6 # To whom correspondence may be addressed: narendra.bairwa@smvdu.ac.in

7 Tel: (91-01991) 285699/285634; Fax: (91-01991)-285694

8 **Running title: SAF1 and RRM3 function in parallel pathways during genotoxic stress**

9 **KEY WORDS: *S. cerevisiae*, F-box motif, E-3 Ligase, Stress**

10 **Abstract:**

11 Unearthing of novel genetic interaction which leads to synthetic growth defects due to
12 inactivation of genes are needed for applications in precision medicine. The genetic
13 interactions among the molecular players involving different biological pathways need to be
14 investigated. The SAF1 gene of *S.cerevisiae* encodes for a protein product which contain N-
15 terminal F-box motif and C-terminal RCC1 domain. The F-box motif interacts with
16 Skp1 subunit of the SCF-E3 ligase and C-terminus with Aah1 (adenine deaminase) for
17 ubiquitination and subsequent degradation by 26S proteasome during phase transition from
18 proliferation state to quiescence phase due to nutrient limitation stress. The replication fork
19 associated protein Rrm3 of *S.cerevisiae* belongs to Pif1 family helicase and function in
20 removal of the non-histone proteins during replication fork movement. Here we have
21 investigated the genetic interaction among both the genes (SAF1 and RRM3) and their role in

22 growth fitness and genome stability. The single and double gene knockout strains of
 23 SAF1 and RRM3 genes was constructed in BY4741 genetic background and checked for the
 24 growth fitness in presence of genotoxic stress causing agents such as hydroxyurea and methyl
 25 methanesulfonate. The strains were also evaluated for nuclear migration defect by DAPI
 26 staining and for HIS3AI marked Ty1 retro-transposition. The *saf1Δrrm3Δ* showed the
 27 extremely slow growth phenotype in rich medium and sensitivity to genotoxic agents such as
 28 HU and MMS in comparison to single gene mutant (*saf1Δ*, *rrm3Δ*) and WT cells. The
 29 *saf1Δrrm3Δ* also showed the defects in nuclear migration as evident by multi-nuclei
 30 phenotype. The *saf1Δrrm3Δ* also showed the elevated frequency of Ty1 retro-transposition in
 31 JC2326 background in comparison to either *saf1Δ* or *rrm3Δ*. Based on these observations we
 32 report that that SAF1 and RRM3 functions in parallel pathway for growth fitness and stability
 33 of the genome.

34 **Introduction:**

35 The proliferation of *Saccharomyces cerevisiae* cells from in and out of quiescence phase due
 36 to nutrients stress or availability remains an active research area for both basic research and
 37 biotechnological purpose. The phase transition process is dependent on the ubiquitin
 38 proteasome system as it was reported that ubiquitin is required for survival during starvation
 39 (1). The SCF E3-ligase component Saf1, of *S.cerevisiae* which is an F-box motif containing
 40 protein recruits the Aah1 for proteasomal mediated degradation upon nutrient deprivation (2).
 41 The *AAH1* gene of *S.cerevisiae*, encodes adenine deaminase (Aah1) which converts adenine
 42 to hypoxanthine. The nutrient deprivation condition induces stress which leads to cell enter
 43 into the quiescence phase. Besides the Aah1, the serine protease B (PRB1), protease C
 44 (PRC1), and Ybr139w, vacuolar origin proteins also have been reported as Saf1 targets for

45 proteasome mediated degradation (3, 4).The serine proteases of the vacuole plays a major
46 role during starvation of a cell.

47 The Pif1 family helicases are well conserved from yeast to humans. In *S. cerevisiae*, RRM3
48 gene encoded product Rrm3 belongs to Pif1 family, shows the 5-3' helicase activity during
49 replication fork movement (5, 6). Rrm3 is involved in the chromosome replication and travels
50 along the replication fork (7).The activity of the Rrm3 counteracted by the replication fork
51 protection complex, Tof1/Csm3 (8). In the absence of Rrm3, replication fork pausing occurs
52 at the ITS1 region of r-DNA, telomeric regions, and difficult to replicate sites in the
53 chromosome (8, 9). Rrm3 was first identified as suppressor of recombination in the repeat
54 region(10).The loss of RRM3 results in the r-DNA circle formation which suggests its role in
55 maintenance of the r-DNA stability. The *rrm3* mutant showed the synthetic fitness defect
56 when combined with mutation in the genome stability regulators such as MRC1, SGS1,
57 RAD53, SRS2, MEC1 and RTT101(11-14). The RRM3 gene has also been implicated in the
58 suppression of the Ty1, Ty2 and Ty3 element (15, 16). Here in this paper we report the
59 genetic interactions among the SAF1 and RRM3 genes. The simultaneous deletion of both
60 gene leads to synthetic growth defects and genomic instability. The growth fitness of the
61 double deletion strain was even further reduced when exposed to MMS and HU in
62 comparison to single gene mutant or WT.

63 **Methods**

64 **Strain and Plasmids**

65 BY4741 (*Mata his3Δ1 leu2Δ0 met15Δ0 ura3Δ0*) *Saccharomyces cerevisiae* strain was used
66 for the construction of the gene deletions. Strain JC2326 (*MAT-ura3, cir0, ura3-167, leu: his,*
67 *his32Ty1his3AI-270, Ty1NEO-588, Ty1 (tub: lacs)-146*)was used for Ty1 retro-transposition

68 assay (Table1).Following, plasmids, pFA6a- KanMX6, pFA6a- His3MX6, and
69 pGADT7(Table 2) were used for PCR amplification of the deletion marker cassette
70 containing 40 bp homology to the upstream and downstream of the target ORFs.

71 **Growth Assay**

72 Growth assessment of the WT and mutants was carried out as mentioned (17). Growth
73 kinetics of strains was analyzed for 14 hrs by measuring optical density at 600nm using
74 TOSHVIN UV- 800 SHIMADZU spectrophotometer. The cultures were also streaked on the
75 YPD plates followed by incubation at 30°C for 2-3 days.

76 **Phase Contrast Microscopy**

77 For microscopy analysis, each strain was grown up to log phase in YPD medium at 30°C and
78 imaged under Leica DM3000 microscope at 100X magnification.

79 **Semi Quantitative Growth Assay**

80 To compare the growth fitness among the WT and mutants, spotting assay was performed.
81 Wild type (BY4741) and its deletion derivatives strains (*saf1Δ,rrm3Δ* and *saf1Δrrm3Δ*) were
82 grown in the 25 ml YPD (Yeast Extract 1% w/v, Peptone 2% w/v, dextrose 2% w/v) medium
83 overnight at 30°C. The next day the cultures were diluted and grown in fresh YPD medium
84 for 3-4hrs so as to reach early log phase (OD600 0.8-1.0). The cultures were equalized by OD
85 at 600nm. A tenfold serial dilution was done and spotted (3μl) onto agar plates containing
86 (YPD and YPD + HU and MMS). The plates were incubated at 30°C for 2-3 days and
87 imaged.

88 **Nuclear Migration Assay using Fluorescence microscopy**

89 For detection of nuclear migration defects, assay described in (18, 19) was adopted. The
90 number of nuclei per cell in yeast strains was determined with nuclear binding dye (DAPI).

Briefly, strains were grown to early log phase ($OD_{600} \sim 0.8$) at 30°C. Yeast cells were washed with distilled water and suspended in 1X PBS (Phosphate Buffer Saline). Further, fixation was done by addition of 70% ethanol before DAPI staining. Cells were washed with 1X PBS then again centrifuged for 1 minute at 2500 rpm. DAPI stain (1mg/ml stock) to final concentration of 2.5µg/ml was added and incubated for 5 minutes at room temperature and visualized under UV light of fluorescent microscope with 100X magnification. A total of 200 cells were counted and grouped according 0, 1, 2 and multi nuclei per cell, more than two nuclei indicated the nuclear migration defect.

Assay for Ty1 retro-transposition

To measure the Ty1 retro-transposition frequency the method mentioned in (16, 20) was followed. Briefly, single colony of JC2326 (reporter strain) and its deletion derivative (*sqf1Δ* and *rrm3Δ* and *sqf1Δrrm3Δ*) strains were inoculated into 10 ml YPD broth and was grown overnight at 30°C. The overnight grown cultures were again inoculated in 5ml YPD at 1:1000 dilutions and grown up to saturation point (144hrs) at 20°C. Then saturated culture was serially diluted and plated on minimal media (SD/His⁻ plates) followed by incubation at 30°C for 3-7 days. The frequency of appearance of His3⁺ colonies considered as measure of Ty1 retro-mobility.

Statistical methods:

Statistical significance of observations was determined using paired student t-test. P-value less than 0.05 indicated significant.

Results

The double mutant *saf1Δrrm3Δ* showed synthetic growth defects

The null mutant of the SAF1 and RRM3 reported to be viable (21). Our data also showed the both single gene mutant (*saf1Δ*, *rrm3Δ*) as viable. The growth of the single gene mutant slightly reduced in comparison to WT (**Figure 1A, C**). The phase contrast imaging of the single gene mutant appeared normal just like WT however the *rrm3Δ* mutant showed the enlargement of the cell (**Figure 1 B**).The growth characteristics of the double mutant *saf1Δrrm3Δ* showed reduced growth phenotype in comparison to *saf1Δ*, *rrm3Δ* and WT when grown in YPD medium (**Figure 1 (A, C)**). It has been reported previously that the growth of the *rrm3Δ* is slower than the WT. The double mutant showed the synthetic growth defects as indicated by streaking. The growth kinetics of the *saf1Δrrm3Δ* indicated that loss of both the gene leads to synthetic growth defects.

The *saf1Δrrm3Δ* cells exhibits Nuclear Migration defects

The analysis of the nucleus using DAPI stain indicated increased percentage of cells showing multi-nuclei phenotype in *saf1Δrrm3Δ* cells in comparison to *saf1Δ*, *rrm3Δ*, and WT (Figure 2A). The multi-nuclei phenotype indicated the defect in the nuclear migration..

The *saf1Δrrm3Δ* cells showed increased Ty1 retro–transposition

Ty1 elements are class of retro-transposons in the *S. cerevisiae* genome which remain in the dormant state due the host encoded genetic factors. In the absence of RRM3, the frequency of Ty1retro-transposition goes up as reported previously (15, 16). In this study the single gene mutant *saf1Δ* showed nearly 8 fold whereas the *rrm3Δ* showed the 85fold change in His⁺ prototroph formation in comparison to WT. However the absence of both the genes (*saf1Δrrm3Δ*) leads to increased frequency (~ 180-fold) of Ty1-retro transposition in the comparison to WT cells. (Figure 3 A, B).

Loss of SAF1 and RRM3 leads to HU and MMS sensitivity

The hydroxyurea (HU) and methyl methanesulfonate (MMS) act as replication checkpoint inhibitor and DNA damaging agent respectively. The Rrm3 mutant reported to have slightly sensitive phenotype in the 200mM hydroxyurea (22). With regard to SAF1 mutant the data is not available. The semi-quantitative growth assay by spot analysis on YPD medium and

YPD+genotoxic agents (MMS and HU) of the WT, *saf1Δ*, *rrm3Δ* and *saf1Δrrm3Δ* indicated the extremely slow growth of the *saf1Δrrm3Δ* when compared with the WT in presence of HU and MMS (Figure 4 A, B). The *saf1Δ*, *rrm3Δ* also showed the reduced growth in presence of the HU and MMS when compared with the WT.

Discussion

The multi-subunit enzymes complex, Skp1-Cul1-F-box (SCF) E-3, recruits the substrate or regulatory proteins through F-box motif containing protein for poly-ubiquitination and subsequent degradation by the 26S proteasome (23, 24). The Saf1 of *S.cerevisiae* is less characterized F-box protein which contains RCC1 repeats (25). The Saf1 has been shown to involved in the phase transition through Aah1p degradation during nutrient deprivation (25). In this paper we have investigated the genetic interaction among the SAF1 and genome stability regulator RRM3. The search for the genetic interactions and their impact on phenotype is crucial for the overall understanding of the biological processes and functioning of molecular player. The investigations of the binary interactions are crucial in unearthing of the pathways for better understanding of the system biology. The null mutant of the SAF1 have been reported as viable (21), exhibited the decrease in cell death (26) phenotype and resistance to histone deacetylase inhibitor CG-1521 drug (27). In our study also SAF1 null mutant showed the decreased growth in comparison to WT. The null mutant of RRM3 reported as viable (21) and showed the abnormal bud morphology (28). Our data with the *rrm3Δ*, showed the phenotype as reported earlier. The RRM3 mutant showed the slightly sensitive phenotype when grown in presence of 200mM HU (22) and MMS (0.01%) (29). In our study we observed that *rrm3Δ* showed slightly reduced growth when grown in presence of the 200mM HU and 0.035% MMS. The retro-transposition frequency in *S.cerevisiae* genome reported to be elevated (16, 30) including Ty1 (31) and Ty3 elements (32) in the null mutant of RRM3. The genetic interaction studies carried out previously reported that

SAF1 interact negatively with CDC 10, CDC11, CDC12, HYP2(33) and DDC1(34). The null mutant of SAF1 showed the synthetic growth defects with HSP82 (35), POL2 (36) and RTT109 (37). The *rrm3Δ* showed the synthetic growth defect with multiple genes such as CCR4, CCS1, CTF18, CTF4, CTF8, DCC1, MMS1, MRE11, PAP2, POP2, RAD51, RAD55, RAD57, RTT107, SRS2, SGS1(38)CTF8, DCC1, RAD52, RAD53 (39) in high throughput studies. However this is the first study which reports the genetic interaction among the SAF1 and RRM3 gene. Our data suggest that both the genes interact genetically to regulate the fitness of the cell and genome stability. The investigated genetic interaction may have implication for the cancer studies as there is higher degree of functional homology of Saf1p, Rrm3 with the mammalian HERC2 and Pif1 helicase respectively. In conclusion, our investigation on binary genetic interaction with the SCF-3 ligase component F- box motif encoding gene SAF1 and genome stability regulator RRM3 suggest that loss of both the genes in *S. cerevisiae* reduces the growth fitness of the WT cells.

Acknowledgment: We thank to Prof. M Joan Curio, Dr. Deepak Sharma, IMTECH, Dr. Ravi Manjithya, NCBS, India for strains and plasmids.

Funding information

This work was supported by a grant (BT/RLF/Re-entry/40/2012) from the Department of Biotechnology, GOI, New Delhi to N.K.B who is recipient of the Ramalingaswami fellowship from DBT, New Delhi.

Conflict of Interest: The authors declare that they have no conflicts of interest with the content of this article.

Author's contributions: NKB conceived and directed the study and wrote the paper with MS and VV. MS performed the experiments and analysed with NKB. VV provided the

bioinformatics facility and analysed the data. All the authors reviewed the results and approved the final version of manuscript.

References:

1. Finley D, Ozkaynak E, & Varshavsky A (1987) The yeast polyubiquitin gene is essential for resistance to high temperatures, starvation, and other stresses. *Cell* 48(6):1035-1046.
2. Escusa S, Camblong J, Galan JM, Pinson B, & Daignan-Fornier B (2006) Proteasome- and SCF-dependent degradation of yeast adenine deaminase upon transition from proliferation to quiescence requires a new F-box protein named Saf1p. *Molecular microbiology* 60(4):1014-1025.
3. Mark KG, Simonetta M, Maiolica A, Seller CA, & Toczyski DP (2014) Ubiquitin ligase trapping identifies an SCF(Saf1) pathway targeting unprocessed vacuolar/lysosomal proteins. *Molecular cell* 53(1):148-161.
4. Mark KG, *et al.* (2015) Prb1 Protease Activity Is Required for Its Recognition by the F-Box Protein Saf1. *Biochemistry* 54(29):4423-4426.
5. Zhou JQ, *et al.* (2002) *Schizosaccharomyces pombe* pfh1+ encodes an essential 5' to 3' DNA helicase that is a member of the PIF1 subfamily of DNA helicases. *Molecular biology of the cell* 13(6):2180-2191.
6. Ivessa AS, Zhou JQ, Schulz VP, Monson EK, & Zakian VA (2002) *Saccharomyces* Rrm3p, a 5' to 3' DNA helicase that promotes replication fork progression through telomeric and subtelomeric DNA. *Genes & development* 16(11):1383-1396.
7. Azvolinsky A, Dunaway S, Torres JZ, Bessler JB, & Zakian VA (2006) The *S. cerevisiae* Rrm3p DNA helicase moves with the replication fork and affects replication of all yeast chromosomes. *Genes & development* 20(22):3104-3116.
8. Mohanty BK, Bairwa NK, & Bastia D (2006) The Tof1p-Csm3p protein complex counteracts the Rrm3p helicase to control replication termination of *Saccharomyces cerevisiae*. *Proceedings of the National Academy of Sciences of the United States of America* 103(4):897-902.
9. Ivessa AS, *et al.* (2003) The *Saccharomyces cerevisiae* helicase Rrm3p facilitates replication past nonhistone protein-DNA complexes. *Molecular cell* 12(6):1525-1536.
10. Keil RL & McWilliams AD (1993) A gene with specific and global effects on recombination of sequences from tandemly repeated genes in *Saccharomyces cerevisiae*. *Genetics* 135(3):711-718.
11. Azam M, *et al.* (2006) Evidence that the *S.cerevisiae* Sgs1 protein facilitates recombinational repair of telomeres during senescence. *Nucleic acids research* 34(2):506-516.
12. Szyjka SJ, Viggiani CJ, & Aparicio OM (2005) Mrc1 is required for normal progression of replication forks throughout chromatin in *S. cerevisiae*. *Molecular cell* 19(5):691-697.
13. Taylor SD, *et al.* (2005) The conserved Mec1/Rad53 nuclear checkpoint pathway regulates mitochondrial DNA copy number in *Saccharomyces cerevisiae*. *Molecular biology of the cell* 16(6):3010-3018.
14. Schmidt KH & Kolodner RD (2004) Requirement of Rrm3 helicase for repair of spontaneous DNA lesions in cells lacking Srs2 or Sgs1 helicase. *Molecular and cellular biology* 24(8):3213-3226.

15. Scholes DT, Banerjee M, Bowen B, & Curcio MJ (2001) Multiple regulators of Ty1 transposition in *Saccharomyces cerevisiae* have conserved roles in genome maintenance. *Genetics* 159(4):1449-1465.
16. Bairwa NK, Mohanty BK, Stamenova R, Curcio MJ, & Bastia D (2011) The intra-S phase checkpoint protein Tof1 collaborates with the helicase Rrm3 and the F-box protein Dia2 to maintain genome stability in *Saccharomyces cerevisiae*. *The Journal of biological chemistry* 286(4):2445-2454.
17. Salari R & Salari R (2017) Investigation of the Best *Saccharomyces cerevisiae* Growth Condition. *Electronic physician* 9(1):3592-3597.
18. Palmer RE, Koval M, & Koshland D (1989) The dynamics of chromosome movement in the budding yeast *Saccharomyces cerevisiae*. *The Journal of cell biology* 109(6 Pt 2):3355-3366.
19. Palmer RE, Sullivan DS, Huffaker T, & Koshland D (1992) Role of astral microtubules and actin in spindle orientation and migration in the budding yeast, *Saccharomyces cerevisiae*. *The Journal of cell biology* 119(3):583-593.
20. Scholes DT, Kenny AE, Gamache ER, Mou Z, & Curcio MJ (2003) Activation of a LTR-retrotransposon by telomere erosion. *Proceedings of the National Academy of Sciences of the United States of America* 100(26):15736-15741.
21. Giaever G, *et al.* (2002) Functional profiling of the *Saccharomyces cerevisiae* genome. *Nature* 418(6896):387-391.
22. Ohya T, Arai H, Kubota Y, Shinagawa H, & Hishida T (2008) A SUMO-like domain protein, Esc2, is required for genome integrity and sister chromatid cohesion in *Saccharomyces cerevisiae*. *Genetics* 180(1):41-50.
23. Hershko A & Ciechanover A (1998) The ubiquitin system. *Annual review of biochemistry* 67:425-479.
24. Bai C, *et al.* (1996) SKP1 connects cell cycle regulators to the ubiquitin proteolysis machinery through a novel motif, the F-box. *Cell* 86(2):263-274.
25. Escusa S, *et al.* (2007) Skp1-Cullin-F-box-dependent degradation of Aah1p requires its interaction with the F-box protein Saf1p. *The Journal of biological chemistry* 282(28):20097-20103.
26. Teng X, *et al.* (2011) Gene-dependent cell death in yeast. *Cell death & disease* 2:e188.
27. Gaupel AC, Begley T, & Tenniswood M (2014) High throughput screening identifies modulators of histone deacetylase inhibitors. *BMC genomics* 15:528.
28. Watanabe M, Watanabe D, Nogami S, Morishita S, & Ohya Y (2009) Comprehensive and quantitative analysis of yeast deletion mutants defective in apical and isotropic bud growth. *Current genetics* 55(4):365-380.
29. Ang JS, Duffy S, Segovia R, Stirling PC, & Hieter P (2016) Dosage Mutator Genes in *Saccharomyces cerevisiae*: A Novel Mutator Mode-of-Action of the Mph1 DNA Helicase. *Genetics* 204(3):975-986.
30. Curcio MJ, *et al.* (2007) S-phase checkpoint pathways stimulate the mobility of the retrovirus-like transposon Ty1. *Molecular and cellular biology* 27(24):8874-8885.
31. Nyswaner KM, Checkley MA, Yi M, Stephens RM, & Garfinkel DJ (2008) Chromatin-associated genes protect the yeast genome from Ty1 insertional mutagenesis. *Genetics* 178(1):197-214.
32. Irwin B, *et al.* (2005) Retroviruses and yeast retrotransposons use overlapping sets of host genes. *Genome research* 15(5):641-654.
33. Costanzo M, *et al.* (2016) A global genetic interaction network maps a wiring diagram of cellular function. *Science (New York, N.Y.)* 353(6306).
34. Costanzo M, *et al.* (2010) The genetic landscape of a cell. *Science (New York, N.Y.)* 327(5964):425-431.
35. Zhao R, *et al.* (2005) Navigating the chaperone network: an integrative map of physical and genetic interactions mediated by the hsp90 chaperone. *Cell* 120(5):715-727.

36. Dubarry M, Lawless C, Banks AP, Cockell S, & Lydall D (2015) Genetic Networks Required to Coordinate Chromosome Replication by DNA Polymerases alpha, delta, and epsilon in *Saccharomyces cerevisiae*. *G3 (Bethesda, Md.)* 5(10):2187-2197.
37. Fillingham J, *et al.* (2008) Chaperone control of the activity and specificity of the histone H3 acetyltransferase Rtt109. *Molecular and cellular biology* 28(13):4342-4353.
38. Pan X, *et al.* (2006) A DNA integrity network in the yeast *Saccharomyces cerevisiae*. *Cell* 124(5):1069-1081.
39. Tong AH, *et al.* (2004) Global mapping of the yeast genetic interaction network. *Science (New York, N.Y.)* 303(5659):808-813.

318 **Tables:**

319 **Table 1: List of primers used for construction of deletion strains**

S.N.	Primer name	Sequence
1	Saf1/ YBR280c	F-5'CCAAAGGATATACTCTCAATTATAAATGGAAAAGCACATC
		CGGATCCCCGGGTAAATTAA-3'
		R-5'ACGGAATCCAAAATGCAAAATCGAAATGACACCTAAAAA TGAATTCGAGCTCGTTTAAAC-3'
2	Rrm3	F-5'AATTATCACTATCATTAGCCCAATAAACAGCTGAAAAGACGGATC
		CCCGGGTTAATTAA-3'
		R-5'AAATAACATAAACAAACAACCTCCGAACAATAACTAAGTACGAAT TCGAGCTCGTTTAAAC-3'
3	Rrm3 Leu	F-5'AGCTCAAAAGTCGAGAGATTGTCTTATAAGACATCCCGCCAACT GTGGAATACTCAG-3'
		R-5'TCAAACCTCAACTAGAGTATATGCATTATTCGTTGCAAGTTGGCCC GAAATCCCCTAC-3'

320

321 **Table 2: List of plasmids used for generating deletion cassette with selectable marker**

S.no.	Plasmid Name	Deletion casttee	PCR product size	Selection Media
1	pFA6a- KanMX6	<i>KanMX6</i>	1559 bp	YPD + G418
2	pFA6a- His3MX6	<i>His3M X6</i>	1403 bp	SC His -

322

323

324

325

326 **Table 3: Yeast strains and their genotype used in the study**

S.no.	Strain	Genotype	Source
1	BY4741 WT	<i>MATa his3Δ1 leu2Δ0 met15Δ0 ura3Δ0</i>	Dr. Deepak Sharma, IMTECH
2	<i>saf1Δ</i>	<i>saf1Δ::His3MX</i>	This study
3	<i>rrm3Δ</i>	<i>rrm3Δ::KanMX</i>	This study
4	<i>saf1Δ rrm3Δ</i>	<i>saf1Δ::His3MX, rrm3Δ::KanMX</i>	This study
5	JC2326	<i>MAT-ura3, cir0, ura3-167, leu: his, his32Ty1his3Δ1-270, Ty1NEO-588, Ty1 (tub: lacs)-146</i>	Prof. M Joan Curcio
6	MS1	<i>saf1Δ::KanMX</i>	This study
7	MS2	<i>rrm3Δ::KanMX</i>	This study
8	MS3	<i>saf1Δ::KanMX, rrm3Δ::LEU2</i>	This study

327
328 **Legends and Figures:**

329 **Figure 1.** Comparative analysis of growth and morphology of WT, *saf1Δ*, *rrm3Δ*,
330 *saf1Δrrm3Δ* strains. A. Growth of streaked strains on YPD plates, streaked culture were
331 incubated for 2-3 days at 30°C and then photographed. B. Phase contrast images of log phase
332 cultures at 100X magnification using Leica DM3000. C. Growth kinetics of strains (WT,
333 *saf1Δ*, *rrm3Δ*, *saf1Δrrm3Δ*). Cells were collected every 2 hour period and cellular growth
334 was measured by optical density (OD) at 600 nm using TOSHVIN UV- 1800 SHIMADZU.
335 The data shown represent the average of three independent experiments the error bars seen
336 represent the standard deviation for each set of data.

337

Figure 2. Assessment of nuclear migration phenotype using DAPI (4',6-diamidino-2-phenylindole) staining of WT, *saf1Δ*, *rrm3Δ*, *saf1Δrrm3Δ*. A. Representative Images of WT, *saf1Δ*, *rrm3Δ*, *saf1Δrrm3Δ* cells showing the status of Nuclear DNA migration. The images were acquired at the 100X magnification using Leica DM3000 fluorescent microscope. B Table showing the percentage from the count of 200 cells as, 0, 1, 2 and multi nucleus in each strain, more than two nuclei indicate the nuclear migration defect.

Figure 3. Assay for genome instability by measuring of *his3AI* marked Ty1 transposition frequency in WT, *saf1Δ*, *rrm3Δ*, *saf1Δrrm3Δ*. A Images of plates showing the Ty1 transposition induced colonies on Synthetic Dropout (SD) plate lacking histidine. B. Bar diagram showing the frequency of Ty1*his3AI* transposition in each strain. The data shown represent the average of three independent experiments. The significance of transposition was determined by using two tailed t-test. P-value (p) less than 0.05 indicates significant difference and the symbol * represent to p<0.05..

Figure 4. Comparative assessment of growth phenotype of WT, *saf1Δ*, *rrm3Δ* and *saf1Δrrm3Δ* strains in presence of 0.035% MMS and 200mM HU genotoxic stress causing agents. Each strain was grown to log phase and was equalized by O.D at 600 nm. The tenfold serially diluted samples of each strains spotted on YPD, YPD+stress agent containing agar plates. The result indicates that *saf1Δrrm3Δ* double mutant exhibits the sensitivity in the presence of HU and MMS.

A**WT*****saf1* Δ** ***rrm3* Δ** ***saf1* Δ *rrm3* Δ** **B Nuclear Migration Phenotype**

Yeast Strain	0 Nucleus per cell	1 Nucleus per cell	2 Nucleus per cell	Multi Nucleus per cell
BY4741 WT	0%	80%	20%	0%
<i>saf1</i> Δ	5%	45%	30%	20%
<i>rrm3</i> Δ	4%	19%	14%	63%
<i>saf1</i> Δ <i>rrm3</i> Δ	2%	12%	9%	78%

saf1 Δ *rrm3* Δ *saf1* Δ *rrm3* Δ