

## **Cofactor selectivity in methylmalonyl-CoA mutase, a model cobamide-dependent enzyme**

Olga M. Sokolovskaya<sup>1,2</sup>, Kenny C. Mok<sup>1</sup>, Jong Duk Park<sup>1,3</sup>, Jennifer L. A. Tran<sup>1,4</sup>, Kathryn A. Quanstrom<sup>1,5</sup>, and Michiko E. Taga<sup>1\*</sup>

<sup>1</sup> Department of Plant & Microbial Biology, University of California, Berkeley, Berkeley, CA 94720 USA

<sup>2</sup> Department of Chemistry, University of California, Berkeley, Berkeley, CA 94720 USA

<sup>3</sup> Present address: Department of Chemistry, Princeton University, Princeton, NJ 08544 USA

<sup>4</sup> Present address: Evidation Health, San Mateo, CA 94401 USA

<sup>5</sup> Present address: University of Michigan School of Medicine, Ann Arbor, MI 48109 USA

\* Correspondence to [taga@berkeley.edu](mailto:taga@berkeley.edu)

## Summary

Cobamides, a uniquely diverse family of enzyme cofactors related to vitamin B<sub>12</sub>, are produced exclusively by bacteria and archaea but used in all domains of life. While it is widely accepted that cobamide-dependent organisms require specific cobamides for their metabolism, the biochemical mechanisms that make cobamides functionally distinct are largely unknown. Here, we examine the effects of cobamide structural variation on a model cobamide-dependent enzyme, methylmalonyl-CoA mutase (MCM). The *in vitro* binding affinity of MCM for cobamides can be dramatically influenced by small changes in the structure of the lower ligand of the cobamide, and binding selectivity differs between bacterial orthologs of MCM. In contrast, variations in the lower ligand have minor effects on MCM catalysis. Bacterial growth assays demonstrate that cobamide requirements of MCM *in vitro* largely correlate with *in vivo* cobamide dependence. This result underscores the importance of enzyme selectivity in the cobamide-dependent physiology of bacteria.

## Introduction

Cobalamin, commonly referred to as vitamin B<sub>12</sub>, is a versatile enzyme cofactor used by organisms in all domains of life. In humans, cobalamin is essential for methionine synthesis and the breakdown of fatty acids, amino acids, and cholesterol (Brodie et al., 1970; Fenton et al., 1982). Bacteria and archaea additionally use cobalamin and related cofactors, cobamides, for deoxyribonucleotide synthesis (Licht et al., 1996), metabolism of various carbon and energy sources (Barker, 1985; Berg et al., 2007; Chang and Frey, 2000; Chen et al., 2001; Erb et al., 2008; Ferguson and Krzycki, 1997; Forage and Foster, 1982; Jeter, 1990; Korotkova et al., 2002; Krasotkina et al., 2001; Ljungdahl, 1986; Scarlett and Turner, 1976; Stupperich and Konle, 1993; Vrijbloed et al., 1999), synthesis of secondary metabolites (Allen and Wang, 2014; Blaszczyk et al., 2016; Bridwell-Rabb et al., 2017; Jung et al., 2014; Kim et al., 2017; Marous et al., 2015; Pierre et al., 2012; Werner et al., 2011), sensing light (Ortiz-Guerrero et al., 2011), and other processes (Barker, 1985; Chang and Frey, 2000; Chen et al., 2001; Cracan and Banerjee, 2012; Gough et al., 2000; Miles et al., 2011; Parks et al., 2013; Romine et al., 2017; Yaneva et al., 2012). The finding that 86% of bacterial species encode at least one cobamide-dependent enzyme in their genome (Shelton et al., 2019) demonstrates the prevalence of cobamide-dependent metabolisms. Widespread use of these cofactors can be attributed to their chemical versatility, as they facilitate challenging chemical reactions including radical-initiated rearrangements, methylation reactions, and reductive cleavage of chemical bonds (Banerjee and Ragsdale, 2003; Bridwell-Rabb and Drennan, 2017).

All cobamides share the same core structure (Figure 1): a corrin ring that coordinates a cobalt ion, a variable “upper” axial ligand (*R* in Figure 1), and a pseudo-nucleotide that is covalently attached to the corrin ring through an aminopropanol linker (Hodgkin et al., 1956) or an ethanolamine linker, in the case of nor-cobamides (Keller et al., 2016; Kräutler et al., 2003). The major differences among cobamides are in the structure of the nucleotide base, more commonly referred to as the lower axial ligand for its ability to coordinate the central cobalt ion. In cobalamin, the lower ligand is 5,6-dimethylbenzimidazole (Figure 1, boxed); in other cobamides, different benzimidazoles, phenolics, and purines constitute the lower ligand (Figure 2D) (Allen and Stabler, 2008; Girard et al., 2009; Men et al., 2015; Renz, 1999; Yan et al., 2018).

While cobamides containing different lower ligands share the same chemically reactive moieties, specifically the cobalt center and methyl or 5'-deoxyadenosyl upper axial ligands, they are nonetheless functionally distinct. Culture-based studies have shown that only a subset of cobamides supports a given bacterial metabolism, and uptake or production of other cobamides can inhibit growth (Helliwell et al., 2016; Keller et al., 2018; Mok and Taga, 2013; Watanabe et al., 1992; Yan et al., 2018; Yan et al., 2016; Yi et al., 2012). The requirements of bacteria for particular cobamides is notable given the diversity of cobamides present in host-associated and environmental samples (Allen and Stabler, 2008; Girard et al., 2009; Men et al., 2015), coupled with the absence of *de novo* cobamide biosynthesis in more than half of bacteria (Shelton et al., 2019). Despite the biological relevance of cobamide structure, and the prevalence of cobamide use among bacteria (Degnan et al., 2014a; Rodionov et al., 2003; Shelton et al., 2019; Zhang et al., 2009), little is understood about the biochemical mechanisms by which cobamides differentially impact microbial physiology.

The effect of lower ligand structure on the biochemistry of cobamide-dependent enzymes has been studied to a limited extent. In “base-on” enzymes, the lower ligand base coordinates the central cobalt ion of the cobamide, as drawn in Figure 1 (Larsson et al., 2010; Shibata et al., 2018; Yamanishi et al., 2002). Because the lower ligand is part of the catalytic center of the enzyme, lower ligand structure can influence catalysis through a variety of mechanisms (Brown, 2006; Conrad et al., 2015; Kozłowski and Zgierski, 2004), and cobamides unable to form an intramolecular coordinate bond are catalytically inactive in base-on enzymes (Poppe et al., 2000; Poppe et al., 1997). In contrast, in “base-off” enzymes the lower ligand is bound by the enzyme more than 10 Å away from the active site (Berkovitch et al., 2004; Bommer et al., 2014; Bridwell-Rabb et al., 2017; Drennan et al., 1994; Gruber et al., 2001; Jost et al., 2015; Kurteva-Yaneva et al., 2015; Mancía et al., 1996; Payne et al., 2015a; Payne et al., 2015b; Wolthers et al., 2010). In a subset of base-off enzymes, referred to as “base-off/His-on,” a histidine residue from the protein coordinates the cobalt ion in place of the lower ligand (Drennan et al., 1994; Mancía et al., 1996). Despite its distance from the reactive center, lower ligand structure affects the activity of base-off enzymes, as evidenced by the cobamide cofactor selectivity of methionine synthase (Tanioka et al., 2010), methylmalonyl-CoA mutase (MCM) (Lengyel et al., 1960; Poppe et al., 1997), reductive dehalogenases (Keller et al., 2018), and other enzymes (Barker et al., 1960; Lengyel et al., 1960; Poppe et al., 2000). However, the mechanisms by which lower ligand structure affects the biochemistry of base-off cobamide-dependent enzymes remain unclear.

As MCM is one of the most abundant cobamide-dependent enzymes in bacterial genomes (Shelton et al., 2019), and one of the two cobamide-dependent enzymes in humans, we have chosen to study the cobamide selectivity of MCM as a model for base-off/His-on enzymes, all of which share a structurally conserved B<sub>12</sub>-binding domain (Dowling et al., 2012; Drennan et al., 1994). MCM catalyzes the interconversion of (*R*)-methylmalonyl-CoA and succinyl-CoA, a bidirectional reaction used in propionate metabolism (Hosotani et al., 1980; Savvi et al., 2008; Vrijbloed et al., 1999), catabolism of branched amino acids and odd-chain fatty acids (Fenton et al., 2014; Savvi et al., 2008), polyhydroxybutyrate degradation (Charles et al., 1997), secondary metabolite biosynthesis (Hunaiti and Kolattukudy, 1984), and autotrophic carbon dioxide fixation (Berg et al., 2007; Han et al., 2012). MCM-dependent pathways have been harnessed industrially for the bioproduction of propionate, bioplastics, biofuels, and antibiotics (Akawi et al., 2015; Aldor et al., 2002; Dayem et al., 2002; Gonzalez-Garcia et al., 2017; Gross et al., 2006; Li et al., 2017; Reeves et al., 2007; Srirangan et al., 2013).

The presence of a cobamide lower ligand is required for MCM activity, as evidenced by the observation that adenosylcobinamide, a cobamide intermediate lacking a lower ligand (Figure 1), does not support MCM activity *in vitro* (Chowdhury and Banerjee, 1999). Three studies provide evidence that MCM is selective for cobamides with particular lower ligands. First, MCM from *Propionibacterium shermanii* was found to have different apparent  $K_M$  values for cobamides, increasing from AdoCbl to Ado[Bza]Cba to Ado[Ade]Cba (refer to Table 1 for full names of cobamides), and MCM from sheep had a higher apparent  $K_M$  for Ado[Bza]Cba than AdoCbl (Lengyel et al., 1960). Second, *P. shermanii* MCM had a lower apparent  $K_M$  for Ado[Cre]Cba than AdoCbl (Poppe et al., 1997). Third, in *Sinorhizobium meliloti* bacteroids MCM activity was highest with AdoCbl, intermediate with Ado[Bza]Cba, and absent with Ado[Ade]Cba (De Hertogh et al., 1964). Each of these studies includes only one or two cobamides other than cobalamin, and understandably so; cobamides are difficult to obtain in high quantities and must be purified from large volumes of bacterial cultures. Because of this, the response of MCM orthologs to the full diversity of cobamides has not been explored, and the mechanistic basis of cobamide selectivity remains unclear.

To investigate the mechanisms by which diverse lower ligands affect MCM function, we conducted *in vitro* binding and activity assays with MCM from *S. meliloti* (*SmMCM*). We discovered major differences in the binding affinities of eight naturally occurring cobamides for *SmMCM*, while cobamide structure affected enzyme activity to a lesser extent. Using six additional cobamides, five of which are novel analogs that have not been observed in nature or described previously, we identified structural elements of lower ligands that are determinants of binding to *SmMCM*. To probe the hypothesis that enzyme selectivity influences bacterial growth, we characterized the cobamide dependence of *S. meliloti* growth *in vivo*. By bridging the results of *in vitro* biochemistry of three bacterial MCM orthologs and the cobamide-dependent growth phenotypes of *S. meliloti*, we have elucidated molecular factors that contribute to the cobamide-dependent physiology of bacteria.

## Results

### Lower ligand structure influences cobamide binding to MCM.

We chose *SmMCM* as a model to examine how lower ligand structure influences MCM function based on previous work demonstrating its activity as a homodimer encoded by a single gene (Charles and Aneja, 1999; Miyamoto et al., 2003). We purified eight naturally occurring cobamides for *in vitro* studies of this protein, and chemically adenosylated each cobamide to produce the biologically active form used by MCM for catalysis. Previous studies showed that binding of cobamides to *P. shermanii* MCM can be detected *in vitro* by measuring quenching of intrinsic protein fluorescence (Chowdhury and Banerjee, 1999). We found that the fluorescence of purified, His-tagged *SmMCM* also decreased in a dose-dependent manner when the protein was reconstituted with increasing concentrations of AdoCbl (Figure 2A). The equilibrium dissociation constant ( $K_d$ ) derived from these measurements,  $0.03 \pm 0.02 \mu\text{M}$  (Figure 2D), is 6-fold lower than the  $K_d$  reported for *P. shermanii* MCM (Chowdhury and Banerjee, 1999). Ado-cobinamide also bound *SmMCM*, as was observed with *P. shermanii* MCM (Chowdhury and Banerjee, 1999), albeit with over 10-fold reduced affinity compared to cobalamin (Figure 2A, D).

We next measured binding of other benzimidazolyl cobamides to *SmMCM* and found that Ado[5-MeBza]Cba and Ado[Bza]Cba, the cobamides most structurally similar to AdoCbl, also bound the enzyme. However, the absence of one or two methyl groups, respectively, in the

lower ligands of these cobamides caused a decrease in binding affinity relative to AdoCbl (Figure 2A, D). Strikingly, no binding of Ado[5-OHBza]Cba to *Sm*MCM was detected at low micromolar concentrations. To rule out the possibility that Ado[5-OHBza]Cba binds *Sm*MCM but does not cause a fluorescence quench, we used an alternative, filtration-based, binding assay and observed little to no binding of Ado[5-OHBza]Cba to *Sm*MCM at micromolar concentrations (Figure S1A, B).

We expanded our analysis of *Sm*MCM-cobamide binding selectivity to include cobamides from other structural classes. Both of the phenolyl cobamides tested, Ado[Cre]Cba and Ado[Phe]Cba, bound *Sm*MCM with affinities similar to those of cobalamin and other benzimidazolyl cobamides (Figure 2B, D). In contrast, the purinyl cobamides Ado[2-MeAde]Cba and Ado[Ade]Cba had lower affinities for *Sm*MCM compared to most benzimidazolyl cobamides (Figure 2B, D): Ado[2-MeAde]Cba bound *Sm*MCM with ~20-fold lower affinity than cobalamin, and Ado[Ade]Cba did not bind to any significant extent at micromolar concentrations (verified by the filtration assay, Figure S1C). Interestingly, for all three classes of lower ligands, the presence of a methyl substituent promoted binding relative to other cobamides of the same structural class.

#### Identification of structural elements that interfere with cobamide binding.

When cobamides are bound to MCM, the lower ligand is surrounded by protein residues (Froese et al., 2010; Mancina et al., 1996). Therefore, the reduced affinity of certain cobamides for the enzyme could be a result of exclusion of their lower ligands from this binding pocket because of steric or electrostatic repulsion. We hypothesized that the poor binding of the purinyl cobamides Ado[Ade]Cba and Ado[2-MeAde]Cba is due to the presence of the exocyclic amine (N10) based on several observations: 1) Ado[5-OHBza]Cba, which also contains a polar functional group, had impaired binding (Figure 2A, D); 2) in a crystal structure of *Homo sapiens* MCM (*Hs*MCM) (Froese et al., 2010), residues Ala731, Thr726, and Phe721, all of which are conserved in *Sm*MCM, would be expected to sterically interfere with the binding of an exocyclic amine (Figure S2A, asterisk); and 3) structural modeling of *Hs*MCM, which shares 59% amino acid identity to *Sm*MCM in the B<sub>12</sub>-binding domain, bound to Ado[Ade]Cba suggests a significant displacement of the adenine lower ligand relative to the position of the lower ligand of AdoCbl, accompanied by significant expansion of the lower ligand binding pocket, which would be an unlikely conformation for the protein to adopt (Figure S2B-D).

To test the importance of the exocyclic amine of adenine in cofactor exclusion, we produced an unsubstituted purinyl cobamide, Ado[Pur]Cba (Yan et al., 2018). Ado[Pur]Cba also had low affinity for *Sm*MCM (Figure 2C, D), suggesting that the exocyclic amine of adenine is not a major cause of binding exclusion. Consistent with this result, two novel benzimidazolyl cobamides, Ado[7-MeBza]Cba and Ado[7-AmBza]Cba, bound *Sm*MCM with comparable affinities to other benzimidazolyl cobamides (Figure 2C, D), despite being functionalized at the position analogous to N10 of adenine. Rather, these results suggest that the presence of nitrogens in the six-membered ring of the lower ligand interferes with binding. To test this hypothesis directly, we produced three novel cobamide analogs that contain at least one nitrogen in the six-membered ring of the lower ligand base. Comparison of the binding of Ado[6-MePur]Cba and Ado[7-MeBza]Cba (Figure 2C) supported a role of ring nitrogens in binding inhibition, and comparison of binding affinities between Ado[Bza]Cba and Ado[5-AzaBza]Cba (Figures 2A, C), and between Ado[7-AmBza]Cba and Ado[3-DeazaAde]Cba (Figure 2C),

revealed that a single nitrogen atom in the six-membered ring of the lower ligand was sufficient to severely impair binding.

#### Bacterial MCM orthologs have distinct selectivity.

To test whether cofactor-binding selectivity is a general phenomenon across bacterial MCM orthologs, we compared the cobamide-binding profile of *Sm*MCM to that of MCM orthologs from *Escherichia coli* (*Ec*MCM) and *Veillonella parvula* (*Vp*MCM). Activity of *Ec*MCM with AdoCbl has been reported both *in vivo* and *in vitro*, although its physiological role in *E. coli* remains unclear (Gonzalez-Garcia et al., 2017; Haller et al., 2000). Annotations for two MCM homologs are present in the genome of *V. parvula*, and we purified the one that exhibits MCM activity when expressed in *S. meliloti* (see Materials and Methods). Because *S. meliloti* produces cobalamin (Kliwer and Evans, 1963), *E. coli* produces [2-MeAde]Cba when provided cobinamide (Hazra et al., 2015), and *V. parvula* produces [Cre]Cba (Crofts et al., 2013), we expected that each ortholog should have distinct cobamide selectivity. As expected, *Ec*MCM had highest affinity for its native cobamide, Ado[2-MeAde]Cba (Figure 3A, C). All other cobamides bound with 2- to 3-fold reduced affinities relative to Ado[2-MeAde]Cba. Similarly, *Vp*MCM had a higher affinity for Ado[Cre]Cba, its native cobamide, than AdoCbl (Figure 3B, C). *Vp*MCM also bound Ado[2-MeAde]Cba and Ado[Bza]Cba with similar affinity.

We constructed a sequence alignment of MCM orthologs from diverse organisms known to produce or use various cobamides, in search of amino acid residues that could account for differences in cobamide binding (Figure S3A). The B<sub>12</sub>-binding domains of diverse MCM orthologs had high overall amino acid identity (38-70%). Cases of low identity correlated with differences in the structural configuration of MCM, which occurs in different organisms as a homodimer (Froese et al., 2010; Haller et al., 2000; Miyamoto et al., 2010; Miyamoto et al., 2003), heterodimer (Birch et al., 1993; Mancina et al., 1996; Miyamoto et al., 2002; Zhang et al., 1999), or heterotetramer (Han et al., 2012; Yabuta et al., 2015) (Figure S3B). We focused our analysis on residues immediately surrounding the lower ligand in the structure of *Hs*MCM (Figure 3A, triangles). For the most part, these residues are highly conserved between orthologs. Interestingly, however, *Hs*MCM residues Phe638, Phe722, and Ala731, which are conserved in *Sm*MCM, are substituted with the more polar residues Tyr, Tyr, and Ser, respectively, in *Ec*MCM (Figure S3A), which has a higher affinity for purinyl cobamides. Whether or not these residues co-vary with cobamide selectivity across other MCM orthologs is difficult to interpret because the cobamide selectivity of MCM from most organisms is unknown.

#### The lower ligand of cobamides modulates MCM reaction kinetics.

We reconstituted *Sm*MCM with saturating amounts of the five cobamides that bound with highest affinity and measured conversion of (*R*)-methylmalonyl-CoA to succinyl-CoA under steady state conditions. Interestingly, the substrate K<sub>M</sub> was nearly invariable among the cobamides tested (Figure 4). Turnover was highest with AdoCbl (26 ± 1 s<sup>-1</sup>) and 2- to 3-fold lower with other cobamides. Thus, all of the cobamides tested, including the unnatural cobamide analogs, supported *Sm*MCM catalysis with modest differences in *k*<sub>cat</sub>. This finding is consistent with a previous observation that adenosylcobinamide-GDP, a cobamide precursor with an extended nucleotide loop and a guanine base, supported activity of *P. shermanii* MCM with only slight catalytic impairment compared to AdoCbl (Chowdhury et al., 2001).

MCM-dependent growth of *S. meliloti* correlates with the binding selectivity of *Sm*MCM for benzimidazolyl and purinyl cobamides, but not phenolyl cobamides.

To assess whether the cobamide-dependent growth of *S. meliloti* reflects MCM selectivity as observed *in vitro*, we cultured *S. meliloti* under conditions that require MCM activity. Examination of metabolic pathways encoded in the *S. meliloti* genome using the KEGG database (Kanehisa and Goto, 2000) suggests that the degradation of branched amino acids isoleucine and valine to succinyl-CoA, an intermediate of the citric acid cycle, requires MCM. Indeed, growth of *S. meliloti* on L-isoleucine and L-valine as the only carbon sources was dependent on the presence of the *bhbA* gene, which encodes MCM (Charles and Aneja, 1999) (Figure S4).

We constructed an *S. meliloti* strain incapable of synthesizing cobalamin and lacking cobamide-dependent enzymes other than MCM to ensure that differential growth could be attributed solely to MCM selectivity for added cobamides (see Materials and Methods). We cultivated this strain with L-isoleucine and L-valine as sole carbon sources in medium supplemented with different cobamides in their cyanylated (CN) forms, which is the form typically used for *in vivo* growth assays. Under these growth conditions, the maximum growth yield (OD<sub>600</sub>) achieved at high concentrations of all of the cobamides was indistinguishable (Figure S5). However, the concentration of cobamides required to achieve half of the maximal OD<sub>600</sub> (EC<sub>50</sub>) differed based on the cobamide provided (Figure 5). Consistent with the binding data, CNCbl had the lowest EC<sub>50</sub> value. EC<sub>50</sub> values for CN[Bza]Cba and CN[2-MeAde]Cba were 5-fold higher than CNCbl, and other cobamides had EC<sub>50</sub> values two orders of magnitude higher than CNCbl.

With the notable exception of the phenolyl cobamides, differences in the EC<sub>50</sub> values of cobamides *in vivo* qualitatively correlated with the binding selectivity that we observed *in vitro* (Figure 2). Among benzimidazolyl cobamides, EC<sub>50</sub> values increased from cobalamin to [Bza]Cba to [5-OHBza]Cba, consistent with the relative binding affinities of these cobamides. Similarly, [2-MeAde]Cba, which had an intermediate binding affinity for *Sm*MCM, had a lower EC<sub>50</sub> value than [Ade]Cba and [Pur]Cba, both of which did not bind to *Sm*MCM at low micromolar concentrations *in vitro*. The ability of [5-OHBza]Cba, [Ade]Cba, and [Pur]Cba to support growth suggests that these cobamides can bind *Sm*MCM at concentrations higher than those tested *in vitro*; a control experiment with an *S. meliloti* strain lacking MCM rules out the possibility that high concentrations of cobamides (10 μM) abiotically enable growth on isoleucine and valine (Figure S4).

We considered the possibility that differences in cobamide internalization by *S. meliloti* could also influence the EC<sub>50</sub> measurements shown in Figure 5. When *S. meliloti* cultures were supplemented with equimolar amounts of CNCbl, CN[Ade]Cba, or CN[Cre]Cba, the concentration of cobalamin extracted from the cellular fraction was 2- to 3-fold higher than [Ade]Cba and 5- to 6-fold higher than [Cre]Cba (Figure S6A). This result suggests that cobamides are differentially internalized or retained by the cells. However, MCM-dependent growth does not correlate with intracellular cobamide concentrations, as intracellular concentrations of cobalamin comparable to those of [Ade]Cba and [Cre]Cba supported *S. meliloti* growth to high densities (Figure S6). Therefore, the high EC<sub>50</sub> value of CN[Ade]Cba relative to CNCbl is more likely attributable to enzyme selectivity. Additional factors that could explain the high EC<sub>50</sub> of CN[Cre]Cba are considered in the Discussion.

## Discussion

Cobamides are distinct from other cofactors in their extensive structural diversity, with over a dozen forms that differ in the lower ligand base and nucleotide loop. How cobamide lower ligand structure influences the activity of cobamide-dependent enzymes has not been extensively explored. Here, we report a systematic analysis of the effects of cobamide lower ligand structure on the function of a model cobamide-dependent enzyme, MCM. Our results show that MCM exhibits varied affinities for different cobamides, and that this selectivity is linked to the physiology of the organism.

Our results show that the major determinant of cobamide selectivity in *Sm*MCM is binding, with small changes in the lower ligand capable of dramatically altering the binding affinity of a cobamide. One explanation for these differences is that the chemical compatibility between the lower ligand base and the binding pocket of the protein strongly influences the binding affinity of cobamides; repulsion of the lower ligand on the basis of electrostatics could reduce the binding affinity of cobamides to MCM. While the structure of *Sm*MCM has not been determined, a model generated by sequence alignment to *Hs*MCM suggested a highly hydrophobic lower ligand binding pocket. Consistent with this, we observed higher affinity of cobamides with hydrophobic lower ligands to *Sm*MCM, as well as interference of ring nitrogens with cobamide binding. Introducing mutations to change the properties of the lower ligand binding pocket of *Sm*MCM to accommodate alternative lower ligands proved challenging, as it resulted in reduced protein solubility and overall impaired cobamide binding (data not shown).

On the other hand, sequence alignments suggested that many of the hydrophobic residues predicted to immediately surround the lower ligand are conserved between diverse MCM orthologs that differ in cobamide selectivity. Assuming that the arrangement of the lower ligand binding pocket is similar across MCM orthologs, this suggests that interactions within the lower ligand binding pocket are not sufficient to account for selectivity. In a similar vein, examination of the residues surrounding the lower ligand in the cobamide-bound structures of reductive dehalogenases does not reveal the basis of exclusion of certain cobamides (Keller et al., 2018). These observations suggest that the lower ligand may have an unknown role in the binding of cobamides to MCM. Consistent with this idea, studies of the kinetics and pH dependence of AdoCbl binding to *P. shermanii* MCM suggest a pre-association step, wherein a cofactor-protein complex is formed prior to displacement of the lower ligand of the cofactor by a histidine residue in the protein (Chowdhury and Banerjee, 1999). The nature of this complex is unknown, but potential interactions between the lower ligand and this conformation of the enzyme could provide an opportunity for lower ligand structure to impact the outcome of binding.

Our analysis of MCM orthologs from *E. coli* and *V. parvula* demonstrates that variations in cobamide selectivity have evolved in organisms with different physiologies. The cobamide selectivity patterns in the three MCM orthologs we examined correlate with the physiologies of the bacteria in two ways. First, in all three cases, each MCM ortholog has highest affinity for the native cobamide produced by the organism, suggesting that cobamide biosynthesis and selectivity of cobamide-dependent enzymes have coevolved. Second, *Sm*MCM is more selective than *Ec*MCM and *Vp*MCM, and these differences in selectivity correlate with differences in cobamide biosynthesis, acquisition, and use in these organisms. *S. meliloti* synthesizes cobalamin *de novo* and is incapable of attaching purinyl and phenolyl lower ligands to cobamide precursors (Crofts et al., 2013). Thus, its cobamide-dependent enzymes have likely



evolved to function best with cobalamin. In contrast, *E. coli* does not synthesize cobamides *de novo* and instead relies on the importer BtuBFCD to acquire cobamides from the environment (Borths et al., 2005; Heller et al., 1985). Alternatively, *E. coli* can produce a variety of benzimidazolyl and purinyl cobamides when provided with precursors (Hazra et al., 2015), making the ability to use multiple cobamides likely advantageous. Like *S. meliloti*, *V. parvula* synthesizes cobamides *de novo*, but can produce both benzimidazolyl and phenolyl cobamides (Crofts et al., 2013; Hazra et al., 2013) and also encodes membrane transport components adjacent to cobalamin riboswitches (Mukherjee et al., 2019), which are likely to be cobamide importers (Nahvi et al., 2004; Rodionov et al., 2003). Thus, the ability of VpMCM to bind diverse cobamides is similarly consistent with its physiology.

Relative to cobamide binding selectivity, our results suggest that effects of lower ligand structure on the catalytic activity of MCM are minor. Among the cobamides we tested, the maximum differences in SmMCM turnover were 3-fold. We did not observe inhibition of MCM activity with any cobamides, in contrast to the strong inhibition that has been observed with analogs containing variations in the upper ligand or central metal, known as antivitamin (Calafat et al., 1995; Fukuoka et al., 2005; Widner et al., 2016).

In addition to elucidating the biochemical basis of cobamide selectivity in MCM, a major aim of our work was to link biochemical selectivity with cobamide-dependent growth. Our results with benzimidazolyl and purinyl cobamides support the hypothesis that enzyme selectivity is a major determinant of cobamide-dependent growth. Interestingly, although phenolyl cobamides bound SmMCM with high affinity and supported catalysis *in vitro*, high concentrations were required to support growth of *S. meliloti*. This discrepancy can be partially explained by poorer internalization or retention of these cofactors as compared to cobalamin (Figure S5). The observation that the intracellular cobamide concentrations were 50- to 190-fold higher than the amount added to the growth medium (Figure S5A) suggests that cobamides could be internalized by an uptake mechanism that favors cobalamin, distinct from both BtuBFCD and ECF-CbrT (Rodionov et al., 2009; Santos et al., 2018), both of which are absent from *S. meliloti*. Thus, we propose a model in which the cobamide-dependent growth of bacteria is influenced not only by binding selectivity of cobamide-dependent enzymes, but also by cobamide import (Figure 6). The lower effectiveness of phenolyl cobamides in supporting growth of *S. meliloti* could additionally be explained by inefficient adenosylation of these cobamides *in vivo*, as MCM requires the adenosyl upper axial ligand for activity. Whether or not adenosyltransferase enzymes, specifically CobA and PduO (Escalante-Semerena et al., 1990; Johnson et al., 2001) in *S. meliloti*, are selective with respect to lower ligand structure is unknown.

We and others have proposed the possibility of manipulating microbial communities using cobamides by taking advantage of the differential cobamide-dependent growth of bacteria (Abreu and Taga, 2016; Degnan et al., 2014b; Seth and Taga, 2014; Yan et al., 2018). Cobamides are predicted to mediate microbial interactions that are critical to the assembly of complex communities (Belzer et al., 2017; Degnan et al., 2014a; Heal et al., 2017; Helliwell et al., 2016; Ma et al., 2017; Men et al., 2015; Shelton et al., 2019; Yan et al., 2012), so the ability to selectively inhibit or promote the growth of particular species using corrinoids with various lower ligands could be applied to alter the composition of microbial communities in ways that could promote environmental and human health. This possibility hinges on the ability to predict which cobamides support or inhibit growth of an organism of interest, which requires an understanding of the major biochemical determinants of growth. We observed here that the cobamide binding selectivity of a model base-off cobamide-dependent enzyme correlates with

growth to a large extent. Thus, uncovering protein residues that confer selectivity would enable prediction of selectivity in cobamide-dependent enzymes, thereby facilitating prediction of the cobamide requirements of organisms of interest. Furthermore, our results suggest that additional steps of cobamide trafficking may be important determinants of cobamide-dependent growth. Future studies to understand how these various steps depend on cobamide structure will ultimately allow us to better understand, predict, and manipulate microbial interactions.

## Materials and Methods

### Chemical reagents

Chemicals were obtained from the following sources: 5'-chloro-5'-deoxyadenosine, Santa Cruz Biotechnology; 4-methylbenzimidazole, Accela; 5-methyl-1H-benzimidazole, ACROS Organics; phenol, J. T. Baker; zinc metal, Fisher Scientific; 5-methoxybenzimidazole, purine, and *para*-cresol, Alfa Aesar; methylmalonyl-CoA, methylmalonic acid, coenzyme A, adenosylcobalamin (coenzyme B<sub>12</sub>), cyanocobalamin, dicyanocobinamide, 6-methylpurine, 1H-imidazo[4,5-c]pyridine-4-amine (3-deazaadenine), benzimidazole, adenine hemisulfate, 5-azabenzimidazole, 1H-benzo[d]imidazol-7-amine (4-aminobenzimidazole), 2-methyl-1H-purine-6-amine (2-methyladenine), and bovine serum albumin (BSA), Sigma.

### Molecular cloning, protein expression and purification

*SmMCM* (locus SM\_b20757, *bhbA*) was expressed from the pET28a vector, with an N-terminal hexahistidine (6xHis) tag, in *E. coli* BL21(DE3)pLysS (cloning primers listed in Table S1). The expression strain was grown to an optical density at 600 nm (OD<sub>600</sub>) of 0.6-0.8 at 37 °C, cooled on ice for 15 min, and induced with 1 mM IPTG for 2.5 h at 37 °C. Cells were lysed by sonication in 25 mM Tris-HCl pH 8.0, 300 mM NaCl, 10 mM imidazole, with 0.5 mM PMSF, 1 µg/mL leupeptin, 1 µg/mL pepstatin, and 1 mg/mL lysozyme. Clarified lysate was treated with 0.05% polyethyleneimine. An ÄKTA Pure 25 Fast Protein Liquid Chromatography (FPLC) system was used to purify the protein over a GE 5 mL HisTrap HF column, using a gradient of 21 to 230 mM imidazole in the lysis buffer. Purified protein was dialyzed into 25 mM Tris-HCl pH 8.0, 300 mM NaCl, 10% glycerol and concentrated with a Vivaspin 10,000 MWCO protein concentrator. Final protein concentration was determined by A<sub>280</sub> using the theoretical extinction coefficient 55810 M<sup>-1</sup> cm<sup>-1</sup> (Fortier et al., 2012). *EcMCM* (locus b2917, *scpA*, previously *sbmA*) was expressed with an N-terminal 6xHis tag from a pET28a vector in *E. coli* BL21(DE3), by induction at OD<sub>600</sub> 0.6-0.8 with 0.1 mM IPTG, for 3.5 h at 30 °C. The protein was purified as described above and the final concentration was determined by Coomassie-stained SDS-PAGE, using BSA as a standard.

The *V. parvula* genome has two MCM annotations: a heterotetramer (loci Vpar\_RS06295, Vpar\_RS06290) and a heterodimer (loci Vpar\_RS09005, Vpar\_RS09000). The functionality of both homologs was tested by complementation in *S. meliloti*. The two putative *VpMCM* enzymes were cloned into the pTH1227 vector and transferred by conjugation into an *S. meliloti* *bhbA::Tn5* mutant. Complementation was assessed by growth in M9 liquid medium containing L-isoleucine and L-valine (see “*S. meliloti* growth assays” for additional details). *S. meliloti* co-expressing Vpar\_RS09005 and Vpar\_RS09000 showed identical growth to a strain expressing *SmMCM* from pTH1227 and was selected for *in vitro* studies.

The α subunit of *VpMCM* (encoded by Vpar\_RS09005) was expressed with an N-terminal 6xHis tag from the pET-Duet expression vector in *E. coli* BL21(DE3). Protein

expression was induced with 520  $\mu$ M IPTG for 6 h at 30 °C. The protein was batch purified by nickel affinity and subsequently purified by FPLC using a HiTrapQ column with an NaCl gradient from 50 to 500 mM in 20 Tris-HCl pH 8.0, 10% glycerol. The  $\beta$  subunit of VpMCM (encoded by Vpar\_RS09000) was expressed separately with an N-terminal 6xHis tag from the pET-Duet expression vector in *E. coli* BL21(DE3). Expression was induced with 1 mM IPTG for 22 h at 16 °C and the protein was purified using nickel-affinity chromatography as described for SmMCM. Purified protein was dialyzed into 25 mM Tris-HCl pH 8.0, 300 mM NaCl, 10% glycerol, and 1 mM  $\beta$ -mercaptoethanol. Concentration of  $\alpha$  and  $\beta$  subunits was determined by absorbance at 280 nm ( $A_{280}$ ) using the theoretically calculated extinction coefficients 75290 M<sup>-1</sup> cm<sup>-1</sup> and 74260 M<sup>-1</sup> cm<sup>-1</sup>, respectively (Fortier et al., 2012). Equimolar amounts of  $\alpha$  and  $\beta$  subunits were combined during the setup of fluorescence binding assays.

*E. coli* thiokinase containing an N-terminal 6xHis tag was expressed from a vector provided by Gregory Campanello from the laboratory of Ruma Banerjee. Expression was induced with 1 mM IPTG in *E. coli* BL21(DE3)pLysS at 28 °C for 3 h. The protein was purified as a heterodimer using nickel-affinity chromatography as described above. His-tagged *Rhodopseudomonas palustris* MatB (Crosby et al., 2012) was expressed from a pET16b expression plasmid provided by Omer Ad from the laboratory of Michelle Chang. The protein was overexpressed in *E. coli* BL21(DE3) at 16 °C overnight, after induction with 1 mM IPTG, and purified by nickel affinity chromatography as indicated above. Thiokinase and MatB concentrations were determined by Coomassie-stained SDS-PAGE, using BSA as a standard.

#### Guided biosynthesis, extraction, and purification of cobamides

*Sporomusa ovata* strain DSM 2662 was used for the production of its native cobamide, [Cre]Cba, and for production of [Phe]Cba, [5-MeBza]Cba, [Bza]Cba, [5-OHBza]Cba, [7-MeBza]Cba, and [7-AmBza]Cba, by guided biosynthesis as previously described (Mok and Taga, 2013). 5-OHBza was synthesized as described (Crofts et al., 2013). *Salmonella enterica* serovar Typhimurium strain LT2 and *Propionibacterium acidipropionici* strain DSM 20273 were used for production of [Ade]Cba (Hoffmann et al., 2000; Yi et al., 2012). [2-MeAde]Cba, [Pur]Cba, [5-AzaBza]Cba, [3-DeazaAde]Cba, and [6-MePur]Cba were produced by guided biosynthesis in *P. acidipropionici*. Cobamides were extracted as previously described (Yi et al., 2012) and purified by High Performance Liquid Chromatography (HPLC) using previously published methods (Crofts et al., 2014; Crofts et al., 2013; Yi et al., 2012) as well as additional methods listed in Table S2. In many cases more than one method was required to achieve high purity. Identity of cobamides was confirmed by Liquid Chromatography (LC) coupled to Mass Spectrometry (MS) using an Agilent 1260 LC/6120 quadrupole MS instrument. Lower ligand orientation in the novel cobamides [7-MeBza]Cba, [7-AmBza]Cba, [3-DeazaAde]Cba, and [6-MePur]Cba was inferred based on their absorbance spectra, which reveal a base-on conformation in the cyanylated form (Figure S7). The orientation of the lower ligands in [Pur]Cba and [5-AzaBza]Cba was not determined.

#### Chemical adenosylation of cobamides

Cobamide adenosylation was performed as previously described (Brown and Zou, 1999; Crofts et al., 2014). Briefly, cobamides at concentrations 0.5 – 1 mM were reduced with activated zinc metal under anaerobic conditions, with vigorous stirring for 0.5 – 2 h. 5'-chloro-5'-deoxyadenosine was added and adenosylation was allowed to proceed for 1 – 3 h in the dark. The progress of the reaction was monitored by HPLC. Following adenosylation, cobamides

were desalted using a C18 SepPak (Waters), purified by HPLC, desalted again, dried, and stored at -20 °C or -80 °C.

### Cobamide quantification

Purified cobamides were dissolved in water and quantified by UV-Vis spectrophotometry on a BioTek Synergy 2 plate reader using the following extinction coefficients: for cyanlated benzimidazolyl cobamides,  $\epsilon_{518} = 7.4 \times 10^3 \text{ M}^{-1} \text{ cm}^{-1}$  (Hogekamp, 1975); for cyanlated purinyl cobamides,  $\epsilon_{548} = 7.94 \times 10^3 \text{ M}^{-1} \text{ cm}^{-1}$  (Schneider and Stroinski, 1987); for cyanlated phenolyl cobamides,  $\epsilon_{495} = 9.523 \times 10^3 \text{ M}^{-1} \text{ cm}^{-1}$  (Stupperich et al., 1988); for adenosylated benzimidazolyl cobamides (AdoCbl, Ado[5-MeBza]Cba, Ado[Bza]Cba, Ado[5-OHBza]Cba, Ado[7-MeBza]Cba, and Ado[7-Ambza]Cba), which are predominantly base-on in water,  $\epsilon_{522} = 8.0 \text{ mM}^{-1} \text{ cm}^{-1}$  (Hogekamp, 1975); for adenosylated purinyl and phenolyl cobamides (Ado[Ade]Cba, Ado[2-MeAde]Cba, Ado[Pur]Cba, Ado[Cre]Cba, and Ado[Phe]Cba), which are predominantly base-off in water,  $\epsilon_{458} = 8.8 \text{ mM}^{-1} \text{ cm}^{-1}$  (Schneider and Stroinski, 1987); for adenosylated azabenzimidazolyl cobamides (Ado[3-DeazaAde]Cba, Ado[5-AzaBza]Cba and Ado[6-MePur]Cba), which are a mixture of base-on and base-off in water, the concentration was estimated from the average of concentrations calculated using the extinction coefficients above.

### Fluorescence Binding Assays

An *in vitro* assay previously described for measuring binding of AdoCbl to *P. shermanii* MCM (Chowdhury and Banerjee, 1999) was adapted to a 96-well format: MCM (0.2  $\mu\text{M}$ ) was combined with a range of cobamide concentrations (as specified in each experiment) in a black 96-well plate in 50 mM potassium phosphate pH 7.5 with 1 mM DTT, on ice. All steps involving cobamides were conducted in the dark. The plate was centrifuged for 1 min at 3800 rpm to level the surface of the liquid in each well. The plate was then incubated for 40 min at 30 °C to allow binding, with a brief shaking step after 30 min. Preliminary experiments showed that this time is sufficient for equilibration. Following incubation, fluorescence emission at 340 nm (5 nm slit width) was measured upon excitation at 282 nm (5 nm slit width) using a Tecan Infinite M1000 PRO Plate Reader. Fluorescence, normalized to the initial value, was plotted as a function of cobamide concentration, and fit to the following equation (Warner and Copley, 2007):

$$\frac{F}{F_0} = 1 + \frac{\Delta F_{max} ([E] + [L] + K_d) - \sqrt{([E] + [L] + K_d)^2 - 4[E][L]}}{2[E]}$$

where  $F$  is fluorescence,  $F_0$  is initial fluorescence,  $[E]$  is total enzyme concentration,  $[L]$  is total ligand concentration, and  $K_d$  is the binding dissociation constant.

### Filtration binding assay

Cobamides (10  $\mu\text{M}$ ) with and without MCM (15  $\mu\text{M}$ ) were incubated in 50 mM Tris-phosphate buffer pH 7.5 at 30 °C for 40 min, transferred to Nanosep 10K Omega centrifugal devices (Pall Corporation), and centrifuged for 5 minutes at 13,900 x g to separate unbound cobamides from enzyme-bound cobamides. The UV-Vis spectra of the filtrates were recorded on a BioTek Synergy 2 plate reader.

### Structural modeling

A model of *Sm*MCM was generated using the Swiss-Model software (Guex et al., 2009) based on the known crystal structure of *Homo sapiens* MCM (*Hs*MCM) (PDB ID: 2XIJ) (Froese et al., 2010). No major differences were observed in the B<sub>12</sub>-binding domain between *Sm*MCM

models generated from *HsMCM* and *Propionibacterium freudenrichii* MCM (PDB ID: 4REQ) (Mancia et al., 1996).

Maestro (Schrödinger Release 2015: Maestro) was used to generate a model of *HsMCM* bound to [Ade]Cba. The initial structure of *HsMCM* bound to cobalamin (PDB ID: 2XIQ) (Froese et al., 2010) was prepared using standard methods. A constrained energy minimization (atoms within 10 Å of cobalamin freely moving; atoms within a second 10 Å shell constrained by a force constant of 200; remaining structure frozen) was performed using MacroModel (Schrödinger Release 2015: MacroModel). The structure of the lower ligand was then modified to adenine, and the constrained energy minimization was repeated to generate a model of the lower ligand binding pocket bound to [Ade]Cba.

### Enzymatic synthesis of (*R*)-methylmalonyl-CoA

(*R*)-methylmalonyl-CoA synthesis reactions contained the following in 10 mL: 100 mM sodium phosphate pH 7.5, 20 mM MgCl<sub>2</sub>, 5 mM ATP, 10 mM methylmalonic acid, 2 mM coenzyme A, 5 mM β-mercaptoethanol, and 1.5 μM purified MatB protein. After combining ingredients on ice, the reaction was incubated at 37 °C for 1 h. The reaction was then frozen in liquid nitrogen and lyophilized. To purify (*R*)-methylmalonyl-CoA, the dried reaction mixture was resuspended in 3.2 mL water and the protein was precipitated with 200 μL trichloroacetic acid; precipitate was pelleted; supernatant was neutralized with 200 μL of 10 M NaOH; and salts and remaining starting materials were removed using a C18 SepPak column (Waters) (loaded in 0.1% formic acid, washed with water, methylmalonyl-CoA eluted with 50% methanol in water). Formation of (*R*)-methylmalonyl-CoA was initially verified by <sup>1</sup>H NMR and in subsequent preparations by HPLC (Table S3). The concentration of (*R*)-methylmalonyl-CoA was determined using an extinction coefficient of 12.2 mM<sup>-1</sup> cm<sup>-1</sup> at 259 nm.

### MCM activity assays

A thiokinase-coupled, spectrophotometric MCM activity assay was adapted from previous work (Taoka et al., 1994), except that ADP was used instead of GDP, and the experiment was conducted in 96-well plates. Final concentrations of reagents in the assays are as follows: Tris-phosphate buffer pH 7.5, 50 mM; DTNB, 400 μM; ADP, 1 mM; MgCl<sub>2</sub>, 10 mM; (*R*)-methylmalonyl-CoA, 0 – 4 mM; thiokinase, 5 μM; MCM, 50 nM; and cobamides, 2 μM. Preliminary experiments were conducted to ensure that concentrations of thiokinase, DTNB, and cobamides were not rate limiting.

Three separate mixes were prepared, all in 1X Tris-phosphate buffer: an assay mix containing DTNB, ADP, and MgCl<sub>2</sub>, a substrate mix containing (*R*)-methylmalonyl-CoA, and an enzyme mix containing thiokinase, MCM, and cobamides. All steps involving cobamides were conducted in the dark. The assay and enzyme mixes were prepared as a master mix and aliquoted into 96-well plates; substrate mixes were prepared in individual wells, in triplicate. All components were incubated at 30 °C for 40 minutes to equilibrate temperature and allow pre-binding of cobamides and MCM. After incubation, one replicate at a time, the substrate mix was added to the assay mix, followed by the enzyme mix. Absorbance at 412 nm ( $A_{412}$ ) was recorded immediately after addition of enzyme and for 1-3 minutes, every 3 seconds, on a BioTek Synergy 2 plate reader. The increase in  $A_{412}$  in reactions lacking substrate was subtracted from all readings, to account for reactivity of DTNB with thiols on protein surfaces.  $A_{412}$  values were converted to concentration of free CoA using a pathlength correction determined for the reaction volume and extinction coefficient of 14150 M<sup>-1</sup> cm<sup>-1</sup>.

### *S. meliloti* growth assays

MCM-dependent growth experiments were performed with *S. meliloti* strain Rm1021 *cobD::gus Gm<sup>R</sup> methH::Tn5 ΔnrdJ pMS03-nrdAB<sub>Ec</sub><sup>+</sup>*, which lacks cobamide-dependent enzymes other than MCM and does not synthesize cobalamin. *cobD* is required for cobalamin biosynthesis (Campbell et al., 2006), *methH* encodes methionine synthase (Banerjee et al., 1989; Campbell et al., 2006; Sato et al., 1974), and *nrdJ* encodes ribonucleotide reductase (Cowles et al., 1969). Because *nrdJ* is essential, the *E. coli* cobamide-independent ribonucleotide reductase encoded by *nrdA* and *nrdB* was expressed from the pMS03 plasmid (Taga and Walker, 2010). The strain was pre-cultured in M9 medium (Maniatis et al., 1989) (modified concentration of MgSO<sub>4</sub>: 1 mM) containing 0.1% sucrose, 2 g/L isoleucine, 2 g/L valine, 1 g/L methionine, and 20 μg/mL gentamycin, shaking at 30 °C. After two days, cells were washed and diluted to an OD<sub>600</sub> of 0.02 into M9 medium containing 4 g/L isoleucine, 4 g/L valine, 1 g/L methionine, 20 μg/mL gentamycin, and cobamides at various concentrations as indicated for each experiment, in 384-well plates. The plates were incubated at 30 °C for 145 h in a Biotek Synergy 2 plate reader with linear shaking at 1140 cpm. OD<sub>600</sub> was measured in 1 h increments.

For quantification of intracellular cobamides in *S. meliloti*, the strain above was pre-cultured as described, diluted into 50 mL of M9 medium containing 0.2% sucrose and various cobamides, and grown for 48 h (until OD<sub>600</sub> 0.6-0.8). Cobamides were extracted from cell pellets as previously described (Yi et al., 2012), using 5 mL of methanol containing 500 μg of potassium cyanide, and including a partial purification by means of a wash step with 20% methanol in water during the SepPak desalting procedure. Extracted cobamides were quantified by HPLC using peak areas at 525 nm and external standard curves, and cellular cobamide concentrations were calculated assuming 8 x 10<sup>8</sup> cells/mL at OD<sub>600</sub> 1.0 and cellular volume of 1 μm<sup>3</sup>.

### **Acknowledgements**

We thank current and past members of the Taga lab and Ruma Banerjee, Judith Klinman, Susan Marqusee, and David Savage for helpful discussions; Amrita Hazra, Amanda Shelton, Alexa Nicolas, Zachary Hallberg, Kristopher Kennedy, Joseph Maa, and Judith Klinman for critical reading of the manuscript; Anna Beatrice Grimaldo for help with guided biosynthesis of cobamide analogs; Amrita Hazra and Florian Widner for providing 5-OHBza; Victoria Innocent for assisting with cobamide accumulation assays; Ruma Banerjee, Gregory Campanello, Michelle Chang, and Omer Ad for providing expression plasmids; Kathleen Durkin for help with molecular modeling; and Krishna Niyogi and Arash Komeili for use of their equipment. This work was supported by NIH grants R01GM114535 and DP2AI117984 to MET. OS was also supported by the NIH Chemical Biology Training Grant T32 GM066698, JT by the UCSF work study program, and modeling work at the UC Berkeley Molecular Graphics and Computation Facility by NIH S10OD023532.

## Figures




Figure 1: The structure of cobalamin. The lower ligand, boxed, varies in other cobamides. Cobinamide, a cobamide precursor, lacks a nucleotide base (delineated by the wavy line).

Table 1: Abbreviations used for cobamides and upper axial ligands.

Abbreviation	Cobamide name
Cbl	cobalamin
[5-MeBza]Cba	5-methylbenzimidazolylcobamide
[Bza]Cba	benzimidazolylcobamide
[5-OHBza]Cba	5-hydroxybenzimidazolylcobamide
[Cre]Cba	<i>para</i> -cresolylcobamide
[Phe]Cba	phenolylcobamide
[Ade]Cba	adeninylcobamide
[2-MeAde]Cba	2-methyladeninylcobamide
[Pur]Cba	purinylcobamide
[7-MeBza]Cba	7-methylbenzimidazolylcobamide
[7-AmBza]Cba	7-aminobenzimidazolylcobamide
[5-AzaBza]Cba	5-azabenzimidazolylcobamide
[3-DeazaAde]Cba	3-deazaadeninylcobamide
[6-MePur]Cba	6-methylpurinylcobamide
Prefix	Upper ligand
Ado	5'-deoxyadenosine
CN	cyanide




Figure 2: Binding of diverse cobamides to *SmMCM* (see also Figures S1, S2). Fluorescence decrease of *SmMCM* when reconstituted with (A) benzimidazolyl cobamides and cobinamide, (B) phenolyl and purinyl cobamides, and (C) additional cobamide analogs (benzimidazolyl cobamides, blue; purinyl cobamides, orange; azabenzimidazolyl cobamides, black). Data points represent the mean and standard deviation of three technical replicates from a single experiment. (D)  $K_d$  values for different cobamides, reported as the average and standard deviation of three or more independent experiments, each consisting of technical triplicates. "n. d.," not determined, indicates that binding was too weak to determine  $K_d$ . \*While it was unreported at the time of our study, [Pur]Cba was discovered to be the cobamide naturally produced by *Desulfitobacterium hafniense* (Yan et al., 2018).






Figure 3: Binding selectivity of diverse MCM orthologs (see also Figure S3). Fluorescence binding assays with (A) *E. coli* MCM and (B) *V. parvula* MCM. Data points and error bars represent the mean and standard deviation, respectively, of technical triplicates from a single experiment; each replicate consisted of an independent cobamide dilution.  $K_d$  values from the fitted curves in (A) and (B) are reported in (C); error values reflect the standard error of the curve fit.  $K_d$  values for *VpMCM* binding to Ado[Cre]Cba and AdoCbl and for *EcMCM* binding to all cobamides were reproduced in independent experiments.




Figure 4: Activity of *Sinorhizobium meliloti* MCM with different cobamide cofactors. (A) Succinyl-CoA formation was detected using a coupled spectrophotometric assay (Taoka et al., 1994). DTNB, dithionitrobenzoate (Ellman's Reagent); TNB, thionitrobenzoate; CoA, Coenzyme A. (B) Michaelis-Menten kinetic analysis of *SmMCM* reconstituted with various cobamides. Data points and error bars represent the mean and standard deviation, respectively, of three technical replicates from one experiment; each replicate consisted of an independent substrate dilution. Kinetic constants are presented in (C).




Figure 5: MCM-dependent growth of *S. meliloti* *cobD::gus* Gm<sup>R</sup> *meth::Tn5*  $\Delta$ *nrdJ* pMS03-*nrdAB*<sub>Ec</sub><sup>+</sup> with various cobamides (see also Figures S4, S5). (A) Dose dependence of growth based on OD<sub>600</sub> at 145 h. Data points and error bars represent the mean and standard deviation, respectively, of three biological replicates from a single experiment. EC<sub>50</sub> values reported in (B) are the average and standard deviation of five or more biological replicates across two or more independent experiments.




Figure 6: Model describing biochemical determinants of cobamide-dependent growth in bacteria. Cobamides differentially impact bacterial growth due to selective cobamide import and biosynthesis, cobamide-binding selectivity of cobamide-dependent enzymes, and cobamide-dependent catalysis. For MCM-dependent growth of *S. meliloti*, cobamide-binding selectivity is most strongly correlated with cobamide-dependent growth of the organism.

## References

- Abreu, N.A., and Taga, M.E. (2016). Decoding molecular interactions in microbial communities. *FEMS Microbiol Rev* **40**, 648-663.
- Akawi, L., Srirangan, K., Liu, X., Moo-Young, M., and Perry Chou, C. (2015). Engineering *Escherichia coli* for high-level production of propionate. *J Ind Microbiol Biotechnol* **42**, 1057-1072.
- Aldor, I.S., Kim, S.W., Prather, K.L., and Keasling, J.D. (2002). Metabolic engineering of a novel propionate-independent pathway for the production of poly(3-hydroxybutyrate-co-3-hydroxyvalerate) in recombinant *Salmonella enterica* serovar Typhimurium. *Appl Environ Microbiol* **68**, 3848-3854.
- Allen, K.D., and Wang, S.C. (2014). Initial characterization of Fom3 from *Streptomyces wedmorensis*: The methyltransferase in fosfomycin biosynthesis. *Archives of biochemistry and biophysics* **543**, 67-73.
- Allen, R.H., and Stabler, S.P. (2008). Identification and quantitation of cobalamin and cobalamin analogues in human feces. *Am J Clin Nutr* **87**, 1324-1335.
- Banerjee, R., and Ragsdale, S.W. (2003). The many faces of vitamin B<sub>12</sub>: Catalysis by cobalamin-dependent enzymes. *Annu Rev Biochem* **72**, 209-247.
- Banerjee, R.V., Johnston, N.L., Sobeski, J.K., Datta, P., and Matthews, R.G. (1989). Cloning and sequence analysis of the *Escherichia coli methH* gene encoding cobalamin-dependent methionine synthase and isolation of a tryptic fragment containing the cobalamin-binding domain. *J Biol Chem* **264**, 13888-13895.
- Barker, H.A. (1985). beta-Methylaspartate-glutamate mutase from *Clostridium tetanomorphum*. *Methods in enzymology* **113**, 121-133.
- Barker, H.A., Smyth, R.D., Weissbach, H., Toohey, J.I., Ladd, J.N., and Volcani, B.E. (1960). Isolation and properties of crystalline cobamide coenzymes containing benzimidazole or 5,6-dimethylbenzimidazole. *J Biol Chem* **235**, 480-488.
- Belzer, C., Chia, L.W., Aalvink, S., Chamlagain, B., Piironen, V., Knol, J., and de Vos, W.M. (2017). Microbial metabolic networks at the mucus layer lead to diet-independent butyrate and vitamin B<sub>12</sub> production by intestinal symbionts. *mBio* **8**, e00770-00717.
- Berg, I.A., Kockelkorn, D., Buckel, W., and Fuchs, G. (2007). A 3-hydroxypropionate/4-hydroxybutyrate autotrophic carbon dioxide assimilation pathway in Archaea. *Sci* **318**, 1782-1786.
- Berkovitch, F., Behshad, E., Tang, K.H., Enns, E.A., Frey, P.A., and Drennan, C.L. (2004). A locking mechanism preventing radical damage in the absence of substrate, as revealed by the x-ray structure of lysine 5,6-aminomutase. *Proceedings of the National Academy of Sciences of the United States of America* **101**, 15870-15875.
- Birch, A., Leiser, A., and Robinson, J.A. (1993). Cloning, sequencing, and expression of the gene encoding methylmalonyl-coenzyme A mutase from *Streptomyces cinnamonensis*. *J Bacteriol* **175**, 3511-3519.
- Blaszczyk, A.J., Silakov, A., Zhang, B., Maiocco, S.J., Lanz, N.D., Kelly, W.L., Elliott, S.J., Krebs, C., and Booker, S.J. (2016). Spectroscopic and electrochemical characterization of the iron-sulfur and cobalamin cofactors of TsrM, an unusual radical S-adenosylmethionine methylase. *Journal of the American Chemical Society* **138**, 3416-3426.

- Bommer, M., Kunze, C., Fessler, J., Schubert, T., Diekert, G., and Dobbek, H. (2014). Structural basis for organohalide respiration. *Science* **346**, 455-458.
- Borths, E.L., Poolman, B., Hovorup, R.N., Locher, K.P., and Rees, D.C. (2005). In vitro functional characterization of BtuCD-F, the *Escherichia coli* ABC transporter for vitamin B<sub>12</sub> uptake. *Biochemistry* **44**, 16301-16309.
- Bridwell-Rabb, J., and Drennan, C.L. (2017). Vitamin B<sub>12</sub> in the spotlight again. *Current opinion in chemical biology* **37**, 63-70.
- Bridwell-Rabb, J., Zhong, A., Sun, H.G., Drennan, C.L., and Liu, H.W. (2017). A B<sub>12</sub>-dependent radical SAM enzyme involved in oxetanocin-A biosynthesis. *Nature* **544**, 322-326.
- Brodie, J.D., Burke, G.T., and Mangum, J.H. (1970). Methylcobalamin as an intermediate in mammalian methionine biosynthesis. *Biochemistry* **9**, 4297-4302.
- Brown, K.L. (2006). The enzymatic activation of coenzyme B<sub>12</sub>. *Dalton Transactions*, 1123-1133.
- Brown, K.L., and Zou, X. (1999). Thermolysis of coenzymes B<sub>12</sub> at physiological temperatures: activation parameters for cobalt-carbon bond homolysis and a quantitative analysis of the perturbation of the homolysis equilibrium by the ribonucleoside triphosphate reductase from *Lactobacillus leichmannii*. *J Inorg Biochem* **77**, 185-195.
- Calafat, A.M., Taoka, S., Puckett, J.M., Semerad, C., Yan, H., Luo, L., Chen, H., Banerjee, R., and Marzilli, L.G. (1995). Structural and electronic similarity but functional difference in methylmalonyl-CoA mutase between coenzyme B<sub>12</sub> and the analog 2',5'-dideoxyadenosylcobalamin. *Biochemistry* **34**, 14125-14130.
- Campbell, G.R., Taga, M.E., Mistry, K., Lloret, J., Anderson, P.J., Roth, J.R., and Walker, G.C. (2006). *Sinorhizobium meliloti bluB* is necessary for production of 5,6-dimethylbenzimidazole, the lower ligand of B<sub>12</sub>. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 4634-4639.
- Chang, C.H., and Frey, P.A. (2000). Cloning, sequencing, heterologous expression, purification, and characterization of adenosylcobalamin-dependent D-lysine 5,6-aminomutase from *Clostridium sticklandii*. *J Biol Chem* **275**, 106-114.
- Charles, T.C., and Aneja, P. (1999). Methylmalonyl-CoA mutase encoding gene of *Sinorhizobium meliloti*. *Gene* **226**, 121-127.
- Charles, T.C., Cai, G.Q., and Aneja, P. (1997). Megaplasmid and chromosomal loci for the PHB degradation pathway in *Rhizobium (Sinorhizobium) meliloti*. *Genetics* **146**, 1211-1220.
- Chen, H.P., Wu, S.H., Lin, Y.L., Chen, C.M., and Tsay, S.S. (2001). Cloning, sequencing, heterologous expression, purification, and characterization of adenosylcobalamin-dependent D-ornithine aminomutase from *Clostridium sticklandii*. *J Biol Chem* **276**, 44744-44750.
- Chowdhury, S., and Banerjee, R. (1999). Role of the dimethylbenzimidazole tail in the reaction catalyzed by coenzyme B<sub>12</sub>-dependent methylmalonyl-CoA mutase. *Biochemistry* **38**, 15287-15294.
- Chowdhury, S., Thomas, M.G., Escalante-Semerena, J.C., and Banerjee, R. (2001). The coenzyme B<sub>12</sub> analog 5'-deoxyadenosylcobinamide-GDP supports catalysis by methylmalonyl-CoA mutase in the absence of trans-ligand coordination. *J Biol Chem* **276**, 1015-1019.
- Conrad, K.S., Jordan, C.D., Brown, K.L., and Brunold, T.C. (2015). Spectroscopic and computational studies of cobalamin species with variable lower axial ligation:

- implications for the mechanism of Co-C bond activation by class I cobalamin-dependent isomerases. *Inorg Chem* **54**, 3736-3747.
- Cowles, J.R., Evans, H.J., and Russell, S.A. (1969). B<sub>12</sub> coenzyme-dependent ribonucleotide reductase in *Rhizobium* species and the effects of cobalt deficiency on the activity of the enzyme. *J Bacteriol* **97**, 1460-1465.
- Cracan, V., and Banerjee, R. (2012). Novel coenzyme B<sub>12</sub>-dependent interconversion of isovaleryl-CoA and pivalyl-CoA. *J Biol Chem* **287**, 3723-3732.
- Crofts, T.S., Hazra, A.B., Tran, J.L., Sokolovskaya, O.M., Osadchiy, V., Ad, O., Pelton, J., Bauer, S., and Taga, M.E. (2014). Regiospecific formation of cobamide isomers is directed by CobT. *Biochemistry* **53**, 7805-7815.
- Crofts, T.S., Seth, E.C., Hazra, A.B., and Taga, M.E. (2013). Cobamide structure depends on both lower ligand availability and CobT substrate specificity. *Chemistry & biology* **20**, 1265-1274.
- Crosby, H.A., Rank, K.C., Rayment, I., and Escalante-Semerena, J.C. (2012). Structure-guided expansion of the substrate range of methylmalonyl coenzyme A synthetase (MatB) of *Rhodopseudomonas palustris*. *Appl Environ Microbiol* **78**, 6619-6629.
- Dayem, L.C., Carney, J.R., Santi, D.V., Pfeifer, B.A., Khosla, C., and Kealey, J.T. (2002). Metabolic engineering of a methylmalonyl-CoA mutase-epimerase pathway for complex polyketide biosynthesis in *Escherichia coli*. *Biochemistry* **41**, 5193-5201.
- De Hertogh, A.A., Mayeux, P.A., and Evans, H.J. (1964). The relationship of cobalt requirement to propionate metabolism in *Rhizobium*. *Journal of Biological Chemistry* **239**, 2446-2453.
- Degnan, P.H., Barry, N.A., Mok, K.C., Taga, M.E., and Goodman, A.L. (2014a). Human gut microbes use multiple transporters to distinguish vitamin B<sub>12</sub> analogs and compete in the gut. *Cell host & microbe* **15**, 47-57.
- Degnan, P.H., Taga, M.E., and Goodman, A.L. (2014b). Vitamin B<sub>12</sub> as a modulator of gut microbial ecology. *Cell metabolism* **20**, 769-778.
- Dowling, D.P., Croft, A.K., and Drennan, C.L. (2012). Radical use of Rossmann and TIM barrel architectures for controlling coenzyme B<sub>12</sub> chemistry. *Annu Rev Biophys* **41**, 403-427.
- Drennan, C.L., Huang, S., Drummond, J.T., Matthews, R.G., and Ludwig, M.L. (1994). How a protein binds B<sub>12</sub>: A 3.0 Å X-ray structure of B<sub>12</sub>-binding domains of methionine synthase. *Science* **266**, 1669-1674.
- Erb, T.J., Retey, J., Fuchs, G., and Alber, B.E. (2008). Ethylmalonyl-CoA mutase from *Rhodobacter sphaeroides* defines a new subclade of coenzyme B<sub>12</sub>-dependent acyl-CoA mutases. *J Biol Chem* **283**, 32283-32293.
- Escalante-Semerena, J.C., Suh, S.J., and Roth, J.R. (1990). *cobA* function is required for both *de novo* cobalamin biosynthesis and assimilation of exogenous corrinoids in *Salmonella typhimurium*. *Journal of Bacteriology* **172**, 273-280.
- Fenton, W.A., Gravel, R.A., and Rosenblatt, D.S. (2014). Disorders of propionate and methylmalonate metabolism. In *The Online Metabolic and Molecular Bases of Inherited Disease*, A.L. Beaudet, B. Vogelstein, K.W. Kinzler, S.E. Antonarakis, A. Ballabio, K.M. Gibson, and G. Mitchell, eds. (New York, NY: The McGraw-Hill Companies, Inc.).
- Fenton, W.A., Hack, A.M., Willard, H.F., Gertler, A., and Rosenberg, L.E. (1982). Purification and properties of methylmalonyl coenzyme A mutase from human liver. *Archives of biochemistry and biophysics* **214**, 815-823.

- Ferguson, D.J., Jr., and Krzycki, J.A. (1997). Reconstitution of trimethylamine-dependent coenzyme M methylation with the trimethylamine corrinoid protein and the isozymes of methyltransferase II from *Methanosarcina barkeri*. *J Bacteriol* **179**, 846-852.
- Forage, R.G., and Foster, M.A. (1982). Glycerol fermentation in *Klebsiella pneumoniae*: functions of the coenzyme B<sub>12</sub>-dependent glycerol and diol dehydratases. *J Bacteriol* **149**, 413-419.
- Fortier, A., Grosdidier, A., Hernandez, C., Baratin, D., Kuznetsov, D., de Castro, E., Gasteiger, E., Csardi, G., Rossier, G., Stockinger, H., *et al.* (2012). ExPASy: SIB bioinformatics resource portal. *Nucleic acids research* **40**, W597-W603.
- Froese, D.S., Kochan, G., Muniz, J.R., Wu, X., Gileadi, C., Ugochukwu, E., Krysztowska, E., Gravel, R.A., Oppermann, U., and Yue, W.W. (2010). Structures of the human GTPase MMAA and vitamin B<sub>12</sub>-dependent methylmalonyl-CoA mutase and insight into their complex formation. *J Biol Chem* **285**, 38204-38213.
- Fukuoka, M., Nakanishi, Y., Hannak, R.B., Krautler, B., and Toraya, T. (2005). Homoadenosylcobalamins as probes for exploring the active sites of coenzyme B<sub>12</sub>-dependent diol dehydratase and ethanolamine ammonia-lyase. *Febs j* **272**, 4787-4796.
- Girard, C.L., Santschi, D.E., Stabler, S.P., and Allen, R.H. (2009). Apparent ruminal synthesis and intestinal disappearance of vitamin B<sub>12</sub> and its analogs in dairy cows. *Journal of dairy science* **92**, 4524-4529.
- Gonzalez-Garcia, R.A., McCubbin, T., Wille, A., Plan, M., Nielsen, L.K., and Marcellin, E. (2017). Awakening sleeping beauty: production of propionic acid in *Escherichia coli* through the *sbm* operon requires the activity of a methylmalonyl-CoA epimerase. *Microbial cell factories* **16**, 121.
- Gough, S.P., Petersen, B.O., and Duus, J.O. (2000). Anaerobic chlorophyll isocyclic ring formation in *Rhodobacter capsulatus* requires a cobalamin cofactor. *Proceedings of the National Academy of Sciences of the United States of America* **97**, 6908-6913.
- Gross, F., Ring, M.W., Perlova, O., Fu, J., Schneider, S., Gerth, K., Kuhlmann, S., Stewart, A.F., Zhang, Y., and Muller, R. (2006). Metabolic engineering of *Pseudomonas putida* for methylmalonyl-CoA biosynthesis to enable complex heterologous secondary metabolite formation. *Chemistry & biology* **13**, 1253-1264.
- Gruber, K., Reitzer, R., and Kratky, C. (2001). Radical shuttling in a protein: ribose pseudorotation controls alkyl-radical transfer in the coenzyme B<sub>12</sub> dependent enzyme glutamate mutase. *Angewandte Chemie (International ed in English)* **40**, 3377-3380.
- Guex, N., Peitsch, M.C., and Schwede, T. (2009). Automated comparative protein structure modeling with SWISS-MODEL and Swiss-PdbViewer: a historical perspective. *Electrophoresis* **30 Suppl 1**, S162-173.
- Haller, T., Buckel, T., Rétey, J., and Gerlt, J.A. (2000). Discovering new enzymes and metabolic pathways: Conversion of succinate to propionate by *Escherichia coli*. *Biochemistry* **39**, 4622-4629.
- Han, Y., Hawkins, A.S., Adams, M.W., and Kelly, R.M. (2012). Epimerase (Msed\_0639) and mutase (Msed\_0638 and Msed\_2055) convert (S)-methylmalonyl-coenzyme A (CoA) to succinyl-CoA in the *Metallosphaera sedula* 3-hydroxypropionate/4-hydroxybutyrate cycle. *Appl Environ Microbiol* **78**, 6194-6202.
- Hazra, A.B., Han, A.W., Mehta, A.P., Mok, K.C., Osadchiy, V., Begley, T.P., and Taga, M.E. (2015). Anaerobic biosynthesis of the lower ligand of vitamin B<sub>12</sub>. *Proceedings of the National Academy of Sciences of the United States of America* **112**, 10792-10797.

- Hazra, Amrita B., Tran, Jennifer L.A., Crofts, Terence S., and Taga, Michiko E. (2013). Analysis of substrate specificity in CobT homologs reveals widespread preference for DMB, the lower axial ligand of vitamin B<sub>12</sub>. *Chemistry & biology* 20, 1275-1285.
- Heal, K.R., Qin, W., Ribalet, F., Bertagnolli, A.D., Coyote-Maestas, W., Hmelo, L.R., Moffett, J.W., Devol, A.H., Armbrust, E.V., Stahl, D.A., *et al.* (2017). Two distinct pools of B<sub>12</sub> analogs reveal community interdependencies in the ocean. *Proceedings of the National Academy of Sciences of the United States of America* 114, 364-369.
- Heller, K., Mann, B.J., and Kadner, R.J. (1985). Cloning and expression of the gene for the vitamin B<sub>12</sub> receptor protein in the outer membrane of *Escherichia coli*. *J Bacteriol* 161, 896-903.
- Helliwell, K.E., Lawrence, A.D., Holzer, A., Kudahl, U.J., Sasso, S., Krautler, B., Scanlan, D.J., Warren, M.J., and Smith, A.G. (2016). Cyanobacteria and eukaryotic algae use different chemical variants of vitamin B<sub>12</sub>. *Curr Biol* 26, 999-1008.
- Hodgkin, D.C., Kamper, J., Mackay, M., Pickworth, J., Trueblood, K.N., and White, J.G. (1956). Structure of vitamin B<sub>12</sub>. *Nature* 178, 64-66.
- Hoffmann, B., Oberhuber, M., Stupperich, E., Bothe, H., Buckel, W., Konrat, R., and Kräutler, B. (2000). Native corrinoids from *Clostridium cochlearium* are adenylcobamides: Spectroscopic analysis and identification of pseudovitamin B<sub>12</sub> and factor A. *J Bacteriol* 182, 4773-4782.
- Hogenkamp, H.P.C. (1975). The chemistry of cobalamins and related compounds. In *Cobalamin Biochemistry and Pathophysiology*, B.M. Babior, ed. (John Wiley and Sons, Inc.), p. 55.
- Hosotani, K., Yokota, A., Nakano, Y., and Kitaoka, S. (1980). The metabolic pathway of propionate in *Euglena gracilis* z grown under illumination. *Agricultural and Biological Chemistry* 44, 1097-1103.
- Hunaiti, A.A., and Kolattukudy, P.E. (1984). Source of methylmalonyl-coenzyme A for erythromycin synthesis: Methylmalonyl-coenzyme A mutase from *Streptomyces erythreus*. *Antimicrobial agents and chemotherapy* 25, 173-178.
- Jeter, R.M. (1990). Cobalamin-dependent 1,2-propanediol utilization by *Salmonella typhimurium*. *Journal of general microbiology* 136, 887-896.
- Johnson, C.L., Pechonick, E., Park, S.D., Havemann, G.D., Leal, N.A., and Bobik, T.A. (2001). Functional genomic, biochemical, and genetic characterization of the *Salmonella pduO* gene, an ATP:cob(I)alamin adenosyltransferase gene. *J Bacteriol* 183, 1577-1584.
- Jost, M., Cracan, V., Hubbard, P.A., Banerjee, R., and Drennan, C.L. (2015). Visualization of a radical B<sub>12</sub> enzyme with its G-protein chaperone. *Proceedings of the National Academy of Sciences of the United States of America* 112, 2419-2424.
- Jung, W.S., Kim, E., Yoo, Y.J., Ban, Y.H., Kim, E.J., and Yoon, Y.J. (2014). Characterization and engineering of the ethylmalonyl-CoA pathway towards the improved heterologous production of polyketides in *Streptomyces venezuelae*. *Applied microbiology and biotechnology* 98, 3701-3713.
- Kanehisa, M., and Goto, S. (2000). KEGG: kyoto encyclopedia of genes and genomes. *Nucleic acids research* 28, 27-30.
- Keller, S., Kunze, C., Bommer, M., Paetz, C., Menezes, R.C., Svatoš, A., Dobbek, H., and Schubert, T. (2018). Selective utilization of benzimidazolyl-norcobamides as cofactors by the tetrachloroethene reductive dehalogenase of *Sulfurospirillum multivorans*. *Journal of Bacteriology*.

- Keller, S., Treder, A., von Reuss, S.H., Escalante-Semerena, J.C., and Schubert, T. (2016). The SMUL\_1544 gene product governs norcobamide biosynthesis in the tetrachloroethene-respiring bacterium *Sulfurospirillum multivorans*. *J Bacteriol* **198**, 2236-2243.
- Kim, H.J., Liu, Y.N., McCarty, R.M., and Liu, H.W. (2017). Reaction catalyzed by GenK, a cobalamin-dependent radical S-adenosyl-L-methionine methyltransferase in the biosynthetic pathway of gentamicin, proceeds with retention of configuration. *Journal of the American Chemical Society* **139**, 16084-16087.
- Kliwer, M., and Evans, H.J. (1963). Identification of cobamide coenzyme in nodules of symbionts & isolation of the B<sub>12</sub> coenzyme from *Rhizobium meliloti*. *Plant physiology* **38**, 55-59.
- Korotkova, N., Chistoserdova, L., Kuksa, V., and Lidstrom, M.E. (2002). Glyoxylate regeneration pathway in the methylotroph *Methylobacterium extorquens* AM1. *Journal of Bacteriology* **184**, 1750-1758.
- Kozlowski, P.M., and Zgierski, M.Z. (2004). Electronic and steric influence of trans axial base on the stereoelectronic properties of cobalamins. *The Journal of Physical Chemistry B* **108**, 14163-14170.
- Krasotkina, J., Walters, T., Maruya, K.A., and Ragsdale, S.W. (2001). Characterization of the B<sub>12</sub>- and iron-sulfur-containing reductive dehalogenase from *Desulfitobacterium chlororespirans*. *J Biol Chem* **276**, 40991-40997.
- Kräutler, B., Fieber, W., Ostermann, S., Fasching, M., Ongania, K.-H., Gruber, K., Kratky, C., Mikl, C., Siebert, A., and Diekert, G. (2003). The cofactor of tetrachloroethene reductive dehalogenase of *Dehalospirillum multivorans* is norpseudob<sub>12</sub>, a new type of a natural corrinoid, Vol 86.
- Kurteva-Yaneva, N., Zahn, M., Weichler, M.T., Starke, R., Harms, H., Muller, R.H., Strater, N., and Rohwerder, T. (2015). Structural basis of the stereospecificity of bacterial B<sub>12</sub>-dependent 2-hydroxyisobutyryl-CoA mutase. *J Biol Chem* **290**, 9727-9737.
- Larsson, K.M., Logan, D.T., and Nordlund, P. (2010). Structural basis for adenosylcobalamin activation in AdoCbl-dependent ribonucleotide reductases. *ACS chemical biology* **5**, 933-942.
- Lengyel, P., Mazumder, R., and Ochoa, S. (1960). Mammalian methylmalonyl isomerase and vitamin B<sub>12</sub> coenzymes. *Proceedings of the National Academy of Sciences of the United States of America* **46**, 1312-1318.
- Li, J., Zhu, X., Chen, J., Zhao, D., Zhang, X., and Bi, C. (2017). Construction of a novel anaerobic pathway in *Escherichia coli* for propionate production. *BMC biotechnology* **17**, 38-38.
- Licht, S., Gerfen, G.J., and Stubbe, J. (1996). Thiyl radicals in ribonucleotide reductases. *Science* **271**, 477-481.
- Ljungdahl, L.G. (1986). The autotrophic pathway of acetate synthesis in acetogenic bacteria. *Annual review of microbiology* **40**, 415-450.
- Ma, A.T., Beld, J., and Brahmsha, B. (2017). An amoebal grazer of cyanobacteria requires cobalamin produced by heterotrophic bacteria. *Applied and Environmental Microbiology* **83**, e00035-00017.
- Mancia, F., Keep, N.H., Nakagawa, A., Leadlay, P.F., McSweeney, S., Rasmussen, B., Bosecke, P., Diat, O., and Evans, P.R. (1996). How coenzyme B<sub>12</sub> radicals are generated: the crystal structure of methylmalonyl-coenzyme A mutase at 2 Å resolution. *Structure (London, England : 1993)* **4**, 339-350.



- Maniatis, T., Fritsch, E.F., and Sambrook, J. (1989). *Molecular cloning: a laboratory manual* (Cold spring harbor laboratory press).
- Marous, D.R., Lloyd, E.P., Buller, A.R., Moshos, K.A., Grove, T.L., Blaszczyk, A.J., Booker, S.J., and Townsend, C.A. (2015). Consecutive radical S-adenosylmethionine methylations form the ethyl side chain in thienamycin biosynthesis. *Proceedings of the National Academy of Sciences of the United States of America* *112*, 10354-10358.
- Men, Y., Seth, E.C., Yi, S., Crofts, T.S., Allen, R.H., Taga, M.E., and Alvarez-Cohen, L. (2015). Identification of specific corrinoids reveals corrinoid modification in dechlorinating microbial communities. *Environmental microbiology* *17*, 4873-4884.
- Miles, Z.D., McCarty, R.M., Molnar, G., and Bandarian, V. (2011). Discovery of epoxyqueuosine (oQ) reductase reveals parallels between halorespiration and tRNA modification. *Proceedings of the National Academy of Sciences of the United States of America* *108*, 7368-7372.
- Miyamoto, E., Tanioka, Y., Nishizawa-Yokoi, A., Yabuta, Y., Ohnishi, K., Misono, H., Shigeoka, S., Nakano, Y., and Watanabe, F. (2010). Characterization of methylmalonyl-CoA mutase involved in the propionate photoassimilation of *Euglena gracilis* Z. *Archives of microbiology* *192*, 437-446.
- Miyamoto, E., Watanabe, F., Charles, T.C., Yamaji, R., Inui, H., and Nakano, Y. (2003). Purification and characterization of homodimeric methylmalonyl-CoA mutase from *Sinorhizobium meliloti*. *Archives of microbiology* *180*, 151-154.
- Miyamoto, E., Watanabe, F., Yamaji, R., Inui, H., Sato, K., and Nakano, Y. (2002). Purification and characterization of methylmalonyl-CoA mutase from a methanol-utilizing bacterium, *Methylobacterium extorquens* NR-1. *Journal of nutritional science and vitaminology* *48*, 242-246.
- Mok, K.C., and Taga, M.E. (2013). Growth inhibition of *Sporomusa ovata* by incorporation of benzimidazole bases into cobamides. *J Bacteriol* *195*, 1902-1911.
- Mukherjee, S., Das Mandal, S., Gupta, N., Drory-Retwitzer, M., Barash, D., and Sengupta, S. (2019). RiboD: A comprehensive database for prokaryotic riboswitches. *Bioinformatics* (Oxford, England).
- Nahvi, A., Barrick, J.E., and Breaker, R.R. (2004). Coenzyme B<sub>12</sub> riboswitches are widespread genetic control elements in prokaryotes. *Nucleic acids research* *32*, 143-150.
- Ortiz-Guerrero, J.M., Polanco, M.C., Murillo, F.J., Padmanabhan, S., and Elias-Arnanz, M. (2011). Light-dependent gene regulation by a coenzyme B<sub>12</sub>-based photoreceptor. *Proceedings of the National Academy of Sciences of the United States of America* *108*, 7565-7570.
- Parks, J.M., Johs, A., Podar, M., Bridou, R., Hurt, R.A., Jr., Smith, S.D., Tomanicek, S.J., Qian, Y., Brown, S.D., Brandt, C.C., *et al.* (2013). The genetic basis for bacterial mercury methylation. *Science* *339*, 1332-1335.
- Payne, K.A., Fisher, K., Sjuts, H., Dunstan, M.S., Bellina, B., Johannissen, L., Barran, P., Hay, S., Rigby, S.E., and Leys, D. (2015a). Epoxyqueuosine reductase structure suggests a mechanism for cobalamin-dependent tRNA modification. *J Biol Chem* *290*, 27572-27581.
- Payne, K.A., Quezada, C.P., Fisher, K., Dunstan, M.S., Collins, F.A., Sjuts, H., Levy, C., Hay, S., Rigby, S.E., and Leys, D. (2015b). Reductive dehalogenase structure suggests a mechanism for B<sub>12</sub>-dependent dehalogenation. *Nature* *517*, 513-516.

- Pierre, S., Guillot, A., Benjdia, A., Sandstrom, C., Langella, P., and Berteau, O. (2012). Thiostrepton tryptophan methyltransferase expands the chemistry of radical SAM enzymes. *Nature chemical biology* 8, 957-959.
- Poppe, L., Bothe, H., Bröker, G., Buckel, W., Stupperich, E., and Rétey, J. (2000). Elucidation of the coenzyme binding mode of further B<sub>12</sub>-dependent enzymes using a base-off analogue of coenzyme B<sub>12</sub>, Vol 10.
- Poppe, L., Stupperich, E., Hull, W.E., Buckel, T., and Retey, J. (1997). A base-off analogue of coenzyme-B<sub>12</sub> with a modified nucleotide loop; <sup>1</sup>H-NMR structure analysis and kinetic studies with (*R*)-methylmalonyl-CoA mutase, glycerol dehydratase, and diol dehydratase. *European journal of biochemistry* 250, 303-307.
- Reeves, A.R., Brikun, I.A., Cernota, W.H., Leach, B.I., Gonzalez, M.C., and Weber, J.M. (2007). Engineering of the methylmalonyl-CoA metabolite node of *Saccharopolyspora erythraea* for increased erythromycin production. *Metabolic engineering* 9, 293-303.
- Renz, P. (1999). Biosynthesis of the 5,6-dimethylbenzimidazole moiety of cobalamin and of the other bases found in natural corrinoids. In *Chemistry and Biochemistry of B12*, R. Banerjee, ed. (New York: John Wiley & Sons, Inc.), pp. 557-566.
- Rodionov, D.A., Hebbeln, P., Eudes, A., ter Beek, J., Rodionova, I.A., Erkens, G.B., Slotboom, D.J., Gelfand, M.S., Osterman, A.L., Hanson, A.D., *et al.* (2009). A novel class of modular transporters for vitamins in prokaryotes. *Journal of Bacteriology* 191, 42.
- Rodionov, D.A., Vitreschak, A.G., Mironov, A.A., and Gelfand, M.S. (2003). Comparative genomics of the vitamin B<sub>12</sub> metabolism and regulation in prokaryotes. *J Biol Chem* 278, 41148-41159.
- Romine, M.F., Rodionov, D.A., Maezato, Y., Anderson, L.N., Nandhikonda, P., Rodionova, I.A., Carre, A., Li, X., Xu, C., Clauss, T.R., *et al.* (2017). Elucidation of roles for vitamin B<sub>12</sub> in regulation of folate, ubiquinone, and methionine metabolism. *Proceedings of the National Academy of Sciences of the United States of America* 114, E1205-e1214.
- Santos, J.A., Rempel, S., Mous, S.T.M., Pereira, C.T., ter Beek, J., de Gier, J.-W., Guskov, A., and Slotboom, D.J. (2018). Functional and structural characterization of an ECF-type ABC transporter for vitamin B<sub>12</sub>. *eLife* 7, e35828.
- Sato, K., Inukai, S., and Shimizu, S. (1974). Vitamin B<sub>12</sub>-dependent methionine synthesis in *Rhizobium meliloti*. *Biochemical and biophysical research communications* 60, 723-728.
- Savvi, S., Warner, D.F., Kana, B.D., McKinney, J.D., Mizrahi, V., and Dawes, S.S. (2008). Functional characterization of a vitamin B<sub>12</sub>-dependent methylmalonyl pathway in *Mycobacterium tuberculosis*: Implications for propionate metabolism during growth on fatty acids. *Journal of Bacteriology* 190, 3886.
- Scarlett, F.A., and Turner, J.M. (1976). Microbial metabolism of amino alcohols. Ethanolamine catabolism mediated by coenzyme B<sub>12</sub>-dependent ethanolamine ammonia-lyase in *Escherichia coli* and *Klebsiella aerogenes*. *Journal of general microbiology* 95, 173-176.
- Schneider, Z., and Stroinski, A. (1987). Comprehensive B<sub>12</sub>: Chemistry, Biochemistry, Nutrition, Ecology, Medicine. In, W.d. Gruyter, ed.
- Schrödinger Release 2015: MacroModel, S., LLC, New York, NY, 2015.
- Schrödinger Release 2015: Maestro, S., LLC, New York, NY, 2015.
- Seth, E.C., and Taga, M.E. (2014). Nutrient cross-feeding in the microbial world. *Frontiers in microbiology* 5, 350.

- Shelton, A.N., Seth, E.C., Mok, K.C., Han, A.W., Jackson, S.N., Haft, D.R., and Taga, M.E. (2019). Uneven distribution of cobamide biosynthesis and dependence in bacteria predicted by comparative genomics. *The ISME journal* **13**, 789-804.
- Shibata, N., Sueyoshi, Y., Higuchi, Y., and Toraya, T. (2018). Direct participation of a peripheral side chain of a corrin ring in coenzyme B<sub>12</sub> catalysis. *Angewandte Chemie (International ed in English)* **57**, 7830-7835.
- Srirangan, K., Akawi, L., Liu, X., Westbrook, A., Blondeel, E.J., Aucoin, M.G., Moo-Young, M., and Chou, C.P. (2013). Manipulating the sleeping beauty mutase operon for the production of 1-propanol in engineered *Escherichia coli*. *Biotechnology for biofuels* **6**, 139.
- Stupperich, E., Eisinger, H.J., and Krautler, B. (1988). Diversity of corrinoids in acetogenic bacteria. *p*-cresolylcobamide from *Sporomusa ovata*, 5-methoxy-6-methylbenzimidazolylcobamide from *Clostridium formicoaceticum* and vitamin B<sub>12</sub> from *Acetobacterium woodii*. *European journal of biochemistry* **172**, 459-464.
- Stupperich, E., and Konle, R. (1993). Corrinoid-dependent methyl transfer reactions are involved in methanol and 3,4-dimethoxybenzoate metabolism by *Sporomusa ovata*. *Appl Environ Microbiol* **59**, 3110-3116.
- Taga, M.E., and Walker, G.C. (2010). *Sinorhizobium meliloti* requires a cobalamin-dependent ribonucleotide reductase for symbiosis with its plant host. *Mol Plant Microbe Interact* **23**, 1643-1654.
- Tanioka, Y., Miyamoto, E., Yabuta, Y., Ohnishi, K., Fujita, T., Yamaji, R., Misono, H., Shigeoka, S., Nakano, Y., Inui, H., *et al.* (2010). Methyladeninylcobamide functions as the cofactor of methionine synthase in a cyanobacterium, *Spirulina platensis* NIES-39. *FEBS Lett* **584**, 3223-3226.
- Taoka, S., Padmakumar, R., Lai, M.T., Liu, H.W., and Banerjee, R. (1994). Inhibition of the human methylmalonyl-CoA mutase by various CoA-esters. *J Biol Chem* **269**, 31630-31634.
- Vrijbloed, J.W., Zerbe-Burkhardt, K., Ratnatilleke, A., Grubelnik-Leiser, A., and Robinson, J.A. (1999). Insertional inactivation of methylmalonyl coenzyme A (CoA) mutase and isobutyryl-CoA mutase genes in *Streptomyces cinnamonensis*: influence on polyketide antibiotic biosynthesis. *J Bacteriol* **181**, 5600-5605.
- Warner, J.R., and Copley, S.D. (2007). Pre-steady-state kinetic studies of the reductive dehalogenation catalyzed by tetrachlorohydroquinone dehalogenase. *Biochemistry* **46**, 13211-13222.
- Watanabe, F., Nakano, Y., and Stupperich, E. (1992). Different corrinoid specificities for cell growth and cobalamin uptake in *Euglena gracilis* Z, Vol 138.
- Werner, W.J., Allen, K.D., Hu, K., Helms, G.L., Chen, B.S., and Wang, S.C. (2011). In vitro phosphinate methylation by PhpK from *Kitasatospora phosalacinea*. *Biochemistry* **50**, 8986-8988.
- Widner, F.J., Lawrence, A.D., Deery, E., Heldt, D., Frank, S., Gruber, K., Wurst, K., Warren, M.J., and Krautler, B. (2016). Total synthesis, structure, and biological activity of adenosylrhodibalamin, the non-natural rhodium homologue of coenzyme B<sub>12</sub>. *Angewandte Chemie (International ed in English)* **55**, 11281-11286.
- Wolthers, K.R., Levy, C., Scrutton, N.S., and Leys, D. (2010). Large-scale domain dynamics and adenosylcobalamin reorientation orchestrate radical catalysis in ornithine 4,5-aminomutase. *J Biol Chem* **285**, 13942-13950.

- Yabuta, Y., Kamei, Y., Bito, T., Arima, J., Yoneda, K., Sakuraba, H., Ohshima, T., Nakano, Y., and Watanabe, F. (2015). Functional and structural characteristics of methylmalonyl-CoA mutase from *Pyrococcus horikoshii*. *Biosci Biotechnol Biochem* 79, 710-717.
- Yamanishi, M., Yunoki, M., Tobimatsu, T., Sato, H., Matsui, J., Dokiya, A., Iuchi, Y., Oe, K., Suto, K., Shibata, N., *et al.* (2002). The crystal structure of coenzyme B<sub>12</sub>-dependent glycerol dehydratase in complex with cobalamin and propane-1,2-diol. *European journal of biochemistry* 269, 4484-4494.
- Yan, J., Bi, M., Bourdon, A.K., Farmer, A.T., Wang, P.H., Molenda, O., Quaile, A.T., Jiang, N., Yang, Y., Yin, Y., *et al.* (2018). Purinyl-cobamide is a native prosthetic group of reductive dehalogenases. *Nature chemical biology* 14, 8-14.
- Yan, J., Ritalahti, K.M., Wagner, D.D., and Löffler, F.E. (2012). Unexpected specificity of interspecies cobamide transfer from *Geobacter* spp. to organohalide-respiring *Dehalococcoides mccartyi* strains. *Appl Environ Microbiol* 78, 6630-6636.
- Yan, J., Simsir, B., Farmer, A.T., Bi, M., Yang, Y., Campagna, S.R., and Löffler, F.E. (2016). The corrinoid cofactor of reductive dehalogenases affects dechlorination rates and extents in organohalide-respiring *Dehalococcoides mccartyi*. *The ISME journal* 10, 1092-1101.
- Yaneva, N., Schuster, J., Schäfer, F., Lede, V., Przybylski, D., Paproth, T., Harms, H., Müller, R.H., and Rohwerder, T. (2012). Bacterial acyl-CoA mutase specifically catalyzes coenzyme B<sub>12</sub>-dependent isomerization of 2-hydroxyisobutyryl-CoA and (S)-3-hydroxybutyryl-CoA. *J Biol Chem* 287, 15502-15511.
- Yi, S., Seth, E.C., Men, Y.J., Stabler, S.P., Allen, R.H., Alvarez-Cohen, L., and Taga, M.E. (2012). Versatility in corrinoid salvaging and remodeling pathways supports corrinoid-dependent metabolism in *Dehalococcoides mccartyi*. *Appl Environ Microbiol* 78, 7745-7752.
- Zhang, W., Yang, L., Jiang, W., Zhao, G., Yang, Y., and Chiao, J. (1999). Molecular analysis and heterologous expression of the gene encoding methylmalonyl-coenzyme A mutase from rifamycin SV-producing strain *Amycolatopsis mediterranei* U32. *Applied Biochemistry and Biotechnology* 82, 209-225.
- Zhang, Y., Rodionov, D.A., Gelfand, M.S., and Gladyshev, V.N. (2009). Comparative genomic analyses of nickel, cobalt and vitamin B<sub>12</sub> utilization. *BMC genomics* 10, 78.