Paraoxonase and acylated homoserine lactones in urine from patients with urinary tract infections

John Lafleur^{1*}, Richard L. Amdur²

¹Department Emergency Medicine, George Washington School of Medicine, Washington, DC, United States of America

²Department of Surgery, George Washington School of Medicine, Washington, DC, United States of America

*Corresponding author

johnlafleur@gmail.com

Abstract

Paraoxonases are mammalian enzymes that have a number of roles including the inhibition of bacterial virulence and biofilm formation by microorganisms that quorum sense with acylated homoserine lactones. Paraoxonases have previously been reported to inhibit *P. aeruginosa* biofilm formation in mammalian airways and skin. An innate immune role for paraoxonases in urinary tract infection has not previously been reported. We performed western blots for paraoxonase1 in urine from patients with urinary tract infection; we also tested urinary tract infection urine for the presence of acylated homoserine lactones using a cellular reporter system. We report here that paraoxonase1 was not found with our western blot assay in the urine of normal control patients; in those with urinary tract infection, paraoxonase1 was associated with *E. coli* UTI. Acylated homoserine lactones, but not paraoxonases, were found in the bulk urine of those with *P. aeruginosa* urinary tract infection. We hypothesize that paraoxonase may play a similar innate immune role in infected urine as has previously described in skin and airways.

Introduction

The paraoxonase PON family of mammalian lactonases are an evolutionarily conserved (1–3) innate immune mechanism that limit bacterial virulence and biofilm formation by degrading quorum sensing (QS) acylated homoserine lactones (AHLs) produced by some microorganisms.(4–13) These bacteria include *P. aeruginosa* as well as other environmental opportunists with large genomes and flexible lifestyles that are frequently found to be occult members of infecting biofilms.(14–16) Many of these belong to the group of non-fermenting gram negative bacilli (NFGNB).(17,18) UTI caused by NFGNB have been reported to more commonly infect those with urinary catheters, diabetes mellitus, and previous hospitalizations.(19) In multiple studies over the course of the last decade, PON have been shown to be protective against infection by *P. aeruginosa* biofilm formation in mammalian airways and skin cells.(9,20) Another body surface that is subject to environmental exposure is the urinary system, and PON has previously been reported in urine from healthy subjects.(21)

AHLs are known to be potent promoters of biofilm formation and virulence expression in gram negative pathogens that quorum sense with them, while at the same time causing tissue inflammation and derangement of host immunity.(22–26) Although they have not previously been reported, the presence of AHLs in urine from patients with UTI would support a role for PON in the defense against infection by *P. aeruginosa*.

Uncomplicated UTI in an immunocompetent host is characterized by single species infection, (27) *E. coli* 80% of the time. It may be hypothesized that the limited spectrum of uncomplicated UTI uropathogens is due to competent immunity, including the inhibition by PON of environmental opportunists that QS with AHLs, such as *P. aeruginosa* and many of the NFGNB.

To further explore these issues, we set out to assay for AHLs and PON in urine from patients with UTI presenting to the busy emergency department of a large city hospital. We hypothesized that 1) UTI patients will have PONS present, while non-UTI patients will not; 2) among UTI patients, the presence of PONS will be significantly associated with the presence of urinary pathogens

Materials and methods

Human subjects enrollment

Study protocol was reviewed/approved by Lifspan IRB

Study site: Anderson emergency department of Rhode Island Hospital

Inclusion criteria:

1. Greater than 18 years of age and able to give informed consent for study participation.
2. 10 or more white cells in urine analysis with symptoms of urinary tract infection.
3. Urine culture sent to the hospital microbiology department (prior to administration of antibiotics).

Control subjects: Emergency department patients with minor complaints unrelated to urinary system and without significant metabolic derangement such as fever, hyperglycemia, renal disease (acute or chronic), significant hypertension.

Urine samples: once enrolled in study subjects were asked to provide 50-100 ml of clean-catch urine in a sterile cup. This was immediately frozen at -80 for further study.

Growth media: Plates and broth were Luria-Bertani (LB).

Bacterial culture. The long chain HSL reporter strain *E. coli* JM109 (pSB1142) (carries *P. aeruginosa lasR* fused to luxCDABE), and *P. aeruginosa* PAO1 carrying PlasB-luxCDABE were grown in LB broth with shaking at 38 deg. C.

Reagents. 3-oxo-C12-HSL stock solution 20 mg/ml (Sigma-Aldrich) was diluted 1:50,000 in water. This dilution was arrived at empirically by testing against luminescence in the long chain HSL reporter strain *E. coli* JM109.

Western blotting. Urine samples from enrolled research subjects with UTI were stored at -80 deg. C., and thawed for use. 25 microliter samples of unprocessed urine were assayed for PON1 using the Bio-Rad iBlot system as previously described.

Antibodies. Primary antibody: polyclonal human PON1 from rabbit (Atlas Antibodies).

Secondary antibody: goat anti-rabbit reporter

Measures

Culture results were recoded into a binary variable (positive/negative). As a sensitivity analysis, we also coded those patients who were positive but with <50k cfu as negative. Positively skewed continuous variables and those with outliers were recoded into ordinal variables.

Data analysis

Associations between diagnosis and categorical variables were analyzed using chi-square or Fishers Exact Test. Comparison of continuous variables across groups was done using 2-tailed independent groups t-tests or the Kruskal-Wallis test for skewed variables. In order to test the independent association of PON1 with being culture-positive in UTI patients, we used multivariable logistic regression, adjusting for variables that might be confounds. These were defined as having an association with PON1 with $p < .10$. We also used a multivariable logistic

regression model to develop an optimal prediction model for being culture-positive in UTI patients. This was based on the patient variables that were associated with being culture-positive with $p < .10$, dropping any for which an odds ratio could not be calculated due to low sample size. SAS version 9.4 (Cary, NC) was used for data analysis, with $p < .05$ considered significant.

Results and discussion

For the 70 patients in the study, mean age was 60 ± 22 , 13 (19%) were black and 44 (63%) were white, and 48 (69%) were female. 11 (16%) were catheterized. Culture was positive in 39/61 cases (64%), while PON1 was positive in 22 cases (36%). There were 61 UTI patients and 9 controls in the sample.

Controls and UTI patients differed significantly on age, with UTI older, on serum creatinine (UTI higher), on highest temperature (UTI higher), and on lowest DBP (UTI lower), and there was a trend-level association for hemoglobin (UTI lower) (Table 1). Patients with UTI were older, and, probably for that reason, have higher average creatinine--due to age-related decline in kidney function (Table 1). Higher temperature in UTI subjects is likely due to some subjects being systemically ill.

Table 1. Patient variables by diagnosis (UTI vs control)

Patient variable	Control (n=9)	UTI (n=61)	p
Age	42 $\pm$ 16	63 $\pm$ 22	.009
Race			.45
Black	3 (33%)	10 (16%)	
White	5 (56%)	39 (64%)	
Other	1 (11%)	12 (20%)	
Gender female	4 (44%)	44 (72%)	.13

Catheterized	0	11 (18%)	.34
WBC	9.3 ± 4.4	11.4 ± 4.4	.25
Hemoglobin	13.7 ± 0.9	12.5 ± 1.9	.07 ^A
Serum creatinine	0.8 ± 0.1	1.2 ± 0.9	.042 ^A
Highest HR	85 ± 14	93 ± 21	.29
Highest temp	97.7 ± 0.6	99.0 ± 1.5	.0001
Lowest systolic bp	123 ± 16	118 ± 20	.40
Lowest diastolic bp	78 ± 11	68 ± 13	.044

^A using Kruskal-Wallis test.

PON1 was significantly associated with UTI diagnosis. Of the 61 UTI patients, 22 (36%) were

PON1 positive, while none of the controls were PON1 positive (Fisher Exact test p=.049).

PON1 was not significantly associated with any demographic or laboratory values (Table 2), but

was significantly associated with higher HR (higher in PON1 positive) (Table 2).

Table 2. Associations between patient variables and PON1 in patients with UTI

Patient variable	PON1 Neg (n=39)	PON1 Pos (n=22)	p
Age	62 ± 22	63 ± 21	.82
Race			.60
Black	5 (13%)	5 (23%)	
White	26 (67%)	13 (59%)	
Other	8 (21%)	4 (18%)	
Gender female	29 (74%)	15 (68%)	.61
Catheterized	5 (13%)	6 (27%)	.18
WBC	11.2 ± 4.4	11.7 ± 4.6	.68
Hemoglobin	12.5 ± 2.0	12.4 ± 1.7	.75
Serum creatinine	1.2 ± 1.0	1.2 ± 0.5	.48 ^A
Highest HR	88 ± 16	101 ± 25	.03
Highest temp	99.0 ± 1.6	98.9 ± 1.3	.66
Lowest systolic bp	120 ± 20	114 ± 20	.27
Lowest diastolic bp	68 ± 12	70 ± 13	.45

PON1 was significantly associated with positive culture in UTI patients: PON was positive in 4/22 with negative culture (18%) versus 18/39 with positive culture (46%; $p=.03$; Table 3). We did a sensitivity analysis coding patients who had culture $< 50k$ cfu as negative (rather than the default coding as positive), and found that the association was still significant (culture negative had 23% PON positive, culture positive had 48% PON positive, $p=.04$). Thus, in UTI patients, presence of PON in urine was associated with urine culture growing out a urinary pathogen, in contrast to urogenital flora, or no growth. However, in the sample of UTI patients, after adjusting for higher HR, the association between PON1 and culture was no longer significant (an OR for PON1 Pos vs Neg was 3.08 [95% ci 0.84-11.20], $p=.09$).

Table 3. Association of Positive Culture with PON1 and other patient variables, in patients with UTI. N and column-% are shown.

Culture	Culture Neg (n=22)	Culture Pos (n=39)	p
PON1 Positive	4 (18%)	18 (46%)	.03
Age	60 $\pm$ 24	64 $\pm$ 20	.45
Race			.17
Black	1 (5%)	9 (23%)	
White	16 (73%)	23 (59%)	
Other	5 (23%)	7 (18%)	
Gender female	16 (73%)	28 (72%)	.94
Catheterized	1 (5%)	10 (26%)	.045
WBC	9.8 $\pm$ 3.5	12.2 $\pm$ 4.6	.046
Hemoglobin	12.7 $\pm$ 1.7	12.3 $\pm$ 2.0	.44
Serum creatinine	1.1 $\pm$ 0.6	1.2 $\pm$ 1.0	.44
Highest HR	86 $\pm$ 16	97 $\pm$ 22	.064
Highest temp	98.8 $\pm$ 1.4	99.0 $\pm$ 1.5	.61
Lowest systolic bp	119 $\pm$ 19	116 $\pm$ 21	.58
Lowest diastolic bp	71 $\pm$ 9	67 $\pm$ 14	.23

In addition to PON1, other patient variables that were associated with being culture-positive, in UTI patients, included WBC (higher with culture positive), and being catheterized (more frequent for culture positive). Highest HR was marginally associated with culture-positive (Table 3). We created an optimal prediction model for being culture-positive, which included PON1, highest HR, and WBC (being catheterized was dropped because OR could not be calculated for this variable due to small sample size), which had an area under the ROC curve of 0.72 for predicting culture-positive.

Using the equation: $\text{risk} = -2.43 + 1.007 \cdot \text{PON1} + .014 \cdot \text{highestHR} + .138 \cdot \text{WBC}$, and then $\text{probability} = \exp(\text{risk}) / (1 + \exp(\text{risk}))$, and then splitting the probabilities into tertiles, we found that the observed incidence of being culture positive in tertiles 1 through 3, respectively, were 39%, 75%, and 83% ($p=.01$).

19 out of 38 positive urine cultures grew out *E. coli* alone (50%) . PON1 was positive in 10 of these (53%). When compared to cultures that grew out multiple organisms (including those with ‘urogenital flora’) PON was significantly associated with cultures that grew out *E. coli* alone, $P=0.05$ (Table 4). Four NFGNB (10%) were cultured from PON negative urines (all of them *P. aeruginosa*). In addition, three other gram-negative environmental opportunists were cultured from PON negative urines: *Serratia marscesens*, *Citrobacter freundii*, and *Klebsiella pneumoniae*. These organisms are lactose fermenters, so do not meet criteria for being NFGNB, however, like NFGNB, they are multi-drug resistant environmental opportunists. Additionally these bacteria have all been reported to produce or QS with AHLs.(28–30) Among PON positive urines there were no NFGNB in culture.

Table 4. Univariable association of presence of PON in urine with *E. coli* alone versus cultures growing out multiple different bacteria (including ‘urogenital flora’)

	E. coli alone (n=21)	Mult. organisms. (n=25)
PON pos.	10 (53%)	6 (24%)
PON neg.	9 (47%)	19 (76%)

^A Chi-square 5.3; P=0.05

Using an *E. coli* luminescent reporter construct, C12 AHL was also assayed for in urine samples. C12 AHLs were only seen in the urines that were culture positive for *P. aeruginosa*. Concentrations of C12 AHL in one sample (patient #9) was about 1.5 micromolar. The other three samples in which *P. aeruginosa* grew out of culture had considerably lower concentrations (see table 5). Biologically relevant concentrations of AHLs for QS are considered to be 1-5 micromolar.(25) One possible interpretation of concentrations of C12 considerably below this in three of four samples suggests that QS and virulence expression in *P. aeruginosa* UTI is not a planktonic phenomenon in the urine but occurring on mucosal surfaces of the bladder/urinary system where surface colonization/biofilm formation can take place, and immune interactions are likely to take place.

Table 5. Subjects whose cultures grew out *P. aeruginosa* had urine that contained C12 AHL (except for subject #61 where it was not detectable).

Subject #	Luminescence (ALU)	Micromolar conc.
9	167,000	1.5
25	22,000	0.2
35	8,000	0.07
61	0	0

These preliminary results indicate a positive association between PON level and positive culture in patients with UTI, although HR may be functioning as a confounding factor. Levels of C12 AHL in planktonic cultures necessary to initiate QS-related *lasB* expression have previously been reported to be about 1 micromolar(31), a concentration of C12 AHLs that is not uncommonly seen in planktonic cultures of *P. aeruginosa*. By contrast, C12 AHL levels associated with *P. aeruginosa* biofilms in flow cells have been found to be hundreds of times higher.(32) C12 AHL levels in urine from subjects with *P. aeruginosa* UTI have not previously been reported, though detection in urine of non-AHL *P. aeruginosa* mediators of QS associated with pulmonary infection has recently been reported.(33) In the current study results are somewhat equivocal as 3 out of 4 urines were found to contain concentrations of C12 AHLs significantly below the 1 micromolar threshold. There are two possible scenarios that may be imagined in the case of *P. aeruginosa* UTI—that QS occurs mostly on bladder mucosa surfaces where concentrations of microorganisms are likely to be higher, and interactions with mediators of host immunity more intense, or that QS is also occurring in bulk urine. Host bladder epithelial response to UTI include urination, exocytosis of intracellular urinary pathogens, and sloughing of bladder epithelial cells with adherent or intracellular urinary pathogens.(34) Since most planktonic *P. aeruginosa* in UTI will be flushed with urination, possible adaptive advantages conferred by QS among planktonic population of *P. aeruginosa* are that production of virulence factors, such as C12 AHL, may be ramped up in the *P. aeruginosa* population as a whole if QS is occurring in bulk urine. The results of the current study are preliminary, but since 3 of 4 samples showed what appear to be significantly sub-threshold levels of C12 AHL, QS at mucosal surfaces seems more likely. Urine has recently been reported to independently promote *P.*

aeruginosa biofilm formation,(35) suggesting that in the absence of the normal QS-mediated mechanisms for biofilm formation, a biofilm may still be formed in *P. aeruginosa* UTI.

We found in the current study that the presence of PON predicts growth of uropathogens as opposed to “urogenital flora” or, “no growth”. The latter two are results of urine culture which are not considered to represent significant infections, but, rather, some source of inflammation resulting in urinary symptoms mimicking acute UTI. One possible explanation for this is that UTI-mediated upregulation of urinary system TLR4/5 may result in the increases expression of PON seen in our PON positive UTI subjects. Urinary TLR signaling has been found to be sensitive to uropathogens,(36,37) resulting in activation of NF-kB and the expression of the pro-inflammatory genes IL-6 and IL-8 with consequent ingress to the bladder mucosa of neutrophils.(38) There is no report in the literature of TLR4/5 mobilization of PON, so at present this remains conjecture. By combining information on PON1 level, higher HR, and WBC, it was possible to obtain an accurate estimate of the probability of having positive cultures.

Our findings include that PON positive subjects had significantly more UTIs caused by *E. coli* alone, rather than multi-species infections, or infections with opportunists such as *P. aeruginosa*. It has previously been reported that the large majority of uncomplicated UTIs in normal hosts are caused by single species—meaning that under normal conditions community richness in UTI is very limited compared to other human microbiomes.(27) Infection associated with impaired immunity is characterized by difficult to eradicate biofilms, polymicrobial infections, and infection with opportunistic organisms that don’t readily infect immune-competent hosts. Viewed in this light, PON positive patients more nearly conform to immune-competent patients with UTIs caused by a single pathogen, while PON negative patients were more likely to have UTIs more characteristic of deranged immune competence. The current study does not make it

possible to draw any causal link between the presence of PON in urine and the immune-competent pattern of uncomplicated UTI caused by a single pathogen, mostly *E. coli*. However, the role of PON in innate immunity of the airway and skin,(9,20) and the role that AHL QS plays in many NFGNB environmental opportunists, along with possible occult roles played by fastidious environmental opportunists in establishing complex multi-species infections,(39) suggests that urinary PON may have a protective role.

Conclusion

The current study reports for the first time AHLs in the urine of subjects with *P. aeruginosa* UTIs. However, the significance of this, and the role that AHLs play in QS among planktonic *P. aeruginosa* remains to be investigated. We found that UTI subjects with PON positive urines were much more likely to have uncomplicated *E. coli* UTI. What mechanism, if any, underlies this finding is at present unclear.

References

1. Bar-Rogovsky H, Hugematter A, Tawfik DS. The evolutionary origins of detoxifying enzymes: the mammalian serum paraoxonases (PONs) relate to bacterial homoserine lactonases. *J Biol Chem* [Internet]. 2013 Aug 16 [cited 2014 Dec 3];288(33):23914–27. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3745338&tool=pmcentrez&rendertype=abstract>
2. Elias M, Tawfik DS. Divergence and convergence in enzyme evolution: parallel evolution of paraoxonases from quorum-quenching lactonases. *J Biol Chem* [Internet]. 2012 Jan 2 [cited 2013 Feb 28];287(1):11–20. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3249062&tool=pmcentrez&rendertype=abstract>
3. Khersonsky O, Tawfik DS. Structure-reactivity studies of serum paraoxonase PON1 suggest that its native activity is lactonase. *Biochemistry*. 2005 Apr 26;44(16):6371–82.
4. Chun CK, Ozer EA, Welsh MJ, Zabner J, Greenberg EP. Inactivation of a *Pseudomonas aeruginosa* quorum-sensing signal by human airway epithelia. *Proc Natl Acad Sci U S A*. 2004 Mar 9;101(10):3587–90.
5. Ozer E a, Pezzulo A, Shih DM, Chun C, Furlong C, Lusis AJ, et al. Human and murine paraoxonase 1 are host modulators of *Pseudomonas aeruginosa* quorum-sensing. *FEMS Microbiol Lett* [Internet]. 2005 Dec 1 [cited 2014 Oct 16];253(1):29–37. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/16260097>
6. Manolescu B. Paraoxonases as Protective Agents Against N-Acyl Homoserine Lactone–Producing Pathogenic Microorganisms. *Maedica (Buchar)* [Internet]. 2013 [cited 2014 Oct 16];8(1):49–52. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3749762/>
7. Stoltz DA, Ozer EA, Taft PJ, Barry M, Liu L, Kiss PJ, et al. *Drosophila* are protected from *Pseudomonas aeruginosa* lethality by transgenic expression of paraoxonase-1. *J Clin Invest*. 2008 Sep;118(9):3123–31.
8. Stoltz DA, Ozer EA, Ng CJ, Janet MY, Reddy ST, Lusis AJ, et al. Paraoxonase-2 deficiency enhances *Pseudomonas aeruginosa* quorum sensing in murine tracheal epithelia. *Am J Physiol Cell Mol Physiol*. 2007;292(4):L852–60.
9. Simanski M, Babucke S, Eberl L, Harder J. Paraoxonase 2 acts as a quorum sensing-quenching factor in human keratinocytes. *J Invest Dermatol* [Internet]. 2012 Sep [cited 2013 Feb 13];132(9):2296–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22534874>
10. Devarajan A, Bourquard N, Grijalva VR, Gao F, Ganapathy E, Verma J, et al. Role of PON2 in innate immune response in an acute infection model. *Mol Genet Metab*

- 276 [Internet]. 2013 Nov [cited 2014 Oct 18];110(3):362–70. Available from:
277 <http://www.ncbi.nlm.nih.gov/pubmed/23911207>
- 278 11. Schweikert E-M, Amort J, Wilgenbus P, Förstermann U, Teiber JF, Horke S.
279 Paraoxonases-2 and -3 Are Important Defense Enzymes against *Pseudomonas aeruginosa*
280 Virulence Factors due to Their Anti-Oxidative and Anti-Inflammatory Properties. *J Lipids*
281 [Internet]. 2012 Jan [cited 2013 Feb 13];2012:352857. Available from:
282 [http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3335252&tool=pmcentrez&re](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3335252&tool=pmcentrez&rendertype=abstract)
283 [ndertype=abstract](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3335252&tool=pmcentrez&rendertype=abstract)
- 284 12. Amara N, Krom BP, Kaufmann GF, Meijler MM. Macromolecular inhibition of quorum
285 sensing: enzymes, antibodies, and beyond. *Chem Rev*. 2011 Jan 12;111(1):195–208.
- 286 13. Gowda LK, Marie MAM. Role of quorum-sensing molecules in infections caused by
287 Gram-negative bacteria and host cell response. *Rev Med Microbiol* [Internet]. 2014 Jul
288 [cited 2014 Dec 3];25(3):66–70. Available from:
289 [http://content.wkhealth.com/linkback/openurl?sid=WKPTLP:landingpage&an=00013542-](http://content.wkhealth.com/linkback/openurl?sid=WKPTLP:landingpage&an=00013542-201407000-00002)
290 [201407000-00002](http://content.wkhealth.com/linkback/openurl?sid=WKPTLP:landingpage&an=00013542-201407000-00002)
- 291 14. Dubern J-F, Cigana C, De Simone M, Lazenby J, Juhas M, Schwager S, et al. Integrated
292 whole genome screening for *Pseudomonas aeruginosa* virulence genes using multiple
293 disease models reveals that pathogenicity is host specific. *Environ Microbiol* [Internet].
294 2015;n/a-n/a. Available from: <http://doi.wiley.com/10.1111/1462-2920.12863>
- 295 15. Tümmler B, Wiehlmann L, Klockgether J, Cramer N. Advances in understanding
296 *Pseudomonas*. *F1000Prime Rep* [Internet]. 2014;6(February):9. Available from:
297 </pmc/articles/PMC3913036/?report=abstract>
- 298 16. Davenport P, Griffin JL, Welch M. Quorum sensing is accompanied by global metabolic
299 changes in the opportunistic human pathogen, *Pseudomonas aeruginosa*. *J*
300 *Bacteriol* [Internet]. 2015;197(12):JB.02557-14. Available from:
301 <http://jb.asm.org/lookup/doi/10.1128/JB.02557-14>
- 302 17. Vincenti S, Quaranta G, De Meo C, Bruno S, Ficarra MG, Carovillano S, et al. Non-
303 fermentative gram-negative bacteria in hospital tap water and water used for
304 haemodialysis and bronchoscope flushing: Prevalence and distribution of antibiotic
305 resistant strains. *Sci Total Environ* [Internet]. 2014;499:47–54. Available from:
306 <http://linkinghub.elsevier.com/retrieve/pii/S004896971401208X>
- 307 18. Brooke JS. *Stenotrophomonas maltophilia*: An emerging global opportunistic pathogen.
308 *Clin Microbiol Rev*. 2012;25:2–41.
- 309 19. Solanki R, Dawar R. Nonfermenting Gram-Negative Bacilli and Urinary Tract Infection -
310 Sorting the Mystery of Infections Caused. *J Med Microbiol Diagnosis* [Internet].
311 2015;04(04):2–5. Available from: [http://www.omicsonline.org/open-](http://www.omicsonline.org/open-access/nonfermenting-gramnegative-bacilli-and-urinary-tract-infection-sortingthe-mystery-of-infections-caused-2161-0703-1000210.php?aid=65584)
312 [access/nonfermenting-gramnegative-bacilli-and-urinary-tract-infection-sortingthe-](http://www.omicsonline.org/open-access/nonfermenting-gramnegative-bacilli-and-urinary-tract-infection-sortingthe-mystery-of-infections-caused-2161-0703-1000210.php?aid=65584)
313 [mystery-of-infections-caused-2161-0703-1000210.php?aid=65584](http://www.omicsonline.org/open-access/nonfermenting-gramnegative-bacilli-and-urinary-tract-infection-sortingthe-mystery-of-infections-caused-2161-0703-1000210.php?aid=65584)
- 314 20. Stoltz DA, Ozer EA, Zabner J. paraoxonases, quorum sensing, and pseudomonas

- 315 aeruginosa.pdf. The Paraoxonases: Their Role in Disease Development and Xenobiotic
316 Metabolism. Springer; 2008. 307-319 p.
- 317 21. Lescuyer P, Pernin A, Hainard A, Bigeire C, Burgess JA, Zimmermann-Ivol C, et al.
318 Proteomic analysis of a podocyte vesicle-enriched fraction from human normal and
319 pathological urine samples. *ProteomicsClinical Appl*. 2008 Jul;2(7–8):1008–18.
- 320 22. Boontham P, Robins A, Chandran P, Pritchard D, Camara M, Williams P, et al.
321 Significant immunomodulatory effects of *Pseudomonas aeruginosa* quorum-sensing signal
322 molecules: possible link in human sepsis. *Clin Sci (Lond)*. 2008 Dec;115(11):343–51.
- 323 23. Hartmann A, Schikora A. Quorum sensing of bacteria and trans-kingdom interactions of
324 N-acyl homoserine lactones with eukaryotes. *J Chem Ecol [Internet]*. 2012 Jun [cited 2013
325 Feb 12];38(6):704–13. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22648507>
- 326 24. Li Y, Zhou H, Zhang Y, Chen C, Huang B, Qu P, et al. N-3-(oxododecanoyl)-L-
327 homoserine lactone promotes the induction of regulatory T-cells by preventing human
328 dendritic cell maturation. *Exp Biol Med [Internet]*. 2015;240:896–903. Available from:
329 <http://ebm.sagepub.com/lookup/doi/10.1177/1535370214564742>
- 330 25. Kravchenko V V., Kaufmann GF, Mathison JC, Scott D a., Katz AZ, Wood MR, et al. N-
331 (3-oxo-acyl)homoserine lactones signal cell activation through a mechanism distinct from
332 the canonical pathogen-associated molecular pattern recognition receptor pathways. *J Biol*
333 *Chem*. 2006;281(39):28822–30.
- 334 26. Kravchenko V, Garner AL, Mathison J, Seit-Nebi A, Yu J, Gileva IP, et al. Facilitating
335 Cytokine-Mediated Cancer Cell Death by Proteobacterial N -Acylhomoserine Lactones.
336 *ACS Chem Biol [Internet]*. 2013;8:1117–20. Available from:
337 <http://dx.doi.org/10.1021/cb4000184>
- 338 27. Willner D, Low S, Steen J a., George N, Nimmo GR, Schembri M a., et al. Single clinical
339 isolates from acute uncomplicated urinary tract infections are representative of dominant
340 In Situ populations. *MBio*. 2014;5(2):1–10.
- 341 28. Annapoorani A, Karim A, Abdul K. Inhibition of Quorum Sensing Mediated Virulence
342 Factors Production in Urinary Pathogen *Serratia marcescens* PS1 by Marine Sponges.
343 2012;52(June):160–6.
- 344 29. Hentzer M, Givskov M. Pharmacological inhibition of quorum sensing for the treatment
345 of chronic bacterial infections. *J Clin Invest*. 2003 Nov;112(9):1300–7.
- 346 30. Yin W, Purmal K, Chin S, Chan X, Koh C, Sam C, et al. N-Acyl Homoserine Lactone
347 Production by *Klebsiella pneumoniae* Isolated from Human Tongue Surface.
348 2012;4:3472–83.
- 349 31. Pearson JP, Passador L, Iglewski BH, Greenberg EP. A second N-acylhomoserine lactone
350 signal produced by *Pseudomonas aeruginosa*. *Proc Natl Acad Sci U S A*.
351 1995;92(February):1490–4.
- 352 32. Charlton TS, de Nys R, Netting a, Kumar N, Hentzer M, Givskov M, et al. A novel and

sensitive method for the quantification of N-3-oxoacyl homoserine lactones using gas chromatography-mass spectrometry: application to a model bacterial biofilm. *Environ Microbiol.* 2000;2:530–41.

33. Barr HL, Halliday N, Camara M, Barrett D a., Williams P, Forrester DL, et al. *Pseudomonas aeruginosa* quorum sensing molecules correlate with clinical status in cystic fibrosis. *Eur Respir J* [Internet]. 2015;1–9. Available from: <http://erj.ersjournals.com/cgi/doi/10.1183/09031936.00225214>
34. Greene SE, Hibbing ME, Janetka J, Chen SL, Hultgren J. Human Urine Decreases Function and Expression of Type 1 Pili in Uropathogenic *Escherichia coli*. *MBio.* 2015;6(4):1–9.
35. Cole SJ, Records AR, Orr MW, Linden SB, Lee VT. Catheter-associated urinary tract infection by *Pseudomonas aeruginosa* is mediated by exopolysaccharide-independent biofilms. *Infect Immun.* 2014;82(5):2048–58.
36. Ragnarsdóttir B, Jönsson K, Urbano A, Grönberg-Hernandez J, Lutay N, Tammi M, et al. Toll-like receptor 4 promoter polymorphisms: common TLR4 variants may protect against severe urinary tract infection. *PLoS One* [Internet]. 2010 Jan [cited 2014 Oct 18];5(5):e10734. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2873976&tool=pmcentrez&rendertype=abstract>
37. Ingersoll M a, Albert ML. From infection to immunotherapy: host immune responses to bacteria at the bladder mucosa. *Mucosal Immunol* [Internet]. 2013 Nov [cited 2014 Oct 20];6(6):1041–53. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24064671>
38. Nielubowicz GR, Mobley HLT. Host-pathogen interactions in urinary tract infection. *Nat Rev Urol* [Internet]. 2010;7(8):430–41. Available from: <http://dx.doi.org/10.1038/nrurrol.2010.101>
39. Zhang L, Gowardman J, Morrison M, Krause L, Playford EG, Rickard CM. Molecular investigation of bacterial communities on intravascular catheters: No longer just *Staphylococcus*. *Eur J Clin Microbiol Infect Dis.* 2014;33:1189–98.