A light-dependent molecular link between competition cues and defense responses in plants

3

4 G. L. Fernández-Milmanda¹, C. D. Crocco¹, M. Reichelt², C. A. Mazza¹, T. G. Köllner², T. Zhang^{3,5},
5 M. D. Cargnel¹, M. Z. Lichy¹, A. J. Koo³, A. T. Austin¹, J. Gershenzon², C. L. Ballaré^{1,4†}

6

7 ¹IFEVA, Consejo Nacional de Investigaciones Científicas y Técnicas – Universidad de Buenos Aires, Ave.
8 San Martín 4453, C1417DSE, Buenos Aires, Argentina

9 ²Max Planck Institute for Chemical Ecology, Hans-Knöll-Str. 8, 07745 Jena, Germany

10 ³Department of Biochemistry, University of Missouri, Columbia 65211, USA

11 ⁴IIBIO, Consejo Nacional de Investigaciones Científicas y Técnicas–Universidad Nacional de San Martín,
12 B1650HMP Buenos Aires, Argentina

13 ⁵Current address: College of Agriculture, South China Agricultural University, Guangdong, China

14

15 † Corresponding author. Email: ballare@ifeva.edu.ar

16

17

18 **ABSTRACT**

19 **One of the principal internal signals controlling plant growth and defense is jasmonate (JA), a**
 20 **potent growth inhibitor that is simultaneously a central regulator of plant immunity to**
 21 **herbivores and pathogens. When shade-intolerant plants perceive the proximity of**
 22 **competitors using the photoreceptor phytochrome B (phyB), they accelerate growth and**
 23 **down-regulate JA responses. However, the mechanisms by which photoreceptors relay light**
 24 **cues to the JA signaling pathway are not understood. Here we identify a sulfotransferase**
 25 **(ST2a) that is strongly up-regulated by plant proximity perceived by phyB via the phyB-**
 26 **Phytochrome Interacting Factor (PIF) signaling module. By catalyzing the formation of a**
 27 **sulfated JA derivative, ST2a acts to degrade bioactive forms of JA and represents a direct**
 28 **molecular link between photoreceptors and hormone signaling in plants. The enzyme**
 29 **provides a molecular mechanism for prioritizing shade avoidance over defense under close**
 30 **plant competition.**

31

32 **RESULTS and DISCUSSION**

33 Growth responses to competition with other plants (1) and defense responses to the attack of
 34 consumer organisms (2) are two paradigmatic examples of adaptive phenotypic plasticity in
 35 plants. However, the mechanistic and functional links between these responses are not well
 36 understood. JAs are potent growth inhibitors (3) and regulators of cell division (4, 5), and their
 37 role in balancing growth and defense is evolutionarily conserved in land plants, from
 38 bryophytes (6) to angiosperms (7). When shade-intolerant plants perceive a high risk of
 39 competition for light with neighboring individuals, they activate the shade-avoidance
 40 syndrome (SAS), which allows them to position their leaves in well-illuminated areas of the
 41 canopy. Under these competitive conditions, plants also often attenuate the expression of JA-
 42 mediated defense responses against pathogens and herbivore (8). This attenuation of defense
 43 presumably allows the plant to efficiently focus its resources and developmental decisions on
 44 escaping shade, sacrificing plant parts that are unlikely to contribute to resource capture.

45 Plants perceive the proximity of competitors using photoreceptors. Low ratios of red (R) to far-
 46 red (FR) radiation (R:FR ratio), which indicate a high risk of competition, result in partial
 47 inactivation of the photoreceptor phyB, which in turn promotes growth-related hormonal
 48 pathways (9), and attenuates signaling mechanisms involved in the activation of defense
 49 responses, such as the JA and salicylic acid signaling pathways (8). The attenuation of defense

50 responses triggered by supplemental FR radiation or mutations in the *PHYB* gene is not merely
51 a consequence of redirecting resources to growth (i.e., a simple reflection of an energetic
52 tradeoff between growth and defense)(10, 11). Attenuation of JA responses under low R:FR
53 ratios has been associated with FR-induced changes in the balance between DELLA and JA-ZIM-
54 domain (JAZ) repressor proteins (12, 13), and decreased stability of key transcription factors,
55 such as MYC2 (13). However, the specific molecular links between the phyB and JA signaling
56 pathways have not been demonstrated.

57 Most JA-upregulated genes and metabolites are down-regulated by low R:FR ratios or phyB
58 inactivation (8, 11). In order to identify signaling elements that could be involved in the
59 suppression of defense in plants exposed to competition cues, we searched for genes with an
60 inverse pattern of regulation (i.e., JA-related genes showing increased expression under
61 conditions that inactivate phyB). In a microarray experiment, we found a small cluster of
62 approximately 100 genes that were positively regulated by JA whose expression was promoted
63 by phyB inactivation; this cluster included various *JAZ* genes, and a gene of the
64 sulfotransferase family annotated as *ST2a/SOT15* (Fig. S1A). Moreover, in an analysis of
65 expression patterns of JA-related genes using publicly-available microarray databases, we
66 discovered that this *ST2a* gene was consistently and strongly upregulated by low R:FR ratios
67 (Fig. S1B). Experimentally, we confirmed a strong upregulation of *ST2a* mRNA by FR radiation
68 treatments that mimicked the effects of plant proximity and mild suppression by UV-B
69 radiation (Fig. S1C), suggesting that this gene could indeed be involved in relaying information
70 on the canopy light conditions to the JA signaling pathway. Chromatin immunoprecipitation
71 sequencing (ChIP-seq) results indicate that *ST2a* is among the direct targets of Phytochrome
72 Interacting Factors (PIFs) (14). PIFs are growth-promoting transcription factors (15), and PIF4,
73 PIF5, and PIF7 have been shown to link phyB inactivation with growth responses to shade
74 signals (16, 17). We tested a *pif4 pif5 pif7* triple knock out mutant (18) and found that *ST2a*
75 mRNA was upregulated by tissue damage just like in Col-0, but the transcriptional response of
76 *ST2a* to supplemental FR radiation was completely lost (Fig. S1D). These results demonstrate
77 that low R:FR ratios upregulate the transcription of *ST2a* via the phyB-PIF transcription
78 module.

79 *ST2a* belongs to a family of 21 sulfotransferase-encoding genes in Arabidopsis (19, 20), and
80 shows sequence similarity to proteins in many dicotyledonous species (Fig. S2). In vitro, the
81 *ST2a* protein catalyzes the sulfation of 12-hydroxy JA (OH-JA) to form JA sulfate (HSO₄-JA) (21).
82 First described in the late 1800s (22), sulfation consists of the transfer of a sulfate residue to a

hydroxyl or amino group. In mammalian systems, sulfation represents an important pathway for the biotransformation of hormones, neurotransmitters, and numerous xenobiotics (23, 24), which in most cases results in increased water solubility and decreased biological activity. Based largely on their activities and substrate specificities in vitro, sulfotransferases have also been proposed to be important in the regulation of hormonal signaling in plants (19, 20). For example, in Arabidopsis, a tyrosylprotein-sulfotransferase was reported to regulate the activity of hormone peptides involved in the control of cell proliferation (25). However, there is very little evidence from functional genetic studies that sulfotransferases are involved in the adaptive modulation of phytohormonal pathways, and no information about their ecological role in the regulation of plant responses to environmental cues.

We reasoned that a plausible physiological role for the activation of *ST2a* transcription via the phyB-PIF module could be the attenuation of JA-mediated responses in plants exposed to competition signals (low R:FR ratios). To test this hypothesis, we treated Arabidopsis plants with methyl JA (MeJA) and evaluated the formation of JA-related metabolites and accompanying changes in the expression of genes related to JA metabolism. Plants were kept under either white light (Amb light treatment) or white light supplemented with FR radiation (FR treatment). MeJA treatment induced rapid (≤ 30 min) increases in concentrations of JA and the bioactive conjugate JA-Ile, which were followed by increases in further metabolites, including OH-JA, OH-JA-Ile, and later-on by a marked increase in COOH-JA-Ile and HSO₄-JA (Fig. 1). Addition of FR to ambient light significantly decreased the abundance of JA and JA-Ile, and their oxidized derivatives. In contrast, the pool of HSO₄-JA was significantly increased by supplemental FR. The increase in HSO₄-JA concentration under FR correlated well with a dramatic increase in *ST2a* mRNA level (Fig. S3). Genes encoding other enzymes involved in the metabolism of JAs, such as *IAR3*, *ILL6*, and *CYP94B3* were also up-regulated by FR but to a much lesser extent than *ST2a* (Fig. 1, Fig. S3). In summary, low R:FR ratios decreased the pools of JA and JA-Ile, and this effect coincided with a massive up-regulation of *ST2a* gene expression.

In the field, shade treatments, compared to full sunlight, have been reported to attenuate the JA burst induced by insect herbivory, which correlates with attenuated production of defense metabolites (26). Are the effects of shade on JA accumulation mediated by phyB inactivation and functionally connected with *ST2a* gene expression? To address this question, we first isolated and characterized two *ST2a* null alleles (*st2a-1* and *st2a-2*), and demonstrated that both knock-out mutants produced only trace levels of HSO₄-JA (Fig. S4). The mutant carrying

the *st2a-1* allele was used for further functional characterization. In Col-0 plants, supplemental FR reduced the JA burst induced by wounding, and this effect correlated with an increase in the pool of HSO₄-JA (Fig. 2). Importantly, in *st2a-1* plants, the effect of FR attenuating the JA response burst was completely missing (Fig. 2). OH-JA was reduced by FR in Col-0, and it was more abundant in *st2a-1* than in wild type plants (Fig. S5). Within the resolution of our sampling, the levels of JA-Ile were very low and variable, and most of the JA-Ile conjugates were present in the carboxylated form (COOH-JA-Ile) 4 h after wounding. The concentrations of the sum of JA-Ile conjugates was significantly lower in FR plants than in plants of the ambient light treatment (Fig. S5), and significantly higher in *st2a-1* than in Col-0 plants at 4 h. Overaccumulation of COOH-JA-Ile has also been reported in lines lacking JOX/JAOs (27), the enzymes responsible for generating OH-JA (the putative substrate of ST2a). Thus, genetic ablation of *JAO/JOX* or *ST2a* results in increased flux through JA-Ile metabolism and catabolite accumulation. Because *ST2a* transcription in response to FR supplementation was minimal in the *pif4 pif5 pif7* triple mutant (Fig. S1D), we used this mutant as a complementary genetic tool to manipulate *ST2a* mRNA levels. Plants of *pif4 pif5 pif7* accumulated HSO₄-JA in response to wounding, but failed to do so in response to supplemental FR (Fig. S6A). Furthermore, when *ST2a* expression was physiologically manipulated in Col-0 plants using various combinations of FR and wounding treatments, the variation in the HSO₄-JA content at 4 h after wounding was largely explained by the variation in the levels of *ST2a* mRNA detected by qPCR (Fig. S6B). Finally, when we overexpressed *ST2a* under a strong promoter, we obtained JA metabolite profiles that were qualitatively similar to those generated in response to FR radiation (HSO₄-JA was greatly increased, with a concomitant reduction in OH-JA, JA, and oxidized JA-Ile conjugates; Fig S7). Neither wounding nor supplemental FR radiation affected the abundance of transcripts of *ST2b*, a gene closely related to *ST2a* (Fig. S2), and a *st2b* null mutant showed normal levels of HSO₄-JA (Fig. S8). These results provide compelling evidence that ST2a is the sole sulfotransferase responsible for the generation of HSO₄-JA in Arabidopsis, and that the increased sulfation of OH-JA under low R:FR ratios is functionally connected with the transcriptional upregulation of the *ST2a* gene mediated by the phyB-PIF module. Pharmacological experiments indicate that exogenous applications of OH-JA, the preferred ST2a substrate in vitro, can enhance some JA-Ile triggered responses in Arabidopsis (27) and induce the expression of certain genes, including *ST2a* (21). Therefore, sulfation of OH-JA by ST2a may be critical for the generation of a genuinely inactive metabolite, channeling JA molecules away from bioactive pools.

149 To define the functionality of ST2a, we measured JA-response markers in plants exposed to
150 mechanical wounding under contrasting light conditions. Genes involved in JA biosynthesis
151 (*LOX2*), JA signaling (*MYC2*), and JA response (*VSP2*) were regulated as expected in Col-0, with
152 FR repressing the response to wounding (Fig. 3A). In the *st2a-1* null mutant, the basal
153 expression of these genes was higher than in Col-0, and the suppressing effect of FR radiation
154 completely disappeared (Fig. 3A). RNAseq analysis of samples from wounded rosettes revealed
155 a statistically significant overlap between the genes downregulated by FR in Col-0 plants and
156 those upregulated by the *st2a-1* mutation under FR radiation. The group of overlapping genes
157 was significantly enriched in the GO terms “Response to JA” and “JA biosynthetic process” (Fig.
158 3B and Data File S1). Consistent with the pattern of expression of JA biosynthetic genes in Col-
159 0 and *st2a-1*, we found that FR reduced the accumulation of *cis*-12-oxo-phytodienoic acid (*cis*-
160 OPDA) in Col-0, particularly at high rates of FR supplementation, but this effect of FR was less
161 marked in *st2a-1* plants (Fig. S9). Glucosinolates (GS) are important defense compounds in
162 Arabidopsis, which are often regulated by JA (28) (Fig. S10). In Col-0, the accumulation of these
163 JA-dependent compounds was attenuated when plants were exposed to supplemental FR
164 radiation (Fig. 3C), as expected (29). In contrast, in *st2a-1* plants, FR failed to inhibit GS
165 accumulation (Fig. 3C). Collectively, these data (Fig. 3) indicate that the sulfation reaction
166 catalyzed by ST2a plays a central role suppressing JA-dependent responses in plants
167 undergoing shade avoidance.

168 To investigate the functional role of changes in JA metabolism caused by ST2a activity, we
169 tested the *st2a-1* null mutant in bioassays with larvae of *Spodoptera littoralis* (a chewing
170 insect) and *Botrytis cinerea* (a necrotrophic pathogen). In Col-0, supplemental FR radiation
171 caused increased growth of *S. littoralis* caterpillars that fed on the plants, and increased the
172 size of necrotic lesions generated by *B. cinerea* (Fig. 4A). These FR effects were missing in
173 plants of *st2a-1*, which correlated strongly with the lack of effect of FR reducing the
174 concentration of JA (Fig. 2), JA marker gene transcripts, and defense compounds (Fig. 3).
175 Furthermore, the *pif4 pif5 pif7* triple mutant, which did not upregulate the transcription of
176 *ST2a* in response to supplemental FR, was significantly more resistant to *B. cinerea* than Col-0
177 under low R:FR ratios (Fig. 4A). These data provide compelling empirical support for a
178 functional connection between *ST2a* transcription, increased JA catabolism, and reduced
179 defense under low R:FR ratios.

180 Rosettes of the *st2a-1* null mutant appeared similar to those of Col-0 under ambient light, and
181 they showed normal morphological responses to supplemental FR radiation (leaf hyponasty

182 and petiole elongation) (Fig. 4B). However, when plants were treated with MeJA, the shade-
 183 avoidance response to supplemental FR radiation was significantly attenuated in *st2a-1*; in
 184 contrast, Col-0 plants were capable of reconfiguring their morphology, and showed a normal
 185 response to FR even under MeJA elicitation (Fig. 4B and Fig. S11). Taken together, these results
 186 suggest that the key function of the sulfotransferase ST2a under low R:FR ratios is to facilitate
 187 the inactivation of JA, thereby allowing the plant to express its full repertoire of shade-
 188 avoidance responses and maximize its competitive ability in crowded stands.

189 Failure to respond to competition signals with a rapid reconfiguration of shoot architecture
 190 and leaf traits carries a disproportionate fitness penalty for plants competing for light in fast
 191 growing stands. Under these conditions, suppression of the 'growth brake' (3, 4) imposed by JA
 192 could be a key determinant of success, even if it comes at the cost of attenuating defense
 193 responses. Our results demonstrate the molecular mechanism that links neighbor perception
 194 via phyB with the attenuation of JA signaling (Fig. 4C), and provide a compelling example of the
 195 role of sulfotransferases in the adaptive modulation of hormonal metabolism in plants. This
 196 phyB-dependent sulfation mechanism generates a metabolic sink for bioactive JA, and allows
 197 the plant to refocus its strategy on rapid growth when the perceived risk of competition for
 198 light is strong.

199

- 200 1. J. Schmitt, J. R. Stinchcombe, M. S. Heschel, H. Huber, The adaptive evolution of
 201 plasticity: phytochrome-mediated shade avoidance responses. *Integr. Comp. Biol.* **43**,
 202 459 (2003).
- 203 2. E. E. Farmer, *Leaf Defence*. (Oxford University Press, Oxford, UK, 2014), pp. 224.
- 204 3. Y. Yan *et al.*, A downstream mediator in the growth repression limb of the jasmonate
 205 pathway. *Plant Cell* **19**, 2470 (2007).
- 206 4. Y. Zhang, J. G. Turner, Wound-Induced endogenous jasmonates stunt plant growth by
 207 inhibiting mitosis. *PLoS ONE* **3**, e3699 <https://doi.org/10.1371/journal.pone.0003699>
 208 (2008).
- 209 5. W. Zhou *et al.*, A jasmonate signaling network activates root stem cells and promotes
 210 regeneration. *Cell*, doi.org/10.1016/j.cell.2019.03.006 (2019).
- 211 6. I. Monte *et al.*, A single JAZ repressor controls the jasmonate pathway in *Marchantia*
 212 *polymorpha*. *Mol. Plant* **12**, 185 (2019).
- 213 7. Q. Guo *et al.*, JAZ repressors of metabolic defense promote growth and reproductive
 214 fitness in *Arabidopsis*. *P. Natl. Acad. Sci. USA* **115**, E10768 (2018).

- 215 8. C. L. Ballaré, Light regulation of plant defense. *Annu. Rev. Plant Biol.* **65**, 335 (2014).
- 216 9. M. de Wit, V. C. Galvão, C. Fankhauser, Light-mediated hormonal regulation of plant
217 growth and development. *Annu. Rev. Plant Biol.* **67**, 613 (2016).
- 218 10. J. E. Moreno, Y. Tao, J. Chory, C. L. Ballaré, Ecological modulation of plant defense via
219 phytochrome control of jasmonate sensitivity. *P. Natl. Acad. Sci. USA* **106**, 4935 (2009).
- 220 11. M. L. Campos *et al.*, Rewiring of jasmonate and phytochrome B signalling uncouples
221 plant growth-defense tradeoffs. *Nat. Comm.* **7**, 12570 | DOI: 10.1038/ncomms12570 |
222 (2016).
- 223 12. M. Leone, M. M. Keller, I. Cerrudo, C. L. Ballaré, To grow or defend? Low red:far-red
224 ratios reduce jasmonate sensitivity in Arabidopsis seedlings by promoting DELLA
225 degradation and increasing JAZ10 stability. *New Phytol.* **204**, 355 (2014).
- 226 13. J. M. Chico *et al.*, Repression of jasmonate-dependent defenses by shade involves
227 differential regulation of protein stability of MYC transcription factors and their JAZ
228 repressors in Arabidopsis. *Plant Cell* **26**, 1967 (2014).
- 229 14. E. Oh, J.-Y. Zhu, Z.-Y. Wang, Interaction between BZR1 and PIF4 integrates
230 brassinosteroid and environmental responses. *Nat. Cell Biol.* **14**, 802 (2012).
- 231 15. P. Leivar, P. H. Quail, PIFs: pivotal components in a cellular signaling hub. *Trends Plant*
232 *Sci.* **16**, 19 (2011).
- 233 16. S. Lorrain, T. Allen, P. D. Duek, G. C. Whitelam, C. Fankhauser, Phytochrome-mediated
234 inhibition of shade avoidance involves degradation of growth-promoting bHLH
235 transcription factors. *Plant J.* **53**, 312 (2008).
- 236 17. L. Li *et al.*, Linking photoreceptor excitation to changes in plant architecture. *Gene*.
237 *Dev.* **26**, 785 (2012).
- 238 18. M. de Wit, K. Ljung, C. Fankhauser, Contrasting growth responses in lamina and petiole
239 during neighbor detection depend on differential auxin responsiveness rather than
240 different auxin levels. *New Phytol.* **208**, 198 (2015).
- 241 19. F. Hirschmann, F. Krause, J. Papenbrock, The multi-protein family of sulfotransferases
242 in plants: composition, occurrence, substrate specificity, and functions. *Front. Plant*
243 *Sci.* **5**, (2014).
- 244 20. A. Koprivova, S. Kopriva, Sulfation pathways in plants. *Chem. Biol. Int.* **259**, 23 (2016).
- 245 21. S. K. Gidda *et al.*, Biochemical and molecular characterization of a hydroxyjasmonate
246 sulfotransferase from Arabidopsis thaliana. *J. Biol. Chem.* **278**, 17895 (2003).
- 247 22. E. Baumann, Ueber Sulfosäuren im Harn. *Ber. Dtsch. Chem. Ges.* **9**, 54 (1876).
- 248 23. K. Nagata, Y. Yamazoe, Pharmacogenetics of sulfotransferase. *Annu. Rev. Pharmacol.*
249 *Toxicol.* **40**, 159 (2000).

- 250 24. N. Gamage *et al.*, Human sulfotransferases and their role in chemical metabolism.
251 *Toxicol. Sci.* **90**, 5 (2006).
- 252 25. R. Komori, Y. Amano, M. Ogawa-Ohnishi, Y. Matsubayashi, Identification of
253 tyrosylprotein sulfotransferase in Arabidopsis. *P. Natl. Acad. Sci. USA* **106**, 15067
254 (2009).
- 255 26. A. Agrawal, E. Kearney, A. Hastings, T. Ramsey, Attenuation of the jasmonate burst,
256 plant defensive traits, and resistance to specialist monarch caterpillars on shaded
257 common milkweed (*Asclepias syriaca*). *J. Chem. Ecol.* **38**, 893 (2012).
- 258 27. E. Smirnova *et al.*, Jasmonic acid oxidase 2 hydroxylates jasmonic acid and represses
259 basal defense and resistance responses against *Botrytis cinerea* infection. *Mol. Plant*
260 **10**, 1159 (2017).
- 261 28. B. A. Halkier, J. Gershenzon, Biology and biochemistry of glucosinolates. *Annu. Rev.*
262 *Plant Biol.* **57**, 303 (2006).
- 263 29. M. D. Cargnel, P. V. Demkura, C. L. Ballaré, Linking phytochrome to plant immunity:
264 low red: far-red ratios increase Arabidopsis susceptibility to *Botrytis cinerea* by
265 reducing the biosynthesis of indolic glucosinolates and camalexin. *New Phytol.* **204**,
266 342 (2014).
- 267 30. C. Wasternack, I. Feussner, The Oxylipin Pathways: Biochemistry and Function. *Annu.*
268 *Rev. Plant Biol.* **69**, 363 (2018).

269

270 ACKNOWLEDGMENTS

271 We thank I. Cerrudo, P. Karssemeijer and B. F. Alliani for help with the initial characterization
272 of *st2a* lines and P. V. Demkura and B. Rothe for technical assistance. **Funding:** Supported by
273 grants of the Agencia Nacional de Promoción Científica y Tecnológica, Universidad de Buenos
274 Aires and The New Phytologist Trust (C.L.B. and A.T.A.), the Max Planck Society (J.G.), National
275 Science Foundation (IOS-1557439; A.J.K.), a Georg Forster Research Award from the Alexander
276 von Humboldt Foundation (to C.L.B.) and a Deutscher Akademischer Austauschdienst
277 Fellowship (to G.L.F.-M.). **Author contributions:** G.L.F.-M. contributed to all aspects of this
278 research; C.D.C. contributed to experimental design, data collection, analysis and
279 interpretation; M.R. designed and executed protocols for metabolite and hormone analyses;
280 T.Z. and A.J.K. designed and performed hormone profiling; M.D.C. carried out bioassays and
281 M.Z.L. screened mutant lines and helped with gene expression analyses; C.A.M. and T.G.K.
282 performed analyses of transcriptomic data and helped with data interpretation; A.T.A. and J.G.
283 contributed to the general conception of the project and data interpretation; C.L.B. conceived

284 the project and contributed to data generation and analysis, and wrote manuscript with input
 285 from all co-authors. **Competing interests:** The authors declare that they have no competing
 286 interests. **Data and materials availability:** All data needed to evaluate the conclusions in the
 287 paper are present in the main text or the supplementary materials.

288

289 **SUPPLEMENTARY MATERIALS**

290 The file includes:

291 Materials and Methods

292 Figs. S1 to S12

293 Tables S1 and S2

294 References

295 Data Files S1 and S2

296

297

Fig. 1. FR supplementation reduces the pool of bioactive JAs and increases *ST2a* transcription and JA sulfation. Col-0 Arabidopsis plants were sprayed with 200 μ M MeJA and harvested at the indicated time points for measurements of JA pools and gene expression. The color of the box outline indicates the direction of the FR effect: Blue = downregulation; Yellow = transient upregulation; Red = upregulation; unboxed genes were not significantly regulated by FR. Metabolic map adapted from Wasternack and Feussner (30). The bar charts show quantitative data for metabolite concentrations (thin bars indicate 1 SE; n = 3 biological replicates). For gene expression data, see Fig. S3.

Fig. 2. FR attenuates the JA burst triggered by mechanical wounding and increases the concentration of HSO₄-JA in a *st2a*-dependent manner. Significant genotype x light (GxL) interaction terms are indicated. For each time point, different letters indicate significant differences between means ($P < 0.05$); thin bars indicate 1 SE ($n = 6$ biological replicates). DW = dry weight. For additional jasmonate pools, see Fig. S5.

Fig. 3. FR downregulates gene and metabolite markers of jasmonate signaling in an *ST2a*-dependent manner. (A) qPCR results for selected markers of JA synthesis, signaling and response. (B) Summary of RNAseq results demonstrating a significant overlap between the genes downregulated by FR in Col-0 plants and those upregulated by the *st2a-1* mutation in wounded plants. The table shows the GO categories overrepresented in the set of overlapping genes (for details on analysis see Data File S1). (C) Suppression by FR of glucosinolate accumulation in plants treated with MeJA, mechanical wounding or insect herbivory (*Spodoptera littoralis*) was missing in a *st2a-1* null mutant. For specific data on 4MSOB and I3M in wounded plants, see Fig. S10 B). For induced plants, the significance of the genotype x light (GxL) interaction term is indicated in panels A and C. Different letters indicate significant ($P < 0.05$) differences between means; thin bars indicate 1 SE ($n = 6$ biological replicates for glucosinolate data or 3 for transcriptomic data).

335

336

337

338 **Fig. 4. Sulfotransferase ST2a is key in regulating the growth/defense balance in response to**
 339 **changes in the R:FR ratio.** (A) Under FR supplementation, plants that do not upregulate *ST2a*
 340 expression are better defended than their Col-0 counterparts. Bioassays were carried out
 341 comparing Col-0 and *st2a-1* rosettes using larvae of *Spodoptera littoralis* (upper panel) and
 342 inoculations with *Botrytis cinerea* spore suspensions (middle panel), and also comparing Col-0
 343 and *pif4pif5pif7* triple mutants inoculated with *B. cinerea* spore suspensions (lower panel). The
 344 significance of the genotype x light interaction term (GxL) is indicated for each factorial
 345 experiment; different letters indicate significant ($p < 0.05$) differences between means. (B)
 346 *st2a-1* rosettes display normal phenotypes under control conditions but, compared with Col-0
 347 rosettes, they display impaired shade-avoidance responses when exposed to low doses of
 348 MeJA (100 μ M). **, $p < 0.01$; for full dataset, see Fig. S10. (C) Conceptual model linking the
 349 perception of low R:FR ratios via phyB with the modulation of jasmonate metabolism and
 350 signaling through regulation of *ST2a* transcription via the phyB-PIF transcription module. Pr,
 351 inactive form of phytochrome; Pfr, active form of phytochrome.