

1 Title: Population Health Genetic Screening for Tier 1 Inherited Diseases in Northern Nevada:  
2 90% of At-Risk Carriers are Missed

3 Authors:

4 Grzymiski JJ\* <sup>1,2</sup>, Elhanan G<sup>2</sup>, Smith E<sup>2</sup>, Rowan C<sup>1</sup>, Slotnick N<sup>1</sup>, Dabe S<sup>2</sup>, Schlauch K<sup>2</sup>, Read R<sup>2</sup>,  
5 Metcalf WJ<sup>2</sup>, Lipp B<sup>2</sup>, Reed H<sup>2</sup>, Levin E<sup>3</sup>, Kao J<sup>3</sup>, Rashkin M<sup>3</sup>, Bowes J<sup>3</sup>, Bolze A<sup>3</sup>, Dunaway K<sup>3</sup>,  
6 Washington N<sup>3</sup>, Slonim A<sup>1</sup>, Lu JT\* <sup>3</sup>

7 Affiliations

- 8 1. Renown Health; Reno, NV  
9 2. Desert Research Institute; Reno, NV  
10 3. Helix; San Carlos, CA

11 \*Corresponding Authors:

12 Joseph Grzymiski, PhD.  
13 Joe.Grzymiski@dri.edu

14  
15 James Lu, M.D. Ph.D.  
16 james.lu@helix.com  
17

18 **Conflicts**

19 EL, JK, MR, JB, AB, KD, NW, JL are employees of Helix.

20 **Data Availability Statement**

21 The data that support the findings of this study are available from the corresponding author  
22 upon reasonable request.

23 **Code Availability Statement**

24 The code that supports the findings of this study are available from the corresponding author  
25 upon reasonable request.

26  
27

## 1 Abstract

2 In an unselected population of 23,713 participants who underwent clinical exome sequencing as  
3 a part of the Healthy Nevada Project (HNP) in Northern Nevada (Renown Health, Reno,  
4 Nevada) from March 15, 2018, to Sept 30, 2018 (Table S1) we find a 1.26% carrier rate for  
5 expected pathogenic and likely pathogenic genetic variants in (FH: *LDLR*, *PCSK9*, *APOB*),  
6 Hereditary Breast and Ovarian Cancer (HBOC: *BRCA1*, *BRCA2*) and Lynch Syndrome (LS:  
7 *MLH1*, *MSH2*, *MSH6*, *PSM2*) with over 90% of carriers undetected under current medical  
8 practice. 26% of carriers were found to have advanced disease with 70% first diagnosed before  
9 the age of 65. Less than 20% of all carriers had any documented suspicion for inherited genetic  
10 disease in the medical record and upon direct follow-up survey under 40% of carriers had family  
11 history of relevant disease. A population preventative genetic screening approach for patients  
12 under 45 may improve outcomes.

## 13 Main

14 In an unselected population of 23,713 participants who underwent clinical exome sequencing as  
15 a part of the Healthy Nevada Project (HNP) in Northern Nevada (Renown Health, Reno,  
16 Nevada) from March 15, 2018, to Sept 30, 2018 (Table S1) we find a 1.26% carrier rate for  
17 expected pathogenic and likely pathogenic genetic variants in (FH: *LDLR*, *PCSK9*, *APOB*),  
18 Hereditary Breast and Ovarian Cancer (HBOC: *BRCA1*, *BRCA2*) and Lynch Syndrome (LS:  
19 *MLH1*, *MSH2*, *MSH6*, *PSM2*) with over 90% of carriers undetected under current medical  
20 practice. 26% of carriers were found to have advanced disease with 70% first diagnosed before  
21 the age of 65. Less than 20% of all carriers had any documented suspicion for inherited genetic  
22 disease in the medical record and upon direct follow-up survey under 40% of carriers had family  
23 history of relevant disease. A population preventative genetic screening approach for patients  
24 under 45 may improve outcomes.

25 Here we report on genetic risk and disease manifestation for the above three inherited  
26 autosomal dominant diseases (9 genes) that have evidence-based recommendations for  
27 genetic testing targeted towards subsets of the population based on family history, ethnic  
28 background, or other demographic characteristics <sup>1</sup>. Interrogating the clinical genetic data for  
29 previously described pathogenic and likely pathogenic (P/LP) variants <sup>2</sup> identified 290  
30 individuals for a population prevalence of 1.26% (Table S1). Although not corrected for familial  
31 relatedness or potential ascertainment bias, the prevalence rate of BRCA related HBOC (~1 in  
32 185), LS (~1 in 520) and FH (~1 in 208) are in line with recent reports <sup>3-5</sup>. Carriers broadly  
33 resemble the overall demographic profile of the HNP with a median age of 47, 63% female, and  
34 78% caucasian (Table S1). We then curated and reviewed available medical records, extracted  
35 and processed in February 2019, and combined this data with surveyed family history disease  
36 risk (See Online Methods) for each of the 243 participants (83.3% of the 290 participants) who  
37 are patients at Renown Health System. We find that 62 (26%) patients had already presented  
38 with disease. First disease diagnosis was ascertainable for only 55 patients (of the 64) as some  
39 patients enter Renown Health as patients with disease. The median age of first diagnosis for  
40 these patients was 60 with 88.6% of disease first presenting after the age of 40 (Figure 1), and  
41 70% before the age of 65. Of these patients, 26 (47.2%) had no documented diagnostic  
42 information that could have raised suspicion of underlying genetic condition prior to diagnosis,  
43 suggesting a somewhat acute presentation.

44 For HBOC carriers, we were able to review 108 medical records. These patients had a median  
45 age of 51 with 91 (84.3%) of the carriers falling outside of NCCN practice guidelines for genetic

1 testing. We also find that 23 of the HBOC carriers (21.2%) had a diagnosis of breast or ovarian  
2 cancer with a median diagnosis age of 60. In comparison, the prevalence of non-BRCA1/2  
3 HBOC in women at age 55 is ~1.47%<sup>6</sup>. 9 of the 23 diseased carriers (39.1%) were within  
4 NCCN guidelines for genetic testing; however, only five of the nine had genetic testing prior to  
5 HNP, likely due to early onset of disease (median age at diagnosis of 45 for tested individuals).  
6 The remaining four had a median age of diagnosis of 64.5, which likely reduced clinical  
7 suspicion of germline disease. We also found that 14 (60.9%) of the carriers with disease were  
8 nonetheless tested after disease diagnosis even though they fell outside of NCCN guidelines  
9 (Table 1). Although we find no differences in disease-free progression between BRCA1 and  
10 BRCA2 carriers (Figure S1) in our cohort but do find that family history provides incremental risk  
11 excess of carrier status alone (Figure S2). This is consistent with previous studies that show  
12 that family history is an independent risk factor for breast/ovarian cancer<sup>7</sup>. Lastly we find that for  
13 15 of the 23 (65.2%) of the diseased patients, there was no diagnostic evidence that could  
14 indicate suspicion for disease prior to diagnosis with breast or ovarian cancer. Instead, these  
15 individuals either presented acutely or through routine clinical screening for breast cancer (i.e.  
16 mammography).

17 For LS carriers, 37 medical records were available and reviewed. These patients had a median  
18 age of 58 with 35 (94.6%) of the carriers falling outside of NCCN practice guidelines (Table 1)  
19 for genetic testing. Carriers of LS P/LP variants had marked increases in oncologic events; 13  
20 of the 37 (35.1%) carriers had a colon, ovarian or other cancer diagnosis with a median age of  
21 first diagnosis of 44. Seven of the 13 had colon cancer while six of the 13 had another type of  
22 malignancy; 4 patients presented with malignancy in two or more locations. In contrast the  
23 prevalence of colon cancer at ages 40-49 is 0.12%<sup>6</sup> (Table 1) in individuals without LS. The  
24 prevalence rate in our unselected population is in-line with the approximately ~40% cumulative  
25 incidence rates of cancer by age 50 in family history-selected LS mutation carriers<sup>8</sup>. Of the  
26 seven patients with colon cancer, six had documentary evidence in the medical record that  
27 could have raised suspicion for an underlying genetic condition, however only 2 of these  
28 patients fell within NCCN guidelines and were tested for Lynch Syndrome. No other patients  
29 had genetic testing. This is perhaps unsurprising given that average risk CRC screening starts  
30 at age 50<sup>9</sup>. In contrast, annual screening in LS patients is recommended at age 20<sup>10</sup>.

31 In FH carriers, 98 medical records were reviewed along with family history ascertained during  
32 the Return of Results (See Online Methods). We found that 63% of FH carriers did not have  
33 known family history of heart disease; in addition, only three individuals were found to have an  
34 indication for genetic testing and only two had testing. The remaining 95 (96.9%) of carriers  
35 were found to be outside of guidelines-based testing<sup>11</sup>. Of the 98 carriers, 26 (26.5%) had  
36 manifested with atherosclerotic cardiovascular disease (ASCVD). We found a relative  
37 enrichment of 2.3 (95% CI: 1.13-3.47, Two Proportion Test) of males to females; 14 of 33  
38 (42.4%) of male carriers (median age of diagnosis 64) and 12 of 65 (18.5%) of female carriers  
39 (median age of diagnosis 63) had manifested ASCVD (Table 1). This is in line with previously  
40 described rates for phenotypic FH<sup>12</sup>. In comparison, the prevalence for non-FH ASCVD in 40-  
41 59 y.o males and females is 6.3% and 5.6%<sup>13</sup>. We also observed no discernable differences in  
42 disease penetrance between APOB, PCSK9 and LDLR (Figure S3). Lastly, there were 8 major  
43 acute events out of the 26 disease cases: five myocardial Infarctions (MI) and three cerebral  
44 vascular accidents (CVA). For six of these eight events, the first documented ASCVD related  
45 event was the acute event. The remaining patients all had evidence of coronary intervention.

46 To our knowledge, the ascertainment of genetic disease in an unselected population using a  
47 population health approach has not previously been performed. While the observed prevalence

1 rates in HNP for carriers of FH, HBOC, and LS are consistent with previously published  
2 biobanking studies<sup>3-5</sup>, we discover that current medical guidelines are largely inadequate for  
3 detecting carriers of these three diseases. Over 90% of P/LP carriers are undetected today  
4 because guidelines are narrowly focused on selected populations. Yet disease penetrance for  
5 LS and FH in our unselected population is similar to that in selected populations recommended  
6 by guidelines, and substantially elevated in HBOC carriers (albeit at a lower rate than those with  
7 additional family history). Our study also illustrates that average risk screening protocols may  
8 not adequately identify at-risk individuals as average risk screening typically is less intensive  
9 and starts later in life: screening using mammography for breast cancer starts at age 50 in  
10 women<sup>14</sup>, for colon cancer at age 50 in both sexes<sup>15</sup>, and for lipid disorders at age 35 and 45  
11 for men and women<sup>16</sup>, respectively.

12 The HNP follows a public health community-based approach with direct participant outreach,  
13 enrollment and return of results. This also includes direct to participant genetic education and  
14 physician CME courses (See Online Methods). Some self-selection for the study is evident  
15 from the relative enrichment of caucasians and the higher educational attainment of the cohort.  
16 However these demographics typically enrich for higher levels of engagement with health  
17 systems<sup>17</sup> and higher compliance with preventative care. Our analysis of medical records is  
18 also limited to 12 years of electronic health records from the Renown Health System, although  
19 this is the dominant health system in Northern Nevada with more than 70% market share.


20 In the over 1.1 million health records at Renown Health, diagnosis of FH, LS or HBOC occurs in  
21 less than 0.2% of patients. Our results suggest that genetic predisposition testing as a first line  
22 preventative genetic screen for these conditions could lower mortality and morbidity related to  
23 inherited disease risks that are today insufficiently detected under current medical practice. A  
24 population health screening approach to genetic medicine requires different scrutiny given the  
25 potential for added costs, over-interpretation of disease risk, and ethical and social factors<sup>18</sup>.  
26 Cost-effectiveness studies have found predispositional genetic screening with return of a limited  
27 panel of genetic information to be cost effective in a population under 45 years old when testing  
28 costs are in the low hundreds of dollars<sup>19,20</sup>. Our experience with the Healthy Nevada Project  
29 suggests that a public health based approach to genetic screening in younger patients can  
30 provide substantial clinical, economic, and patient benefit.

31

32

## 1 Figures and Tables

- 2 **Figure 1:** Empirical cumulative distribution for age of first diagnosis for FH related  
3 Atherosclerotic cardiovascular disease (ASCVD), BRCA1/2 related Hereditary Breast and  
4 Ovarian Cancer (HBOC) and Lynch Syndrome related malignancy (LS Ca).




5  
6

7

	HBOC				LS						FH			
	BRCA1	BRCA2	Total		MLH1	MSH2	MSH6	PSM2	Total		APOB	LDLR	PCKS9	Total
# of Carriers	51	84	135	# of Carriers	8	4	17	17	46	# of Carriers	26	79	12	117
# w/ Medical Records	42	66	108	# w/ Medical Records	5	3	16	13	37	# w/ Medical Records	20	67	11	98
Sex				Sex						Sex				
Female	26	36	62	Female	5	1	10	8	24	Female	13	46	6	65
Male	16	30	46	Male	0	2	6	5	13	Male	7	21	5	33
Median Age [Range]	45 [18-81]	52 [19-81]	51 [18-81]	Median Age [Range]	51 [26-66]	28 [28-31]	57 [26-84]	68 [26-84]	58 [26-84]	Median Age [Range]	46 [19-79]	47 [18-77]	56 [27-81]	47 [18-81]
# of Patients with indication for genetic testing	17			# of Patients with indication for genetic testing	2					# of Patients with indication for genetic testing	3			
# w/ Genetic Testing Completed (July 2018)	8 (47.9%)			# w/ Genetic Testing Completed (July 2018)	2 (100%)					# w/ Genetic Testing Completed (July 2018)	2 (67%)			
# of Patients without indication for genetic testing	91			# of Patients without indication for genetic testing	35					# of Patients without indication for genetic testing	95			
# w/ Genetic Testing Completed (July 2018)	5			# w/ Genetic Testing Completed (July 2018)	1					# w/ Genetic Testing Completed (July 2018)	2			
# with breast/ovarian CA	10	13	23	# w/ LS related malignancy	3	3	6	3	15	# with ASVCD	6	18	2	26
Sex				Sex						Sex				
Female	9	13	22	Female	3	1	4	3	11	Female	4	8	0	12
Male	1	0	1	Male	0	2	2	0	4	Male	2	10	2	14
Median age [Range] at time of diagnosis	62.5 [40-67]	52 [23-71]	60 [23-71]	Median age [Range] at time of diagnosis	45 [26-65]					Median age [Range] at time of diagnosis	53 [32-72]	65 [32-70]	57	62 [32-72]
Prior Clinical Suspicion of Genetic Disease	3	5	8	Prior Clinical Suspicion of Genetic Disease	1	2	2	0	5	Prior Clinical Suspicion of Genetic Disease	5	6	1	12


**Table 1:** Detailed information about HBOC, LS, and FH including carriers per gene and disease prevalence

**Figure S1: No difference in disease free progression between carriers of BRCA1 and BRCA2 P/LP variants.**






**Figure S2: Disease-free progression in HBOC carriers that are within and outside of NCCN guidelines for testing.**



**Figure S3: No difference in disease free progression between carriers of APOB, LDLR, and PCSK9**



**Table S1: Demographic Profile of Healthy Nevada Project Participants.**

<b>Race/Ethnicity:</b>	<b>HBOC carriers</b>	<b>FH Carriers</b>	<b>LS Carriers</b>	<b>Healthy Nevada Project</b>	<b>Renown Health System</b>	<b>Washoe County</b>
Count	135	117	46	23,709	~1.03M	421,407
Prevalence	0.57%	0.49%	0.19%			
Sex						
Male (%)	39%	35%	33%	34%	51%	50%
Female (%)	61%	65%	67%	66%	49%	50%
Age (Median, Stdev)	44 [16.9]	47.2 [18.2]	50.5 [19]	44.5 [19]	49 [24]	38
Ethnicity (Count, %)						
Caucasian	107 (79.3%)	87 (74.4%)	38 (82.6%)	81%	72%	63%
Hispanic/Latino	17 (12.6%)	6 (5.1%)	1 (2.2%)	10%	10%	25%
Asian	4 (3%)	11 (9.4%)	3 (6.5%)	3%	3%	5%
Native American	3 (2.2%)	0 (0%)	2 (4.3%)	1%	1%	1%
Pacific Islander	0 (0%)	0 (0%)	0 (0%)	1%	0%	1%
African American	2 (1.5%)	8 (6.8%)	1 (2.2%)	1%	3%	2%
Other/Unknown	2 (1.5%)	5 (4.3%)	1 (2.2%)	3%	11%	3%

**Table S2: List of P/LP Variants for all cases**

Disease	Gene	# of Cases	Relative Contribution	Population Prevalence
HBOC	BRCA1	51	37.78%	0.215%
HBOC	BRCA2	84	62.22%	0.354%
LS	MLH1	8	17.39%	0.034%
LS	MSH2	4	8.70%	0.017%
LS	MSH6	17	36.96%	0.072%
LS	PMS2	17	36.96%	0.072%
FH	APOB	26	22.22%	0.110%
FH	LDLR	79	67.52%	0.333%
FH	PCKS9	12	10.26%	0.051%

## Online Methods

### Healthy Nevada Project

The Healthy Nevada Project (<https://healthynv.org/>) is an all-comers human subject research study on health determinants in Northern Nevada with specific recruitment foci in rural and socio-economically depressed areas in Northern Nevada. This ongoing program, with a targeted enrollment of 250,000, targets a diverse demographic: 23.8% of individuals are recruited from rural zip codes and ~20% of the study consists of minority participants, predominantly hispanic and native american (Table S1). Participants consent to a research protocol that includes (1) clinical "Exome+" sequencing, (2) linking of sequencing data with electronic health records, (3) return of secondary findings and (4) recontact for health surveys and participation in future studies. 99.2% agreed to return of secondary findings and 96.1% agreed to recontact for future studies. The project 1106618-15, was approved by the University of Nevada, Reno Institutional Review Board.

### Return of Secondary Findings

The return of secondary findings process started in October 2018 and was completed for these patients by February 2019. For each positive secondary finding the de-identified participant ID was cross-matched with a de-identified consent. If the participant consented to return of results, the data was re-identified by an authorized member of the research team. Contact information was then linked by a designated research coordinator at the Office of Research and Education using Renown Health's secure email platform. Linked contact list and individual reports were stored in a secure folder on an encrypted server only accessible to designated research staff.

Clinical Research Coordinators scheduled a 15 minute phone appointment to speak to a clinician regarding their results. Participants were called 1-2 days prior to the first available appointment to minimize wait time for the delivery of their results over the phone. If there was no answer, a voicemail was left with the participant. A phone call was also made to any secondary number on file, and a message left if applicable. The voicemail was then followed up with an email to the same participant with similar information to call back and schedule an appointment. The date of scheduling or unsuccessful attempts to contact each participant was recorded on the contact list. Every effort was made to minimize the significance of the call to the greatest extent possible until the day of the scheduled appointment to avoid undue stress to the participant.

All medically significant results were delivered by a clinical provider with appropriate credentialing to deliver genetic results. To ensure consistency between participants, a generalized script was used by the clinician when delivering results. At the time of the appointment, the clinician verified that the participant still wanted to be informed of any findings that might be relevant to their health prior to continuing the call. Once confirmed, the clinician informed the individual of their secondary finding, what it meant for their health, and the importance of following up with a medical professional upon receiving their results. Family history pertaining to their results was collected for research purposes using Renown's family history questionnaire. This questionnaire was designed to survey family medical history specific to (1) NCCN guidelines for Lynch Syndrome, (2) Tyrer-Cuzick<sup>21</sup>, or Penn II empirical risk calculators<sup>22</sup> for HBOC, and (3) family disease risk for cardiovascular disease<sup>12,23</sup>. Prior to finishing the call, participants were asked if they would like a copy of their results delivered to them either by secure email and/or through the postal service.

Once a participant was notified of their results, a copy of their signed report was sent to their preferred method of delivery (email or letter), along with supporting documents. Supporting documentation included: a cover letter with contact information for the study, templated letters they can provide to their physicians and family members, a copy of available primary care physicians in the area if requested by the participant, and any other educational or informational material pertaining to their findings as deemed applicable. Source documents for the return of results were maintained by research staff per research policy. In addition, a copy of returned results is retained in our Clinical Trial Management System for compliance with study protocols.

## **Electronic health record analysis**

HNP participant IDs were mapped to medical record IDs. For all HNP participants with valid medical record IDs, clinical data were retrieved from the Renown Epic Clarity database (February 2019) utilizing a script. The script was designed to collect patient diagnoses from multiple Clarity tables including, but not limited to, encounter diagnoses, problem lists, medical histories and claims, and invoices. The script also de-identified records by removing all PHI. The diagnoses retrieved were encoded in Renown's proprietary codes and were mapped to ICD10-CM and ICD9-CM diagnosis codes utilizing a mapping schema. For each patient, each unique ICD diagnosis code was tagged with the date of its first and last occurrence across all the Clarity tables that were queried by the script. Data regarding lipid profile results and evidence of prescriptions of lipid lowering therapies and their types were retrieved in a similar manner.

For each patient, diagnosis codes, lab data and medication orders were reviewed by a clinician. Diagnosis codes that were not deemed relevant to the underlying FH, LS and HBOC conditions were removed. The remaining codes, laboratory tests, medication orders and timestamps were used to determine the status of the condition itself, family history and other risk factors. This includes determining if 1) the patient had genetic testing prior to Return of Results from the HNP, 2) if the patient's family history and patient conditions supported clinical suspicion for an underlying genetic condition and 3) if the patient had a diagnosis of a disease that could reasonably be associated with underlying genetic risk such as hereditary breast & hereditary ovarian cancer, Lynch Syndrome associated malignancies and /or ASCVD related diagnoses and the earliest date of diagnosis.

## **Assay and Sequencing**

Sequencing for all participants was performed at Helix's CLIA (#05D2117342) and CAP Accredited (#9382893) facility in San Diego, CA. The Exome+ (Helix) is a proprietary exome that combines a highly performant medical exome with a microarray backbone into a single sequencing assay ([www.helix.com](http://www.helix.com)). The Exome+ provides highly uniform and complete coverage across medical regions of the genome, with a target coverage >20x for >99% of all targeted bases in this study, for all participants. All returned variants meet Helix's validation criteria for clinical return. Performance specification can be found in Helix's Exome+ Performance White Paper (<https://cdn.helix.com/wp-content/uploads/2017/07/Helix-Personal-Genomics-Platform-White-Paper.pdf>).

## **Selection of Pathogenic/ Likely Pathogenic Alleles**

The list of pathogenic/ likely pathogenic were filtered from CLINVAR using filters for pathogenic or likely pathogenic variants. We then removed any variants with conflicting interpretations and any variants whose origin was somatic. These data were generated from the June 3, 2018 CLINVAR database (<https://www.ncbi.nlm.nih.gov/clinvar/>).

## References

1. Khoury, M. J. *et al.* A collaborative translational research framework for evaluating and implementing the appropriate use of human genome sequencing to improve health. *PLoS Med.* **15**, e1002631 (2018).
2. Richards, S. *et al.* Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genetics in Medicine* **17**, 405–423 (2015).
3. Dewey, F. E. *et al.* Distribution and clinical impact of functional variants in 50,726 whole-exome sequences from the DiscovEHR study. *Science* **354**, (2016).
4. Abul-Husn, N. S. *et al.* Genetic identification of familial hypercholesterolemia within a single U.S. health care system. *Science* **354**, (2016).
5. Manickam, K. *et al.* Exome Sequencing–Based Screening for BRCA1/2 Expected Pathogenic Variants Among Adult Biobank Participants. *JAMA Network Open* **1**, e182140 (2018).
6. Noone AM, Howlader N, Krapcho M, Miller D, Brest A, Yu M, Ruhl J, Tatalovich Z, Mariotto A, Lewis DR, Chen HS, Feuer EJ, Cronin KA. SEER Cancer Statistics Review, 1975-2015. *National Cancer Institute. Bethesda, MD*, (2018). Available at: [https://seer.cancer.gov/csr/1975\\_2015/index.html](https://seer.cancer.gov/csr/1975_2015/index.html). (Accessed: 17th April 2019)
7. Kuchenbaecker, K. B. *et al.* Risks of Breast, Ovarian, and Contralateral Breast Cancer for BRCA1 and BRCA2 Mutation Carriers. *JAMA* **317**, 2402–2416 (2017).
8. Møller, P. *et al.* Cancer incidence and survival in Lynch syndrome patients receiving colonoscopic and gynaecological surveillance: first report from the prospective Lynch syndrome database. *Gut* **66**, 464–472 (2017).
9. Final Update Summary: Colorectal Cancer: Screening - US Preventive Services Task Force. (1AD). Available at: <https://www.uspreventiveservicestaskforce.org/Page/Document/UpdateSummaryFinal/colorectal-cancer-screening>. (Accessed: 12th May 2019)
10. Giardiello, F. M. *et al.* Guidelines on genetic evaluation and management of Lynch syndrome: a consensus statement by the US Multi-society Task Force on colorectal cancer. *Am. J. Gastroenterol.* **109**, 1159–1179 (2014).
11. Reiner, Ž. Management of patients with familial hypercholesterolaemia. *Nature Reviews Cardiology* **12**, 565–575 (2015).
12. Perak, A. M. *et al.* Long-Term Risk of Atherosclerotic Cardiovascular Disease in US Adults With the Familial Hypercholesterolemia Phenotype. *Circulation* **134**, 9–19 (2016).
13. Mozaffarian, D. *et al.* Heart disease and stroke statistics--2015 update: a report from the American Heart Association. *Circulation* **131**, e29–322 (2015).
14. Final Recommendation Statement: Breast Cancer: Screening - US Preventive Services Task Force. (1AD). Available at: <https://www.uspreventiveservicestaskforce.org/Page/Document/RecommendationStatementFinal/breast-cancer-screening1>. (Accessed: 12th May 2019)
15. Final Update Summary: Colorectal Cancer: Screening - US Preventive Services Task Force. (1AD). Available at: <https://www.uspreventiveservicestaskforce.org/Page/Document/UpdateSummaryFinal/colorectal-cancer-screening2>. (Accessed: 12th May 2019)
16. Screening for Lipid Disorders in Adults: Recommendation Statement. *AFP* **80**, 1273 (2009).
17. National Academies of Sciences, Engineering, and Medicine, Health and Medicine Division, Board on Population Health and Public Health Practice & Committee on Community-Based



- Solutions to Promote Health Equity in the United States. *Communities in Action: Pathways to Health Equity*. (National Academies Press (US), 2017).
18. Andermann, A., Blancquaert, I., Beauchamp, S. & Déry, V. Revisiting Wilson and Jungner in the genomic age: a review of screening criteria over the past 40 years. *Bull. World Health Organ.* **86**, 317–319 (2008).
  19. Bennette, C. S., Gallego, C. J., Burke, W., Jarvik, G. P. & Veenstra, D. L. The cost-effectiveness of returning incidental findings from next-generation genomic sequencing. *Genet. Med.* **17**, 587–595 (2015).
  20. Zhang, L. *et al.* Population genomic screening of all young adults in a health-care system: a cost-effectiveness analysis. *Genetics in Medicine* (2019). doi:10.1038/s41436-019-0457-6
  21. Tyrer, J., Duffy, S. W. & Cuzick, J. A breast cancer prediction model incorporating familial and personal risk factors. *Stat. Med.* **23**, 1111–1130 (2004).
  22. Lindor, N. M. *et al.* Predicting BRCA1 and BRCA2 gene mutation carriers: comparison of PENN II model to previous study. *Familial Cancer* **9**, 495–502 (2010).
  23. Bertuzzi, M., Negri, E., Tavani, A. & La Vecchia, C. Family history of ischemic heart disease and risk of acute myocardial infarction. *Preventive Medicine* **37**, 183–187 (2003).