Type-3 Secretion System–induced pyroptosis protects *Pseudomonas* against cell-autonomous immunity

Elif Eren^{1¶}, Rémi Planès^{1¶}, Julien Buyck^{2,3¶}, Pierre-Jean Bordignon¹, André Colom¹, Olivier Cunrath², Roland F. Dreier², José C. Santos², Valérie Duplan-Eche⁴, Emmanuelle Näser¹, Antonio Peixoto¹, Dirk Bumann², Céline Cougoule¹, Agnès Coste⁵, Olivier Neyrolles¹, Petr Broz^{2,6*} & Etienne Meunier^{1,2*#}

¹ Institute of Pharmacology and Structural Biology (IPBS), University of Toulouse, CNRS, Toulouse; France

² University of Basel, Biozentrum, Focal Area Infection Biology, Basel; Switzerland

³ UFR Medicine and Pharmacy, INSERM U1070, University of Poitiers, France

⁴ Center of Physiopathology of Toulouse Purpan (CPTP), INSERM, CNRS, University of Toulouse, France.

⁵ UMR152, University of Toulouse, IRD, Toulouse; France

⁶ Department of Biochemistry, University of Lausanne, Epalinges; Switzerland

Running Head: *Pseudomonas*-exploited pyroptosis drives infection

Address correspondence to: etienne.meunier@ipbs.fr (EM)

Present Address: Institute of Pharmacology and Structural Biology (IPBS), University of Toulouse, CNRS, Toulouse; France

*E. Meunier and *P. Broz contributed equally

¶ These authors contributed equally to this work.

28 **Abstract (139 words)**

29 Inflammasome-induced pyroptosis comprises a key cell-autonomous immune process
30 against intracellular bacteria, namely the generation of dying cell structures. These so-
31 called pore-induced intracellular traps (PITs) entrap and weaken intracellular microbes.
32 However, the immune importance of pyroptosis against extracellular pathogens remains
33 unclear. Here, we report that Type-3 secretion system (T3SS)-expressing *Pseudomonas*
34 *aeruginosa* (*P. aeruginosa*) escaped PIT immunity by inducing a NLRC4 inflammasome-
35 dependent macrophage pyroptosis response in the extracellular environment. To the
36 contrary, phagocytosis of *Salmonella* Typhimurium promoted NLRC4-dependent PIT
37 formation and the subsequent bacterial caging. Remarkably, T3SS-deficient
38 *Pseudomonas* were efficiently sequestered within PIT-dependent caging, which favored
39 exposure to neutrophils. Conversely, both NLRC4 and caspase-11 deficient mice
40 presented increased susceptibility to T3SS-deficient *P. aeruginosa* challenge, but not to
41 T3SS-expressing *P. aeruginosa*. Overall, our results uncovered that *P. aeruginosa* uses its
42 T3SS to overcome inflammasome-triggered pyroptosis, which is primarily effective against
43 intracellular invaders.

55 **Importance (119 words)**

56 Although innate immune components confer host protection against infections, the
 57 opportunistic bacterial pathogen *Pseudomonas aeruginosa* (*P. aeruginosa*) exploits the
 58 inflammatory reaction to thrive. Specifically the NLRC4 inflammasome, a crucial immune
 59 complex, triggers an Interleukin (IL)-1 β and -18 deleterious host response to *P.*
 60 *aeruginosa*. Here, we provide evidence that, in addition to IL-1 cytokines, *P. aeruginosa*
 61 also exploits the NLRC4 inflammasome-induced pro-inflammatory cell death, namely
 62 pyroptosis, to avoid efficient uptake and killing by macrophages. Therefore, our study
 63 reveals that pyroptosis-driven immune effectiveness mainly depends on *P. aeruginosa*
 64 localization. This paves the way toward our comprehension of the mechanistic
 65 requirements for pyroptosis effectiveness upon microbial infections and may initiate
 66 targeted approaches in order to ameliorate the innate immune functions to infections.

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82 Manuscript length (5621 words)

83 Introduction

84 Inflammasomes are pro-inflammatory cytosolic complexes whose activation leads
85 to auto-processing of the protease caspase-1. Caspase-1 activation triggers the cleavage
86 and release of the pro-inflammatory cytokines, interleukin (IL)-1 β and IL-18 (1), as well as
87 a pro-inflammatory form of cell death, called pyroptosis. Cleavage and activation of the
88 pore forming effector gasdermin D (GSDMD) by caspase-1 or -11 can induce pyroptosis
89 (2, 3). Several sensors contribute to inflammasome formation, including AIM2-like
90 receptors (AIM2), PYRIN, and a subset of NOD-like receptors (NLRs), namely
91 NLRP1/NLRP1B, NLRP3, NLRP6, NLRP7 and NLRC4, and the caspase-11-induced non-
92 canonical inflammasome pathway (1).

93 Activation of the NLRC4 inflammasome follows an original path as it requires
94 additional helper NLRs – the neuronal apoptosis inhibitory proteins (NAIPs) – that, upon
95 ligand recognition, form hetero-oligomeric complexes with NLRC4 and promote its
96 activation (4–10). Both NAIP5 and NAIP6 directly recognize cytosolic flagellin while NAIP-
97 1 and -2 detect the type-3 secretion system (T3SS) apparatus needle and rod subunits,
98 respectively (1, 6). Importantly, resulting IL-1 β and IL-18 cytokine secretion mediates
99 intracellular bacterial clearance by inducing, respectively, phagocyte recruitment and the
100 production of the microbicidal cytokine interferon (IFN)- γ (1, 6). A recently discovered and
101 understudied cell-autonomous immune process known as pyroptosis promotes
102 intracellular bacteria entrapment and weakening in pore-induced intracellular trap
103 structures (PITs) (11–13), which facilitates subsequent bacterial elimination through
104 efferocytosis (13–16). However, host-adapted bacteria can inhibit PITs formation, which
105 enables bacterial proliferation (14).

106 While NLRC4 activation confers protection to various pathogens, including
107 *Salmonella* Typhimurium, *Burkholderia pseudomallei* and *Legionella pneumophila* (1, 6), it

also drives host susceptibility to the opportunist bacteria *Pseudomonas aeruginosa* (*P. aeruginosa*) (17, 18). In particular, *P. aeruginosa* T3SS and flagellin components induce NLRC4-mediated IL-1 β and IL-18 secretion, which inhibits protection mediated by both Th17 cells and anti-microbial peptides produced by airway epithelial cells (17, 18). This apparent paradox underlies complex host-microbe interactions, as NLRC4 confers protection against other pathogens that express T3SS and flagellin. Here, we examined the underlying host-microbe mechanisms through which NLRC4 specifically drives susceptibility to *P. aeruginosa*. We used *Salmonella enterica* serovar Typhimurium (SL1344) and PAO1, a *P. aeruginosa* strain, that expresses a functional T3SS (T3SS⁺) in parallel with an isogenic mutant of this strain, inactivated in the T3SS transcriptional regulator *exsA*, unable to express a functional T3SS apparatus (T3SS⁻).

Results

Bacterial localization induces differential pyroptosis-dependent cell-autonomous response in macrophages.

To decipher the molecular mechanisms underlying NLRC4-mediated sensing of the bacterial pathogen *P. aeruginosa*, we infected bone marrow-derived macrophages (BMDMs) from wild-type (WT) and *Nlrc4*^{-/-} mice with either T3SS-expressing (T3SS⁺) *P. aeruginosa* and *S. Typhimurium* strains (19) and we measured the capacity of both strains to trigger NLRC4 inflammasome-dependent pyroptotic cell death and IL-1 β release. In agreement with previous published reports, these data indicate that both bacteria triggered cell-death in a T3SS and NLRC4-dependent manner within 3 hours of infection (MOI 15) (**Fig. 1A, Fig. S1A**).

While *S. Typhimurium* and *P. aeruginosa* have a different niche tropism (intracellular for *Salmonella* and extracellular for *Pseudomonas*), we hypothesized that the NLRC4 response might lead to different bacterial fate in macrophages. Consequently, CFU assays

134 after 1 hour of infection (MOI 15) found more intracellular *S. Typhimurium* than *P.*
135 *aeruginosa* in WT BMDMs (**Fig. 1B**). Strikingly, such differences in intracellular numbers of
136 *P. aeruginosa* was partially lost in *Nlrc4*^{-/-} macrophages and independent from *P.*
137 *aeruginosa* T3SS-injected exoenzymes STY (**Fig. 1B, Fig. S1B**), suggesting that the
138 NLRC4 inflammasome response contributed to *P. aeruginosa* avoiding macrophage
139 uptake. Whereas *Salmonella* could efficiently establish an intracellular replicative niche in
140 *Nlrc4*^{-/-} BMDMs, intracellular *P. aeruginosa* failed to do so after 24 hours of infection (**Fig.**
141 **1C**), which suggests that *P. aeruginosa* exploits NLRC4-dependent response to avoid
142 macrophage intracellular-autonomous immunity. Phagocytosis is the main process by
143 which macrophages engulf bacteria. Hence, inhibiting phagocytosis by cytochalasin D
144 reduced NLRC4-dependent cell death and IL-1 β release upon infection with *S.*
145 *Typhimurium* but not with *P. aeruginosa* (**Fig. 1D**). Remarkably, T3SS⁺-induced
146 phagocytosis-independent activation of the NLRC4 inflammasome was specific to only *P.*
147 *aeruginosa*, as other T3SS-expressing bacteria triggered NLRC4 inflammasome
148 responses in a phagocytosis-dependent manner (**Fig. S1C**). To rule out that actin
149 polymerization might directly control NLRC4 activation, we electroporated purified flagellin
150 in WT and *Nlrc4*^{-/-} BMDMs in presence or absence of cytochalasin D. Cell death evaluation
151 showed that cytochalasin D did not modify cytosolic flagellin-induced NLRC4
152 inflammasome response (**Fig. S1D**). Thus, these results identified the unique capability of
153 T3SS-expressing *P. aeruginosa* to promote NLRC4-inflammasome activation in a
154 phagocytosis-independent manner.

155 “Pore-induced Intracellular Traps” (PITs) entrap intracellular *S. Typhimurium* (11–13). As
156 *P.aeruginosa* induced phagocytosis-independent activation of the NLRC4 inflammasome,
157 we reasoned that such process might be a virulence strategy to escape bacterial caging
158 into PITs. Therefore, we infected primary WT BMDMs with both *P. aeruginosa* and *S.*
159 *Typhimurium* strains (MOI 3) for 1h to induce PIT formation and evaluated the number of
160 bacteria associated with pyroptotic structures (e.g. ASC⁺ cells). Strikingly, our results

found higher amount of *Salmonella* associated within PIT structures than *P. aeruginosa*, showing that *P. aeruginosa*-activated NLRC4 inflammasome allowed escape from PIT-driven intracellular trapping (**Fig. 1E, F**).

Then, we aimed to determine whether pyroptosis have a differential *in vivo* regulatory function after *P. aeruginosa* and *S. Typhimurium* exposure. Using a peritoneal mouse model of infection (3×10^6 CFUs, 6h) with either strains, we evaluated the contribution of pyroptosis on the early clearance of each bacterial strain in WT or in *gasderminD*^{-/-} (*gsdmD*^{-/-}) mice, which are deficient at inducing pyroptosis. Therefore, mice lacking *gsdmD* showed an early (6h) susceptibility to STm challenge whereas they presented improved resistance to *P. aeruginosa* infection (**Fig. S1F**). Overall, these results demonstrated that *P. aeruginosa* exploited the NLRC4 inflammasome to escape pyroptosis-mediated capture and sequestration, a process that was reversed in absence of *Nlrc4*.

***Pseudomonas* triggers T3SS-independent NLRC4 and Caspase11 inflammasome activation in macrophages.**

Our results showed that both *S. Typhimurium* and *P. aeruginosa*-induced differential PIT-dependent bacterial trapping. Yet, we wanted to rule out that the observed phenotype could be driven by intrinsic properties of each bacterial strain, such as flagellin and T3SS immunogenicity and/or expression levels.

Therefore, we relied on a surprising observation where we found that infection of macrophages with a T3SS-deficient strain of *P. aeruginosa* still triggered early (3 hours) residual NLRC4 response and, late (4-10 hours) caspase11 non-canonical inflammasome pathway (**Fig. 2A; Figs. S2A, B**). Even though high doses (MOI 25, 50 or 100) of T3SS⁻ *P. aeruginosa* were necessary to induce reduced levels of NLRC4 activation, residual inflammasome response was both still NLRC4- and -flagellin dependent, as T3SS⁻/*fliC* strains failed to elicit measurable IL-1 β release as well as caspase-1 and gasdermin-D

(GSDMD) processing (**Figs. 2A, B; Fig. S2C**). Furthermore, we noted that flagellin complementation in both T3SS⁺ and T3SS⁻ strains restored NLRC4-driven cell death and IL-1 β release (**Figs. S2D, E**). However, only live T3SS-deficient *P. aeruginosa* induced NLRC4-dependent IL-1 β release as it was not detected using heat-killed (HK) and PFA-killed bacteria (20) (**Fig. S2F**). To exclude that genetic inactivation of *exsA* allowed residual T3SS expression, we tested *P. aeruginosa* lacking PscC, a key T3SS structural component (21). Importantly, *pscC* and *exsA*-deficient bacteria triggered comparable levels of NLRC4-dependent cell death and IL-1 β release (**Fig. S2G**). These results indicate that activation of the NLRC4 inflammasome was not caused by residual T3SS expression.

In the host, cytosolic NAIP proteins bind flagellin or T3SS components and initiate NLRC4 oligomerization (6). To confirm direct involvement of flagellin in NLRC4 activation, we infected macrophages lacking either *Naip5* (*Naip5*^{-/-}), the main flagellin sensor, or the 5 different *Naips* (*Naip* ^{Δ 1-6}), with either T3SS⁺ or T3SS⁻ strain. Both *Naip5*^{-/-} and *Naip* ^{Δ 1-6} BMDMs failed to undergo pyroptosis and release IL-1 β when infected with either strain (**Fig. 2C**). Based on these results, we conclude that flagellin directly triggered the NLRC4 inflammasome response during T3SS⁻ *P. aeruginosa* infection.

Altogether, these findings reveal, in addition to the Caspase-11 non canonical inflammasome, an unrecognized and unexpected capacity of *P. aeruginosa* to alternatively activate the NAIP5/NLRC4 inflammasome in a T3SS-independent yet flagellin-dependent manner. These findings further suggest that flagellin can reach the host cell cytosol independently of bacterial secretion systems.

210

T3SS⁻ *P. aeruginosa*-containing damaged phagosomes associate to sequential activation of NLRC4 inflammasome and Caspase-11.

While our results indicated that flagellin and LPS reach the host cytosol in absence of a functional T3SS, the underlying mechanism still remained unknown. We hypothesized

215 that a key candidate involves direct entry of *P. aeruginosa* into the intracellular
 216 compartment. Therefore we evaluated whether T3SS-independent activation of both
 217 NAIP5/NLRC4 and Caspase-11 inflammasomes required *P. aeruginosa* phagocytosis.
 218 Consequently, infection of macrophages with T3SS⁻ *P. aeruginosa* showed that
 219 phagocytosis inhibition abrogated cell death, IL-1 β release and Caspase-1 processing
 220 (**Fig. 3A-C**). Hence, these results show that, in absence of a functional T3SS, *P.*
 221 *aeruginosa* induces inflammasome response in a phagocytosis-dependent manner. As
 222 phagocytosis of T3SS-deficient *P. aeruginosa* is a prerequisite for inflammasome
 223 response, we hypothesized that NAIP5/NLRC4 and Caspase11 responses required that
 224 both flagellin and LPS reached the cytosol by leaking from *P. aeruginosa*-containing
 225 phagosomes. To visualize phagosomal membrane alterations, we probed for galectin-3
 226 (GAL3), a lectin that binds galactosides on permeabilized host cell endovesicles (22). If *P.*
 227 *aeruginosa*-containing phagosomes were compromised, we expected GAL3 recruitment
 228 around bacteria. To avoid unwanted GAL3 recruitment to permeabilized phagosomes from
 229 pyroptosis and not *P. aeruginosa* specifically, we infected immortalized *Casp1*^{-/-}/*11*^{-/-}
 230 macrophages with two T3SS mutants (e.g. T3SS⁻ and T3SS/*fliC*) for 3 hours. Confocal
 231 microscopy revealed that GAL3 did target a small proportion (~ 6-10%) of intracellular
 232 T3SS⁻ and T3SS/*fliC* bacteria (**Fig. 3D**). We then reasoned that cells with active
 233 inflammasomes, *i.e.* containing ASC specks, should also contain at least one GAL3-
 234 positive bacterium. Using macrophages expressing an active inflammasome (ASC
 235 specks⁺), we quantified the percentage of cells that had at least one GAL3- positive *P.*
 236 *aeruginosa*. We consistently found that approximately 45% of *iCasp1*^{-/-}/*11*^{-/-}
 237 macrophages presenting an ASC speck were also positive for GAL3-stained T3SS⁻ *P.*
 238 *aeruginosa* (**Figs. 3D, E**), but we did not detect any specks using T3SS/*fliC* strain (**Fig.**
 239 **3E**). *Casp1*^{-/-}/*11*^{-/-} deficient BMDMs can form a NLRC4 inflammasome that recruit
 240 caspase-8 that also triggers cell death. To verify that in *Casp1*^{-/-}/*11*^{-/-} iBMDMs, GAL3
 241 recruitment to intracellular bacteria did not require caspase-8, we removed *Nlrc4* in GAL3-

242 mcherry expressing *Casp1^{-/-}/11^{-/-}* iBMDMs (**Fig. S3A**) and quantified GAL3 recruitment to
243 T3SS⁻ *P. aeruginosa*. We did not detect any defect for bacteria targeted by GAL3-
244 mecherry in the *Nlrc4^{-/-}/Casp1^{-/-}/11^{-/-}* iBMDMs, hence confirming that T3SS⁻ *P.*
245 *aeruginosa* is targeted by GAL3 independently of the NLRC4-dependent pathway (**Fig.**
246 **S3B**). Overall, these results show that T3SS⁻ *P. aeruginosa* trigger NLRC4 inflammasome
247 from altered phagosomes.

248 Given that T3SS⁻ *P. aeruginosa* were exposed to the cytosol, we speculated that
249 phagosome alterations could also expose *P. aeruginosa* LPS to the host cell cytosol,
250 which might promote caspase11 recruitment and activation. Thurston et al. showed that
251 caspase11 was recruited to cytosolic Salmonella in infected epithelial cells (25). So, we
252 sought to determine whether caspase11 associated with compromised *P. aeruginosa*-
253 containing phagosomes. Microscopy observations of galectin-3+ (GAL3+) bacteria in IFN γ -
254 primed *Nlrc4^{-/-}* iBMDMs showed recruitment of both GAL3 and active caspase on T3SS⁻
255 *P. aeruginosa* after 3 hours of infection, which was not observed in *Nlrc4^{-/-}/Casp11^{-/-}*
256 CRISPR-Cas9 iBMDMs (**Fig. 3F**). These results confirm that bacteria accessible to the cell
257 cytosol directly recruited caspase-11.

258 Altogether, these results indicate that both NLRC4 and Caspase11 directly detected
259 intracellular T3SS⁻ *P. aeruginosa* flagellin and LPS from compromised phagosomes.

260

261 **Pyroptosis-induced PITs is efficient only against T3SS-deficient *Pseudomonas*.**

262 As we previously showed that T3SS-expressing *P. aeruginosa* and *S. Typhimurium*
263 triggered differential PIT responses in macrophages, we evaluated whether pyroptosis-
264 induced PITs also promoted restriction of intracellular *P. aeruginosa*. Primary WT BMDMs
265 were infected with T3SS⁺ or T3SS⁻ for 3 h to induce PIT formation. Time-lapse microscopy
266 revealed that cells with compromised plasma membranes (e.g. positive for the DNA
267 impermeant binding probe TO-PRO-3) contained a high number of intracellular T3SS⁻ *P.*
268 *aeruginosa*, while few T3SS⁺ *P. aeruginosa* were associated with dead cells (**Movie 1 that**

269 **refers to T3SS⁺ PAO1-infected cells, Movie 2 that refers to T3SS⁻PAO1-infected**
 270 **cells).** In addition, confocal microscopy experiments showed that PITs (*i.e.* ASC⁺ cells)
 271 were enriched specifically with T3SS⁻ bacteria (**Fig. 4A, B**). Based on these results, we
 272 conclude that T3SS-expressing *P. aeruginosa* hijacked PIT-mediated intracellular bacterial
 273 entrapment and weakening, a process reversed in the absence of T3SS expression.

274 The capability of PITs to entrap mostly T3SS-deficient bacteria suggested that
 275 these structures might also have a direct or an indirect microbicidal function against these
 276 strains, as suggested by *S. Typhimurium* infection (13, 15). We then infected primary WT
 277 BMDMs with either T3SS⁺ or T3SS⁻ *P. aeruginosa* for 3h and monitored the microbicidal
 278 potential of PITs. PITs-associated or –unbound bacteria were harvested and exposed to
 279 purified neutrophils or PBS for 1h30. We quantified CFUs, which showed that PITs-
 280 associated bacteria specifically were more susceptible to neutrophil killing than the PITs-
 281 unbound fraction (**Fig. 4C**). Since phagocytosis of both T3SS⁺ and T3SS⁻ *P. aeruginosa*
 282 was comparable in *Nlrc4* deficient BMDMs (**Fig. 4D**), we hypothesized that pyroptosis
 283 induction in these cells would weaken both bacteria to the same extent. We infected *Nlrc4*
 284 ^{-/-} BMDMs with both T3SS⁺ and T3SS⁻ *P. aeruginosa* strains and subjected them to ATP
 285 stimulation to ensure similar NLRP3 inflammasome-dependent pyroptosis induction in both
 286 conditions. Intracellular bacteria were harvested, counted and exposed to a secondary
 287 stress signal (*i.e.* H₂O₂). Both intracellular *P. aeruginosa* strains showed the same
 288 increased susceptibility to H₂O₂ (**Fig. 4E**), thus demonstrating that PIT-driven intracellular
 289 trapping was only efficient against T3SS-deficient *P. aeruginosa*. Finally, we performed a
 290 peritoneal mouse model of infection (3x10⁶ CFUs, 4h) with GFP-expressing T3SS⁺ or
 291 T3SS⁻ *P. aeruginosa*, and evaluated bacteria-containing dead macrophages efferocytosis
 292 by neutrophils (13). Strickingly, recruited neutrophils (Ly6G⁺) having efferocytosed
 293 bacteria-containing macrophages (Ly6G⁺/F4/80⁺/GFP⁺) was more efficient against T3SS⁻
 294 *P. aeruginosa* than to T3SS⁺ bacteria (**Fig. 4E; Fig. S4A**). Overall, these results

295 demonstrated that *P. aeruginosa* used its T3SS to escape PIT-mediated capture and
 296 sequestration. This process was reversed in absence of T3SS-expression or of *Nlrc4*.
 297
 298 **Both NLRC4 and Caspase-11 specifically protect against acute infection with T3SS-**
 299 **deficient *P. aeruginosa*.**

300 To evaluate the *in vivo* relevance of our findings, we infected WT and *Nlrc4*^{-/-} mice
 301 with either T3SS⁺ or T3SS⁻ *P. aeruginosa*. Because T3SS⁻ *P. aeruginosa* is highly
 302 susceptible to immune defenses (23), we infected mice with a higher dose of this strain.
 303 When infected for 18h with T3SS⁺ *P. aeruginosa*, *Nlrc4*^{-/-} mice presented lowered
 304 bacterial loads in the bronchoalveolar lavage fluid (BALF) than their WT counterparts (**Fig.**
 305 **5A**), consistent with a prior report (18). Remarkably, *Nlrc4*^{-/-} mice infected with the T3SS⁻
 306 strain showed an increased BALF bacterial loads compared to WT controls after 18h of
 307 infection (**Fig. 5A**). The use of T3SS⁻/*fliC*⁻ *P. aeruginosa* did not show any involvement of
 308 NLRC4 on bacterial loads in BALFs, confirming that flagellin was the principal component
 309 that mediates *in vivo* NLRC4 response to T3SS-deficient *P. aeruginosa* (**Fig. S5A**).

310 While strongly reduced, the residual IL-1 β levels found in BALFs of mice infected
 311 with the T3SS-deficient *P. aeruginosa* strain remained partially dependent on the NLRC4
 312 inflammasome (**Fig. 5B**), whereas infection with T3SS⁻/*fliC*⁻ bacteria showed no NLRC4-
 313 dependent IL-1 β release (**Fig. S5B**). Since we previously showed that caspase-11 could
 314 also detect intracellular *P. aeruginosa*, we speculated about the potential immune
 315 importance of this pathway *in vivo*. Although *Casp11* deficient mice did not show
 316 significant involvement of caspase-11 at controlling the T3SS⁺ strain, T3SS⁻ and T3SS⁻
 317 /*fliC*⁻ bacterial infection induced a higher susceptibility in these mice than their WT
 318 counterparts (**Figs. 5C, D; Figs. S5C, SD**). This result verified that both NLRC4 and
 319 caspase-11 play paired protective roles only against T3SS-deficient *P. aeruginosa*.

320 After airway infection, *P. aeruginosa* will encounter alveolar macrophages as the
 321 first phagocytic cells. So, we evaluated the importance of alveolar macrophages on

inflammasome-triggered differential host responses to *P. aeruginosa*. Mice infected with T3SS⁺ and T3SS⁻ *P. aeruginosa* showed that alveolar phagocyte depletion (**Fig. 5E**) increased alveolar loads of T3SS⁻ *P. aeruginosa* (**Fig. S5E**) and strongly impaired IL-1 β release in the alveoli (**Fig. 5F**). These results demonstrated that alveolar macrophages controlled inflammasome-mediated immune responses to *P. aeruginosa* in mice.

Overall, these results demonstrate that T3SS-expressing *P. aeruginosa* exploited the NLRC4-dependent response to their own advantage, but T3SS deficiency uncovered host protective NLRC4- and caspase-11 responses.

Discussion

Although the NAIP-NLRC4 inflammasome primarily controls intracellularly adapted bacteria *Salmonella* or *Legionella* spp, several studies indicate the critical function of the T3SS-flagellin complex in mediating NLRC4-dependent host susceptibility to *P. aeruginosa* infection is still discussed (17, 18, 24)(25). Regarding *P. aeruginosa* infection, the deleterious functions of both IL-1 β and IL-18 on immune control of T3SS-expressing *P. aeruginosa* infection is well documented (18, 26), yet the role of pyroptosis in this process remains unclear.

Here, we provide evidence that NLRC4-dependent pyroptosis of macrophages drive differential outcomes of T3SS-expressing *P. aeruginosa* and *S. Typhimurium*. Whereas *S. Typhimurium* remains entrapped into PITs, as previously demonstrated (13, 16), *P. aeruginosa*-induced extracellular activation of the NLRC4 inflammasome allows bacterial escape from macrophage and pyroptosis-driven cell-autonomous immunity. To our knowledge, such response is unique to *P. aeruginosa*, as other T3SS-expressing bacteria also required uptake to activate the NLRC4 inflammasome. A recent study found that *P. aeruginosa* T3SS triggered a Caspase-3/-7 deleterious host response in an ASC-dependent manner (27). Therefore, it is tempting to speculate that a similar mechanism

underlying T3SS-expressing *P. aeruginosa*-induced cell apoptosis could also mediate bacterial escape from host immune responses.

In addition, we uncovered that a T3SS-deficient strain of *P. aeruginosa* also triggered residual, but of immune importance, NAIP5/NLRC4 inflammasome, in a T3SS-independent manner, which is in agreement with findings of Faure et al., where lethal challenge of T3SS-deficient *P. aeruginosa* still induce a NLRC4-dependent response in mice. In addition, T3SS-expressing *P. aeruginosa* triggered inflammasome assembly through a process that did not heavily utilize phagocytosis (28–30). We do not believe that the presence of a genetically encoded phagosomal permeabilization system in *P. aeruginosa* is likely, as only a minority (~6-10%) of bacteria were accessible to cytosolic galectin-3. Another explanation could involve phagosome maturation-induced *P. aeruginosa* local production of outer-membrane vesicles (OMVs), which could expose bacterial ligands to host cell cytosol sensors (31–34). Conversely, *Nlrc4* deficiency or long-term infection of macrophages promoted a switch in the NLRC4 response to a LPS-induced caspase-11 non-canonical inflammasome path. Although previous results from others and our group indicated that OMVs from *P. aeruginosa* induced non-canonical inflammasome pathway in a phagocytosis-independent, yet endocytosis-dependent manner, our results here demonstrated that T3SS-deficient *P. aeruginosa*-triggered caspase-11 non-canonical inflammasome responses required bacterial uptake (31–34). Future investigations will be critical to determine the specific roles of OMVs and intracellular *P. aeruginosa* for nucleation of the non-canonical inflammasome route. Immunodeficiency, burn-, nosocomial- and ventilator-associated injuries render patients susceptible to chronic *P. aeruginosa* infection, in which most strains are deficient for T3SS and/or flagellin expression (35, 36). A recent report suggested that T3SS⁻ *P. aeruginosa* promoted the non-canonical inflammasome pathway in an IFN- and GBP-dependent manner (37). Consistent with these data, we also showed that caspase-11 protected mice from acute infection with lethal doses of T3SS-deficient *P. aeruginosa*.

375 The *in vivo* function of NLRC4 to *P. aeruginosa* infection remains unresolved as it
 376 can be either protective or deleterious to the host in mouse models (24, 38). Our mouse
 377 model of acute infection revealed a deleterious role of NLRC4 on host defenses during
 378 exposure to T3SS-expressing *P. aeruginosa* (17, 18). Importantly, we demonstrated that
 379 *P. aeruginosa* T3SS deficiency induced both NLRC4- and caspase-11-driven host
 380 protection during infection. Attree's group analyzed isolated naturally T3SS deficient
 381 clinical isolates from patients (39). Some strains expressed flagellin and were motile (40).
 382 These findings warrant investigating whether these bacteria also induce a NLRC4
 383 dependent response.

384 Overall, our study uncovered that T3SS-expressing *P. aeruginosa* remained
 385 extracellular by triggering cell pyroptosis, a process that enabled bacteria to overwhelm
 386 macrophage PITs-induced host immunity. We further demonstrated that phagocytosis of
 387 *P. aeruginosa* that lacked T3SS expression triggered host protective pyroptosis in a T3SS-
 388 dependent manner through both NLRC4 and caspase-11 inflammasomes. In summary, we
 389 propose that bacterial localization drives the effectiveness of the inflammasome response
 390 during bacterial infection (**Graphical abstract**). Future studies will determine if this is a
 391 conserved response among bacterial pathogens.

392

393

394

395

396

397

398

399

400 **Methods**

401 All reagents and biological samples used in this study are listed in the **table S1**

402 **Mice**

403 *Casp11^{-/-}*, *Casp1^{-/-}/Casp11^{-/-}*, *Nlr4^{-/-}*, *Nlrp3^{-/-}* and *Gsdmd^{-/-}* mice have been
404 previously described (41–43). Mice were bred in the animal facilities of the University of
405 Basel (Basel, Switzerland) or at the IPBS institute (Toulouse, France). Bones from *Naip5^{-/-}*
406 and *Naip^{Δ1-6}* mice were a kind gift from R. E Vance (UC Berkeley, USA) (43). Janvier and
407 Charles Rivers companies provided WT mice.

409 **Animal infections**

410 6-8 age and sex-matched animals (8–10 weeks old) per group were infected
411 intranasally with 5×10^6 (PAO1 T3SS⁺) or 1.5×10^7 (PAO1 T3SS⁻ or T3SS⁻/*fliC*) CFUs of
412 mid-late exponential phase *Pseudomonas aeruginosa* in 40μl PBS. Animals were
413 sacrificed 18h after infection and bronchoalveolar fluids (BALFs) were collected in PBS.
414 When specified, cellular contents (flow cytometry), bacterial loads (serial dilutions and
415 CFU plating) and cytokine levels (ELISA) were evaluated. No randomization or blinding
416 were done.

418 **Clodronate-induced alveolar phagocyte depletion**

419 C57BL/6 mice received intranasal instillation of 40μL of either clodronate or PBS
420 loaded liposomes to deplete alveolar macrophages (18). 48 h later, mice were intranasally
421 infected with either 5×10^6 (PAO1 T3SS⁺) or 1.5×10^7 (PAO1 T3SS⁻) CFUs of mid-late
422 exponential phase *Pseudomonas aeruginosa* (OD of 1.0-1.6) in 40μl PBS. 18 h after
423 infection, BALFs were collected and bacterial loads (serial dilutions and CFU plating),
424 cytokine levels (ELISA) and immune cell contents were evaluated. Briefly, cells were
425 pelleted (1000 rpm, 5 minutes) and alveolar macrophages were subsequently stained with

a cocktail of fluorochrome-conjugated antibodies detailed in the material section. Cells were then fixed in 4% PFA before fluorescence associated cell sorting (FACS) analysis using a LSRII instrument (BD Biosciences). Data analysis and processing were performed using FlowJo.10 software. The plot FSC-A vs FSC-H was used to discriminate doublets from single cells. Alveolar macrophages were defined as CD11c⁺/F480⁺ in Ly6C⁻/Ly6G⁻/CD19⁻/TCRβ⁻ population.

432

433 **Genetic invalidation of *Nlr4* and *Caspase11* genes in *iCasp1^{-/-}/Casp11^{-/-}* and *iNlr4^{-/-}*** 434 ***immortalized BMDMs***

Nlr4 was knocked-out using the crispr/cas9 system in *iCasp1^{-/-}/Casp11^{-/-}* macrophages and *Casp11* was knocked-out from *iNlr4^{-/-}* macrophages. Single guide RNAs (sgRNA) specifically targeting *Nlr4* exon3 (Forward: 5'CACCGTTACTGTGAGCCCTTGGAGC3' reverse: 5'AAACGCTCCAAGGGCTCACAGTAA-C3') and *Caspase-11* exon 2 forward (5'CACCGCTTAAGGTGTTGGAACAGCT3') reverse (5'AAACAGCTGTTCCAACACCTTAAGC3') were designed using Benchling tool (Benchling.com), and oligonucleotides were synthesized by Sigma-Aldrich. The gene-specific crispr guide RNA oligonucleotides were hybridized and cloned in Lenti-gRNA-Puromycin vector using BsmBI restriction sites (lentiGuide-Puro, Addgene 52963, from Feng Zhang lab). These constructs were transfected (using lipofectamine 2000) into HEK293T cells together with the lentiviral packaging vector p8.91 (from Didier Trono lab, EPFL, Switzerland) and a envelop VSVg-encoding vector (pMD.2G, Addgene 12259, from Didier Trono lab) for 48 h. Then, viral supernatants were harvested, filtered on 0.45 μm filter. Cas9-expressing recipient-cells either *iCasp1^{-/-}/Casp11^{-/-}* or *iNlr4^{-/-}* (1,000,000 cells/well in 6-well plates) were generated by lentiviral transduction with Cas9-expressing lentiviral vector (lentiCas9-Blast, Addgene 52962, from Feng Zhang lab) and then infected with the lenti-Guide viral particles in presence of 8μg/ml polybrene and centrifugated for 2 h at 2900 rpm at 32°C. 48 h later, medium was replaced and Puromycin selection

453 (10µg/mL) was applied to select positive clones for two weeks. Puromycin-resistant cells
454 were sorted at the single cell level by FACS (Aria cell sorter). Individual clones were
455 subjected to western blotting to confirm the absence of either *Nlrc4* or *Caspase-11* gene
456 products. To ensure clonal reproducibility, at least 2 positive clones were compared for
457 inflammasome response.

458

459 **Cloning and cell transduction**

460 Galectin-3-morange construct was a kind gift from J. Enninga (Institut Pasteur,
461 Paris, France). Galectin-3-morange coding sequence was sub-cloned into the pMSCV2.2
462 plasmid by first excising EGFP at the EcoRI sites followed by insertion using NotI/XhoI
463 sites. Immortalized WT, *Nlrc4*^{-/-}, *Casp1*^{-/-}/*Casp11*^{-/-} or *Nlrc4*^{-/-}/*Casp11*^{-/-} BMDMs were
464 then transduced with retroviral particles, positive cells for morange fluorescence were
465 FACS to allow clonal selection.

466

467 **Cell culture and infections.**

468 Bone-marrow derived macrophages (BMDMs) were differentiated in DMEM
469 (Invitrogen) with 10% v/v FCS (Thermo Fisher Scientific), 10% MCSF (L929 cell
470 supernatant), 10 mM HEPES (Invitrogen), and nonessential amino acids (Invitrogen).
471 BMDMs were seeded in 6-, 24-, or 96-well-plates at a density of 1.25x10⁶, 2.5x10⁵, or
472 5x10⁴ per well. When required BMDMs were pre-stimulated overnight with 100ng/mL of
473 PAM3CSK4 (InvivoGen). For infections with *Pseudomonas* strains, bacteria were grown
474 overnight in Luria Broth (LB) at 37 °C with aeration. Bacterial cultures were diluted 1/50
475 and grew until mid-late exponential phase (OD of 1.0-1.6) and added to the macrophages
476 at multiplicity of infection (MOI) of 25, 50 and 100, in serum and antibiotic-free medium
477 (OPTIMEM), or as otherwise indicated. Then, to ensure homogenous infections, plates
478 were centrifuged for 1 minute, 800 rpm. 1 h after infection, extracellular bacteria were
479 killed by adding gentamicin (100µg/ml, Invitrogen). When specified, paraformaldehyde

(PFA)-killed (PFA 4%, 20 minutes) or heat-killed (95°C, 15 minutes) of *P. aeruginosa* strains were used to infect BMDMs at MOI of 50. When required, *Shigella flexneri* (M90T), *Chromobacter violaceum* or various *Salmonella* strains (SL1344) were grown in LB in the presence or absence of antibiotics (specified in the resource table) at 37°C with constant agitation overnight. To ensure proper T3SS and flagellin expression, bacteria were sub-cultured the next day in LB media for 3 h until reaching an OD of 0,6-1.

When required, Flagellin was electroporated with Neon™ Transfection System (ThermoFischer) according manufacturer's protocol. Briefly, 5 x 10⁵ cells were resuspended in Buffer R and 0.5 ug/ml Flagellin was electroporated in 10 µl tips using 2 pulses of 1720V and 10 width. Cells were then plated in 24 well-plates.

Transfection of cells with Fagellin (Invivogen, 0, 5µg/mL/2,5x10⁵ cells) or LPS (O111:B4, *E. coli*, 1µg/2,5x10⁵ cells Invivogen) was achieved using FuGeneHD (Promega) transfection reagent in Opti-MEM culture medium (44).

493

494 **Bone marrow Neutrophil isolation**

Neutrophils were isolated from bone-marrow cells of WT mice by positive selection using Anti-Ly6G MicroBeads UltraPure isolation kit (Miltenyi-Biotec, Anti-Ly6G MicroBeads UltraPure mouse) according to the manufacturer's instructions. Characterization of the purified population by Fluorescence Associated Cell Sorting showed an enrichment of more than 98% of Ly6G^{high} cells.

500

501 **Cytokine and pyroptosis measurement**

IL-1β (eBioscience) was measured by ELISA. Cell-death was assayed using LDH Cytotoxicity Detection Kit (Takara). To normalize for spontaneous lysis, the percentage of LDH release was calculated as follows: (LDH infected – LDH uninfected)/(LDH total lysis – LDH uninfected)*100.

506

507 **Western blotting**

508 Cell and supernatant protein lysates were prepared as previously described. Antibodies
509 used were mouse anti-mouse caspase-1 antibody (Casper, Addipogen), goat anti-mouse
510 IL-1 β antibody (AF-401-NA; R&D Systems), anti- β -actin antibody (A1978; Sigma), anti-
511 GAPDH (GTX100118; GeneTex), anti gasdermin-D (Abcam, ab209845), anti-caspase-11
512 (Novus, NB120-10454), anti-NLRC4 (Abcam, ab201792) and anti-NLRP3 (AdipoGen, AG-
513 20B-0014). After incubation with primary antibody the membrane was washed 3 times with
514 TTBS, and then Immunoreactive bands were detected by incubation for 2 h with the
515 appropriate secondary antibodies conjugated with horseradish peroxidase (Diagomics).
516 Proteins of interest were visualized using ECL substrate (Biorad) and images were
517 acquired using ChemiDoc Imaging System (Biorad). Working dilutions of the antibodies
518 are listed in **table 1**.

519

520 **Intracellular CFU experiments**

521 2.5×10^5 BMDMs were infected with various PAO1 strains at indicated MOIs for
522 various times. For CFU assays after 24 hours stimulation, gentamicin protection assay
523 was performed using gentamicin at 50 μ g/mL after 2 hours infections of macrophages.
524 Supernatants were removed, and cells washed 5 times with PBS. Cells were then lysed in
525 triton x-100 (Sigma), 0.1% and plated at 37°C overnight for CFUs numeration on LB-agar.

526

527 **Microscopy**

528 2.5×10^5 BMDMs were seeded on glass coverslips and infected as described above.
529 When desired, wells were washed three times with PBS and fixed with 4% PFA for 10
530 minutes at 37°C. Excess of PFA was removed by PBS washes and quenching using 0,1M
531 Glycine for 10 min at room temperature. Permeabilization and primary antibody staining
532 were performed O/N at 4°C using Saponin 0.1%/BSA 3% solution. Stainings were
533 performed using Hoescht (DNA labeling), rabbit anti-ASC (SCBT, 1/500), chicken anti-

534 *P.aeruginosa* LPS (1/500, Agrisera) or anti-CD45-2-APC (1/250, Biolegends). Coverslips
535 were washed with Saponin/BSA solution and incubated with the appropriate secondary
536 antibodies (1/500, Diagnostics). Cells were then washed 3 times with PBS and mounted on
537 glass slides using Vectashield (Vectalabs). Coverslips were imaged using confocal Zeiss
538 LSM 710 (Image Core facility, Biozentrum, Basel and CPTP, Toulouse) or an
539 Olympus/Andor CSU-X1 Spinning disk microscope (Image core Facility, IPBS, Toulouse)
540 using a 63x objective. Unless specified, for each experiment, 5-10 fields (~50-250 cells)
541 were manually counted using Image J software.

542

543 **Time-lapse microscopy**

544 5x10⁵ iBMDMs Gal3-morange expressing cells were seeded in a 35 mm microdish
545 (Ibidi). 1 hour before infection, cells were pre-stained with 0.1 µM CFSE (Biolegend) and
546 infected with T3SS⁺ or T3SS⁻ *P. aeruginosa* with a MOI of 25 or 50. 1µM of membrane
547 impermeant fluorescent DNA binding probe TO-PRO[®]-3 (ThermoFisher) was also added
548 to the culture medium. Live cell imaging was performed using an Olympus/Andor CSU-X1
549 spinning disk microscope with a picture frequency of 5 minutes for 3-4 hours and videos
550 were reconstructed using Image J software.

551

552 **PIT experiments**

553 We have set up a modified form of a published protocol (13). Unless otherwise
554 specified, WT BMDMs were infected using an MOI of 50 PAO1 T3SS⁺ or T3SS⁻ for 3 h.
555 Intracellular bacteria-associated to PITs were visualized and manually quantified by
556 microscopy (Olympus/Andor CSU-X1 spinning disk or inverted confocal microscope Zeiss
557 LSM 710, ~50 cells counted, 5-10 fields). For bacterial colony forming units number
558 enumeration, cells were lysed in Triton X-100, 0.1% and plated on Luria Broth agar.

559

560

561 **Bacterial response to a secondary stress signal**

562 A modified version of published protocol was used (13). WT BMDMs were infected
563 for 2 h with PAO1 T3SS⁺ or T3SS⁻ using MOI of 50. Then, both extracellular and PIT-
564 associated bacteria were harvested. As intracellular bacteria recovery required cell lysis
565 with TritonX-100 0.1%, extracellular bacteria were also incubated with TritonX-100 0.1%
566 for 2 minutes. When specified, bacteria from cells and supernatants were added to 1mL
567 LB supplemented with 50μM Hydrogen Peroxide (H₂O₂) or PBS for 45 minutes, at 37°C
568 and then plated on LB agar plates.

569 For neutrophil killing assays, WT BMDMs (2.5x10⁵ cells) were infected for 2 h with
570 PAO1 T3SS⁺ or T3SS⁻ using MOI of 50. Both supernatant- and bacteria-containing PITs
571 were exposed to neutrophils (1x10⁶ cells) for 1.5 h. Then, cells were treated with Triton X-
572 100 0.1% for 2 minutes, and CFU counts were enumerated on LB agar plates.

573

574 **ImageStreamX experiments**

575 WT mice (n=3 for each bacterial strain) were infected intraperitoneally with 3x10⁶ CFUs in
576 PBS (100μL/mouse) of either T3SS⁺ or T3SS⁻ GFP-expressing bacteria. 4 hours later,
577 peritoneal lavages were collected in 2,5mL of PBS. Neutrophils were stained prior to
578 fixation with anti-Ly6G (APC-Vio770, Miltenyi-Biotec Clone: REA526 | Dilution: 1:50).
579 Then, cells were fixed and permeabilized with BD Fixation/Permeabilisation Kit according
580 to the manufacturer's instructions, and macrophages were labeled with anti-F4/80 (BV421)
581 (Biolegend Clone: BM8 | Dilution: 1:100). Data were acquired on ImageStreamX^{MKII}
582 (Amnis) device (CPTP Imaging and Cytometry core facility) and analyzed using IDEAS
583 software v2.6 (Amnis). The gating strategy used to evaluate efferocytosis of bacteria-
584 containing macrophages by Neutrophils was performed as follows: (i) gate was set on cells
585 in focus and (ii) sub-gated on single cells. Then, we gated both on (iii) Ly6G⁺ Neutrophils
586 and on (iv) F4/80⁺ macrophages within Ly6G⁺ population. (v) To distinguish efferocytosis
587 of intact from fragmented macrophages, we created a mask based on the Surface Area of

588 F4/80+ signal. This was applied to Ly6G⁺/F4/80⁺ gate. (vi) The percentage of GFP⁺
589 bacteria within the Ly6G⁺/F4/80⁺ population was visualized and quantified.

590

591 **Bacterial KO generation and complementation**

592 The knockout vector pEXG2 was constructed and used based on the protocol
593 described by Rietsch et al. (45) with the following modifications. Briefly, 700-bp sequences
594 of the flanking regions of the selected gene were amplified by PCR with Q5 high fidelity
595 polymerase (New England Biolabs). Then, the flanking regions were gel purified and
596 inserted into pEXG2 plasmid by Gibson assembly (46). The assembled plasmid was
597 directly transformed into competent SM10λpir using Mix&Go competent cells (Zymo
598 Research Corporation) and plated on selective LB plates containing 50 µg/mL kanamycin
599 and 15 µg/mL gentamicin. The resulting clones were sequenced, and mating was allowed
600 for 4 h with PAO1 strain at 37°C. The mated strains were selected for single cross over on
601 plates containing 15 µg/mL gentamicin and 20 µg/mL Irgasan (removal of *E.coli* SM10
602 strains). The next day, some clones were grown in LB for 4 hours and streaked on 5%
603 sucrose LB plates overnight at 30°C. *P. aeruginosa* clones were then checked by PCR for
604 mutations. For flagellin complementation experiments, JBOC plasmid (homemade) was
605 used to re-express flagellin gene (*fliC*) under its endogenous promoter in *fliC* deficient
606 PAO1 strains. Briefly, PAO1 *fliC* and its promoter were PCR-amplified, purified and
607 integrated in JBOC plasmid. SM10 *E.coli* were transformed with *fliC* plasmid and allowed
608 to conjugate with PAO1 mutant strains as described above. Positive PAO1 strains were
609 selected on Gentamicin-Irgasan plates and checked by subsequent PCRs. All primers
610 were designed with Snapgene software (GSL Biotech LLC).

611

612 **Statistical analysis**

613 Statistical data analysis was performed using Prism 5.0a (GraphPad Software,
614 Inc.). For comparison of two groups (cell death, cytokine secretion, CFU and microscopy-

based counts), we used two-tailed t-test. Bonferroni correction was applied for multiple comparisons to two-way ANOVA statistical analysis. Data are reported as mean with standard deviation (s.d). For animal experiments Mann-Whitney tests were performed. P values are given in figures, NS means non-significant, unless otherwise specified. Significance is specified as *, ** or *** for P-values <0.05, <0.01 or <0.001 respectively.

Data and software

Fiji and Image J software was used for time-lapse and microscopy images processing and analysis. Snappgene software (GSL Biotech LLC) was used to design all primers. Cytometry data analysis and processing were performed using FlowJo.10 software. ImageStreamX data were analyzed using IDEAS software v2.6 (Amnis). Benchling online software was used to design Crispr guides.

Ethics Statement

All animal experiments were approved (License 2535, Kantonales Veterinärämte Basel-Stadt and License APAFIS#8521-2017041008135771, Minister of Research, France) and performed according to local guidelines (Tierschutz-Verordnung, Basel-Stadt and French ethical laws) and the European Union animal protection directive (Directive 2010/63/EU).

Acknowledgements

We thank the animal facilities, image core facilities and FACS core facility from Biozentrum (Univ. of Basel, Switzerland), IPBS (CNRS, Univ. of Toulouse, France) and CPTP (INSERM, Univ. of Toulouse, France) institutes. Specifically, we acknowledge (S. Allard and D. Daviaud from the CPTP imaging facility. Authors acknowledge V.M Dixit (Genentech) (*Nlrc4*^{-/-}, *Casp11*^{-/-}), R. Flavell (*Casp1*^{-/-} *11*^{-/-}), F. Sutterwala (*Nlrc4*^{-/-}) for sharing mice. We also acknowledge J. Enninga (Institut Pasteur, Paris, France) for galectin-3 plasmid share and R.E. Vance and I. Rauch (University of California at Berkeley) for providing bones from mice deficient for *Naip5*^{-/-} and *Naip*^{A1-6}. We also thank Y. Rombouts and G.L. Villarino for manuscript reading and comments. We also thank funding partners. E.E. and E.M. are funded by a FRM “Amorçage Jeunes Equipes” number AJE20151034460. E.M. is also funded by an ATIP-AVENIR grant and an ERC StG INFLAME 804249. P.B. is funded by the Swiss National Science Foundation (310030_165893). We thank Life Science Editors for editing assistance. Authors declare no conflict of interest.

Author Contribution

E.E, R.P, J.B, P.B and E.M designed the study; D.B, A.Coste and O.N provided essential mouse lines, reagents and expertize. E.E, R.P, J.B, P.J.B, A.Colom, O.C, R.F.D, J.V.S, C.C and E.M performed experiments; E.M wrote the manuscript, with contribution from O.N.

References

1. Broz P, Dixit VM. 2016. Inflammasomes: mechanism of assembly, regulation and signalling. *Nat Rev Immunol* 16:407–420.
2. Kayagaki N, Stowe IB, Lee BL, O'Rourke K, Anderson K, Warming S, Cuellar T, Haley B, Roose-Girma M, Phung QT, Liu PS, Lill JR, Li H, Wu J, Kummerfeld S, Zhang J, Lee WP, Snipas SJ, Salvesen GS, Morris LX, Fitzgerald L, Zhang Y, Bertram EM, Goodnow CC, Dixit VM. 2015. Caspase-11 cleaves gasdermin D for non-canonical inflammasome signalling. *Nature* 526:666–671.
3. Shi J, Zhao Y, Wang K, Shi X, Wang Y, Huang H, Zhuang Y, Cai T, Wang F, Shao F. 2015. Cleavage of GSDMD by inflammatory caspases determines pyroptotic cell death. *Nature* 526:660–665.
4. Broz P, von Moltke J, Jones JW, Vance RE, Monack DM. 2010. Differential requirement for Caspase-1 autoproteolysis in pathogen-induced cell death and cytokine processing. *Cell Host Microbe* 8:471–83.
5. Tentorey JL, Haloupek N, López-Blanco JR, Grob P, Adamson E, Hartenian E, Lind NA, Bourgeois NM, Chacón P, Nogales E, Vance RE. 2017. The structural basis of flagellin detection by NAIP5: A strategy to limit pathogen immune evasion. *Science* 358:888–893.
6. Zhao Y, Shao F. 2015. The NAIP-NLRC4 inflammasome in innate immune detection of bacterial flagellin and type III secretion apparatus. *Immunol Rev* 265:85–102.
7. Zhao Y, Yang J, Shi J, Gong Y-N, Lu Q, Xu H, Liu L, Shao F. 2011. The NLRC4 inflammasome receptors for bacterial flagellin and type III secretion apparatus. *Nature* 477:596–600.
8. Tentorey JL, Kofoed EM, Daugherty MD, Malik HS, Vance RE. 2014. Molecular Basis for Specific Recognition of Bacterial Ligands by NAIP/NLRC4 Inflammasomes. *Mol Cell* 54:17–29.

- 694 9. Hu Z, Yan C, Liu P, Huang Z, Ma R, Zhang C, Wang R, Zhang Y, Martinon F, Miao
695 D, Deng H, Wang J, Chang J, Chai J. 2013. Crystal Structure of NLRC4 Reveals Its
696 Autoinhibition Mechanism. *Science* (80-) 341:172–175.
- 697 10. Diebolder CA, Halff EF, Koster AJ, Huizinga EG, Koning RI. 2015. Cryoelectron
698 Tomography of the NAIP5/NLRC4 Inflammasome: Implications for NLR
699 ActivationStructure.
- 700 11. Davis MA, Fairgrieve MR, Den Hartigh A, Yakovenko O, Duvvuri B, Lood C, Thomas
701 WE, Fink SL, Gale M. 2019. Calpain drives pyroptotic vimentin cleavage,
702 intermediate filament loss, and cell rupture that mediates immunostimulation. *Proc*
703 *Natl Acad Sci U S A* 116:5061–5070.
- 704 12. Wang J, Deobald K, Re F. 2019. Gasdermin D Protects from Melioidosis through
705 Pyroptosis and Direct Killing of Bacteria. *J Immunol* ji1900045.
- 706 13. Jorgensen I, Zhang Y, Krantz BA, Miao EA. 2016. Pyroptosis triggers pore-induced
707 intracellular traps (PITs) that capture bacteria and lead to their clearance by
708 efferocytosis. *J Exp Med* 213:2113–2128.
- 709 14. Codo AC, Saraiva AC, dos Santos LL, Visconde MF, Gales AC, Zamboni DS,
710 Medeiros AI. 2018. Inhibition of inflammasome activation by a clinical strain of
711 *Klebsiella pneumoniae* impairs efferocytosis and leads to bacterial dissemination.
712 *Cell Death Dis* 9:1182.
- 713 15. Jorgensen I, Lopez JP, Laufer SA, Miao EA. 2016. IL-1 β , IL-18, and eicosanoids
714 promote neutrophil recruitment to pore-induced intracellular traps following
715 pyroptosis. *Eur J Immunol* 46:2761–2766.
- 716 16. Miao EA, Leaf IA, Treuting PM, Mao DP, Dors M, Sarkar A, Warren SE, Wewers
717 MD, Aderem A. 2010. Caspase-1-induced pyroptosis is an innate immune effector
718 mechanism against intracellular bacteria. *Nat Immunol* 11:1136–1142.
- 719 17. Faure E, Mear J-B, Faure K, Normand S, Couturier-Maillard A, Grandjean T, Balloy
720 V, Ryffel B, Dessein R, Chignard M, Uyttenhove C, Guery B, Gosset P, Chamaillard

721 M, Kipnis E. 2014. *Pseudomonas aeruginosa* Type-3 Secretion System Dampens
722 Host Defense by Exploiting the NLRC4-coupled Inflammasome. Am J Respir Crit
723 Care Med 189:799–811.

724 18. Cohen TS, Prince AS. 2013. Activation of inflammasome signaling mediates
725 pathology of acute P. aeruginosa pneumonia. J Clin Invest 123:1630–1637.

726 19. Brutinel ED, Vakulskas CA, Brady KM, Yahr TL. 2008. Characterization of ExsA and
727 of ExsA-dependent promoters required for expression of the Pseudomonas
728 aeruginosa type III secretion system. Mol Microbiol 68:657–671.

729 20. Kupz A, Guarda G, Gebhardt T, Sander LE, Short KR, Diavatopoulos DA, Wijburg
730 OLC, Cao H, Waithman JC, Chen W, Fernandez-Ruiz D, Whitney PG, Heath WR,
731 Curtiss R, Tschopp J, Strugnell RA, Bedoui S. 2012. NLRC4 inflammasomes in
732 dendritic cells regulate noncognate effector function by memory CD8+ T cells. Nat
733 Immunol 13:162–169.

734 21. Hauser AR. 2009. The type III secretion system of Pseudomonas aeruginosa:
735 infection by injection. Nat Rev Microbiol 7:654–65.

736 22. Thurston TLM, Wandel MP, von Muhlinen N, Foeglein A, Randow F. 2012. Galectin
737 8 targets damaged vesicles for autophagy to defend cells against bacterial invasion.
738 Nature 482:414–8.

739 23. Jyot J, Balloy V, Jouvion G, Verma A, Touqui L, Huerre M, Chignard M, Ramphal R.
740 2011. Type II secretion system of Pseudomonas aeruginosa: in vivo evidence of a
741 significant role in death due to lung infection. J Infect Dis 203:1369–77.

742 24. Sutterwala FS, Mijares LA, Li L, Ogura Y, Kazmierczak BI, Flavell RA. 2007.
743 Immune recognition of Pseudomonas aeruginosa mediated by the IPAF/NLRC4
744 inflammasome. J Exp Med 204:3235–45.

745 25. Tolle L, Yu F, Kovach MA, Ballinger MN, Newstead MW, Zeng X, Nunez G,
746 Standiford TJ. 2015. Redundant and cooperative interactions between TLR5 and
747 NLRC4 in protective lung mucosal immunity against Pseudomonas aeruginosa. J

748 Innate Immun 7:177–86.

749 26. Schultz MJ, Knapp S, Florquin S, Pater J, Takeda K, Akira S, van der Poll T. 2003.

750 Interleukin-18 impairs the pulmonary host response to *Pseudomonas aeruginosa*.

751 Infect Immun 71:1630–4.

752 27. Hughes AJ, Knoten CA, Morris AR, Hauser AR. 2018. ASC acts in a caspase-1-

753 independent manner to worsen acute pneumonia caused by *Pseudomonas*

754 *aeruginosa*. J Med Microbiol 67:1168–1180.

755 28. Amiel E, Lovewell RR, O'Toole GA, Hogan DA, Berwin B. 2010. *Pseudomonas*

756 *aeruginosa* Evasion of Phagocytosis Is Mediated by Loss of Swimming Motility and

757 Is Independent of Flagellum Expression. Infect Immun 78:2937–2945.

758 29. Lovewell RR, Patankar YR, Berwin B. 2014. Mechanisms of phagocytosis and host

759 clearance of *Pseudomonas aeruginosa*. Am J Physiol Lung Cell Mol Physiol

760 306:L591-603.

761 30. Patankar YR, Lovewell RR, Poynter ME, Jyot J, Kazmierczak BI, Berwin B. 2013.

762 Flagellar motility is a key determinant of the magnitude of the inflammasome

763 response to *Pseudomonas aeruginosa*. Infect Immun 81:2043–52.

764 31. Bitto NJ, Baker PJ, Dowling JK, Wray-McCann G, De Paoli A, Tran LS, Leung PL,

765 Stacey KJ, Mansell A, Masters SL, Ferrero RL. 2018. Membrane vesicles from

766 *Pseudomonas aeruginosa* activate the noncanonical inflammasome through

767 caspase-5 in human monocytes. Immunol Cell Biol 96:1120–1130.

768 32. Finethy R, Luoma S, Orench-Rivera N, Feeley EM, Haldar AK, Yamamoto M,

769 Kanneganti T-D, Kuehn MJ, Coers J. 2017. Inflammasome Activation by Bacterial

770 Outer Membrane Vesicles Requires Guanylate Binding Proteins. MBio 8:e01188-17.

771 33. Santos JC, Dick MS, Lagrange B, Degrandi D, Pfeffer K, Yamamoto M, Meunier E,

772 Pelczar P, Henry T, Broz P. 2018. LPS targets host guanylate-binding proteins to the

773 bacterial outer membrane for non-canonical inflammasome activation. EMBO J

774 37:e98089.

- 775 34. Vanaja SK, Russo AJ, Behl B, Banerjee I, Yankova M, Deshmukh SD, Rathinam
776 VAK. 2016. Bacterial Outer Membrane Vesicles Mediate Cytosolic Localization of
777 LPS and Caspase-11 Activation. *Cell* 165:1106–1119.
- 778 35. Mahenthiralingam E, Campbell ME, Speert DP. 1994. Nonmotility and phagocytic
779 resistance of *Pseudomonas aeruginosa* isolates from chronically colonized patients
780 with cystic fibrosis. *Infect Immun* 62:596–605.
- 781 36. Jain M, Ramirez D, Seshadri R, Cullina JF, Powers CA, Schulert GS, Bar-Meir M,
782 Sullivan CL, McColley SA, Hauser AR. 2004. Type III Secretion Phenotypes of
783 *Pseudomonas aeruginosa* Strains Change during Infection of Individuals with Cystic
784 Fibrosis. *J Clin Microbiol* 42:5229–5237.
- 785 37. Balakrishnan A, Karki R, Berwin B, Yamamoto M, Kanneganti T-D. 2018. Guanylate
786 binding proteins facilitate caspase-11-dependent pyroptosis in response to type 3
787 secretion system-negative *Pseudomonas aeruginosa*. *Cell death Discov* 4:3.
- 788 38. Iannitti RG, Napolioni V, Oikonomou V, De Luca A, Galosi C, Pariano M, Massi-
789 Benedetti C, Borghi M, Puccetti M, Lucidi V, Colombo C, Fiscarelli E, Lass-Flörl C,
790 Majo F, Cariani L, Russo M, Porcaro L, Ricciotti G, Ellemunter H, Ratclif L, De
791 Benedictis FM, Talesa VN, Dinarello CA, van de Veerdonk FL, Romani L. 2016. IL-1
792 receptor antagonist ameliorates inflammasome-dependent inflammation in murine
793 and human cystic fibrosis. *Nat Commun* 7:10791.
- 794 39. Elsen S, Huber P, Bouillot S, Couté Y, Fournier P, Dubois Y, Timsit JF, Maurin M,
795 Attrée I. 2014. A type III secretion negative clinical strain of *Pseudomonas*
796 *aeruginosa* employs a two-partner secreted exolysin to induce hemorrhagic
797 pneumonia. *Cell Host Microbe*.
- 798 40. Reboud E, Elsen S, Bouillot S, Golovkine G, Basso P, Jeannot K, Attrée I, Huber P.
799 2016. Phenotype and toxicity of the recently discovered *exlA* -positive *Pseudomonas*
800 *aeruginosa* strains collected worldwide. *Environ Microbiol* 18:3425–3439.
- 801 41. Broz P, Newton K, Lamkanfi M, Mariathasan S, Dixit VM, Monack DM. 2010.

- 802 Redundant roles for inflammasome receptors NLRP3 and NLRC4 in host defense
- 803 against *Salmonella*. J Exp Med 207:1745–1755.
- 804 42. Broz P, Ruby T, Belhocine K, Bouley DM, Kayagaki N, Dixit VM, Monack DM. 2012.
- 805 Caspase-11 increases susceptibility to *Salmonella* infection in the absence of
- 806 caspase-1. Nature 490:288–291.
- 807 43. Rauch I, Tenthorey JL, Nichols RD, Al Moussawi K, Kang JJ, Kang C, Kazmierczak
- 808 BI, Vance RE. 2016. NAIP proteins are required for cytosolic detection of specific
- 809 bacterial ligands in vivo. J Exp Med.
- 810 44. Rühl S, Broz P. 2015. Caspase-11 activates a canonical NLRP3 inflammasome by
- 811 promoting K⁺ efflux. Eur J Immunol 45:2927–2936.
- 812 45. Rietsch A, Vallet-Gely I, Dove SL, Mekalanos JJ. 2005. ExsE, a secreted regulator
- 813 of type III secretion genes in *Pseudomonas aeruginosa*. Proc Natl Acad Sci
- 814 102:8006–8011.
- 815 46. Gibson DG, Young L, Chuang R-Y, Venter JC, Hutchison CA, Smith HO. 2009.
- 816 Enzymatic assembly of DNA molecules up to several hundred kilobases. Nat
- 817 Methods 6:343–345.
- 818 47. Olson M V., Stover CK, Pham XQ, Erwin AL, Mizoguchi SD, Warren P, Hickey MJ,
- 819 Brinkman FSL, Hufnagle WO, Kowalik DJ, Lagrou M, Garber RL, Goltry L, Tolentino
- 820 E, Westbrook-Wadman S, Yuan Y, Brody LL, Coulter SN, Folger KR, Kas A, Larbig
- 821 K, Lim R, Smith K, Spencer D, Wong GK-S, Wu Z, Paulsen IT, Reizer J, Saier MH,
- 822 Hancock REW, Lory S. 2000. Complete genome sequence of *Pseudomonas*
- 823 aeruginosa PAO1, an opportunistic pathogen. Nature 406:959–964.
- 824 48. Santos JC, Dick MS, Lagrange B, Degrandi D, Pfeffer K, Yamamoto M, Meunier E,
- 825 Pelczar P, Henry T, Broz P. 2018. LPS targets host guanylate-binding proteins to the
- 826 bacterial outer membrane for non-canonical inflammasome activation. EMBO J
- 827 e98089.

Figures & legends

Fig. 1: Efficient pyroptosis-induced PIT response in macrophages depends on bacterial localization.

BMDMs were primed with 100 ng/mL of the TLR2 ligand Pam₃CSK₄ for 16 h to induce pro-IL-1 β expression and then infected with *S. Typhimurium* and *P. aeruginosa* for various times.

837 **(A)** Measurement of LDH and IL-1 β release in WT and *Nlrc4*^{-/-} BMDMs infected for 3 h
 838 with PAO1 and *S. Typhimurium* at an MOI of 15.

839 **(B)** Phagocytosis scoring of WT or *Nlrc4*^{-/-} BMDMs infected for 1 hour with PAO1 and *S.*
 840 *Typhimurium* at an MOI of 15.

841 **(C)** CFU evaluation of *Nlrc4*^{-/-} BMDMs infected for various time (1-24h) with PAO1 and *S.*
 842 *Typhimurium* at an MOI of 15.

843 **(D)** Cell death (LDH) and IL-1 β release evaluation in WT BMDMs, pre-incubated or not
 844 with cytochalasin D (0.2 μ g/mL) for 30 minutes, and then infected with either *P. aeruginosa*,
 845 *S. Typhimurium* (MOI 15).

846 **(E, F)** Microscopy illustrations and quantifications of PIT (ASC⁺ cells)-associated *P.*
 847 *aeruginosa* and *S. Typhimurium* in WT BMDMs infected for 1H with an MOI 3.

848 Graphs show mean and s.d of quadruplicate wells (A and D) from three independent
 849 experiments pooled together. CFUs scoring are representative of one experiment
 850 performed 5 times.

851 Regarding microscopy quantification (E, F), 10 fields containing approximately 200 cells
 852 were quantified using the Image J software.

853

854

Fig. 2: T3SS⁻ *P. aeruginosa* activate both NLRC4 and Caspase11 inflammasomes.

PAM3CSK4-primed (100 ng/mL) BMDMs were infected for various times with bacterial strains PAO1 T3SS⁺, T3SS⁻ or T3SS⁻/*fliC*⁻ at an MOI of 25, unless otherwise stated.

(A) Kinetics of IL-1β released by immortalized WT, *Nlrc4*^{-/-} or *Nlrc4*^{-/-}/*Casp11*^{-/-} BMDMs infected with PAO1 T3SS⁻ or T3SS⁻/*fliC*⁻.

(B) Western blot examination of processed caspase-1 (p20) and gasdermin-D (p30) in supernatants and pro-caspase-1 (p45), pro-gasdermin-D (p55), NLRP3, NLRC4 and GAPDH in cell lysates of WT, *Nlrc4*^{-/-}, *Nlrp3*^{-/-} and *Nlrc4*^{-/-}/*Nlrp3*^{-/-} BMDMs infected for 3 h with PAO1 T3SS⁺, T3SS⁻ or T3SS⁻/*fliC*⁻.

(C) Measurement of LDH and IL-1β release from WT, *Naip5*^{-/-} or *Naip*^{Δ1-6} BMDMs infected for 3 h with PAO1 T3SS⁺ or T3SS⁻.

Graphs show mean and s.d of quadruplicate wells **(A, C)** pooled from three independent experiments. Immunoblotting **(B)** is representative of one experiment performed two times.

874

875 **Fig. 3: T3SS⁻ *P. aeruginosa* associate to damaged phagosomes to promote**
 876 **inflammasome response.**

877 PAM3CSK4-primed BMDMs were infected with various *Pseudomonas aeruginosa* strains
 878 at an MOI of 25, unless specified.

879 **(A)** LDH and IL-1 β released by WT and *Nlrc4*^{-/-} BMDMs, pre-incubated or not with
 880 cytochalasin D (0.2 μ g/mL) for 30 minutes, and then infected for 3 h with PAO1 T3SS⁺ or
 881 T3SS⁻.

882 **(B)** Western blot analysis of cleaved CASP1 (p20) in cell supernatants and of pro-CASP1
 883 and β -actin in cell extracts of WT and *Casp1*^{-/-}/*11*^{-/-} BMDMs infected for 3 h with T3SS⁺ or
 884 T3SS⁻ PAO1.

885 **(C)** LDH and IL-1 β release in WT, *Nlrc4*^{-/-} or *Casp11*^{-/-} BMDMs, pre-incubated or not with
 886 cytochalasin D (0.2 μ g/mL) for 30 minutes, and then infected for 10 h with PAO1 T3SS⁺,
 887 **(D, E)** Fluorescence microscopy observations and quantifications of galectin-3 targeted
 888 PAO1 T3SS⁻ or T3SS⁻/*fliC* (MOI 25, 3 hours infection) in *Casp-1*^{-/-}/*11*^{-/-} immortalized
 889 BMDMs transduced with galectin-3-morange.

890 **(F, G)** Observations and quantifications of FAM-VAD-FMK FLICA probe recruitment on
 891 GAL-3-mOrange⁺ *Pseudomonas aeruginosa* (T3SS⁻) in IFN γ -primed immortalized (i)
 892 i*Nlrc4*^{-/-}/GAL3-morange and i*Nlrc4*^{-/-}/*Casp11*^{-/-}/GAL3-mOrange BMDMs 3 h after
 893 infection. Caspase-11 is in green, galectin-3 in red, bacteria in grey and nuclei in blue.
 894 Graphs show mean and s.d. **(A, C)** represent pooled data from at least three independent
 895 experiments performed in quadruplicate. **(B)** is representative of one experiment performed
 896 in duplicate. **(D, F)** Microscopy quantification of 10 fields containing approximately 200
 897 cells performed using the Image J software.

898

Fig. 4: Pyroptosis-induced *P. aeruginosa* entrapment and elimination is mostly efficient against T3SS-deficient bacteria.

Unprimed BMDMs were infected with various *Pseudomonas aeruginosa* strains at an MOI of 25, unless otherwise specified.

(A,B) Microscopy illustrations and quantifications of PIT (ASC⁺ cells)-associated *P. aeruginosa* T3SS⁺ or T3SS⁻ in WT BMDMs infected for 3h.

(C) Bacterial weakness induced by pyroptosis evaluated by LB agar plating of intracellular and extracellular PAO1 T3SS⁺ and T3SS⁻ from WT infected BMDMs after PBS or neutrophil exposure (1 h 30).

(D) CFU scoring of WT or *Nlrc4*^{-/-} BMDMs infected for 1 hour with PAO1 T3SS⁺ or T3SS⁻ at an MOI of 25.

(E) Bacterial weakness induced by pyroptosis (ATP, 5 mM, 2 h) evaluated by LB agar plating of intracellular *P. aeruginosa* T3SS⁺ and T3SS⁻ (MOI 25, 1h) from *Nlrc4*^{-/-} infected BMDMs after PBS or H₂O₂ exposure (50 μM).

914 **(F)** ImagestreamX quantification of the % of (i) Neutrophils (Ly6G⁺)/macrophage (F4/80⁺)
 915 and, (ii) Neutrophils (Ly6G⁺)/macrophages (F4/80⁺)/bacteria (GFP⁺) in the peritoneal cavity
 916 of WT mice infected for 4 h with 3.10⁶ CFUs of T3SS⁺ or T3SS⁻ *P. aeruginosa*. Single dots
 917 are representative of each individual mouse infected with either strain of *P. aeruginosa*.
 918 Images represent the acquisition of more than 100 000 total events. Upper image panel
 919 and lower image panel show representatives images from the Neutrophils
 920 (Ly6G⁺)/macrophages (F4/80⁺) gate obtained from mice infected with T3SS⁺ or T3SS⁻ *P.*
 921 *aeruginosa* respectively. Here, three independent experiments were conducted.
 922 All data **(A-F)** are expressed as mean and s.d. Data are representative of three
 923 independent experiments.

924

925 **Movie 1** shows T3SS⁺ PAO1 mcherry entrapment into dead cells and refers to **Fig. 4**.

926 **Movie 2** shows T3SS⁻ PAO1 mcherry entrapment into dead cells and refers to **Fig. 4**.

927

928

929

930

931

932

933

Fig. 5: Both NLRC4 and Caspase-11 only protect against acute infection with T3SS-deficient *P. aeruginosa*.

(A, C) BALF PAO1 CFU scoring after 18 h of WT, *Nlrc4*^{-/-} **(A)** or *Casp11*^{-/-} **(C)** mice infected with either T3SS+ or T3SS- *P. aeruginosa* strains at 5x10⁶ (T3SS+) and 1.5x10⁷ (T3SS-) CFUs respectively.

(B, D) BALF IL-1β release assay in mice infected as described in **(A, C)**.

(E, F) Role of alveolar macrophages on the immune response to PAO1. **(E)** PAO1 CFUs scoring in BALFs after 18 h of infection with either the T3SS+ or T3SS- *P. aeruginosa* strains in control or clodronate treated mice as in **(A, C)**. **(F)** IL-1β release in BALFs of control or clodronate treated mice infected with either the PAO1 T3SS+ or T3SS- strains as in **(B, D)**.

All data are representative results of 2 **(C-F)** and 3 **(A, B)** independent experiments.

947 Supplementary information

948

949 **Figure S1: *Pseudomonas aeruginosa* T3SS promotes phagocytosis-independent**
 950 **activation of the NLRC4 inflammasome.**

951 BMDMs were primed with 100 ng/mL of the TLR2 ligand Pam₃CSK₄ for 16 h to induce pro-
 952 IL-1 β expression and then infected for 3 hours (h) with various bacteria (MOI 15) or
 953 inflammasome inducers.

954 **(A)** Measurement of cell death (LDH release) from WT and *Nlrc4*^{-/-} BMDMs after 3 h of
 955 infection with either T3SS⁺ or T3SS⁻ PAO1 or STm.

956 **(B)** CFU scoring in WT and *Nlrc4*^{-/-} BMDMs infected for 1h with either PAO1 or *STY*-
 957 deficient *P. aeruginosa*.

958 **(C)** LDH and IL-1 β release quantifications of WT and *Nlrc4*^{-/-} BMDMs infected for 3 h with
 959 either PAO1 or *STY*-deficient *P. aeruginosa* in presence or not of Cytochalasin D (0,2
 960 μ g/mL).

961 **(D)** LDH release quantifications of WT and *Nlrc4*^{-/-} BMDMs, pretreated or not with
 962 Cytochalasin D (0,2 μ g/mL), electroporated (Neon™ Transfection System) with 0.5
 963 ug/ml and then plated for 1 h.

964 **(E)** LDH and IL-1 β released by WT BMDMs, pre-incubated or not with cytochalasin D
 965 (0.2 μ g/mL) for 30 minutes, and then infected with either *P. aeruginosa*, *S. Typhimurium*, *S.*
 966 *flexneri* or *C. violaceum*.

967 **(F)** CFU scoring in the peritoneal cavity of WT and *GsdmD*^{-/-} mice infected for 6 h with
 968 3.10⁶ CFUs of *P. aeruginosa* and *S. Typhimurium*.

969 All graphs **(A-E)** show mean and s.d of quadruplicate wells from three pooled experiments.

970 **(F)** represents one experiment from three independent experiments.

971

972

Figure S2: *P. aeruginosa* triggers T3SS-independent activation of both NLRC4 and Caspase11 inflammasomes.

(A, B) Biochemical **(A)** and functional **(B)** characterization of two iCas9*Nlr4*^{-/-}/sgCasp11^{-/-} CRISPR KO clones. **(B)**, LDH release measurement of CRISPR clones (A) transfected with either 0,5 µg/mL of Flagellin (2 hours) or 5µg/mL of LPS (10 hours).

(C) Cell death evaluation (LDH) in WT and *Nlr4*^{-/-} BMDMs infected with various MOIs of PAO1 T3SS⁺ or T3SS⁻ for 3 h.

(D, E) LDH measurement in WT and *Nlr4*^{-/-} BMDMs infected by PAO1 T3SS⁺, T3SS⁻ or T3SS⁻/*fliC*, complemented or not for flagellin expression (*fliC*::*fliC*) for 3h with an MOI of

50.

984 **(F)** Measurement of IL-1 β release from WT and *Nlr4*^{-/-} BMDMs after 3 h of infection with
 985 live, PFA-killed or heat killed (HK) T3SS deficient-PAO1 at an MOI of 50.
 986 **(G)** Measurement of LDH and IL-1 β release from WT, *Nlr4*^{-/-} and *Casp1*^{-/-}/*11*^{-/-} BMDMs
 987 after 3 h of infection with T3SS deficient-PAO1 *exsA*⁻ or -PAO1 *pscC* at an MOI of 50.
 988 Graphs show mean and s.d of quadruplicate wells pooled from three independent
 989 experiments.
 990

992 **Figure S3: T3SS- *P. aeruginosa* associate to altered phagosomes.**

993 **(A)** Biochemical characterization of two iCas9*Nlr4*^{-/-}/*sgCasp11*^{-/-} CRISPR KO clones.
 994 **(B)** Fluorescence microscopy observations and quantifications of galectin-3 targeted
 995 PAO1 T3SS⁻ (MOI 25, 3 h of infection) in iWT and iCasp1^{-/-}/*11*^{-/-}/*sgNlr4*^{-/-} BMDMs
 996 transduced with galectin-3-morange.
 997 Graphs show mean and s.d of quadruplicate wells from three independent pooled
 998 experiments. Immunoblotting representative of one experiment performed on various
 999 clones. Microscopy analysis visualized 10 fields containing approximately 200 cells using
 1000 Image J software.
 1001

Figure S4: ImagestreamX gating strategy used.

(A) ImagestreamX gating strategy used to visualize and quantify neutrophil-mediated efferocytosis of bacteria-containing macrophages.

(i) A gate was set on cells in focus [Cells in Focus], then (ii) single cells [Single Cells] were gated. (iii) First, gate was put on Ly6G⁺ Neutrophils [Ly6G⁺] and then we selected (iv) F4/80⁺ macrophages [F4/80⁺] within Ly6G⁺ population. (v) A mask based on the surface Area of F4/80⁺ signal was applied to Ly6G⁺/F4/80⁺ gate to discriminate efferocytosis of intact macrophages from fragmented macrophages. As a control, dotted gray line shows surface area of F4/80⁺ signal in the [Ly6G⁻] gate. Then, (vi) we gated on [F4/80⁺ low Area] and measured the intensity of the GFP signal. Red and Blue lines show the intensity of GFP signal measured in mice challenged with either *P. aeruginosa* T3SS⁺ or T3SS⁻ strains respectively. Finally, GFP⁺ bacteria percentage within the gate (vi) was visualized. Upper image panel and lower image panel show representative images from the Neutrophils (Ly6G⁺)/macrophages (F4/80⁺) gate obtained in GFP⁻ gate or GFP⁺ gate respectively.

Figure S5: NLRC4 and Caspase-11 only protect mice against T3SS-deficient *P. aeruginosa*.

(A-D) PAO1 CFUs scoring and IL-1β levels in BALFs from WT, *Nlrc4*^{-/-} **(A, B)** or *Casp11*^{-/-} **(C, D)** mice after 18 h of infection with T3SS-*fliC*⁻ *P. aeruginosa* strain with 1.5x10⁷ CFUs.

(E) FACS representation and quantification of depleted alveolar macrophages (F4/80⁺, CD11c⁺) in control (PBS) and clodronate instilled mice.

1045 **Table S1:** All products, software and biological samples used in the current study,
 1046 including their references and concentrations, are listed in table 1.

Reagent	Provider	Reference
Antibodies		
Mouse anti-mouse caspase-1 1/4000 WB	Adipogen	AG20B-0042
Rabbit anti-IL-18 1/2500 WB	Biovision	5180R-100
Goat anti-mouse IL-1 β 1/1000 WB	R&D	AF-401-NA
Rabbit anti-mouse Gasdemin D 1/1000 WB	Abcam	ab209845
Rat anti-mouse Caspase-11 (17D9) 1/1000 WB	Novus	NB120-10454
Rabbit anti-mouse Nlrc4 1/1000 WB	Abcam	ab201792
Mouse anti-mouse NLRP3 1/1000 WB	AdipoGen	AG-20B-0014
Mouse anti- β -actin 1/10000 WB	Sigma	A1978
Rabbit anti-GAPDH 1/10000 WB	GeneTex	GTX100118
Rabbit anti-ASC 1/1000 IF	SantaCruz	Sc-22514-R
Chicken anti- <i>Pseudomonas</i> LPS 1/5000 IF	Agrisera	IMS01-133-332
Anti-CD45-2-APC IF	Biolegends	109818
Anti-CD19-PE-Cf594 0.2 μ g/mL FACS	Biolegends	115554
Anti-TCR β -APC-Cy7 0.2 μ g/mL FACS	Biolegends	109220
Anti-F4/80-BV421 50 μ g/mL FACS	Biolegends	123132
Anti-CD11b-PE-Cy7 0.2 μ g/mL FACS	Biolegends	101216
Anti-CD11c-A488 0.5 μ g/mL FACS	Biolegends	117311
Anti-Ly6C-PerCP-Cy5 0.2 μ g/mL FACS	Biolegends	128012
Anti-MHCII-PE 0.2 μ g/mL FACS	Biolegends	107608
Anti-CD86-BV510 50 μ g/mL FACS	Biolegends	105039
Anti-Ly6G-A647 0.5 μ g/mL FACS	Biolegends	127610
Brilliant Violet 421™ anti-mouse F4/80 Antibody	Biolegends	123131

anti-LY6G (APC-Vio770)	Miltenyi-Biotech	130-107-916
Sec antibodies		
Goat anti-Rat IgG (H&L) DyLight 488 IF 1/20 1/2000 IF	Diagomics	GtxRt-003- D488NHSX
Chicken anti-Goat IgG (H&L) DyLight 488 1/500 IF	Diagomics	CkxGt-003- D488NHSX
Goat anti-Mouse IgG (H&L) DyLight 488 1/500 IF	Diagomics	GtxMu-003- D488NHSX
Goat anti-Rabbit IgG (H&L) DyLight 488 1/500 IF	Diagomics	GtxRb-003- D488NHSX
Donkey anti-Rabbit IgG (H&L) DyLight 550 1/500 IF	Diagomics	DkxRb-003- D550NHSX
Goat anti-Rat IgG (H&L) DyLight 550 1/500 IF	Diagomics	GtxRt-003- D550NHSX
Donkey anti-Goat IgG (H&L) DyLight 550 1/500 IF	Diagomics	DkxGt-003- D550NHSX
Goat anti-Mouse IgG (H&L) DyLight 550 1/500 IF	Diagomics	GtxMu-003- D550NHSX
Goat anti-Chicken IgG (H&L) DyLight 405 1/500 IF	Diagomics	GtxCk-003- D405NHSX
Goat anti-Rabbit IgG (H&L) DyLight 633 1/500 IF	Diagomics	GtxRb-003- D633NHSX
Goat anti-Rat IgG H&L) DyLight 633 1/500 IF	Diagomics	GtxRt-003- D633NHSX
Hoescht	Sigma	H6024
Rabbit anti-Goat IgG (H&L) HRP Conjugate 1/3000	Diagomics	RbxGt-003- DHRPX

Rabbit anti-Mouse IgG (H&L) HRP Conjugate 1/3000	Diagomics	RbxMu-003- DHRPX
Goat-anti-mouse IgG (H+L) HRP conjugate 1/3000	Diagomics	R-05071-500
Goat-anti-rabbit IgG (H+L) HRP conjugate 1/3000	Diagomics	R-05072-500
Goat-anti-rat IgG (H+L) HRP conjugate 1/3000	Diagomics	R-05075-500
CSFE	Biolegend	423801
TO-PRO [®] -3 Iodide (1/10000)	Thermo Fischer Scientific	T3605
Bacterial and Virus Strains		
PAO1	(47)	
PAO1 Δ exsA (referred as T3SS ⁻)	This study	N.A
PAO1 Δ pscC	This study	N.A
PAO1 Δ exsA, Δ fliC	This study	N.A
PAO1 Δ fliC (<i>fliC</i>)	This study	N.A
PAO1 Δ fliC:: <i>fliC</i>	This study	N.A
PAO1 Δ exsA, Δ fliC:: <i>fliC</i>	This study	N.A
PAO1 mCherry	This study	N.A
PAO1 Δ exsA mCherry (referred as T3SS ⁻)	This study	N.A
PAO1 GFP	This study	N.A
PAO1 Δ exsA GFP (referred as T3SS ⁻)	This study	N.A
STm SL1344 (<i>orgA</i> / <i>ssaV</i>), Tet ^R /Km ^R (grown in LB supplemented with 10µg/ml Tet and 50µg/ml Kan)	(41, 42, 48)	N.A

STm SL1344 <i>orgA</i> ⁻ <i>flaG/B</i> ⁻ <i>fliC</i> ⁻ , Tet ^R /Cm ^R /Km ^R (grown in LB supplemented with 10µg/ml Tet, 25µg/ml Cm and 50µg/ml Kan)	(41, 42, 48)	N.A
<i>E. coli</i> Stbl4	Thermo Fisher	N.A
<i>E. coli</i> Sm10	Conjugative strains (Kanamycin) Km ^R	N.A
Biological Samples		
Chemicals, Peptides, and Recombinant Proteins		
FCS	Fisher Scientific	16010-159
MCSF	L929 cell supernatant	NA
HEPES	Fisher Scientific	SH30237.01
Non-essential amino acids	Invitrogen	
PAM3CSK4	Invivogen	tlrl-pms
LPS-EB ULTRAPURE	Invivogen	tlrl-3pelps
ATP (NLRP3 inflammasome inducer)	invivogen	tlrl-atp
Gentamycin	Invitrogen	15710-049
Kanamycin	Sigma	60615-5G

Irgasan	Sigma	72779-25G
LS Columns	Miltenyi Biotec	130-042-401
Anti-Ly-6G MicroBeads UltraPure mouse	Miltenyi Biotec	30-120-337
BD Fixation/Permeabilisation Kit	BD	554714
WB reagents		
ECL Clarity Max Substrate	BioRad	1705060
ECL Clarity Max Substrate	BioRad	1705062
Western Blot Strip Buffer	Diagomics	R-03722-D50
Tris base	euromedex	200923-A
SDS ultra-pure (4x)	Euromedex	1012
Acrylamide / Bisacrylamide 37.5/1 30%	Euromedex	EU0088-B
Temed	Sigma	T9281-25ML
Ammonium persulfate	Sigma	248614-100g248614-100g
Page Ruler 10-180 kDa	Fisher Scientific	15744052
Vectashield	Vectalabs	H-1000

Triton X-100	Euromedex	2000
Hydrogen Peroxide	Sigma	21763-500ml
DMEM	Fisher Scientific	41965-039
LB	Fisher Scientific	BP1426-2
LB Agar	INVITROGEN	22700025
Cytochalasin D (0.2µg/mL)	Sigma	C8273-1Mg
Roche protease inhibitor cocktail	Sigma	0000000116974 98001
Saponin	Bio Basic canada	SB4521
BSA	Sigma	A9647-100G
Sucrose	Sigma	S7903
Clodronate Liposomes	Liposoma	283539
Critical Commercial Assays		
IL-1β ELISA kit	ebiosciences	12354003
LDH Cytotoxicity Detection Kit	Takara	MK401
FAM-VAD-FMK FLICA Poly Caspase Assay	BioRad	ICT091
Neutrophil Isolation Columns	MACS Miltenyi Biotech	130-097-658
Experimental Models: Cell Lines		
<i>Naip5^{-/-}</i> BMDMs	R.E.Vance	N.A

<i>Naip1-6^{-/-}</i> BMDMs	R.E.Vance	N.A
C57BL/6 J primary BMDMs	Janvier	N.A
C57BL/6 N primary BMDMs	Charles River	N.A
Immortalized <i>Casp-1^{-/-}/11^{-/-}</i> BMDMs	This study	N.A
Immortalized <i>GBP^{Chr3-/-}</i> BMDMs	This study	N.A
Immortalized iCas9 <i>Nlrc4^{-/-}</i>	This study	N.A
Immortalized iCas9 <i>Casp-1^{-/-}/11^{-/-}</i>	This study	N.A
Immortalized iCas9 <i>Casp-1^{-/-}/11^{-/-}/sgNlrc4^{-/-}</i>	This study	N.A
Immortalized iCas9 <i>Nlrc4^{-/-}/sgCasp-11^{-/-}</i>	This study	N.A
Experimental Models: Organisms/Strains		
C57BL/6 J mice	Janvier	N.A
C57BL/6 N mice	Charles River	N.A
<i>Nlrc4^{-/-}</i> and <i>Casp11^{-/-}</i> mice	V.M Dixit (Genentech)	N.A
<i>Nlrp3^{-/-}</i> mice	P. Broz (Univ. of Lausanne)	N.A
<i>Casp-1^{-/-}/11^{-/-}</i> mice	R. Flavell	N.A
<i>Nlrc4^{-/-}</i> mice	F. Sutterwala	N.A
Oligonucleotides		
exsA Rup Fw	SIGMA: ttccacacattatacga gccggaagcataaat gtaaagcaagcttGC TGGCGTTGCTG CTCGG	N.A

exsA Rup Rv	SIGMA: TACCGGGCTTT CAAAAAACGAT TATAAGAACCC CAACACTTCCC G	N.A
exsA Rdown Fw	SIGMA: AGTGTTGGGGT TCTTATAATCGT TTTTTGAAAGC CCGGTAGC	N.A
exsA Rdown Rv	SIGMA: ggaaattaattaaggt accgaattcgagctcg agcccggggatccC GTCTCCAGCTT GCCG	N.A
exsA Rup check	SIGMA: TCGCCGACGGT GACGGCAC	N.A
exsA Rdown check	SIGMA: GAGCAGCGTCG CCATGCCCC	N.A

<i>fliC</i> Rup Fw	SIGMA: ttccacacattatacga gccggaagcataaat gtaaagcaagcttAC GCTATCCCGCC TACCT	N.A
<i>fliC</i> Rup Rv	SIGMA: GTGAGTGACCG TTCCCGGGCGG TGATTTCTCC AAAGGACCTAT TTCG	N.A
<i>fliC</i> Rdown Fw	SIGMA: GTCCTTTGGAG GAAATCACCGC CCGGGAACGGT CAC	N.A
<i>fliC</i> Rdown Rv	SIGMA: ggaaattaattaaggt accgaattcgagctcg agcccggggatccC GCGCTGATCGC ACTCT	N.A
<i>fliC</i> Rup check	SIGMA: ATTCTGCCCGG ATGCCTTGC	N.A

<i>fliC</i> Rdown check	SIGMA: CTTCGTTGGAA GACTTGGCGG	N.A
<i>pscC</i> Rup fw	SIGMA: ttccacacattatacga gccggaagcataaat gtaaagcaagcttCG GCTTCGGCACT GGCGA	N.A
<i>pscC</i> Rup Rv	SIGMA: CCATGCTAATT CCCGCGCTCCA GCAGGCGGCG CATCAG	N.A
<i>pscC</i> Rdown fw	SIGMA: CCCTGATGCGC CGCCTGCTGGA GCGCGGGAATT AGCATGGCC	N.A
<i>pscC</i> Rdown Rv	SIGMA: ggaaattaattaaggt accgaattcgagctcg agcccggggatccG ATGAAATCCAC GCCCTGGC	N.A
<i>pscC</i> Rup check	SIGMA: AGGAACTGGCC AGGGTCGC	N.A

<i>pscC</i> Rdown check	SIGMA: CCCAGCCAGGC TTGTCCG	N.A
JBOC FliC Fw	SIGMA: agccattcTCAGTG AAGCATCAAGA CTAACAAATCct acggggcgcgcatctcg agg	N.A
JBOC FliC rv	SIGMA: GAAAATCTTCT CTCATCCGCCA AAACAGCCAAG CTAATTCTtagcgc agcaggctcag	N.A
Gal-3 up	Sigma: GATCCTCGAGG CCGCCACCATG GTGAGCA	N.A
Gal-3 down	Sigma: GATCGCGGCC GCTTATATCAT GGTATATGAAG CACTG	N.A

<i>Actin</i>	Sigma Fw 5'- agccatgtacgtagcc atcc-3' Rv 5'- ctctcagctgtggtggt gaa-3'	N.A
Recombinant DNA		
pMSCV-Galectin-3-mOrange	This study	
JBOC- <i>fliC</i> plasmid	This study	Addgene deposit number 74611
pMSCV2.2	R.E. Vance	
LentiGuide-Puro	Feng Zhang lab	addgen ref 52963
pMD.2G	Didier Trono lab	addgene ref 12259
p8.91	Didier Trono lab	N.A.
LentiCas9-Blast	Feng Zhang lab	addgen ref 52962
<i>Nlrc4</i> targeting primer Forward:	Sigma-Aldrich 5'CACCG- TTACTGTGAGC CCTTGGAGC3'	

<i>Nlrc4</i> targetting primer reverse:	Sigma-Aldrich 5'AAAC- GCTCCAAGGGC TCACAGTAA-C3'	
<i>Caspase 11</i> targetting primer forward	Sigma-Aldrich 5'CACCG- CTTAAGGTGTT GGAACAGCT3'	
<i>Caspase 11</i> targetting primer reverse	Sigma-Aldrich 5'AAAC- AGCTGTTCCAA CACCTTAAG-C3'	
Software and Algorithms		
Graph Pad Prism 5.0		N.A
Image J		N.A
Snappene	GSL Biotech LLC, Chicago, U.S.A	N.A
<i>FlowJO</i>	FlowJo LLC	N.A
<i>Fiji (Image J)</i>		N.A
<i>Benchling Software</i>		N.A
IDEAS software v2.6	Amnis	

1047

1048 Information and requests for reagents may be sent to Etienne Meunier
1049 (Etienne.meunier@ipbs.fr) or Petr Broz (Petr.broz@unil.ch).