Polygenic risk score of alcohol consumption predicts alcohol-related morbidity and all-cause mortality

Tuomo Kiiskinen¹, Nina J. Mars¹, Teemu Palviainen¹, Jukka Koskela¹, Pietari Ripatti¹, Joel T. Rämö¹, Sanni Ruotsalainen¹, FinnGen, GSCAN Consortium, Aarno Palotie^{1,2}, Pamela A.F. Madden³, Richard J. Rose⁴, Jaakko Kaprio^{1,6}, Veikko Salomaa⁵, Pia Mäkelä⁵, Aki S. Havulinna^{1,5}, Samuli Ripatti^{1,2,6}

Author affiliations

¹Institute for Molecular Medicine Finland (FIMM), HiLIFE, University of Helsinki, Helsinki, Finland

²The Broad Institute of MIT and Harvard, Cambridge, Massachusetts, USA

³Department of Psychiatry, Washington University School of Medicine in St.Louis, St.Louis, Missouri, USA

⁴Department of Psychological and Brain Sciences, Indiana University Bloomington, Bloomington, Indiana, USA

⁵National Institute for Health and Welfare, Helsinki, Finland

⁶Department of Public Health, Clinicum, Faculty of Medicine, University of Helsinki, Helsinki, Finland

Corresponding author: Samuli Ripatti

Email: samuli.ripatti@helsinki.fi

Abstract

Objective: To develop a highly polygenic risk score (PRS) for alcohol consumption and study whether it predicts alcohol-related morbidity and all-cause mortality.

Design: Biobank-based prospective cohort study

Setting: FinnGen Study (Finland)

Participants: 96,499 genotyped participants from the nationwide prospective FinnGen study and 36,499 participants from prospective cohorts (Health 2000, FINRISK, Twin Cohort) with detailed baseline data and up to 25 years of follow-up time.

Main outcome measures: Incident alcohol-related morbidity and alcohol-related or all-cause mortality, based on hospitalizations, outpatient specialist care, drug purchases, and death reports.

Results: In 96,499 FinnGen participants there were in total 4,785 first-observed incident alcohol-related health events. The PRS of alcohol consumption was associated with alcohol-related morbidity and the risk estimate (hazard ratio, HR) between the highest and lowest quintiles of the PRS was 1.67 [95 % confidence interval: 1.52-1.84], $p=3.2 \times 10^{-27}$). In 28,639 participants with comprehensive baseline data from prospective Health 2000 and FINRISK cohorts, 911 incident first alcohol-related events were observed. When adjusted for self-reported alcohol consumption, education, marital status, and gamma-glutamyl transferase blood levels, the risk estimate between the highest and lowest quintiles of the PRS was 1.58 (CI=[1.26-1.99], $p=8.2 \times 10^{-5}$). The PRS was also associated with all-cause mortality with a risk estimate of 1.33 between the highest and lowest quintiles (CI=[1.2-1.47], $p=4.5 \times 10^{-8}$) in the adjusted model. In all 39,695 participants with self-reported alcohol consumption available, a 1 SD increase in the PRS was associated with 11.2 g (=0.93 drinks) higher weekly alcohol consumption ($\beta=11.2$ [9.85-12.58 g], $p = 2.3 \times 10^{-58}$).

Conclusions: The PRS for alcohol consumption associates for both alcohol-related morbidity and all-cause mortality. These findings underline the importance of heritable factors in alcohol-related behavior and the related health burden. The results highlight how measured genetic risk for an important behavioral risk factor can be used to predict related health outcomes.

Introduction

Alcohol drinking is a major dose-dependent contributor to morbidity and mortality. Globally, 3 million annual deaths (5 % of all deaths) result from alcohol consumption, and is also linked to more than 200 disease and injury outcomes.(1) As ethanol is a psychoactive substance with addictive properties,(2) alcohol consumption can lead to the development of alcohol use disorders (AUDs), globally prevalent mental disorders of pathological addictive or abusive drinking patterns, which are linked to worse health outcomes, negative socioeconomic effects, and increased mortality.(3) There is a strong connection between the health burden and the level of alcohol consumed, (4) and in total alcohol has been estimated to be the most damaging of all substances of abuse, in terms of harm caused to self and others.(5)

Alcohol-related behaviors are also affected by genetic factors and the estimated heritability of alcohol consumption in twin studies has ranged between 35% and 65% (weighted average 37%) (6) and its single nucleotide polymorphism -based heritability has been estimated to be 10%. (7) Recent large-scale genome-wide association studies (GWAS) have identified multiple loci associated with alcohol consumption, underlining the importance of large study populations for unraveling the genetic architecture underlying alcohol-related traits. (7, 8) Similarly, GWAS of alcohol dependence and the Alcohol Use Disorders Identification Test (AUDIT) scores have shown the traits to be genetically distinct but positively correlated.(9,10)

Polygenic risk scores (PRS) derived from GWAS summary statistics have showcased improved performance in disease prediction. (11) PRSs for known risk factors have also been shown to associate with the related disease (12) and recently associations between multiple risk factor PRSs and related traits were confirmed and reported. (13, 14) However, the link between PRSs for behavioral traits and associated health outcomes remains poorly understood.

The assessment of potential health risks related to alcohol has so far relied on traditional risk factors, including family history, without explicit measurement of genetic risk. Here we developed a highly polygenic risk score for alcohol consumption and studied whether alcohol-related polygenic burden predicts alcohol-use disorders and other alcohol-related morbidity and mortality in Finnish biobank

cohorts (n=96,499) linked to electronic health records. Furthermore, we studied whether the PRS for alcohol consumption predicts alcohol-related outcomes beyond self-reported alcohol consumption and other related risk factors, thus providing more objective information independent of individual reporting bias or temporal fluctuations.

Materials and methods

Study sample and definition of alcohol-related morbidity

The data is comprised of 96,499 Finnish individuals from FinnGen Data Freeze 2, which includes prospective epidemiological and disease-based cohorts as well as hospital biobank samples (Supplementary Table 1). The data were linked by the unique national personal identification numbers to national hospital discharge, death, and medication reimbursement registries. Additional details and information on the genotyping and imputation are provided in the online-only Supplementary Information.

Alcohol-related baseline measures were available for a subset of the FinnGen dataset consisting of national population survey cohorts: FINRISK, collected in 1992, 1997, 2002, 2007 and 2012 and Health 2000, collected in 2000. The baseline data included self-reported information assessed by questionnaires, anthropometric measures, and blood samples. More detailed descriptions of the FINRISK and Health 2000 studies have been published previously.(15,16)

Additionally, three Finnish twin cohorts, FinnTwin12, NAG-FIN, and Old Twin were pooled and analyzed as one dataset. For these datasets, cohort baseline data was available, but the cohorts were not linked to electronic health records. For details regarding the twin datasets, see the online descriptions (<https://wiki.helsinki.fi/display/twineng/Twinstudy>). (17,18)

Using nationwide registries for deaths (1969-2016), hospital discharges (1969-2016), outpatient specialist appointments (1998-2016) and drug purchases (1995-2016), we combined 21 somatic and psychiatric alcohol-related diagnoses into a composite disease endpoint, harmonizing the International Classification of Diseases (ICD) revisions 8, 9 and 10, and ATC-codes (**Supplementary Table 1**). These registries spanning decades were electronically linked to the cohort baseline data

using the unique national personal identification numbers assigned to all Finnish citizens and residents.

Genotyping and imputation

FinnGen, FINRISK, Health2000, and Finnish Twin Cohort samples were genotyped with Illumina and Affymetrix genomewide SNP arrays. Individuals with non-European ancestry or uncertain sex were excluded. The details about the genotype calling, quality controls and imputation are provided in the Supplementary Information.

Polygenic risk scores

Summary statistics from the largest existing GWAS meta-analysis on alcohol consumption (8) were used for constructing the PRS. To avoid overfitting, a separate ad hoc meta-analysis was performed by GSCAN, excluding all Finnish and 23andMe samples ($n=527,282$ after exclusions). The LDpred-inf method (19) was used to account for linkage disequilibrium (LD) among loci, with whole-genome sequencing data on 2,690 Finns serving as the external LD reference panel. The final scores were generated with PLINK2 (20) by calculating the weighted sum of risk allele dosages for each variant. The number of variants in the final scores was 1.1×10^6 .

Statistical analysis

The Cox proportional hazard model was used to estimate survival curves, hazard ratios (HRs) and 95 % confidence interval (95% CI) in the survival analyses where age was used as the time scale. R's cox.zph function was used to test whether the proportional assumption criteria applied in our models. Linear regression in FINRISK and Health 2000 and linear mixed model in the Twin Cohort was used for estimating the relationship between the PRS and alcohol consumption. Logistic regression in the FINRISK and Health 2000 cohorts and linear mixed model in the Twin Cohort was used to estimate the relationship between alcohol abstinence and the PRS.

Age, sex, genotyping array, and the first ten principal components of ancestry were used as core covariates. Additionally, body mass was used as a covariate in the model estimating the PRS-alcohol

consumption relationship. Self-reported weekly average alcohol consumption from the past year (when unavailable, the past week's consumption) was used as the estimate for alcohol consumption. In the fully adjusted survival model analyses, $\log(x+1)$ -transformed alcohol consumption-estimate, current smoking status, binary higher education status, binary marital/cohabitation status and GGT (Gamma Glutamyl Transferase) blood levels at baseline served as covariates.

In the survival analyses, all prevalent cases and individuals with covariate missingness were excluded. The PRS was normalized and included as a continuous variable in the models. In the survival analysis the highest and lowest genetic risk for alcohol consumption were compared using PRS quintiles.

In analyses using baseline consumption data, the analyses were performed separately in the Health 2000, FINRISK Study, and Twin Cohorts and then meta-analyzed using fixed effects model.

In risk prediction, FINRISK cohorts with at least 10 years of follow-up (from 1992 to 2002) were used to train the model, and the predictive performance was tested in the Health 2000 cohort. The maximal follow-up window was restricted to 10 years. The change in the predictive performance was assessed by comparing models with and without the PRS using the correlated C-index approach (21) along with calculating the continuous reclassification improvement (NRI) (22) and integrated discrimination improvement (IDI).(23) The Hosmer-Lemeshow goodness-of-fit test was used to test model calibration.

Patient and public involvement

No patients were involved in setting the research question or the outcome measures, nor were they involved in developing plans for recruitment, design, or implementation of the study. No patients were asked to advise on interpretation or writing up of results. There are no plans to disseminate the results of the research to study participants or the relevant patient community.

Results

Cohorts

Our primary dataset (FinnGen) is comprised of 96,499 unrelated individuals (54,262 women) with a total of 55,484,114 person-years of registry-based follow-up and 4,785 first-observed alcohol-related major health events. Alcohol consumption estimates were available for a total of 39,695 individuals from the prospective cohorts (FINRISK, Health 2000 and Twin Cohort). Two cohorts, FINRISK and Health 2000, have full registry data and information on self-reported alcohol consumption and related baseline data and consist of 28,639 individuals (94.5% of the participants after excluding 964 prevalent alcohol-related morbidity cases), with 424,053 person-years of registry-based follow-up and 988 first ever alcohol-related events. The interview-based DSM-IV AUD-status was available in a subset of the Twin cohort for 713 cases and 1460 controls.

Table 1. Population characteristics of FinnGen, FINRISK, Health 2000, and Twin Cohort -datasets.

	FinnGen	FINRISK	Health 2000	Twin Cohort
N	96,499 (54,262 women)	23,824 (12,513)	5,945 (3,260)	9,926 (5,036)
N (incident events)	4,785	817	171	NA
Age (years)	57.5(end of follow-up)	48.6 (baseline)	54.0 (baseline)	49.3 (baseline)
Alcohol drinking (g/week)	NA	76.5	73.9	84.8
Non-drinkers	NA	2,874(12%)	1,298 (22%)	30(0.3%)
Current smokers	NA	5,929 (25%)	1,538 (26%)	3,478 (35%)
Higher education	NA	8,612 (36%)	1,721 (28%)	1,192 (12%)
Marriage or co-habitation	NA	17,468 (73%)	4,151 (68%)	6,661 (67%)
GGT (U/I)	NA	33.8	36.6	NA

Alcohol consumption

In a meta-analysis of the three cohorts with alcohol consumption estimates available ($n=39,888$), the PRS for alcohol consumption was strongly associated with self-reported alcohol consumption. A one SD increase in the PRS was associated with an 11.2 g (= 0.93 drinks $\times$ 12g) increase in weekly pure alcohol intake (beta=11.2 [9.85-12.6 g], $p = 2.3 \times 10^{-58}$) (**Fig 1, cohort-specific figures in the supplementary material**). Adding the PRS to the model improved r^2 by ~ 0.6 percentage points (from 9.17 % to 9.80 %). In addition, the PRS was negatively associated with alcohol abstinence (reported alcohol consumption 0). In FINRISK and Health2000, a 1 SD increase in the PRS for alcohol consumption was associated with a 13.7% reduced likelihood of being a nondrinker (OR=0.863 [0.833-0.895], $p=6.1 \times 10^{-16}$) while this was not case in the Twin Cohort where there were only 30 nondrinkers (OR=0.999[0.998-1.00] $p=0.31$). Cohort-specific figures for sex-specific alcohol consumption are in the supplementary material.

Figure 1. Alcohol drinking (g/week) for the deciles of the alcohol consumption polygenic risk score shown for males ($n= 18,887$) and females ($n= 20,808$) with 95% confidence interval error bars ($n=39,695$)

219

220 Alcohol-related morbidity

221 The PRS for alcohol consumption was strongly associated with increased risk for lifelong major
 222 alcohol-related events derived from electronic health-records in the FinnGen dataset (n=96,499,
 223 cases = 4,785) (**Figure 2**). The difference in the risk for alcohol-related morbidity events between
 224 the lowest and highest risk quintiles in the PRS was 67 % (HR=1.67 [1.52-1.84], $p=3.2 \times 10^{-27}$) and a 1
 225 SD increase in the PRS was associated with a 21 % increase in risk (HR=1.21 [1.18-1.25], $p=1.4 \times 10^{-40}$).
 226 The association was similar in both males (HR=1.21 [1.17-1.26], $p=1.8 \times 10^{-28}$) and females
 227 (HR=1.22 [1.16-1.29], $p=4.3 \times 10^{-14}$).

228

229 **Figure 2.** The FinnGen dataset was divided into three groups consisting of the lowest quintile, three
 230 middle quintiles and the highest quintile of the alcohol consumption PRS. The cumulative disease
 231 rate of alcohol-related morbidity is displayed as a function of age (n=96,499).

In the cohorts where alcohol consumption estimates and other related baseline data were available at the cohort entry time, the PRS was associated with an increased risk of incident major alcohol-related events and the association was maintained also in the fully adjusted model ($n = 28,639$, cases = 911). In a meta-analysis of the two cohorts 1 PRS SD was associated with a 26% increased risk of incident alcohol-related events when the consumption-estimate was not in the model ($HR=1.26$ [1.18-1.34], $p=1.1 \times 10^{-12}$) and with a 15 % increase when alcohol consumption was in the model ($HR=1.15$ [1.08-1.22], $p=2.1 \times 10^{-5}$). In a fully adjusted model, including marital status, education, smoking status and GGT, the estimate were unchanged ($HR=1.15$ [1.08-1.22], $p=2.0 \times 10^{-5}$) (Table 2).

Mortality

We observed a similar increase in the risk of alcohol-related and all-cause mortality. In FinnGen with 7,249 deaths one SD increase in the PRS for alcohol consumption was associated with 7 % increase in the risk of death ($HR= 1.07$ [1.05-1.10], $p = 4.3 \times 10^{-9}$). The risk estimate between the highest and lowest 20 % in the PRS was 1.23 ($HR = 1.23$ [1.14-1.23], $p = 1.22 \times 10^{-8}$). In our prospective cohorts, with cause-of-death information available, 4,125 deaths were recorded (Table 2). For all-cause mortality there was 11 % increase in the risk of death per 1 PRS SD in the basic model ($HR=1.11$ [1.07-1.14], $p = 3.2 \times 10^{-10}$) and 9 % in the fully adjusted model ($HR = 1.09$ [1.06-1.12], $p=1.1 \times 10^{-7}$). The risk difference between the highest and lowest quintiles of the PRS was 33 % ($HR=1.33$ [1.2-1.47], $p=4.5e-08$) in the fully adjusted model.

Of the 4,125 deaths 335 were known to be alcohol-related. Without alcohol consumption in the model, the increase in alcohol-related mortality was 26 % per 1 PRS SD ($HR=1.26$ [1.13-1.4], $p=3.7 \times 10^{-5}$). When alcohol consumption was included in the model, the increase was 13 % ($HR=1.13$ [1.01-1.26], $p=0.027$) and in a model with all co-variates, 11 % ($HR=1.11$ [0.996-1.24], $p=0.058$) (Table 2). Similarly, the PRS was associated with a higher risk of death from other than alcohol-related causes ($n=3,790$) when fully adjusted for all covariates ($HR=1.08$ [1.05-1.12], $p=1.4 \times 10^{-6}$).

Table 2. Cohort specific and meta-analyzed associations between the alcohol consumption polygenic risk score and alcohol-related **a)** morbidity and **b)** mortality. In the fully adjusted model age, sex, alcohol consumption, smoking, education, marital status and GGT (U/l) were used as non-genetic covariates.

	FINRISK	Health 2000	Meta-analysis
a) Alcohol-related morbidity	Cases=817	Cases=171	Cases=988
Basic model with age and sex	HR=1.25 [1.16-1.34], p=5.9*10 ⁻¹⁰	HR=1.32 [1.13-1.53], p=0.00036	HR=1.26 [1.18-1.34], p=1.1*10 ⁻¹²
Model with alcohol consumption	HR=1.13 [1.06-1.21], p=0.00053	HR=1.23 [1.06-1.43], p=0.0081	HR=1.15 [1.08-1.22], p=2.1*10 ⁻⁵
Fully adjusted model	HR=1.14 [1.06-1.22], p=0.00027	HR=1.20 [1.03-1.4], p=0.022	HR=1.15 [1.08-1.22], p=2.0*10 ⁻⁵
b) Alcohol-related mortality	Deaths=264	Deaths=71	Deaths=335
Basic model with age and sex	HR=1.21 [1.07-1.37], p=0.0022	HR=1.41 [1.11-1.8], p=0.0045	HR=1.25 [1.12-1.4], p=5.9*10 ⁻⁵
Model with alcohol consumption	HR=1.08 [0.952-1.22], p=0.24	HR=1.34 [1.05-1.71], p=0.017	HR=1.13 [1.01-1.26], p=0.033
Fully adjusted model	HR=1.08 [0.957-1.23], p=0.21	HR=1.22 [0.965-1.55], p=0.096	HR=1.11 [0.996-1.24], p=0.058

DSM-IV Alcohol-use disorder

The PRS was also associated with an interview-based DSM-IV alcohol use disorder diagnosis in the Nicotine Addiction Genetics Family cohort (440 cases, 1,140 controls) and a subset of FinnTwin16 cohort (273 cases, 320 controls). A meta-analysis of the two cohorts (713 cases) resulted in a combined 20 % increase in the prevalence of AUD per 1 PRS SD (OR = 1.20 [1.11-1.31], p = 2.29*10⁻⁵) in the unadjusted model. Adjusting for marital status, education and smoking explained part of

the effect (OR=1.14 [1.02-1.28], p=0.023) and further adjusting with maximal amount of drinks taken explained most of the effect (OR=1.06 [0.94-1.19], p=0.35).

Prediction

The predictive performance of the PRS was evaluated in the Health 2000 cohort (5,732 complete cases, 110 events) with a follow-up-window of 10 years based on the Cox model trained in the FINRISK cohort (18427 complete cases with ≥ 10 years of follow-up, 628 events). In a model not including the alcohol consumption estimate, adding the PRS to the model increased the C-index by 0.020, from 0.69 to 0.71 ($p = 0.017$). Both IDI (0.00242 [0.00102-0.00383], $p = 7.3 \times 10^{-4}$) and NRI (0.335 [0.146-0.523], $p = 5.1 \times 10^{-3}$) shifts were positive and statistically significant. When the log-transformed alcohol consumption estimate was included, a modest improvement of prediction was observed (C-index=0.0022 from 0.812 to 0.814, p-value=0.30; NRI=0.308 [0.119-0.497], $p = 0.0014$ and IDI=0.00173 [0.000726-0.00305], $p = 0.017$). Similarly, a modest gain was observed when adding PRS to a model with all available covariates including also marital status, education status, smoking status and GGT (C-index=0.00183 from 0.847 to 0.849, $p=0.44$; NRI=0.235 [0.0461-0.423], $p=0.015$; IDI =0.00331[0.0000254-0.00659], $p = 0.048$).

Discussion

We developed a highly polygenic risk score for alcohol consumption by obtaining weights from a recently published large-scale discovery sample and showed that the PRS was strongly associated with alcohol consumption in independent biobank cohort samples. An increased polygenic burden for alcohol consumption was associated with higher incidence of major alcohol-induced health events. The associations remained significant when we accounted for self-reported alcohol consumption and other relevant covariates; in a fully adjusted model the relative risk-estimate between the highest and lowest quintiles of the polygenic risk score was 1.6. Furthermore, the PRS was also associated with both alcohol-related, non-alcohol related and all-cause mortality.

Comparison with other studies

Our PRS shows the utility of genetic information for prediction of alcohol-related harm. The PRS, developed from a genetic analysis of cross-sectional self-reported alcohol consumption, was

associated with future risk of major alcohol-related health events. While a large number of PRSs have already been established for various traits and diseases (11), the development of PRSs for behavioral traits, such as substance use, has until now been limited (24-27) and the studies have not assessed their impact on future major health events.

Implications

Our results show that using a large sample size with long follow-up, we were able to build a PRS of alcohol consumption that is associated not only with alcohol consumption in independent samples, but also with future incident alcohol-related health events. In line with the knowledge that alcohol consumption is a major contributor to the worldwide burden of death, especially among working-age adults (1), we found the PRS to be associated also with all-cause mortality, further highlighting the importance of alcohol drinking as a cause of premature death.

Our score provides a genetic basis for potentially identifying a subset of high-risk individuals even early on in life, with potential for more targeted prevention of AUDs and other alcohol-related morbidity. Prevention is a cost-effective and efficient strategy to reduce alcohol related harms (28) and it is labeled one of the United Nations main health-related worldwide strategies of sustainable development (<https://sustainabledevelopment.un.org/sdg3>). A higher genetic predisposition for alcohol-related harms was detected both in the presence and absence of alcohol consumption data, as our PRS predicted alcohol-related harms beyond self-reported alcohol consumption. Health services are encouraged to support initiatives for screening and brief interventions for harmful drinking (29) as an effective strategy for tackling alcohol-related harm.(30) Thus, genetic information could potentially be used to improve the arsenal of possible strategies to detect high-risk individuals for targets of brief interventions. The fact that individuals in the highest PRS quintile showed an elevated risk for alcohol-related health events even in fully adjusted models could justify the use of genetic information even in clinical settings where a detailed history of alcohol consumption estimates, AUDIT-scores, or similar information are attainable. Communicating the information of higher risk for alcohol-related harm to patients could serve as a motivator for reducing drinking or committing to abstinence. The effect of being aware of one's negative expectations could also be unwanted, as is thought to be the case in the stereotype threat phenomenon. (31-34)

339

340 Self-reported alcohol consumption is known to be biased and problematic in terms of reliability and
341 validity for predicting alcohol-related risks (35,36). Some bias derives from true measurement error,
342 but another source is the lifelong temporal fluctuation of alcohol-drinking patterns not captured by
343 a measure at one single timepoint. Our PRS was associated with alcohol-related harms even when
344 adjusting for self-reported alcohol consumption estimate. One potential reason for this is that the
345 PRS contains information from the latent genetic predisposition for alcohol consumption, thus
346 overriding both the true measurement error and temporal fluctuations in alcohol drinking volume.

347

348 It has been hypothesized that alcohol consumption-based genetic discovery might inform more
349 about low-level drinking than about problematic drinking and AUDs.(37) However, we built a
350 polygenic risk score for alcohol consumption and successfully used it to predict alcohol-related
351 harms. Due to the robustness of a self-reported single timepoint alcohol consumption estimate and
352 the fact that different alcohol-related traits are to some degree genetically distinct,(9,10) it is
353 expected that a PRS developed directly for alcohol-related morbidity will outperform our PRS in
354 predicting alcohol-related health burden. Supporting this assumption, the general pattern is that
355 PRSs are more strongly associated with their respective diseases than with related phenotypes.
356 (13,14) Unfortunately, no high-quality summary statistics for alcohol-related harms including both
357 somatic and psychiatric outcomes yet exist; the performed GWAS have only covered alcohol
358 dependence (10) and been far smaller in size than our discovery sample of choice, thus making
359 future efforts for large-scale GWAS discovery based on alcohol-related harms more than necessary.

360

361 **Strengths and limitations**

362

363 Our polygenic risk score was derived using European ancestry discovery samples and tested in the
364 Finnish population. Its applicability in other populations therefore needs further evaluation as the
365 alcohol-related genetic mechanisms may vary between populations. However, it has to be noted
366 that the PRS derived from a non-Finnish sample performed well in the Finnish dataset, even though
367 Finns are somewhat genetically different from the rest of the Europeans (38).

368

369 Our design allowed us to study outcomes prospectively. Our registry-based follow-up captures
370 alcohol-related outpatient and inpatient visits, withdrawal treatment prescription for alcoholism,

and deaths, thus covering major alcohol-related health events over several decades. Nonetheless, some of the milder cases of alcohol-related health problems could have gone undetected.

Conclusions

In conclusion, a polygenic risk score for alcohol consumption was associated with elevated risk for incident alcohol-related health events and all-cause mortality. These findings underline the importance of heritable factors driving alcohol-related behavior. A successful attempt to predict alcohol-related health outcomes with a polygenic risk score shows promise in possible future utilization of genetic information in risk estimation and prediction of alcohol-related harms.

Footnotes

Contributors

All authors contributed to the study concept and design, analysis and interpretation of the data, as well as to the critical revision of the manuscript for important intellectual content or additionally to data acquisition. GSCAN provided the GWAS summary statistics. TK, TP, NJM and JKa performed the statistical analyses with support from SRi, ASH, JKo and SRu. ASH connected the data to the registries. TK drafted the manuscript. The corresponding author attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted.

Funding

SR was supported by the Academy of Finland Center of Excellence in Complex Disease Genetics (Grant No 312062), Academy of Finland (Grant No 285380), the Finnish Foundation for Cardiovascular Research, the Sigrid Juselius Foundation and University of Helsinki HiLIFE Fellow grant. The funding agencies had no role in the design and conduct of the study; collection, analysis, and interpretation of data; or the writing of the manuscript or the decision to submit it for publication. The FinnGen project is funded by two grants from Business Finland (HUS 4685/31/2016 and UH 4386/31/2016) and nine industry partners (AbbVie, AstraZeneca, Biogen, Celgene, Genentech, GSK, Merck/MSD, Pfizer and Sanofi). The Finnish Twin Cohort Nicotine Addictions Genetics family study has been supported by NIH DA12854 to PAFM, genotyping in the twin cohort by Global Research Awards for Nicotine Dependence (GRAND) funded by Pfizer, Inc to JK and the Welcome Trust Sanger Institute, and the Finntwin16 study by NIH AA-12502, AA-00145, and AA-09203 to RJR. JKa has been supported by the Academy of Finland (grants 265240, 263278, 308248, 312073). TK, JTR, SRu and PR were supported by the Doctoral Programme in Population Health, University of Helsinki. JTR was supported by the MD/PhD Program of the Faculty of Medicine, University of Helsinki.

Ethical approval

The study was approved by the Coordinating Ethical Committee of the Helsinki and Uusimaa Hospital District, reference numbers HUS/990/2017 (FinnGen), 246/13/03/00/15, 113/E3/2001 and HUS/1169/2016 (The Twin Cohort). The FINRISK and Health 2000 data are stored in the THL Biobank.

The transfer of the FINRISK and Health 2000 sample collections to the THL biobank has been approved by the Coordinating Ethics Committee of Helsinki University Hospital on 10th October 2014 and by the Ministry of Social Affairs and Health on 9th March 2015. This study was conducted under the THL biobank permission BB2017_64 (FINRISK and Health 2000). No additional ethical approval was needed for meta-analysing the results.

Competing financial interests

All authors have completed the ICMJE uniform disclosure form at http://www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author). AP reports that part of his salary is received from the FinnGen project that is partially funded by nine industry partners AbbVie, AstraZeneca, Biogen, Celgene, Genentech, GSK, Merck/MSD, Pfizer and Sanofi. JKa reports grants from Academy of Finland, grants from US-PHS NIH/NIDA, grants from US-PHS NIH/NIAAA, grants from Pfizer Inc/GRAND program grant, grants from EU FP7, during the conduct of the study. VS reports a conference trip and an honorarium for participating in an advisory board meeting from Novo Nordisk (unrelated to the present study) and a grant to his institute for research collaboration from Bayer (unrelated to the present study); no other relationships or activities that could appear to have influenced the submitted work.

Acknowledgements

We would like to Lea Urpa for proofreading, and Sari Kivikko, Huei-Yi Shen and Ulla Tuomainen for management assistance. We thank all participants of the study cohorts for their generous participation.

References

- (1) World Health Organization. Global status report on alcohol and health 2018. 2018.
- (2) Vengeliene V, Bilbao A, Molander A, Spanagel R. Neuropharmacology of alcohol addiction. *Br J Pharmacol* 2008;154(2):299-315.
- (3) Grant BF, Goldstein RB, Saha TD, Chou SP, Jung J, Zhang H, et al. Epidemiology of DSM-5 alcohol use disorder: results from the National Epidemiologic Survey on Alcohol and Related Conditions III. *JAMA psychiatry* 2015;72(8):757-766.
- (4) Wood AM, Kaptoge S, Butterworth AS, Willeit P, Warnakula S, Bolton T, et al. Risk thresholds for alcohol consumption: combined analysis of individual-participant data for 599 912 current drinkers in 83 prospective studies. *The Lancet* 2018;391(10129):1513-1523.
- (5) Nutt DJ, King LA, Phillips LD. Drug harms in the UK: a multicriteria decision analysis. *The Lancet* 2010;376(9752):1558-1565.
- (6) Dick, DM, et al. The Genetics of Substance Use and Substance Use Disorders, in: Kim Y, Kim Y (editor). *Handbook of behavior genetics*. Springer; 2009; pp 436
- (7) Clarke T, Adams MJ, Davies G, Howard DM, Hall LS, Padmanabhan S, et al. Genome-wide association study of alcohol consumption and genetic overlap with other health-related traits in UK Biobank (N= 112 117). *Mol Psychiatry* 2017;22(10):1376.
- (8) Liu M, Jiang Y, Wedow R, Li Y, Brazel DM, Chen F, et al. Association studies of up to 1.2 million individuals yield new insights into the genetic etiology of tobacco and alcohol use. *Nat Genet* 2019;1.
- (9) Sanchez-Roige S, Palmer AA, Fontanillas P, Elson SL, 23andMe Research Team, Substance Use Disorder Working Group of the Psychiatric Genomics Consortium, et al. Genome-wide association study meta-analysis of the Alcohol Use Disorders Identification Test (AUDIT) in two population-based cohorts. *Am J Psychiatry* 2018;appi. ajp. 2018.18040369.
- (10) Walters RK, Polimanti R, Johnson EC, McClintick JN, Adams MJ, Adkins AE, et al. Transancestral GWAS of alcohol dependence reveals common genetic underpinnings with psychiatric disorders. *Nat Neurosci* 2018;21(12):1656.
- (11) Khera AV, Chaffin M, Aragam KG, Haas ME, Roselli C, Choi SH, et al. Genome-wide polygenic scores for common diseases identify individuals with risk equivalent to monogenic mutations. *Nat Genet* 2018;50(9):1219.
- (12) Holmes MV, Asselbergs FW, Palmer TM, Drenos F, Lanktree MB, Nelson CP, et al. Mendelian randomization of blood lipids for coronary heart disease. *Eur Heart J* 2014;36(9):539-550.
- (13) Richardson TG, Harrison S, Hemani G, Smith GD. An atlas of polygenic risk score associations to highlight putative causal relationships across the human phenome. *eLife* 2019;8:e43657.
- (14) Khera AV, Chaffin M, Wade KH, Zahid S, Brancale J, Xia R, Distefano M, Senol-Cosar O, Haas ME, Bick A, Aragam KG. Polygenic prediction of weight and obesity trajectories from birth to adulthood. *Cell*. 2019 Apr 18;177(3):587-96.
- (15) Borodulin K, Vartiainen E, Peltonen M, Jousilahti P, Juolevi A, Laatikainen T, et al. Forty-year trends in cardiovascular risk factors in Finland. *The European Journal of Public Health* 2014;25(3):539-546.
- (16) Heistaro S. Methodology report: Health 2000 Survey. Helsinki: Publications of the National Health Institute 2008.
- (17) Kaprio J. The Finnish twin cohort study: an update. *Twin Research and Human Genetics*. 2013 Feb;16(1):157-62.
- (18) Kaprio J. Twin studies in Finland 2006. *Twin research and human genetics*. 2006 Dec;9(6):772-7.
- (19) Vilhjálmsson BJ, Yang J, Finucane HK, Gusev A, Lindström S, Ripke S, et al. Modeling linkage disequilibrium increases accuracy of polygenic risk scores. *The American Journal of Human Genetics* 2015;97(4):576-592.
- (20) Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ. Second-generation PLINK: rising to the challenge of larger and richer datasets. *Gigascience* 2015;4(1):7.
- (21) Antolini L, Nam B, D'Agostino RB. Inference on correlated discrimination measures in survival analysis: a nonparametric approach. *Communications in statistics-Theory and Methods* 2004;33(9):2117-2135.
- (22) Pencina MJ, D'Agostino Sr RB, D'Agostino Jr RB, Vasan RS. Evaluating the added predictive ability of a new marker: from area under the ROC curve to reclassification and beyond. *Stat Med* 2008;27(2):157-172.
- (23) Pencina MJ, D'Agostino Sr RB, Steyerberg EW. Extensions of net reclassification improvement calculations to measure usefulness of new biomarkers. *Stat Med* 2011;30(1):11-21.
- (24) Clarke T, Smith AH, Gelernter J, Kranzler HR, Farrer LA, Hall LS, et al. Polygenic risk for alcohol dependence associates with alcohol consumption, cognitive function and social deprivation in a population-based cohort. *Addict Biol* 2016;21(2):469-480.
- (25) Belsky DW, Moffitt TE, Baker TB, Biddle AK, Evans JP, Harrington H, et al. Polygenic risk and the developmental progression to heavy, persistent smoking and nicotine dependence: evidence from a 4-decade longitudinal study. *JAMA psychiatry* 2013;70(5):534-542.

- (26) Vink JM, Hottenga JJ, de Geus EJ, Willemsen G, Neale MC, Furberg H, et al. Polygenic risk scores for smoking: predictors for alcohol and cannabis use? *Addiction* 2014;109(7):1141-1151.
- (27) Marioni RE, Ritchie SJ, Joshi PK, Hagenaars SP, Okbay A, Fischer K, Adams MJ, Hill WD, Davies G, Nagy R, Amador C. Genetic variants linked to education predict longevity. *Proceedings of the National Academy of Sciences*. 2016 Nov 22;113(47):13366-71.
- (28) Spoth RL, Gyuill M, Day SX. Universal family-focused interventions in alcohol-use disorder prevention: cost-effectiveness and cost-benefit analyses of two interventions. *J Stud Alcohol* 2002;63(2):219-228.
- (29) World Health Organization. Global strategy to reduce the harmful use of alcohol. 2010.
- (30) Vandenberg B. Tackling Harmful Alcohol Use: Economics and Public Health Policy FRANCO SASSI (Ed) Paris, France: OECD Publishing, 2015ISBN: 978-92-64-18106-9 (pdf), 240 pp. Paperback and E-book. Available free online. *Drug Alcohol Rev* 2016;3(35):367-368.
- (31) Aronson J, Burgess D, Phelan SM, Juarez L. Unhealthy interactions: The role of stereotype threat in health disparities. *Am J Public Health* 2013;103(1):50-56.
- (32) Burgess DJ, Warren J, Phelan S, Dovidio J, Van Ryn M. Stereotype threat and health disparities: what medical educators and future physicians need to know. *Journal of general internal medicine* 2010;25(2):169-177.
- (33) Steele CM, Aronson J. Stereotype threat and the intellectual test performance of African Americans. *J Pers Soc Psychol* 1995;69(5):797.
- (34) Spencer SJ, Steele CM, Quinn DM. Stereotype threat and women's math performance. *J Exp Soc Psychol* 1999;35(1):4-28.
- (35) Gmel G, Rehm J. Measuring alcohol consumption. *Contemporary Drug Problems* 2004;31(3):467-540.
- (36) Litten RZ, Allen JP. Measuring alcohol consumption: Psychosocial and biochemical methods. : Springer Science & Business Media; 2012.
- (37) Edenberg HJ, Gelernter J, Agrawal A. Genetics of Alcoholism. *Curr Psychiatry Rep* 2019;21(4):26.
- (38) Lek M, Karczewski KJ, Minikel EV, Samocha KE, Banks E, Fennell T, et al. Analysis of protein-coding genetic variation in 60,706 humans. *Nature* 2016;536(7616):285.