Title: Parent and offspring genotypes influence gene expression in early life

Running Title: Parental effects on offspring gene expression

Authors & Affiliations:

Daniel J. Newhouse^{1*}, Margarida Barcelo-Serra², Elaina M. Tuttle², Rusty A. Gonser²,
Christopher N. Balakrishnan¹

¹East Carolina University, ²Indiana State University

*Corresponding Author

Abstract

Parents can have profound effects on offspring fitness. Little, however, is known about the mechanisms through which parental genetic variation influences offspring physiology in natural systems. White-throated sparrows (*Zonotrichia albicollis*, WTSP) exist in two genetic morphs, tan and white, controlled by a large polymorphic supergene. Morphs mate disassortatively, resulting in two pair types: tan male x white female (TxW) pairs, which provide biparental care and white male x tan female (WxT) pairs, which provide female-biased care. To investigate how parental composition impacts offspring, we performed RNA-seq on whole blood of WTSP nestlings sampled from nests of both pair types. Parental pair type had a large effect on nestling gene expression, with 881 genes differentially expressed (DE) and seven correlated gene co-expression modules. The DE genes and modules up-regulated in WxT nests with female-biased parental care primarily function in metabolism and stress-related pathways resulting from the overrepresentation of proteolysis and stress response genes (e.g. SOD2, NR3C1). These results show that parental genotypes and/or associated behaviors influence nestling physiology, and highlight avenues of further research investigating the ultimate implications for the maintenance of this polymorphism. Nestlings also exhibited morph-specific gene expression, with 92 differentially expressed genes, comprising innate immunity genes and genes encompassed by the supergene. Remarkably, we identified the same regulatory hub genes in these blood-derived expression networks as were previously identified in adult WTSP brains (EPM2A, BPNT1, TAF5L). These hub genes were located within the supergene, highlighting the importance of this gene complex in structuring regulatory networks across diverse tissues.

Keywords

Transcriptome, parental effects, early life stress, nestling, RNAseq, ornithology

Introduction

Parents can have profound impacts on offspring development and fitness. Parental effects can manifest throughout the developmental period, both pre- and post-natally (reviewed in Meaney 2001, Lupien et al. 2009) and can be mediated through parental behaviors, genetics and physiology during early development (Trivers 1972). Parents play a substantial role in establishing the early life environment of offspring. For example in birds, parental decisions on nest placement, incubation behavior, and nest defense could strongly impact developmental conditions of the egg. These parental behaviors will impact exposure to sunlight, humidity, temperature, and other environmental impacts of the eggs, which can influence developmental physiology (e.g. Nord & Nilsson 2011). In addition to parental behaviors, prenatal effects often arise via physiological maternal effects. Developing offspring are susceptible to the maternally created environment (e.g. maternal hormones, immune state, nutrition), which influence offspring physiology (Mousseau & Fox 1998, Jacquin et al. 2012; reviewed in Gluckman et al. 2008, Wolf & Wade 2009, Cottrell & Secki 2009).

The magnitude of parental effects, particularly in altricial species, is likely largest during the postnatal period, when offspring rely entirely on the parents for provisioning and growth (Royle et al. 2012). Provisioning plays a prominent role in offspring development, with the quality and quantity of food items crucial for offspring development (van Oers et al. 2015, Griebel et al. 2019). Similar to the prenatal stage, parental behaviors could also have strong impacts on offspring physiology. In many species, offspring are left alone during parental foraging trips, increasing environmental exposure (Lloyd and Martin 2004) and predation risk

(Lima 2009). Parental separation can also increase offspring anxiety (Millstein & Holmes 2007). Siblings must also compete to optimize provisioning, brooding warmth, and preening (Mock & Parker 1997). Thus, this postnatal environment, largely mediated through parental effects, can be a potential source of early life stress (ELS) in offspring, which may result in life-long fitness effects (reviewed in Monaghan 2014).

ELS has broad effects on organisms, including impaired neural development, neuroendocrine signaling, behavior, and physiology (McEwen 2007, Monaghan 2014). For example, ELS is associated with impaired neuroendocrine function and corresponding impaired hypothalamic-pituitary-adrenal (HPA) development, which leads to increase stress response sensitivity later in life (e.g. Heim et al. 2008, Spencer et al. 2009, Crespi et al. 2012, Spencer 2017). ELS can exacerbate behavioral alterations as organisms develop and mature including symptoms of anxiety and depression in the postnatal environment (Noguera et al. 2017) and result in impaired behavior as reproductive adults (e.g. Krause et al. 2009, reviewed in Bolton et al. 2017). While the organismal effects of ELS are well studied, the genetic underpinnings are relatively underexplored. Much of the genetic work in the context of ELS has focused on gene regulatory impacts, particularly in mammalian biomedical models (reviewed in Szyf et al. 2007, Szyf 2009, Silberman et al. 2016, Alyamani & Murgatroyd 2018). In particular, the quality of parental care can have strong impacts on offspring health resulting from epigenetic modifications (Liu et al. 1997, Meaney 2001, Weaver et al. 2004). These gene regulation studies primarily use changes in DNA methylation as an indicator of ELS (Murgatroyd et al. 2009, Kinnally et al. 2011, Lewis & Olive 2014) and recent work has expanded these approaches into non-mammalian organisms (e.g. Rubenstein et al. 2016, Moghadam et al. 2017, Pértille et al. 2017, Gott 2018, Sheldon et al. 2018). DNA methylation studies of ELS investigate changes to the structure of

DNA, but are often limited in the functional implications of ELS (i.e. transcription and translation). In general, these modifications are thought to alter transcriptional activity of genes in the modified genomic region (Berger 2007, Lowdon et al. 2016). Indeed, several studies have also taken candidate gene approaches to investigating gene expression in the context of ELS (Marco et al. 2014, Diaz-Real et al. 2017, Anastasiadi et al. 2018, Reshetnikov et al. 2018). However, very few studies assess genome-wide transcription under ELS (Moghadam et al. 2017), particularly in the context of parental effects (but see: Weaver et al. 2006).

In this study, we examined the white-throated sparrow (*Zonotrichia albicollis*, WTSP) to assess the role of parental genotype on offspring gene expression. WTSPs exist in two plumage morphs, tan (T) and white (W), that are found in both sexes and in roughly equal frequencies (Lowther 1961). These morphs are genetically determined by alternative alleles of a supergene, a group of linked genes that are inherited together, show limited recombination, and maintain complex behavioral traits (i.e. WTSP morphs; Schwander et al. 2014, Taylor & Campagna 2016). The WTSP supergene resulted from a complex chromosomal rearrangement comprising multiple inversions (hereafter referred to as “inversion” or “inverted”). This inversion contains ~1,100 genes on chromosome two, termed ZAL2^m (Throneycroft 1975, Thomas et al. 2008, Romanov et al. 2009, Tuttle et al. 2016). W morphs are nearly always heterozygous for the inversion (ZAL2/ZAL2^m) and T morphs are always homozygous (ZAL2/ZAL2; Thorneycroft 1966, 1975).

This unusual polymorphism in WTSPs influences hormonal profiles and the behavior of both sexes, and thus has the potential to influence pre- and post-natal environments for the offspring of different morphs. W morph males maintain higher levels of testosterone during the pre-laying, incubation, and brooding stages and oestradiol during the laying and brooding stages (Horton et al. 2014). Only oestradiol has been shown to differ between adult female morphs

during the breeding season and is higher in W morph females during the pre-laying and laying stages (Horton et al. 2014). These genetic and hormonal differences also translate into striking behavioral differences. W morphs, for example, are highly territorial and sing frequently whereas T morphs are far less territorial and aggressive (Lowther 1962, Kopachena & Falls 1993, Tuttle 2003, Horton & Holberton 2010, Horton et al. 2014). More importantly from the perspective of offspring, males of each morph also differ in paternal investment (Knapton & Falls 1983, Horton et al. 2014). W morph males are promiscuous and provision nestlings very little. T morph males defend their within-pair paternity through mate guarding and are highly paternal. Females tend to provision at intermediate levels, but T morph females may compensate for unassisted care from W morph males and provision more than W morph females (Knapton & Falls 1983). A final wrinkle in this complex mating system is that morphs nearly always mate with the opposite morph (98.5%, Tuttle et al. 2016), resulting in two stable pair types: T male x W female (TxW) and W male x T female (WxT) (Lowther 1961, Tuttle 2003, Tuttle et al. 2016). Because males differ in paternal investment, this results in two distinct parental care strategies. TxW pairs provide biparental care and WxT pairs provide female-biased parental care. In this study we examined gene expression profiles of offspring from both pair-types in order to assess the physiological consequences of variation in parental genotype.

Methods

Field based sample collection

All nestling whole blood samples in this study came from a breeding population of WTSPs at the Cranberry Lake Biological Station in northern New York, USA (SUNY-ESF, 44.15°N, 74.78°W) and were collected during the 2015 breeding season. We only utilized

samples collected during the first clutch (June 6 - June 14, 2015), as WTSP males may increase paternal investment in replacement broods (Horton et al. 2014). We collected ~80µL blood in capillary tubes via brachial venipuncture on days 5-7 post-hatch. Approximately 60µL blood was preserved in Longmire's lysis buffer (Longmire et al. 1992) for genotyping and ~20µL was immediately placed in RNeasy. Within six hours of collection, samples were placed temporarily into liquid nitrogen, before being shipped overnight on dry ice to -80°C storage until RNA extraction. All animal sampling protocols were approved by the Indiana State University Institutional Animal Care and Use Committee (IACUC 562158-1:ET/RG, 562192-1:ET/RG).

Molecular sexing & genotyping

Nestling DNA was extracted from erythrocytes using the DNA IQ® magnetic extraction system (Promega Corp, Madison, WI USA). To determine sex and morph, we used PCR to fluorescently label and amplify a region of the chromo-helicase-DNA-binding gene, and a region of the vasoactive intestinal peptide following Griffiths et al. (1998) and Michopolous et al. (2007). The PCR products were run and analyzed on an ABI PRISM™ 310 genetic analyzer.

RNA extraction, library preparation, & sequencing

We sampled a total of 52 nestlings for RNA extraction, but due to issues with RNA quality after extraction, only 32 were used for sequencing. These samples represent 23 nestlings from eight TxW pairs and nine nestlings from three WxT pairs. Additionally, these data represent 18 females, 14 males, 15 T morph, and 17 W morph individuals.

We removed RNeasy and homogenized whole blood tissue samples with Tri-Reagent (Molecular Research Company). Total RNA was purified with a Qiagen RNeasy mini kit

(Valencia, CA, USA), followed by DNase treatment and further purification. We quality assessed RNA with an Agilent Bioanalyzer (RIN > 7) (Wilmington, DE, USA). Both library preparation and sequencing were performed at the University of Illinois Roy J. Carver Biotechnology Center. A library was prepared for each RNA sample using the Illumina HT TruSeq (San Diego, CA, USA) stranded RNA sample prep kit. Libraries were distributed into four pools with equimolar concentrations and quantitated via qPCR. Each of the pools was sequenced on an individual lane of an Illumina HiSeq 2500 using the Illumina TruSeq SBS sequencing kit v4 producing 100-nucleotide single-end reads.

Creation of masked reference genome

The WTSP reference genome was generated from a male T morph individual (Tuttle et al. 2016). Thus, the reference genome does not contain any sequence data from the ZAL2^m inversion. To avoid any potential bias in mapping reads derived from W morph individuals onto a T morph genome, we generated a masked reference genome for this study. To do so, we used previously published whole genome sequences from three W morph adults (Tuttle et al. 2016). Reads were adapter trimmed with *Trim Galore!* v0.3.8 (<https://github.com/FelixKrueger/TrimGalore>) and aligned to the WTSP reference genome with *bwa mem* v 0.7.10-r789 (Li 2013). We converted and sorted the resulting SAM alignment to BAM format with *samtools view* and *samtools sort*, respectively (*samtools* v1.2, Li et al. 2009). We then merged all genomic scaffolds corresponding to the ZAL2^m inversion, as identified in Tuttle et al. (2016), with *samtools merge*. We called SNPs within the inversion using *samtools mpileup* and *bcftools call* v 1.2 (Li et al. 2009, Li 2011). We only kept SNPs that were heterozygous in each of the three individuals with *SnpSift* v 4.3p (Cingolani et al. 2012) and used

these SNPs to mask the reference genome with *bedtools maskfasta* v 2.21.0 (Quinlan & Hall 2010).

Quality control, read mapping, differential expression, & gene ontology

We trimmed Illumina sequencing adapters from each of the 32 libraries with *Trim Galore!* v0.3.8 which uses *Cutadapt* v1.7.1 (Martin 2011). Trimmed reads were then mapped to the masked reference genome with *STAR* v2.5.3a (Dobin et al. 2013). The mapping results were then quantified and assigned gene IDs with *htseq-count* v0.6.0 (Anders et al. 2015) specifying ‘-s reverse’ and ‘-i gene’. We then removed lowly expressed genes by summing the counts for each gene across all 32 samples, dividing by 32 to obtain the study average, and removing genes with an average read count of < 5.

All statistical analyses were performed with R v3.5.0 (R Core Team 2013). We first identified outlier samples based on visual inspection of sample distance in a dendrogram within *WGCNA* (Horvath 2011). Two samples, one T female and one T male representing an entire TxW nest, were identified as outliers and removed from all future analyses (Figure S1). Using the remaining 30 samples, we normalized reads accounting for sequencing depth and assessed differential expression with *DEseq2* (Love et al. 2014). We performed variance stabilizing transformation of reads in *DEseq2* and performed PCA and hierarchical clustering based on Euclidean distance of gene expression profiles with *pcaExplorer* v2.6.0 (Marini & Binder 2016). Differential expression analyses utilized pairwise comparisons between nestling morph and pair type (i.e. parental morphs). We controlled for sex in morph comparisons and sex, morph, and nest ID for pair type comparisons. To include nest ID in the pair type comparison, we followed the “individuals nested within groups” guide in the *DEseq2* manual. We did not include nestling

age in analyses, as most samples were 6 days old (n=21), limiting comparisons with nestlings aged Day 5 (n=3) or Day 7 (n=6). Network analysis (see below) did not reveal any effect of age on variables of interest (morph, pair type; data not shown). We also tested for an interaction between nestling morph and pair type utilizing a grouping variable as outlined in the *DEseq2* manual. *DEseq2* determines differential expression with a Wald test followed by Benjamini & Hochberg (1995) FDR correction. Genes were considered differentially expressed (DE) if the FDR corrected p-value was < 0.10. Details for each model run, including the R code used, are in this project's GitHub repository.

We next tested for gene ontology (GO) enrichment among DE genes with *GOrilla* (Eden et al. 2007, 2009). For each *DEseq2* comparison, we ordered the list of genes based on ascending FDR values, excluding any genes in which *DEseq2* did not assign a FDR value. The WTSP genome is not completely annotated, so any loci without a gene symbol were excluded from GO analyses (n=1,926). *GOrilla* places greater weight on genes located at the top of the list (i.e. DE genes), while accounting for the contribution of each gene in the given comparison. GO categories were considered significantly enriched if the FDR corrected p-value < 0.05. *GOrilla* does not support WTSP annotation; so, all analyses were based on homology to human gene symbols.

Weighted gene co-expression network analysis (WGCNA)

We used the *WGCNA* package in R (Zhang & Horvath 2005, Langfelder & Horvath 2008) to identify modules of co-expressed genes in our dataset. We first exported variance stabilizing transformed (vst) read counts from *DEseq2*, removed genes with an average vst < 5 averaged across all 30 samples, and imported the subsequent list of 8,982 genes into *WGCNA*. To

build the co-expression matrix, we chose a soft thresholding power (β) value of 12, at which the network reaches scale-free topology (Figure S2). We generated a signed network with minimum module size of 30 genes and merged highly correlated modules (dissimilarity threshold = 0.20, which corresponds to $R^2 = 0.80$). We then correlated the eigengene, which is the first principal component of a module, of these merged modules with external traits (pair type, morph, sex, nest ID). Modules with $p < 0.05$ were considered significantly correlated with a given trait. For all morph-specific results, we tested for an enrichment of inversion genes with a chi-squared test using a Fisher's exact test ($p < 0.05$).

To visualize the interaction of genes within a module, we generated the intramodular connectivity (IM) score for each gene, which represents the interconnection of module genes. We exported all IM scores for modules of interest and imported into *VisAnt* v5.51 (Hu et al. 2013) for visualization. To maximize network clarity, we only plotted the top 300 interactions based on IM scores. Thus, we only visualized the most connected genes. To identify hub genes, we visualized the Degree Distribution (DD) for the network and selected the most connected genes above a natural break in the distribution. This resulted in one to nine hub genes per module.

To understand the biological function of modules correlated with traits of interest, we performed a target vs background GO analysis in *GOrilla*. For each module, we tested the assigned genes for each module against the entire list of 8,982 genes used for the *WGCNA* analysis. GO categories were significant with a FDR corrected p-value < 0.05 .

Results

Sequencing results

We sequenced each sample to an average depth of 29.4 million reads (range = 16.2-58.5 million reads). The 32 libraries were distributed into four pools in equimolar concentration. One pool contained only four samples, which corresponded to the four samples with lowest RNA concentrations. This pool was sequenced to an average depth of 56.17 million reads per library. The remaining three pools were sequenced to an average depth of 25.62 million reads per library. Samples mapped to our masked genome at an average rate of 91.08% (range = 88.19%-92.87%) (Table S1). A total of 8,982 genes had count values ≥ 5 across all samples, which included 641 located in the W morph inversion. Samples did not segregate by pair type or morph in PCA or hierarchical clustering (Figures S3, S4).

Differential Expression – Morph

Ninety-two genes were differentially expressed between morphs. Sixty-five of these genes (71%) were located in the inversion, representing a significant enrichment ($\chi^2=553.73$, $df=1$, $p<0.00001$) (Table S2). The inversion represents only 641 out the 8,892 genes (7%) sampled here. Additionally, expression of many of these 92 genes was elevated in W morph nestlings and a number of these genes had well-known functions in innate immunity (e.g. IFIT5, IL20RA, EIF2AK2, RSAD2). There was GO enrichment of four categories, two of which are immunity related: “immune response” ($p = 0.019$) and “defense response to virus” ($p = 0.049$) (Table S3).

Differential Expression – Pair Type

Pair type had the largest effect on gene expression, with 881 genes DE between offspring from the two different pair types ($FDR < 0.10$, Table S2). Many genes associated with stress

responses were elevated in nestlings in WxT nests, including the glucocorticoid receptor (NR3C1), superoxide dismutase (SOD)1 & SOD2, DEP domain-containing mTOR-interacting protein (DEPTOR), and several ubiquitin-mediated proteolysis pathway genes (e.g. UBE2D3, PSMD3, PSMD6). Additionally, several immune system related genes were also elevated in WxT nests, including cytokines (e.g. IL2RA, IL7R), suppressor of cytokine signaling 1 (SOCS1), and five putative major histocompatibility complex (MHC) class I loci. No GO categories were significantly enriched, however.

We next tested for a morph-specific response to pair type. Within WxT nests, 40 genes were DE ($p < 0.10$) between T and W morph nestlings. Twelve of these genes (30%) are located within the inversion, again reflecting an enrichment of inversion genes among those differentially expressed between morph ($\chi^2=34.44$, $df=1$, $p<0.00001$). Only two genes (THSD7B & CFAP44) were DE between morphs within TxW nests, both of which are uniquely DE between morphs in TxW nests. No GO categories were enriched in either comparison.

WGCNA – Morph

WGCNA revealed 26 modules, five of which were correlated with morph (Table 1, Figure 1). The light cyan module (183 genes, $R^2=0.67$, $p=5 \times 10^{-5}$) and ivory module (72 genes, $R^2=-0.66$, $p=9 \times 10^{-5}$) contained genes elevated and suppressed, respectively, in W morph nestlings relative to T morph nestlings. These modules are both enriched for genes located within the chromosomal inversion (light cyan module = 70/183 (38%) genes, $\chi^2=266.49$, $df=1$, $p<0.00001$; ivory module = 40/72 (56%), $\chi^2=261.60$, $df=1$, $p<0.00001$) (Figure S5). The hubs of each of these modules are also located in the chromosomal inversion (Table 1, Figure S5). Additionally, the sky blue module (58 genes, $R^2=0.53$, $p=0.003$) and dark red module (102 genes,

$R^2=0.47$, $p=0.009$) (Figure S6) contained genes elevated in W morph nestlings and many of these genes overlap with the immune related genes described in the morph DE tests above. The hubs of these networks (e.g. sky blue: EIF2AK2, IFIT5, OASL; dark red: TRAF5) (Table 1) reflect a conserved innate immunity network structure in avian blood (Kernbach et al., in review) (Figure S6).

WGCNA – Pair Type

We found seven modules correlated with pair type (Table 2, Figure 1). The blue module represented genes that are elevated in nestlings from WxT nests (1,142 genes, $R^2 = -0.45$, $p=0.01$). This module contained both the largest number of genes and correspondingly strongest functional enrichment. Many of these GO enrichments were related to protein function, resulting from the presence of ribosomal genes. Interestingly, several GO categories for metabolism, catabolism, and proteolysis were also enriched, driven by genes encoding ubiquitin-conjugating enzymes and proteasome subunits (e.g. “proteasomal protein catabolic process”, $p=2.34 \times 10^{-4}$; “proteasome-mediated ubiquitin-dependent protein catabolic process”, $p=5.32 \times 10^{-4}$) (Table S4). Many of these (e.g. PSMF1, PSMD3, PSMD6, UBE2D2, UBE2D3, UBE3C) were also DE between offspring of the two pair types (Figure 2). Lastly, the blue module contains one hub gene, NDUFB3 (DD=42) (Figure 2), which is involved in the mitochondrial electron transport chain.

The beige and light green modules represented candidate stress response networks. These modules showed contrasting expression patterns in nestlings from WxT nests (Figures 4 & 5). Although not significantly enriched for any GO categories, the beige module comprised 335 genes that were upregulated in WxT nests relative to TxW nests ($R^2=-0.61$, $p=3 \times 10^{-4}$). DEPTOR,

which functions as an inhibitor of the mTOR pathway in response to stress (e.g. Desantis et al. 2015), was the single hub in the beige module (DD=39, Figure 3). The beige module also contained NR3C1, which is activated in response to increased glucocorticoid secretion. Lastly, the light green module (116 genes, $R^2=0.60$, $p=4 \times 10^{-4}$) contained genes with low expression in TxW nests relative to WxT nests. There were three hub genes (DD > 28), CDK19, CHD4, and EPG5, each with previously described roles in the stress response (Figure 4).

For each pair type module, the correlation was stronger for the overall effect of pair type than any individual nest, indicating that one nest did not drive the correlation. This trend was reflected in gene expression plots of hub genes and candidate genes described above (Figure S7). We did not observe modules correlated with pair type that were also correlated with nestling morph or sex, suggesting there is no morph or sex-specific response to a given pair type at the network level.

Discussion

By assessing genome-wide transcription in nestlings raised by different WTSP pair types we have identified distinct transcriptomic signatures that suggest WxT pairs induce a stress response in developing nestlings relative to TxW pairs. This is reflected both by differential expression of several genes involved in protein degradation as well as networks of co-expressed genes with stress response hubs. Additionally, we identified morph-specific gene expression driven by innate immunity genes and genes located in the chromosome 2 inversion. As adults, the genes within the inversion strongly influence the WTSP neural transcriptome (Balakrishnan et al. 2014, Zinzow-Kramer et al. 2015). Our results here suggest that as nestlings, parental genotypes and

associated behaviors, rather than nestling genotype, have the strongest influence on the nestling transcriptome.

Gene expression differences resulting from pair type

We find 881 genes DE between nestlings raised under the two pair types. Many of these genes function in the proteasome or ubiquitin-mediated proteolysis. Cells naturally use the proteasome for degradation of proteins targeted by the ubiquitination process, but genes involved in proteasome formation (e.g. PSMD6, PSMD11) and ubiquitination (e.g. UBE2B) are up-regulated in cells experiencing mild oxidative stress (Aiken et al. 2011, Shang & Taylor 2011, Livneh et al. 2016) or organisms experiencing abiotic stress (Dhanasiri et al. 2013, Tomalty et al. 2015). Thus, increased expression of these genes in nestlings from WxT nests suggests they are responding to oxidative stress. As a result, there may be a cost to having a W morph father and T morph mother at the nestling stage.

To complement our differential expression approach, we also constructed co-expression networks with *WGCNA*. *WGCNA* identifies modules of co-regulated genes blind to the experimental design. These modules are then correlated with external traits, offering a systems-level view into how conditions impact transcriptional networks. Within these networks, we can then perform GO analyses as described above and identify network hubs, which are the most highly connected genes within that network. Using this approach, we identified 26 modules of co-regulated genes in this dataset (Figure 1), seven of which were significantly correlated with parental pair type. The blue module contains genes that are elevated in nestlings in WxT nests. The blue module hub gene was NDUF3 (Module Membership [MM]=0.938, DD=42) (Figure 2), which encodes a subunit of the mitochondrial membrane respiratory chain. Interestingly,

many of the same proteolysis-related genes highlighted in the differential expression results are also present in this module, resulting in the enrichment of several metabolism and stress-related GO categories (Table S4).

Two modules, light green and beige, contained stress responsive hub genes. The light green module contains genes that are suppressed in nestlings in WxT nests, with three hub genes: CDK19, CHD4, and EPG5 (Figure 4). The absence of EPG5 expression (via knockout) and reduction in CHD4 expression (via knockdown) has been associated with increased DNA damage (Zhao et al. 2013, Larsen et al. 2010). Similarly, down-regulation of CDK19 following knockdown is associated with an increased stress response (Audetat et al. 2017). Suppression of these genes in these nestlings could be indicative of increased cellular damage. The beige module contains genes whose expression is elevated in nestlings from WxT nests and contains one hub gene, DEPTOR, which is an inhibitor of mTOR signaling (Figure 3). The exact role of DEPTOR remains unclear, but up-regulation likely inhibits the mTORC1 pathway to reduce endoplasmic reticulum stress, promote cell survival, and avoid apoptosis (Peterson et al. 2009, Desantis et al. 2015, Catena et al. 2016).

Increased expression of genes in the beige module in these nestlings and the high connectivity of DEPTOR to other co-expressed genes provide further support for a transcriptional stress response within WxT nests. The beige module also contains two well-studied stress responsive genes, superoxide dismutase 2 (SOD2) and the glucocorticoid receptor (NR3C1). SOD2 mitigates the effects of exposure to reactive oxygen species by scavenging free radicals (Zelko et al. 2002). NR3C1 binds glucocorticoids and has primarily been studied in the context of ELS and methylation of an upstream promoter. NRC3C1 methylation is often associated with down-regulation of NR3C1 (e.g. McGowan et al. 2009) and impairment of the

HPA axis, but up-regulation following methylation has also been observed as part of the stress response (Turner et al. 2006, Bockmühl et al. 2015). Increased expression observed here directly implicates the HPA axis and suggests these nestlings may be activating SOD2 and NR3C1 to cope with elevated levels of reactive oxygen species and corticosterone, respectively (Wang et al. 2018, Finsterwald & Alberini 2014). However, further work is needed to investigate stress physiology, corticosterone levels, and uncover the epigenetic state of NR3C1 in these nestlings and how this may relate to ELS (Banerjee et al. 2011, McCoy et al. 2016, Rubenstein et al. 2016, Quirici et al. 2016, Greggor et al. 2017).

How does parental genotype influence offspring gene expression?

In a non-experimental study, we have limited power to make inferences about the mechanism by which parental genotype impacted offspring gene expression. Given the well-studied reproductive biology of WTSPs, however, two mechanisms seem especially likely: hormone-mediated maternal effects and/or differences in parental provisioning. In weighing the evidence for these two non-mutually exclusive possibilities, we conclude that the difference in parental provisioning is the most plausible explanation for the observed gene expression differences. As described above, WTSP morphs differ in hormone levels. Only oestradiol, however, has been shown to differ between adult female morphs during the breeding season and is higher in W morph females during the pre-laying and laying stages (Horton et al. 2014). No baseline differences in any other hormone measured to date (corticosterone, testosterone, DHEA, DHT) have been described during the breeding season (Spinney et al. 2006, Swett & Breuner 2009, Horton & Holberton 2010, Horton et al. 2014). Taken together this suggests that hormone deposition into eggs may not differ dramatically between the morphs. By contrast, there is strong

evidence of differences in provisioning among morph types (Knapton & Falls 1983, Kopachna & Falls 1993, Horton & Holberton 2010, Horton et al. 2014). Reduced provisioning by W morph males appears to be stable across populations resulting in female-biased parental care in WxT nests (Knapton & Falls 1983, Horton et al. 2014). Therefore, parental care variation is a likely source of behaviorally mediated maternal or paternal effects (see Crean & Bonduriansky 2014) that would explain the strong signature of stress exposure in the expression data.

Previous work revealed no difference in clutch size between pair types (Knapton et al. 1984, Formica et al. 2004) and no effect of pair type on nestling mass (Knapton et al. 1984, Tuttle et al. 2017). Also, nestlings did not differ in mass at time of sampling between the TxW and WxT nests used in this study (Smith et al. in review). Increased provisioning by females to compensate for reduced care by males could explain this observation, and this has been observed previously in a separate WTSP population (Knapton & Falls 1983). In this scenario reduced brooding and increased maternal separation could also negatively impact nestling physiology and act as a source of ELS (reviewed in Ledón-Rettig et al. 2013). Somewhat surprisingly, given the gene expression findings described here, a recent study in our study population did not detect differences in reactive oxygen metabolites in plasma of offspring of the two different pair types (Grunst et al. 2019). ROM, however, only provides a limited overview of the stress response and the RNA-seq response we observed could even mitigate long-term consequences of ELS. The results here further highlight the utility of blood RNA-seq as a highly sensitive measure of environmental exposures (Louder et al. 2018).

Our study was carried out in the field as part of a long-term study and is limited by the fact that we did not perform a cross-fostering experiment. We aimed to mitigate potential environmental confounds by restricting sampling of nestlings to a short time period of nine days.

Certainly the environment may influence gene expression in our samples, but consistent changes among the samples in the two pair types suggest the role of parents is a significant driver of nestling gene expression, rather than temporal or spatial environmental variation.

Morph-specific gene expression

We were also interested in morph-specific gene expression and how nestling morph may respond to differences in parental pair type. WTSPs have been studied extensively as adults, but very rarely in other life stages. W morph males and T morph females exhibit earlier reproductive and actuarial senescence, potentially resulting from the high energy expenditure lifestyle of W morph males and biased parental care given by T morph females (Grunst et al. 2018a, Grunst et al. 2018b). There also appears to be seasonal variation in fitness between the morphs as adults. Following cold, wet winters, W morph males exhibit lower recruitment in the breeding grounds, leading to an overproduction of W morph male nestlings, potentially to stabilize morph frequencies in the population (Tuttle et al. 2017). Thus, morph specific differences may arise in early life. We found 92 genes DE between morphs, including many innate immune-related genes and genes located within the inversion (65/92 genes, Table S2). *WGCNA* revealed five modules correlated with morph (Figure 1). These included two innate immunity-related modules with increased expression in W morphs (Dark Red & Sky Blue) and two modules enriched with genes located in the inversion (Ivory = 40/72, Light Cyan = 70/183) (Figures S5, S6). The sky blue module contains nine hub genes and the dark red module contains one hub gene, both of which include well-studied anti-viral genes (e.g. sky blue: OASL, RSAD2; dark red: TRAF5). These genes also form a co-expression module in avian blood following West Nile virus infection (Kernbach et al., in review). Adult WTSP morphs differ in their ability to clear infection (Boyd et

al. 2018), so the immune activation here may be indicative of an increased parasite load in W morph nestlings, although further investigation is required. The light cyan module contains genes elevated in W morph nestlings and contains eight hub genes, each located in the inversion (Table 1). Three of these, EPM2A, BPNT1, and TAF5L, were also identified as hub genes in brain tissues of adult W morph males (Zinzow-Kramer et al. 2015). These nestlings thus exhibit expression differences in inversion genes prior to any phenotypic or behavioral differences, revealing the importance of the inversion in maintaining morph phenotypes throughout life. Additionally, the conservation of network hub genes in a different tissue and life stage highlights avenues for further investigation into WTSP gene regulation.

Despite broad gene expression differences between the morphs, within pair types morph-specific expression was limited. In part due to small sample size, nestlings in TxW nests only have two genes DE between morphs. There is a larger effect of morph within WxT nests, where the number of DE genes increased to 40. These genes encompassed a wide range of gene functions without any obvious stress-related candidate genes. Of these 40 genes, 34 are uniquely DE within WxT nests and do not overlap with the overall list of 92 genes DE between morphs using all samples. Interestingly, glucocorticoid-induced transcript 1 (GLCCI1) is elevated in W morph nestlings in WxT nests. The function of GLCCI1 remains unclear (Kim et al. 2016), but expression differences between morphs observed here implicates the role of glucocorticoids in response to pair type. This suggests that nestling morphs may respond differently to the parental pair type though larger sample sizes will be needed to explore this further.

Conclusions

Using the WTSP, a system with alternative parental care strategies, we show that nestlings in WxT nests (female-biased parental care) have increased expression of stress-related genes, and parental genotypes may act as a source of ELS in the species. Nestling morph also influences transcription, but parental pair type appears to have the greatest effect on their transcriptome. Combined, this supports the parental effects hypothesis (Wade 1998, Schrader et al. 2018), where offspring phenotypes are primarily a result of the nest environment and care received, rather than from offspring genotypes (i.e. T vs. W). Nearly 54% of observed pairs have been WxT (Tuttle et al. 2016). Thus, roughly half of the nestlings in every population will experience female-biased parental care. Our results suggest that these differences in parental pair type have at least short-term consequences on offspring physiology. While we have identified impacts at the level of transcription, an integrative approach assessing nestling WTSP physiology and performing cross-fostering experiments will further elucidate the consequences of variation in parental pair type. Importantly, it remains unclear whether female-biased parental care or differences in maternal effects translate into long-term fitness consequences for offspring. There appears to be a cost associated with parental genotype, as less cooperative reproductive strategy (WxT pairs) accelerates senescence (Grunst et al. 2018a, Grunst et al. 2018b). We show here that this cost is also translated into nestlings within WxT nests via increased stress-related gene expression. This work sets the stage to further explore morph-specific fitness consequences in nestlings experiencing alternative parental care strategies.

Acknowledgements

We acknowledge Lindsay Forrette, Andrea Grunst, and Melissa Grunst for assistance in the field, Sarah Ford for assistance with molecular work, Rachel Wright for WGCNA code, and Cranberry

Lake Biological Station. Funding provided by East Carolina University, Indiana State University, The National Science Foundation (grant no. DUE-0934648) and the National Institutes of Health (grant no. 1R01Gm084229 to E.M.T and R.A.G.) and a Sigma Xi Grants in Aid of Research award to DJN. Birds were banded with color bands and a Fish and Wildlife band (Master Banding Permit 22296 to EMT and permit 24105 to RAG). Dr. Alvaro Hernandez and Chris Wright provided guidance and oversight on sequencing carried out at the University of Illinois. All methods were conducted in accordance with legal and ethical standards and were approved by Indiana State University's Institutional Animal Care and Use Committee (protocols 562158-1:ET/RG and 562192-1:ET/RG).

References

- Aiken, C. T., Kaake, R. M., Wang, X., & Huang, L. (2011). Oxidative Stress-Mediated Regulation of Proteasome Complexes. *Molecular & Cellular Proteomics*, 10(5). Retrieved from <http://www.mcponline.org/content/10/5/R110.006924.abstract>
- Alyamani, R. A. S., & Murgatroyd, C. (2018). Epigenetic Programming by Early-Life Stress. In *Progress in Molecular Biology and Translational Science* (Vol. 157, pp. 133–150). doi: [10.1016/bs.pmbts.2018.01.004](https://doi.org/10.1016/bs.pmbts.2018.01.004)
- Anastasiadi, D., Esteve-Codina, A., & Piferrer, F. (2018). Consistent inverse correlation between DNA methylation of the first intron and gene expression across tissues and species. *Epigenetics & Chromatin*, 11(1). doi: [10.1186/s13072-018-0205-1](https://doi.org/10.1186/s13072-018-0205-1)
- Anders, S., Pyl, P. T., & Huber, W. (2015). HTSeq-A Python framework to work with high-throughput sequencing data. *Bioinformatics*, 31(2), 166–169. doi: [10.1093/bioinformatics/btu638](https://doi.org/10.1093/bioinformatics/btu638)
- Audetat, K. A., Galbraith, M. D., Odell, A. T., Lee, T., Pandey, A., Espinosa, J. M., ... Taatjes, D. J. (2017). A kinase-independent role for CDK19 in p53 response. *Molecular and Cellular Biology*. 37(13), e00626-16 Retrieved from <http://mcb.asm.org/content/early/2017/04/12/MCB.00626-16.abstract>
- Balakrishnan, C. N., Mukai, M., Gonser, R. A., Wingfield, J. C., London, S. E., Tuttle, E. M., & Clayton, D. F. (2014). Brain transcriptome sequencing and assembly of three songbird model systems for the study of social behavior. *PeerJ*, 2, e396–e396. doi: [10.7717/peerj.396](https://doi.org/10.7717/peerj.396)
- Banerjee, S. B., Arterbery, A. S., Fergus, D. J., & Adkins-Regan, E. (2011). Deprivation of maternal care has long-lasting consequences for the hypothalamic-pituitary-adrenal axis of zebra finches. *Proceedings of the Royal Society B: Biological Sciences*, 279(1729), 759–766. doi: [10.1098/rspb.2011.1265](https://doi.org/10.1098/rspb.2011.1265)
- Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate - a practical and powerful approach to multiple testing. *J R Stat Soc B Method*, 57(1), 289-300.
- Berger, S. L. (2007). The complex language of chromatin regulation during transcription. *Nature*, 447(7143), 407–412. doi: [10.1038/nature05915](https://doi.org/10.1038/nature05915)
- Bockmühl, Y., Patchev, A. V., Madejska, A., Hoffmann, A., Sousa, J. C., Sousa, N., ... Spengler, D. (2015). Methylation at the CpG island shore region upregulates Nr3c1 promoter activity after early-life stress. *Epigenetics*, 10(3), 247–257. doi: [10.1080/15592294.2015.1017199](https://doi.org/10.1080/15592294.2015.1017199)
- Bolton, J. L., Molet, J., Ivy, A., & Baram, T. Z. (2017). New insights into early-life stress and behavioral outcomes. *Current Opinion in Behavioral Sciences*, 14, 133–139. doi: [10.1016/j.cobeha.2016.12.012](https://doi.org/10.1016/j.cobeha.2016.12.012)

- Boyd, R. J., Kelly, T. R., MacDougall-Shackleton, S. A., & MacDougall-Shackleton, E. A. (2018). Alternative reproductive strategies in white-throated sparrows are associated with differences in parasite load following experimental infection. *Biology letters*, 14(7), 20180194.
- Catena, V., Bruno, T., De Nicola, F., Goeman, F., Pallocca, M., Iezzi, S., ... Fanciulli, M. (2016). Deptor transcriptionally regulates endoplasmic reticulum homeostasis in multiple myeloma cells. *Oncotarget*, 7(43), 70546–70558. doi: [10.18632/oncotarget.12060](https://doi.org/10.18632/oncotarget.12060)
- Cingolani, P., Patel, V., Coon, M., Nguyen, T., Land, S., Ruden, D., & Lu, X. (2012). Using *Drosophila melanogaster* as a Model for Genotoxic Chemical Mutational Studies with a New Program, SnpSift. *Frontiers in Genetics*, 3, 35–35.
- Cottrell, E. C., & Secki, J. R. (2009). Prenatal stress, glucocorticoids and the programming of adult disease. *Frontiers in Behavioral Neuroscience*, 3, 19. doi: [10.3389/neuro.08.019.2009](https://doi.org/10.3389/neuro.08.019.2009)
- Crean, A. J., & Bonduriansky, R. (2014). What is a paternal effect? *Trends in Ecology & Evolution*, 29(10), 554–559. doi: [10.1016/j.tree.2014.07.009](https://doi.org/10.1016/j.tree.2014.07.009)
- Crespi, E. J., Williams, T. D., Jessop, T. S., & Delehanty, B. (2012). Life history and the ecology of stress: how do glucocorticoid hormones influence life-history variation in animals? *Functional Ecology*, 27(1), 93–106. doi: [10.1111/1365-2435.12009](https://doi.org/10.1111/1365-2435.12009)
- Desantis, A., Bruno, T., Catena, V., De Nicola, F., Goeman, F., Iezzi, S., ... Fanciulli, M. (2015). Chemically induced inhibition of mTOR pathway enables stress-induced autophagy. *The EMBO Journal*, 34(9), 1214–1230.
- Dhanasiri, A. K. S., Fernandes, J. M. O., & Kiron, V. (2013). Liver Transcriptome Changes in Zebrafish during Acclimation to Transport-Associated Stress. *PLOS ONE*, 8(6), e65028–e65028.
- Diaz-Real, J., Kim, S.-Y., & Velando, A. (2017). Plumage colour and the expression of stress-related genes in gull chicks. *Journal of Avian Biology*, 48(9), 1216–1225. doi: [10.1111/jav.01460](https://doi.org/10.1111/jav.01460)
- Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., ... Gingeras, T. R. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*, 29(1), 15–21.
- Eden, E., Lipson, D., Yogev, S., & Yakhini, Z. (2007). Discovering Motifs in Ranked Lists of DNA Sequences. *PLOS Computational Biology*, 3(3), e39–e39.
- Eden, E., Navon, R., Steinfeld, I., Lipson, D., & Yakhini, Z. (2009). GOrilla: a tool for discovery and visualization of enriched GO terms in ranked gene lists. *BMC Bioinformatics*, 10(1), 48–48. doi: [10.1186/1471-2105-10-48](https://doi.org/10.1186/1471-2105-10-48)
- Finsterwald, C., & Alberini, C. M. (2014). Stress and glucocorticoid receptor-dependent mechanisms in long-term memory: From adaptive responses to psychopathologies. *Neurobiology of Learning and Memory*, 112, 17–29. doi: [10.1016/j.nlm.2013.09.017](https://doi.org/10.1016/j.nlm.2013.09.017)
- Formica, V. A., Gonser, R. A., Ramsay, S., & Tuttle, E. M. (2004). Spatial dynamics of alternative reproductive strategies: the role of neighbors. *Ecology*, 85(4), 1125–1136.
- Gluckman, P. D., Hanson, M. A., Cooper, C., & Thornburg, K. L. (2008). Effect of In Utero and Early-Life Conditions on Adult Health and Disease. *New England Journal of Medicine*, 359(1), 61–73. doi: [10.1056/NEJMr0708473](https://doi.org/10.1056/NEJMr0708473)
- Gott, A. J. (2018). Effects of early-life adversity on the adult phenotype in European starlings. Doctoral dissertation, Newcastle University.
- Greggor, A. L., Spencer, K. A., Clayton, N. S., & Thornton, A. (2017). Wild jackdaws' reproductive success and their offspring's stress hormones are connected to provisioning rate and brood size, not to parental neophobia. *General and Comparative Endocrinology*, 243, 70–77. doi: [10.1016/j.ygcen.2016.11.006](https://doi.org/10.1016/j.ygcen.2016.11.006)
- Griebel, I. A., Fairhurst, G. D., Marchant, T. A., & Clark, R. G. (2019). Effects of parental and nest-site characteristics on nestling quality in the Tree Swallow (*Tachycineta bicolor*). *Canadian Journal of Zoology*, 97(1), 63–71. doi: [10.1139/cjz-2018-0109](https://doi.org/10.1139/cjz-2018-0109)
- Griffiths, R., Double, M. C., Orr, K., & Dawson, R. J. G. (1998). A DNA test to sex most birds. *Molecular Ecology*, 7(8), 1071–1075. doi: [10.1046/j.1365-294x.1998.00389.x](https://doi.org/10.1046/j.1365-294x.1998.00389.x)
- Grunst, A. S., Grunst, M. L., Formica, V. A., Korody, M. L., Betuel, A. M., Barcelo-Serra, M., ... Tuttle, E. M. (2018a). Morph-Specific Patterns of Reproductive Senescence: Connections to Discrete Reproductive Strategies. *The American Naturalist*, 191(6), 744–755. doi: [10.1086/697377](https://doi.org/10.1086/697377)
- Grunst, A. S., Grunst, M. L., Gonser, R. A., & Tuttle, E. M. (2019). Developmental stress and telomere dynamics in a genetically polymorphic species. *Journal of Evolutionary Biology*, 32(2), 134–143. doi: [10.1111/jeb.13400](https://doi.org/10.1111/jeb.13400)

- Grunst, M. L., Grunst, A. S., Formica, V. A., Korody, M. L., Betuel, A. M., Barcelo-Serra, M., ... Tuttle, E. M. (2018b). Actuarial senescence in a dimorphic bird: different rates of ageing in morphs with discrete reproductive strategies. *Proceedings of the Royal Society B: Biological Sciences*, 285(1892), 20182053. doi: [10.1098/rspb.2018.2053](https://doi.org/10.1098/rspb.2018.2053)
- Heim, C., Newport, D. J., Mletzko, T., Miller, A. H., & Nemeroff, C. B. (2008). The link between childhood trauma and depression: Insights from HPA axis studies in humans. *Psychoneuroendocrinology*, 33(6), 693–710. doi: [10.1016/j.psyneuen.2008.03.008](https://doi.org/10.1016/j.psyneuen.2008.03.008)
- Horton, B. M., & Holberton, R. L. (2010). Morph-Specific Variation in Baseline Corticosterone and the Adrenocortical Response in Breeding White-Throated Sparrows (*Zonotrichia albicollis*). *The Auk*, 127(3), 540–548. doi: [10.1525/auk.2010.09096](https://doi.org/10.1525/auk.2010.09096)
- Horton, B. M., Moore, I. T., & Maney, D. L. (2014). New insights into the hormonal and behavioural correlates of polymorphism in white-throated sparrows, *Zonotrichia albicollis*. *Animal Behaviour*, 93, 207–219. doi: [10.1016/j.anbehav.2014.04.015](https://doi.org/10.1016/j.anbehav.2014.04.015)
- Horvath, S. (2011). *Weighted network analysis: application in genomics and systems biology*. New York, NY: Springer.
- Hu, Z., Chang, Y.-C., Wang, Y., Huang, C.-L., Liu, Y., Tian, F., ... DeLisi, C. (2013). VisANT 4.0: Integrative network platform to connect genes, drugs, diseases and therapies. *Nucleic Acids Research*, 41(W1), W225–W231.
- Jacquin, L., Blottière, L., Haussy, C., Perret, S., & Gasparini, J. (2012). Prenatal and postnatal parental effects on immunity and growth in ‘lactating’ pigeons: Prenatal and postnatal parental effects. *Functional Ecology*, 26(4), 866–875. doi: [10.1111/j.1365-2435.2012.01988.x](https://doi.org/10.1111/j.1365-2435.2012.01988.x)
- Kernbach, M.E., Newhouse, D.J., Miller, J.M., Hall, R.J., Gibbons, J., Oberstaller, J., Selechnik, D., Jiang, R., Unnasch, T.R., Balakrishnan, C.N., & L.B. Martin. Light pollution increases West Nile virus competence of a ubiquitous passerine reservoir species. *In review*.
- Kim, S.-H., Kim, H.-J., & Kim, C.-W. (2016). GLCCI1 is a novel component associated with the PI3K signaling pathway in podocyte foot processes. *Experimental & Molecular Medicine*, 48(5), e233–e233. doi: [10.1038/emm.2016.28](https://doi.org/10.1038/emm.2016.28)
- Kinnally, E. L., Feinberg, C., Kim, D., Ferguson, K., Leibel, R., Coplan, J. D., & John Mann, J. (2011). DNA methylation as a risk factor in the effects of early life stress. *Brain, Behavior, and Immunity*, 25(8), 1548–1553. doi: [10.1016/j.bbi.2011.05.001](https://doi.org/10.1016/j.bbi.2011.05.001)
- Knapton, R. W., Cartar, R. V., & Falls, J. B. (1984). A comparison of breeding ecology and reproductive success between morphs of the white-throated sparrow. *The Wilson Bulletin*, 60-71.
- Knapton, R. W., & Falls, J. B. (1983). Differences in parental contribution among pair types in the polymorphic White-throated Sparrow. *Canadian Journal of Zoology*, 61(6), 1288–1292. doi: [10.1139/z83-173](https://doi.org/10.1139/z83-173)
- Kopachena, J. G., & Falls, J. B. (1993). Aggressive Performance as a Behavioral Correlate of Plumage Polymorphism in the White-Throated Sparrow (*Zonotrichia albicollis*). *Behaviour*, 124(3/4), 249–266.
- Krause, E. T., Honarmand, M., Wetzel, J., & Naguib, M. (2009). Early Fasting Is Long Lasting: Differences in Early Nutritional Conditions Reappear under Stressful Conditions in Adult Female Zebra Finches. *PLoS ONE*, 4(3), e5015. doi: [10.1371/journal.pone.0005015](https://doi.org/10.1371/journal.pone.0005015)
- Langfelder, P., & Horvath, S. (2008). WGCNA: an R package for weighted correlation network analysis. *BMC Bioinformatics*, 9(1), 559–559. doi: [10.1186/1471-2105-9-559](https://doi.org/10.1186/1471-2105-9-559)
- Larsen, D. H., Poinsignon, C., Gudjonsson, T., Dinant, C., Payne, M. R., Hari, F. J., ... Lukas, J. (2010). The chromatin-remodeling factor CHD4 coordinates signaling and repair after DNA damage. *The Journal of Cell Biology*, 190(5), 731–740.
- Ledón-Rettig, C. C., Richards, C. L., & Martin, L. B. (2013). Epigenetics for behavioral ecologists. *Behavioral Ecology*, 24(2), 311–324. doi: [10.1093/beheco/ars145](https://doi.org/10.1093/beheco/ars145)
- Lewis, C. R., & Olive, M. F. (2014). Early-life stress interactions with the epigenome: potential mechanisms driving vulnerability toward psychiatric illness. *Behavioural Pharmacology*, 1. doi: [10.1097/FBP.0000000000000057](https://doi.org/10.1097/FBP.0000000000000057)
- Li, H. (2011). A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. *Bioinformatics*, 27(21), 2987–2993.
- Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *ArXiv:1303.3997 [q-Bio]*. Retrieved from <http://arxiv.org/abs/1303.3997>

- Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., ... Subgroup, 1000 Genome Project Data Processing. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics*, 25(16), 2078–2079.
- Lima, S. L. (2009). Predators and the breeding bird: behavioral and reproductive flexibility under the risk of predation. *Biological Reviews*, 84(3), 485–513. doi: [10.1111/j.1469-185X.2009.00085.x](https://doi.org/10.1111/j.1469-185X.2009.00085.x)
- Liu, D., Diorio, J., Tannenbaum, B., Caldji, C., Francis, D., Freedman, A., ... Meaney, M. J. (1997). Maternal Care, Hippocampal Glucocorticoid Receptors, and Hypothalamic-Pituitary-Adrenal Responses to Stress. *Science*, 277(5332), 1659–1662.
- Livneh, I., Cohen-Kaplan, V., Cohen-Rosenzweig, C., Avni, N., & Ciechanover, A. (2016). The life cycle of the 26S proteasome: from birth, through regulation and function, and onto its death. *Cell Research*, 26, 869–869.
- Lloyd, J. D., & Martin, T. E. (2004). Nest-site preference and maternal effects on offspring growth. *Behavioral Ecology*, 15(5), 816–823. doi: [10.1093/beheco/arh085](https://doi.org/10.1093/beheco/arh085)
- Longmire, J. L., Gee, G. F., Hardekopf, C. L., & Mark, G. A. (1992). Establishing paternity in whooping cranes (*Grus americana*) by DNA analysis. *The Auk*, 109(3), 522–529.
- Louder, M. I. M., Hauber, M. E., & Balakrishnan, C. N. (2018). Early social experience alters transcriptomic responses to species-specific song stimuli in female songbirds. *Behavioural Brain Research*, 347, 69–76. doi: [10.1016/j.bbr.2018.02.034](https://doi.org/10.1016/j.bbr.2018.02.034)
- Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, 15(12), 550–550. doi: [10.1186/s13059-014-0550-8](https://doi.org/10.1186/s13059-014-0550-8)
- Lowdon, R. F., Jang, H. S., & Wang, T. (2016). Evolution of Epigenetic Regulation in Vertebrate Genomes. *Trends in Genetics*, 32(5), 269–283. doi: [10.1016/j.tig.2016.03.001](https://doi.org/10.1016/j.tig.2016.03.001)
- Lowther, J. K. (1961). Polymorphism in the white-throated sparrow, *Zonotrichia albicollis* (Gmelin). *Canadian Journal of Zoology*, 39(3), 281–292. doi: [10.1139/z61-031](https://doi.org/10.1139/z61-031)
- Lowther, J. K. (1962). *Colour and Behavioural Polymorphism in the White-throated Sparrow, Zonotrichia Albicollis (Gmelin)*. Doctoral dissertation, University of Toronto.
- Lupien, S. J., McEwen, B. S., Gunnar, M. R., & Heim, C. (2009). Effects of stress throughout the lifespan on the brain, behaviour and cognition. *Nature Reviews Neuroscience*, 10(6), 434–445. doi: [10.1038/nrn2639](https://doi.org/10.1038/nrn2639)
- Marco, E. M., Echeverry-Alzate, V., López-Moreno, J. A., Giné, E., Peñasco, S., & Viveros, M. P. (2014). Consequences of early life stress on the expression of endocannabinoid-related genes in the rat brain. *Behavioural Pharmacology*, 25(5–6), 547–556. doi: [10.1097/FBP.0000000000000068](https://doi.org/10.1097/FBP.0000000000000068)
- Marini, F., & Binder, H. (2016). Development of Applications for Interactive and Reproducible Research: a Case Study. *Genomics and Computational Biology*, 3(1), 39. doi: [10.18547/gcb.2017.vol3.iss1.e39](https://doi.org/10.18547/gcb.2017.vol3.iss1.e39)
- Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.Journal*, 17(1), 10–10. doi: [10.14806/ej.17.1.200](https://doi.org/10.14806/ej.17.1.200)
- McCoy, C. R., Rana, S., Stringfellow, S. A., Day, J. J., Wyss, J. M., Clinton, S. M., & Kerman, I. A. (2016). Neonatal maternal separation stress elicits lasting DNA methylation changes in the hippocampus of stress-reactive Wistar Kyoto rats. *European Journal of Neuroscience*, 44(10), 2829–2845. doi: [10.1111/ejn.13404](https://doi.org/10.1111/ejn.13404)
- McEwen, B. S. (2007). Physiology and Neurobiology of Stress and Adaptation: Central Role of the Brain. *Physiological Reviews*, 87(3), 873–904. doi: [10.1152/physrev.00041.2006](https://doi.org/10.1152/physrev.00041.2006)
- McGowan, P. O., Sasaki, A., D'Alessio, A. C., Dymov, S., Labonté, B., Szyf, M., ... Meaney, M. J. (2009). Epigenetic regulation of the glucocorticoid receptor in human brain associates with childhood abuse. *Nature Neuroscience*, 12, 342–342.
- Meaney, M. J. (2001). Maternal Care, Gene Expression, and the Transmission of Individual Differences in Stress Reactivity Across Generations. *Annual Review of Neuroscience*, 24(1), 1161–1192. doi: [10.1146/annurev.neuro.24.1.1161](https://doi.org/10.1146/annurev.neuro.24.1.1161)
- Michopoulos, V., Maney, D. L., Morehouse, C. B., & Thomas, J. W. (2007). A genotyping assay to determine plumage morph in the white-throated sparrow (*Zonotrichia albicollis*). *The Auk*, 124(4), 1330. doi: [10.1642/0004-8038\(2007\)124\[1330:AGATDP\]2.0.CO;2](https://doi.org/10.1642/0004-8038(2007)124[1330:AGATDP]2.0.CO;2)
- Millstein, R. A., & Holmes, A. (2007). Effects of repeated maternal separation on anxiety- and depression-related phenotypes in different mouse strains. *Neuroscience & Biobehavioral Reviews*, 31(1), 3–17. doi: [10.1016/j.neubiorev.2006.05.003](https://doi.org/10.1016/j.neubiorev.2006.05.003)
- Mock, D. W., & Parker, G. A. (1997). *The evolution of sibling rivalry*. Oxford; New York: Oxford University Press.

- Moghadam, H. K., Johnsen, H., Robinson, N., Andersen, Ø., H. Jørgensen, E., Johnsen, H. K., ... Tveiten, H. (2017). Impacts of Early Life Stress on the Methylome and Transcriptome of Atlantic Salmon. *Scientific Reports*, 7(1). doi: [10.1038/s41598-017-05222-2](https://doi.org/10.1038/s41598-017-05222-2)
- Monaghan, P. (2014). Organismal stress, telomeres and life histories. *Journal of Experimental Biology*, 217(1), 57–66. doi: [10.1242/jeb.090043](https://doi.org/10.1242/jeb.090043)
- Mousseau, T. A., & Fox, C. W. (Eds.). (1998). *Maternal effects as adaptations*. New York: Oxford University Press.
- Murgatroyd, C., Patchev, A. V., Wu, Y., Micale, V., Bockmühl, Y., Fischer, D., ... Spengler, D. (2009). Dynamic DNA methylation programs persistent adverse effects of early-life stress. *Nature Neuroscience*, 12(12), 1559–1566. doi: [10.1038/nn.2436](https://doi.org/10.1038/nn.2436)
- Noguera, J. C., Kim, S.-Y., & Velando, A. (2017). Family-transmitted stress in a wild bird. *Proceedings of the National Academy of Sciences*, 114(26), 6794–6799. Retrieved from <http://www.pnas.org/content/early/2017/06/06/1706164114.abstract>
- Nord, A., & Nilsson, J.-Å. (2011). Incubation Temperature Affects Growth and Energy Metabolism in Blue Tit Nestlings. *The American Naturalist*, 178(5), 639–651. doi: [10.1086/662172](https://doi.org/10.1086/662172)
- Pétille, F., Brantsæter, M., Nordgreen, J., Coutinho, L. L., Janczak, A. M., Jensen, P., & Guerrero-Bosagna, C. (2017). DNA methylation profiles in red blood cells of adult hens correlate with their rearing conditions. *The Journal of Experimental Biology*, 220(19), 3579–3587. doi: [10.1242/jeb.157891](https://doi.org/10.1242/jeb.157891)
- Peterson, T. R., Laplante, M., Thoreen, C. C., Sancak, Y., Kang, S. A., Kuehl, W. M., ... Sabatini, D. M. (2009). DEPTOR Is an mTOR Inhibitor Frequently Overexpressed in Multiple Myeloma Cells and Required for Their Survival. *Cell*, 137(5), 873–886. doi: [10.1016/j.cell.2009.03.046](https://doi.org/10.1016/j.cell.2009.03.046)
- Quinlan, A. R., & Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics*, 26(6), 841–842.
- Quirici, V., Guerrero, C. J., Krause, J. S., Wingfield, J. C., & Vásquez, R. A. (2016). The relationship of telomere length to baseline corticosterone levels in nestlings of an altricial passerine bird in natural populations. *Frontiers in Zoology*, 13(1). doi: [10.1186/s12983-016-0133-5](https://doi.org/10.1186/s12983-016-0133-5)
- R Core Team (2013). R: A language and environment for statistical computing.
- Reshetnikov, V. V., Studenikina, A. A., Ryabushkina, J. A., Merkulova, T. I., & Bondar, N. P. (2018). The impact of early-life stress on the expression of HPA-associated genes in the adult murine brain. *Behaviour*, 155(2–3), 181–203. doi: [10.1163/1568539X-00003482](https://doi.org/10.1163/1568539X-00003482)
- Romanov, M. N., Tuttle, E. M., Houck, M. L., Modi, W. S., Chemnick, L. G., Korody, M. L., ... Ryder, O. A. (2009). The value of avian genomics to the conservation of wildlife. *BMC Genomics*, 10(Suppl 2), S10. doi: [10.1186/1471-2164-10-S2-S10](https://doi.org/10.1186/1471-2164-10-S2-S10)
- Royce, N. J., Smiseth, P. T., & Kölliker, M. (Eds.). (2012). *The evolution of parental care*. Oxford University Press.
- Rubenstein, D. R., Skolnik, H., Berrio, A., Champagne, F. A., Phelps, S., & Solomon, J. (2016). Sex-specific fitness effects of unpredictable early life conditions are associated with DNA methylation in the avian glucocorticoid receptor. *Molecular Ecology*, 25(8), 1714–1728. doi: [10.1111/mec.13483](https://doi.org/10.1111/mec.13483)
- Schrader, M., Jarrett, B. J. M., & Kilner, R. M. (2018). Parental care and sibling competition independently increase phenotypic variation among burying beetle siblings. *Evolution*, 72(11), 2546–2552. doi: [10.1111/evo.13607](https://doi.org/10.1111/evo.13607)
- Schwander, T., Libbrecht, R., & Keller, L. (2014). Supergenes and Complex Phenotypes. *Current Biology*, 24(7), R288–R294. doi: [10.1016/j.cub.2014.01.056](https://doi.org/10.1016/j.cub.2014.01.056)
- Shang, F., & Taylor, A. (2011). Ubiquitin–proteasome pathway and cellular responses to oxidative stress. *Free Radical Biology and Medicine*, 51(1), 5–16. doi: [10.1016/j.freeradbiomed.2011.03.031](https://doi.org/10.1016/j.freeradbiomed.2011.03.031)
- Sheldon, E. L., Schrey, A. W., Ragsdale, A. K., & Griffith, S. C. (2018). Brood size influences patterns of DNA methylation in wild Zebra Finches (*Taeniopygia guttata*). *The Auk*, 135(4), 1113–1122. doi: [10.1642/AUK-18-61.1](https://doi.org/10.1642/AUK-18-61.1)
- Silberman, D. M., Acosta, G. B., & Zorrilla Zubilete, M. A. (2016). Long-term effects of early life stress exposure: Role of epigenetic mechanisms. *Pharmacological Research*, 109, 64–73. doi: [10.1016/j.phrs.2015.12.033](https://doi.org/10.1016/j.phrs.2015.12.033)
- Smith, J.N., Newhouse, D.J., Balakrishnan, C.N., Tuttle, E.M., & Gonser, R.A. Differential morph gene expression in chromosome three of the white-throated sparrow (*Zonotrichia albicollis*). *In review*.
- Spencer, K. A. (2017). Developmental stress and social phenotypes: integrating neuroendocrine, behavioural and evolutionary perspectives. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 372(1727). Retrieved from <http://rstb.royalsocietypublishing.org/content/372/1727/20160242.abstract>

- Spencer, K. A., Evans, N. P., & Monaghan, P. (2009). Postnatal Stress in Birds: A Novel Model of Glucocorticoid Programming of the Hypothalamic-Pituitary-Adrenal Axis. *Endocrinology*, 150(4), 1931–1934. doi: [10.1210/en.2008-1471](https://doi.org/10.1210/en.2008-1471)
- Spinney, L. H., Bentley, G. E., & Hau, M. (2006). Endocrine correlates of alternative phenotypes in the white-throated sparrow (*Zonotrichia albicollis*). *Hormones and Behavior*, 50(5), 762–771. doi: [10.1016/j.yhbeh.2006.06.034](https://doi.org/10.1016/j.yhbeh.2006.06.034)
- Swett, M. B., & Breuner, C. W. (2009). Plasma Testosterone Correlates with Morph Type across Breeding Substages in Male White-Throated Sparrows. *Physiological and Biochemical Zoology*, 82(5), 572–579. doi: [10.1086/605392](https://doi.org/10.1086/605392)
- Szyf, M. (2009). The early life environment and the epigenome. *Biochimica et Biophysica Acta (BBA) - General Subjects*, 1790(9), 878–885. doi: [10.1016/j.bbagen.2009.01.009](https://doi.org/10.1016/j.bbagen.2009.01.009)
- Szyf, M., Weaver, I., & Meaney, M. (2007). Maternal care, the epigenome and phenotypic differences in behavior. *Reproductive Toxicology*, 24(1), 9–19. doi: [10.1016/j.reprotox.2007.05.001](https://doi.org/10.1016/j.reprotox.2007.05.001)
- Taylor, S., & Campagna, L. (2016). Avian supergenes. *Science*, 351(6272), 446–447. doi: [10.1126/science.aae0389](https://doi.org/10.1126/science.aae0389)
- Thomas, J. W., Cáceres, M., Lowman, J. J., Morehouse, C. B., Short, M. E., Baldwin, E. L., ... Martin, C. L. (2008). The Chromosomal Polymorphism Linked to Variation in Social Behavior in the White-Throated Sparrow (*Zonotrichia albicollis*) Is a Complex Rearrangement and Suppressor of Recombination. *Genetics*, 179(3), 1455 LP – 1468.
- Thorncroft, H. B. (1966). Chromosomal Polymorphism in the White-Throated Sparrow, *Zonotrichia albicollis* (Gmelin). *Science*, 154(3756), 1571–1572.
- Thorncroft, H. B. (1975). A cytogenetic study of the white-throated sparrow, *Zonotrichia albicollis* (Gmelin). *Evolution*, 29(4), 611–621. doi: [10.1111/j.1558-5646.1975.tb00855.x](https://doi.org/10.1111/j.1558-5646.1975.tb00855.x)
- Tomalty, K. M. H., Meek, M. H., Stephens, M. R., Rincón, G., Fangue, N. A., May, B. P., & Baerwald, M. R. (2015). Transcriptional Response to Acute Thermal Exposure in Juvenile Chinook Salmon Determined by RNAseq. *G3: Genes/Genomes/Genetics*, 5(7), 1335 LP – 1349.
- Trivers, R.L. (1972). Parental investment and sexual selection. In: *Sexual Selection and the Descent of Man* (B. Campbell, ed), pp. 136–179. Aldine, Chicago, IL.
- Turner, J. D., Schote, A. B., Macedo, J. A., Pelascini, L. P. L., & Muller, C. P. (2006). Tissue specific glucocorticoid receptor expression, a role for alternative first exon usage? *Biochemical Pharmacology*, 72(11), 1529–1537. doi: [10.1016/j.bcp.2006.07.005](https://doi.org/10.1016/j.bcp.2006.07.005)
- Tuttle, E. (2003). Alternative reproductive strategies in the white-throated sparrow: behavioral and genetic evidence. *Behavioral Ecology*, 14(3), 425–432.
- Tuttle, E. M., Grunst, A. S., Grunst, M. L., Korody, M. L., Betuel, A. M., Barcelo-Serra, M., ... Gonser, R. A. (2017). Climatically driven changes in population composition and offspring sex-morph ratio in a polymorphic species. *Ecosphere*, 8(4), e01762–e01762. doi: [10.1002/ecs2.1762](https://doi.org/10.1002/ecs2.1762)
- Tuttle, E. M., Bergland, A. O., Korody, M. L., Brewer, M. S., Newhouse, D. J., Minx, P., ... Balakrishnan, C. N. (2016). Divergence and Functional Degradation of a Sex Chromosome-like Supergene. *Current Biology*, 26(3), 344–350. doi: [10.1016/j.cub.2015.11.069](https://doi.org/10.1016/j.cub.2015.11.069)
- van Oers, K., Kohn, G. M., Hinde, C. A., & Naguib, M. (2015). Parental food provisioning is related to nestling stress response in wild great tit nestlings: implications for the development of personality. *Frontiers in Zoology*, 12(Suppl 1), S10. doi: [10.1186/1742-9994-12-S1-S10](https://doi.org/10.1186/1742-9994-12-S1-S10)
- Wade, M. J. (1998). The evolutionary genetics of maternal effects Pp. 5–21 in Mousseau T. A & Fox C. W., eds. Maternal effects as adaptations. Oxford Univ. Press, New York.
- Wang, Y., Branicky, R., Noë, A., & Hekimi, S. (2018). Superoxide dismutases: Dual roles in controlling ROS damage and regulating ROS signaling. *The Journal of Cell Biology*, 217(6), 1915–1928. doi: [10.1083/jcb.201708007](https://doi.org/10.1083/jcb.201708007)
- Weaver, I. C. G., Cervoni, N., Champagne, F. A., D'Alessio, A. C., Sharma, S., Seckl, J. R., ... Meaney, M. J. (2004). Epigenetic programming by maternal behavior. *Nature Neuroscience*, 7, 847–847.
- Weaver, I. C. G., Meaney, M. J., & Szyf, M. (2006). Maternal care effects on the hippocampal transcriptome and anxiety-mediated behaviors in the offspring that are reversible in adulthood. *Proceedings of the National Academy of Sciences of the United States of America*, 103(9), 3480 LP – 3485.
- Wolf, J. B., & Wade, M. J. (2009). What are maternal effects (and what are they not)? *Philosophical Transactions of the Royal Society B: Biological Sciences*, 364(1520), 1107–1115. doi: [10.1098/rstb.2008.0238](https://doi.org/10.1098/rstb.2008.0238)

- Zelko, I. N., Mariani, T. J., & Folz, R. J. (2002). Superoxide dismutase multigene family: a comparison of the CuZn-SOD (SOD1), Mn-SOD (SOD2), and EC-SOD (SOD3) gene structures, evolution, and expression. *Free Radical Biology and Medicine*, 33(3), 337–349. doi: [10.1016/S0891-5849\(02\)00905-X](https://doi.org/10.1016/S0891-5849(02)00905-X)
- Zhang, B., & Horvath, S. (2005). A General Framework for Weighted Gene Co-Expression Network Analysis. *Statistical Applications in Genetics and Molecular Biology*, 4. doi: [10.2202/1544-6115.1128](https://doi.org/10.2202/1544-6115.1128)
- Zhao, H., Zhao, Y. G., Wang, X., Xu, L., Miao, L., Feng, D., ... Zhang, H. (2013). Mice deficient in *Epg5* exhibit selective neuronal vulnerability to degeneration. *The Journal of Cell Biology*, 200(6), 731 LP – 741.
- Zinzow-Kramer, W. M., Horton, B. M., McKee, C. D., Michaud, J. M., Tharp, G. K., Thomas, J. W., ... Maney, D. L. (2015). Genes located in a chromosomal inversion are correlated with territorial song in white-throated sparrows. *Genes, Brain and Behavior*, 14(8), 641–654. doi: [10.1111/gbb.12252](https://doi.org/10.1111/gbb.12252)

Data Accessibility

The 32 RNAseq libraries used in this study will be submitted to the NCBI Sequence Read Archive (SRA). All files needed to produce these results, including code and counts files, will be uploaded to this project's GitHub page: <https://github.com/danielnewhouse/wtsp>

Author Contributions

DJN designed and performed research, analyzed the data, and wrote the paper. MBS performed research, contributed samples, and reviewed drafts of the paper. EMT designed and performed research and contributed samples. RAG designed and performed research, contributed samples, and reviewed drafts of the paper. CNB designed and performed research, contributed reagents, and reviewed drafts of the paper.

816 Tables and Figures

817 Table 1. WGCNA modules correlated with morph, strength of correlation (R^2), p-value, hub
818 gene(s) of module, and the degree distribution of hub gene(s).

Module	R^2	p-value	Hub genes	DD of hub gene(s)
Dark Red	0.47	0.009	TRAF5	32
Ivory	-0.66	9×10^{-5}	GOPC, HDAC2, HINT3, TAF5L, TRMT61B, MARC2	>29
Light Cyan	0.67	5×10^{-5}	BPNT1, EPM2A, LOC102066536 (GST-like), MAN1A1, MEI4, RNASET2, SLC18B1, TTC32	>27
Salmon	-0.5	0.005	NSL1	39
Sky Blue	0.53	0.003	DTX3L, EIF2AK2, IFIT5, LOC102064521 (OASL), LOC102065196 (IFI27L2), PARP9, PARP14, RSAD2, ZNFX1	>22

819

820 Table 2. WGCNA modules correlated with pair type, strength of correlation (R^2), p-value, hub
821 gene(s) of module, and the degree distribution of hub gene(s).

Module	R^2	p-value	Hub genes	DD of hub gene(s)
Beige	-0.61	3×10^{-4}	DEPTOR	39
Blue	-0.45	0.01	NDUFB3	42

Cyan	0.46	0.01	HELZ	36
Dark Orange	0.7	1×10^{-5}	NCOA6	45
Light Green	0.6	4×10^{-4}	CDK19, CHD4, EPG5	>28
Orange	-0.45	0.01	ZFX	31
Pink	0.37	0.04	LOC102060916 (C12orf4)	19

822

823

Figure 1. WGCNA module-trait correlation matrix. Each box contains the R^2 correlation value followed by p-value in parentheses of a given trait with the module. Correlation values range from -1 to 1, with orange colors representing positive correlation and blue colors representing negative correlation.

Figure 2. (A) Network of blue module, highlighting hub gene NDUFB3, along with normalized expression plots of (B) NDUFB3, as well as ubiquitin-mediated proteolysis-related genes (C) PSMD6 and (D) UBE2D3. TxW represents samples from nests sired by a T male and a W female. WxT represents samples from nests sired by a W male and a T female. Each circle represents a gene and diamonds represent hub genes described in Table 2.

Figure 3. (A) Network of beige module, highlighting hub gene DEPTOR, along with normalized expression plots of hub gene (B) DEPTOR, as well as stress responsive genes (C) SOD2 and (D) NR3C1. TxW represents samples from nests sired by a T male and a W female. WxT represents samples from nests sired by a W male and a T female.

Figure 4. (A) Network of light green module and normalized expression plots of hub genes (B) CDK19, (C) CHD4, and (D) EPG5. TxW represents samples from nests sired by a T male and a W female. WxT represent samples from nests sired by a W male and a T female. Each circle represents a gene and diamonds represent hub genes described in Table 2.

Module-trait relationships

A **B** **C** **D**

A **B** **C** **D**

A **B****CDK19** **C****CHD4** **D****EPG5**