Predator-induced selection on urchin activity level depends on urchin body size

Justin Pretorius¹, James L.L. Lichtenstein^{1*}, Erika J. Eliason¹, Adrian C. Stier¹, and Jonathan N. Pruitt^{1,2}

¹ Department of Ecology, Evolution, and Marine Biology, University of California Santa Barbara, Santa Barbara, California 93106 USA

² Department of Psychology, Neuroscience, & Behaviour, McMaster University, Hamilton, Ontario, L8S 4K1 Canada

*Denotes corresponding author. Email: jlllichtenstein@gmail.com; Phone: 1 503 553 9353

11 ABSTRACT

12 Temporally consistent individual differences in behavior impact many ecological processes. We
 13 simultaneously examined the effects of individual variation in prey activity level, covering behavior, and
 14 body size on prey survival with predators using an urchin-lobster system. Specifically, we tested the
 15 hypothesis that slow-moving purple sea urchins (*Strongylocentrotus purpuratus*) and urchins who deploy
 16 extensive substrate (pebbles and stones) covering behavior will out-survive active urchins that deploy
 17 little to no covering behavior when pitted against a predator, the California spiny lobster (*Panulirus*
 18 *interruptus*). We evaluated this hypothesis by first confirming whether individual urchins exhibit
 19 temporally consistent differences in activity level and covering behavior, which they did. Next, we placed
 20 groups of four urchins in mesocosms with single lobster and monitored urchin survival for 108 hours.
 21 High activity level was negatively associated with survival, whereas urchin size and covering behavior
 22 independently did not influence survival. The negative effect of urchin activity level on urchin survival
 23 was strong for smaller urchins and weaker for large urchins. Taken together, these results suggest that
 24 purple urchin activity level and size jointly determine their susceptibility to predation by lobsters. This is
 25 potentially of great interest, because predation by recovering lobster populations could alter the stability
 26 of kelp forests by culling specific phenotypes, like foraging phenotypes, from urchin populations.

27
 28 **Key words:** activity, covering behavior, behavioral type, predator-prey interactions, predation risk,
 29 temperament

30
 31

32
 33
 34
 35

36 1 | INRODUCTION

37

38 Intraspecific differences in morphology and behavior are predicted to shape the outcome of species
 39 interactions. For example, prior studies have shown that larger and faster prey species are more likely to
 40 escape predation (Einfalt & Wahl, 1997; Hoyle & Keast, 1987; Werner & Hall, 1974). Likewise,
 41 individual variation in predator jaw morphology can shape the effect of predators on community
 42 composition at both the producer and primary consumer levels (Post, Palkovacs, Schielke, & Dodson,
 43 2008). Parallel lines of evidence suggest that among individual variation in key traits of both predators
 44 and primary producers can affect ecosystem structure and function (Bolnick et al., 2011; Bolnick et al.,
 45 2003; Wolf & Weissing, 2012). For instance, genetic diversity within primary producer species is
 46 correlated with consumer diversity and community resilience, seemingly by increasing the diversity of
 47 interaction types (Crutsinger et al., 2006; Hughes & Stachowicz, 2004; Johnson, Lajeunesse, & Agrawal,
 48 2006; Reusch, Ehlers, Hämmerli, & Worm, 2005; Rudman et al., 2015). Understanding how intraspecific
 49 variation in morphology and behavior shape the outcome of predator-prey interactions is of specific
 50 interest, because these interactions can change consumer abundance and behavior, and therefore, alter the
 51 role of predation in driving the stability, function, and resilience of ecosystems (Brown, Laundré, &
 52 Gurung, 1999; Miller, Ament, & Schmitz, 2014; Preisser & Bolnick, 2008).

53 The past few decades have seen a rise in the number of studies focused on intraspecific behavioral
 54 variation (Bell, Hankison, & Laskowski, 2009; Sih, Cote, Evans, Fogarty, & Pruitt, 2012). In particular,
 55 much literature has examined the effects of temporally consistent individual differences in behavior, or so
 56 called “animal personalities” (Carter, Feeney, Marshall, Cowlshaw, & Heinsohn, 2013; Gosling, 2001)
 57 and their effect on species interactions (Chang, Teo, Norma-Rashid, & Li, 2017; Lichtenstein, Wright,
 58 McEwen, Pinter-Wollman, & Pruitt, 2017; Nannini, Parkos III, & Wahl, 2012). Animal personalities
 59 imply that there exists a level of constraint to the behavioral plasticity individuals display in any given
 60 context (Dall, Houston, & McNamara, 2004; Dingemanse, Kazem, Réale, & Wright, 2010; Dochtermann

61 & Dingemanse, 2013), meaning that individuals may exhibit behavioral consistency through time and
 62 across contexts, occasionally to their detriment (Jones & Godin, 2010; Pearish, Hostert, & Bell, 2013;
 63 Pruitt, Riechert, & Jones, 2008). For instance, some prey animals ceaselessly seek shelter regardless of the
 64 contextual risk, which can result in decreased opportunities to forage or mate, while other animals may
 65 continuously forage regardless of the level of predation risk (Dall et al., 2004; Wilson, Coleman, Clark, &
 66 Biederman, 1993). Yet, the available empirical evidence suggests that the effects of prey behavior on the
 67 outcome of predator-prey interactions varies considerably among systems (Belgrad & Griffen, 2016;
 68 Keiser, Ingley, Toscano, Scharf, & Pruitt, 2017), and even varies based on the traits of the individuals
 69 (predator and prey) involved (DiRienzo, Pruitt, & Hedrick, 2013; McGhee, Pintor, Suhr, & Bell, 2012;
 70 Sweeney et al., 2013).

71 Here, we use the purple urchin (*Strongylocentrotus purpuratus*), and the California spiny lobster
 72 (*Panulirus interruptus*), to evaluate the effect of prey behavioral tendencies and size on prey survival.
 73 Urchins are ecologically notable because of their ability to consume large quantities of giant kelp
 74 (*Macrocystis pyrifera*) and therefore alter the state of coastal kelp communities (Estes & Duggins, 1995;
 75 Harrold & Reed, 1985; Pearse, 2006). Across the wide geographic distribution of purple urchins along the
 76 west coast of North America (Ebert, 2010), urchins are eaten by a diversity of predatory species including
 77 the sunflower sea star (*Pycnopodia helianthoides*), otters (*Enhydra lutris*), sheephead (*Semicossyphus*
 78 *pulcher*), and California spiny lobsters (*Panulirus interruptus*). These predators can play key roles in
 79 regulating herbivory by urchins on giant kelp (Burt et al., 2018; Caselle, Davis, & Marks, 2018).
 80 However, the role of variation in urchin size structure and individual behavioral variation in their
 81 susceptibility to predation remains mysterious, even though such variation in mortality could explain
 82 variation in community susceptibility to urchin-driven state shifts, like between kelp forests vs. urchin
 83 barrens, a process often referred to as a “tipping point” (Pruitt et al., 2018; Selkoe et al., 2015). We
 84 focused on how two behavioral traits in might determine prey survival in the presence of spiny lobsters:
 85 propensity to move (activity level), which can increase or decrease encounter rates with predators (Huey

& Pianka, 1981; Skelly, 1994), and urchin's tendency to conceal themselves with substrate (covering behavior). Although the function of urchin covering behavior appears to vary among species (i.e., protection from wave action; Dumont, Drolet, Deschênes, & Himmelman, 2007), we predicted that urchins would be safer from predators while buried in substrate.

We tested the hypothesis that slow-moving purple urchins and urchins with a greater tendency to cover themselves and/or larger urchins will enjoy greater survival during staged encounters with sit-and-pursue predators. The former is predicted by Locomotor Crossover Hypothesis proposed by Huey and Pianka (1981), which posits that predators should tend to encounter and consume prey of the opposing activity level (active predators consume sedentary prey, whereas sit-and-pursue predators consume active prey). To probe the role of size, we tested whether urchin size increases urchin survival jointly with urchin behavioral traits. We further evaluated whether prey behavior measured in isolation is correlated with prey behavior during real encounters with predators. Consistency across testing conditions is a commonly untested assumption in animal personality studies, usually overlooked because of logistical limitations. However, the conspicuousness of the urchin covering behavior and the slow movements of predator and prey in this study made ongoing assessment of this behavior achievable. Together, this system allowed us to evaluate how the behavioral and morphological traits of a prey determine their susceptibility to predation.

2 | MATERIALS AND METHODS

2.1 | Animal collection and maintenance

Fifty *Strongylocentrotus purpuratus* were opportunistically collected by hand from the Goleta sewer pipe (34° 24.851 N; 119° 49.749 W) and 120 others were collected from Goleta Pier (34° 24.847 N; 119° 49.718 W, Goleta, CA, U.S.A.) in the summer of 2017. A total of 170 *S. purpuratus* were used for the

experiment. The three lobsters (*Panulirus interruptus*) used in the experiment were transferred into our possession from another research facility on the campus of the University of California at Santa Barbara (UCSB) on 16 August 2017. The three *P. interruptus* used in the study were female, 10.5-11 cm in carapace length, and were collected by hand from the Santa Barbara Channel from May to June 2017.

All *S. purpuratus* were provided brown algae (*Nereocystis luetkeana*) *ad libitum* every other day from 01 July 2017 to 01 November 2017. Purple urchins were housed in four separate 90.0L cm x 141.0W cm x 40.5H cm seawater (9-11° C) flow through A-frame tanks until their behavioral assays. During this time, we measured the total diameter (test and spines) of each urchin using digital calipers. They had average diameter of 7.7 ± 0.20 SE cm and a range of 4.5-11.5 cm. After purple urchins had been assayed for covering behavior, individuals were transferred into numerically labeled 17.15L cm x 11.90W cm x 7.00H cm plastic containers in the same seawater flow through tanks. Plastic containers housing individual purple urchins had fourteen holes of 16.0 mm diameter drilled into their sides to allow seawater flow through. *P. interruptus* were housed in similar seawater flow-through A-frame tanks bisected by plastic dividers with twelve 16mm holes, creating two identical lobster enclosures (90.0L cm x 70.5W cm x 40.1H cm). Each lobster enclosure contained three cinder blocks, two of which supported the plastic dividers. All animals were exposed to open air conditions and natural day-night cycles for the duration of the experiment. This experiment was conducted at the UCSB Marine Laboratory from June to November 2017.

2.2 | Repeatability overview

To estimate short-term behavioral repeatability, we ran ten urchins through 6 trials assessing movement velocity (activity) and covering percentage (covering behavior) over the span of six days. We ran the urchins through two covering behavior tests separated by an hour, and the next day, we ran the urchins through two activity level tests separated by an hour. We repeated this two-day cycle for six days.

Therefore, we assessed the repeatability of activity level and covering behavior with six iterations across three testing days for each assay type in ten focal individuals. These ten testing urchins were not used in predator-prey trials.

The purple urchins not selected for repeatability estimates underwent only two trials of each type, conducted under identical conditions ($n_{\text{individuals}} = 160$) separated by an hour. The average value of the two activity and covering behavior measurements for each urchin served as our activity and covering behavior predictor variables in our statistical analyses predicting prey survival.

2.3 | Measuring urchin activity level

To determine urchin activity level, we estimated their average velocity in open field arenas. The arena was a 40.60L cm x 32.40W cm x 15.25H cm plastic container fashioned with 6 holes of 16.0 mm diameter drilled into its sides and submerged in flow through seawater. The container was completely submerged in seawater within a chilled seawater flow through plastic A-frame tank. We placed a 28.0 cm x 30.5 cm particleboard with a 4 cm grid within each activity container to track urchin movement distance. The board was covered with transparent saran wrap to help the urchins cling to the board. We used movements between squares to calculate an average movement velocity (mm/s) for each individual, one urchin at a time, across a ten-minute trial. We deemed urchins with higher velocities to be more active. This test is very similar to an open field test, a classic metric of activity level (Carter et al., 2013). We performed two such activity trials on a single testing day separated by an hour on each of 160 urchins. We averaged the two scores to obtain a single estimate of each urchin's activity level in a novel environment. Also, two activity containers were run at a time.

2.4 | Measuring sea urchin covering behavior

To estimate urchins' shelter seeking tendencies, we measured their propensity to cover themselves with pebble substrate. Covering behavior resembles common metrics of "boldness" in the personality literature and has antipredator benefits in some species of urchins (Amsler, McClintock, & Baker, 1999). However, the behavior can be cued by wave action and sunlight in other urchin species (Dumont et al., 2007), although urchins deploy these behaviors in the absence of both (Pawson & Pawson, 2013). We examine here whether covering behavior might provide an antipredator benefit to purple urchins with predators during staged encounters.

Individual purple urchins ($n = 170$) were haphazardly placed into pairs and then tested under social conditions. Pairs of asymmetrically sized urchins (difference in test [spherical] diameter of 10.0 mm or greater) were created to track individual identity during these trials. Urchins were tested in pairs because of the time constraints of testing so many urchins. Pairs of purple urchins were placed into 40.60L cm x 32.40W cm x 15.25H cm plastic containers with transparent lids fashioned with 6 holes of 16.0 mm diameter drilled into their sides to allow chilled seawater to flow through. These were the same containers used for the activity level test, but for covering behavior trials, the bottoms of these containers were layered with 2.0-3.0 cm of fresh composite pebbles of 0.3-1.5 cm diameter, rather than a particle board grid. These containers were then completely submerged in chilled seawater flow within plastic A-frame tanks entirely separate from their home A-frame tank.

Once the pair of purple urchins were placed into the container and submerged, they were permitted one hour to move about their arenas and cover themselves. After this time, we opened the arena and estimated the percent cover (test + spines) of each urchin by eye to the nearest 5%. These estimates were taken by a single observer (Pretorius J). We performed two trials in a testing day separated by an hour. We then averaged each urchin's scores to obtain a single estimate of urchins' covering behavior.

2.5 Spiny lobster vs. sea urchin staged interactions

Each staged predator-prey interaction consisted of a cohort of 4 haphazardly selected urchins and a single predator lobster. Lobsters were maintained under identical feeding conditions (*ad libitum* mussels interspersed with purple urchins to ensure they did eat urchins) and then starved for four days prior to the initiation of our trials, and all purple urchins had already been assayed for activity and covering behavior. Because we had three lobsters, we ran three of these trials at once. Predators were permitted to interact with prey for 108 hours during our trials, because this duration resulted in an urchin mortality rate over 50%.

Staged interactions were conducted in natural seawater (9-11° C) flow-through tanks divided by plastic barriers perforated by twelve 16 cm holes. The bottom of each arena (90.0L cm x 40.5W cm x 70.5H cm) was layered with 2.5-4.0 cm of pebbles of diameter 0.3-1.5 cm and contained three submerged cinderblocks to permit prey to hide from predators in physical retreats or, most commonly, within the substrate. The sides of the tank likewise served as physical refuge for prey, allowing them to climb up the sides of the container and out of reach of the lobsters.

To distinguish individual purple urchins throughout staged interactions, one of four colors (yellow, green, pink, or blue) of acrylic nail polish was lightly dotted on the side of a few of each individual's peripheral spines. Once acrylic nail polish had been applied to every purple urchin within a group of four, all four purple urchins were simultaneously placed by hand into four corners (quadrants) of each arena containing a lobster. Care was taken to place urchins outside of the lobster's field of view. For each occasion, staged interactions began at 14:30 hours. We recorded each urchin's covering percentage and the quadrant they occupied every hour for the first 24 hours and every six hours for the next 84 hours. We estimated activity during trials by using movement between quadrants. After 108-hours, a thorough search of every mesocosm was conducted. We recorded which urchins had survived by their color markings. We repeated these trials four times separated by a week using entirely new urchins, so each lobster was used in four trials. The flow-through A-frame tanks for predation trials were housed on a separate concrete patio surrounded by chain link caging to prevent mammalian predators (raccoons, etc.) from eating

captive marine fauna, which is required for the vertebrates housed in other nearby enclosures. Thus, urchins that went missing during our trials had few escape options and no other likely predators.

2.6 | Statistical Methods

2.6.1 | Repeatability of behavior

The repeatability of each behavioral trait was determined using the rptR package (Stoffel, Nakagawa, & Schielzeth, 2017) in R version 3.4.3 (R Development Core Team, 2010). We used this package to fit mixed models with ‘individual ID’ and “testing day” (1-3) as random effects, “trial number” (1/2) as a random effect nested within testing day, and each behavioral test was set as the response variable. These models estimate repeatability as the proportion of variance determined by ‘individual ID’ to the total behavioral variance. The rptR package constructs 95% CI of the repeatability estimates using 1000 bootstrap iterations of these repeatability estimations. The activity level repeatability models were linear mixed model, because a Gaussian distribution best fit the data. The covering behavior repeatability models were generalized linear mixed effects models fit with a Poisson distribution, because a Poisson distribution best fit the data. Repeatability estimates are deemed significant if their 95% confidence interval do not overlap 0.

2.6.2 | Spiny lobster-sea urchin staged interactions

To investigate the factors predicting urchin survival, we deployed a generalized linear mixed model (GLMM) fit with a binomial distribution in the lme4 package in R (Bates, Mächler, Bolker, & Walker, 2014). Our full GLMM had (0/1) urchin survival as its response variable, and its predictor variables were urchin average activity level, average urchin covering behavior, urchin diameter, and all of their interaction terms. Lobster ID and trial ID were included as random effects in our analysis. To build

the model with the greatest likelihood, we ran this full model through backwards stepwise BIC model selection (Chambers & Hastie, 1992).

238

2.6.3 | Cross contextual covering behavior and activity

We tested (1) whether urchins' pre-mesocosm average covering percentage predicted covering behavior in the staged interactions, and (2) whether pre-mesocosm average velocity (activity level) correlated with the frequency of movement between quadrants of staged interactions. However, urchins varied in their number of behavioral observations collected during staged interactions, because urchins varied in how long they survived. Therefore, we weighted urchins' average covering percentage and frequency of movement between quadrants during staged interactions by probability of survival estimated. Survival probability was estimated using a binomial logistic regression with survival as the response variable and activity level, covering behavior, diameter, activity by covering behavior, activity by diameter, and covering behavior by diameter interaction terms as explanatory variables. To calculate weighted average covering percentage and frequency of movement between quadrants, we multiplied centered individuals' average covering percentage and frequency of movement by individuals' probability of survival respectively. Weighted average covering percentage and weighted frequency of movement between quadrants were non-Gaussian. Therefore, we used Spearman's rank-order tests to assess the correlation between pre-interaction covering percentage and weighted in situ covering percentage, and the correlation between pre-interaction velocity and weighted in situ movement between quadrants.

255

3 | RESULTS

257

3.1 | Repeatability of behavior

259

Urchins behaved consistently across time in their average velocity/activity level (Repeatability [95% CIs] = 0.30 [0.03, 0.57]) and average covering percentage (Repeatability [95% CIs] = 0.73 [0.38, 0.87]). Between-individual differences in covering behavior are notably very high in this test population (Bell et al., 2009). Both of these behaviors are repeatable across a period longer than the lobster predation trials.

3.2 | Spiny lobster-sea urchin staged interactions

Urchins had a survival rate of 33.33 ± 6.88 % SE. The best model contained urchin diameter, activity level, and covering behavior (BIC = 75.3, Δ BIC = 3.5). Slow-moving purple urchins survived more frequently than fast-moving urchins (Figure 1; Activity Level: $\beta = -2.99 \pm 1.30$ SE, $z = -2.30$, $p = 0.02$). However, the effect (Figure 2A; Activity Level x Diameter: $\beta = 0.16 \pm 0.08$ SE, $z = 1.97$, $p < 0.05$). Urchin covering behavior, on its own, had no direct effect on urchin survival (Covering behavior: $\beta = -0.01 \pm 2.37$ SE, $z = -0.04$, $p = 0.98$). However, smaller urchins that exhibited greater covering behavior were more likely to survive, whereas urchin covering behavior appeared unrelated to survival among large urchins (Figure 2B; Covering behavior x Diameter: $\beta = -0.38 \pm 0.18$ SE, $z = -2.12$, $p = 0.03$). Larger urchins appeared to be slightly more likely to survive than small urchins, but not significantly so (Diameter: $\beta = 0.06 \pm 0.03$ SE, $z = 1.68$, $p = 0.09$), and Lobster ID and trial iteration failed to explain a significant component of the variation in urchin survival, suggesting that survival rates were homogenous across mesocosms.

3.3 | Cross contextual covering behavior and activity

Individual-level differences in average covering percentage were correlated with weighted average covering percentages during staged interactions (Kendall's tau: $\tau = 0.26$, $p = 0.01$). Individual activity, however, seemingly did not persist across contexts, as individual-level differences in average velocity were not correlated with weighted frequency of movement between quadrants during staged interactions

(Kendall's tau: $\tau = 0.04$, $p = 0.69$). However, activity level was measured somewhat differently in pre-trial assays during the trial itself, which obscures the interpretability of this result.

4 | DISCUSSION

Here we evaluated the effects of individual size (test diameter), activity (velocity), and covering behavior on prey survival using California spiny lobsters and purple sea urchins. We found that slow-moving urchins were more likely to survive encounters with lobsters (Figure 1), whereas covering behavior did not predict urchin survival on its own. This is consistent with findings suggesting that active prey are more likely to succumb to predation (Huey & Pianka, 1981; Perry, 2007), particularly with sedentary predators, and that urchin covering behavior does not regularly predict survival in a direct manner (Zhao et al., 2014). Urchin size also did not predict survival directly. Instead, urchin size interacted with urchin activity level and covering behavior to determine urchin survival. Specifically, when purple sea urchins were small, high activity level and low covering behavior reduced their chances of survival, but when urchins were large, their behavior predicted their survival less clearly (Figure 2). Taken together, these findings convey that, even in confined experimental conditions, urchin covering and movement behavior helps to determine survival with predators, but that the ideal behavioral phenotype for survival is contingent on size.

Classical theory on predator-prey interactions predicts that more active animals will be more susceptible to predation (Lima & Dill, 1990). However, more nuanced theoretical frameworks predict that the relationship between prey activity patterns and survival will depend on the foraging mode of the predator (Huey & Pianka, 1981; Perry, 2007; Sweeney et al., 2013). In particular, high activity levels are predicted to be especially costly when prey interact with sit-and-wait or sit-and-pursue predators, which rely on prey movement for encounters. Our results here lend support to both predictions. We found that fast-moving purple urchins were more likely to succumb to lobsters than sedentary urchins. This outcome

is consistent with classical optimal foraging and predator-prey interaction theory. However, the fact that spiny lobsters deploy sit-and-pursue foraging tactics (MacDiarmid, Hickey, & Maller, 1991; Tegner & Levin, 1983) suggests that this outcome should be even more likely. The effect of urchin movement speed also deteriorated with urchin size. This finding suggests that activity and covering behavior only mediates survival for smaller and easier to eat urchins. When urchins reach a certain size, they can safely move about with less risk of being eaten (Brooks & Dodson, 1965). It is possible that the detrimental effect of activity level may be driven by the relatively small size of our arenas. Taken together, the age and size structure of urchin populations may determine whether and how lobster predation applies selection pressure on urchin behavior.

Our findings suggest that urchin behavior may potentially play a role in driving the dynamics of kelp forest ecosystems at fine spatial scales. Specifically, smaller (e.g., younger or resource deprived) and more active sea urchin populations may be more susceptible to predation by spiny lobsters, which may alter the role that lobsters play in driving the dynamics of urchin prey. Recent studies have emphasized the importance of size-structured interactions between lobsters and urchins in driving the dynamics of trophically-linked species within the kelp forest (Dunn, Baskett, & Hovel, 2017). However, the role of intraspecific variation in urchin behavior in mediating these size-structured interactions and the resulting effects on kelp forest ecosystem dynamics is not well-studied. A comprehensive integration of intraspecific variation in urchin behavior in driving interactions between urchins and their predators, as well as urchins and kelp, will require a deeper understanding of how urchin behavior shapes their impacts on kelp (amount consumed, foraging mode, etc.) and how conflicting pressures, like the introduction or reintroduction of other predators, including sea otters, or competitors, might generate conflicting selection pressures on prey behavior.

Conclusions

We found that purple urchin size and behavioral tendencies synergistically determined their susceptibility to predation by spiny lobsters, which are known predators of urchins in nature (Lafferty, 2004; Tegner & Levin, 1983). Understanding the links between prey traits and performance is central to understanding trait evolution and, in the case of this system, potentially important for predicting community level outcomes. Urchin herbivory can defoliate kelp forests, thereby precipitating local loss of biodiversity (Graham, 2004) and altering the value of fisheries (Hamilton, Caselle, Malone, & Carr, 2010; Pearse, 2006). Our results here are a tentative first step towards leveraging the predictive value of intraspecific trait variation to predict how key players in kelp forest ecosystems interact. Moving forward, the challenge is to document how other environmental pressures cause alterations to the trait distributions of predator and prey populations, either in the form of selection or via phenotypic plasticity, and either trophic level's effects on community level outcomes. We reason that a comprehensive understanding of the role of predator and prey traits in this system will ultimately enable us to better predict or even mitigate urchin generated transitions in kelp forest states. There remains much to be learned, but our results here hint that the effects of individual variation in this system are likely to be strong.

ACKNOWLEDGEMENTS

We are indebted to the California Coastal Commission for issuing research and collection permits (SCP). We would also like to thank Christoph Pierre for collecting the animals for these studies and assisting in their laboratory maintenance. Funds for this work were generously provided by the University of California, Santa Barbara as start-up to J.N.P., NSF grant awards to J.N.P. (No. 1352705 and No. 1455895), and an NIH grant award to J.N.P. (No. R01GM115509).

5 | REFERENCES

- Amsler, C. D., McClintock, J. B., & Baker, B. J. (1999). An Antarctic feeding triangle: defensive interactions between macroalgae, sea urchins, and sea anemones. *Marine Ecology Progress Series*, 183, 105-114.
- Bates, D., Mächler, M., Bolker, B., & Walker, S. (2014). Fitting linear mixed-effects models using lme4. *arXiv preprint arXiv:1406.5823*.
- Belgrad, B. A., & Griffen, B. D. (2016). Predator–prey interactions mediated by prey personality and predator hunting mode. *Proceedings of the Royal Society of London B: Biological Sciences*, 283(1828), 20160408.
- Bell, A. M., Hankison, S. J., & Laskowski, K. L. (2009). The repeatability of behaviour: a meta-analysis. *Animal Behaviour*, 77(4), 771-783.
- Bolnick, D. I., Amarasekare, P., Araújo, M. S., Bürger, R., Levine, J. M., Novak, M., . . . Vasseur, D. A. (2011). Why intraspecific trait variation matters in community ecology. *Trends in Ecology & Evolution*, 26(4), 183-192.
- Bolnick, D. I., Svanbäck, R., Fordyce, J. A., Yang, L. H., Davis, J. M., Hulsey, C. D., & Forister, M. L. (2003). The ecology of individuals: incidence and implications of individual specialization. *The American Naturalist*, 161(1), 1-28.
- Brooks, J. L., & Dodson, S. I. (1965). Predation, body size, and composition of plankton. *Science*, 150(3692), 28-35.
- Brown, J. S., Laundré, J. W., & Gurung, M. (1999). The ecology of fear: optimal foraging, game theory, and trophic interactions. *Journal of Mammalogy*, 80(2), 385-399.
- Burt, J. M., Tinker, M. T., Okamoto, D. K., Demes, K. W., Holmes, K., & Salomon, A. K. (2018). Sudden collapse of a mesopredator reveals its complementary role in mediating rocky reef regime shifts. *Proceedings of the Royal Society of London B: Biological Sciences*, 285(1883), 20180553.
- Carter, A. J., Feeney, W. E., Marshall, H. H., Cowlishaw, G., & Heinsohn, R. (2013). Animal personality: what are behavioural ecologists measuring? *Biological Reviews*, 88(2), 465-475.
- Caselle, J. E., Davis, K., & Marks, L. M. (2018). Marine management affects the invasion success of a non-native species in a temperate reef system in California, USA. *Ecology letters*, 21(1), 43-53.
- Chambers, J. M., & Hastie, T. J. (1992). *Statistical models in S* (Vol. 251): Wadsworth & Brooks/Cole Advanced Books & Software Pacific Grove, CA.
- Chang, C.-c., Teo, H. Y., Norma-Rashid, Y., & Li, D. (2017). Predator personality and prey behavioural predictability jointly determine foraging performance. *Scientific reports*, 7.
- Crutsinger, G. M., Collins, M. D., Fordyce, J. A., Gompert, Z., Nice, C. C., & Sanders, N. J. (2006). Plant genotypic diversity predicts community structure and governs an ecosystem process. *Science*, 313(5789), 966-968.
- Dall, S. R., Houston, A. I., & McNamara, J. M. (2004). The behavioural ecology of personality: consistent individual differences from an adaptive perspective. *Ecology letters*, 7(8), 734-739.
- Dingemanse, N. J., Kazem, A. J., Réale, D., & Wright, J. (2010). Behavioural reaction norms: animal personality meets individual plasticity. *Trends in Ecology & Evolution*, 25(2), 81-89.
- DiRienzo, N., Pruitt, J. N., & Hedrick, A. V. (2013). The combined behavioural tendencies of predator and prey mediate the outcome of their interaction. *Animal Behaviour*, 86(2), 317-322.
- Dochtermann, N. A., & Dingemanse, N. J. (2013). Behavioral syndromes as evolutionary constraints. *Behavioral Ecology*, 24(4), 806-811.
- Dumont, C. P., Drolet, D., Deschênes, I., & Himmelman, J. H. (2007). Multiple factors explain the covering behaviour in the green sea urchin, *Strongylocentrotus droebachiensis*. *Animal Behaviour*, 73(6), 979-986.
- Dunn, R. P., Baskett, M. L., & Hovel, K. A. (2017). Interactive effects of predator and prey harvest on ecological resilience of rocky reefs. *Ecological Applications*, 27(6), 1718-1730.
- Ebert, T. A. (2010). Demographic patterns of the purple sea urchin *Strongylocentrotus purpuratus* along a latitudinal gradient, 1985–1987. *Marine Ecology Progress Series*, 406, 105-120.

- 408 Einfalt, L. M., & Wahl, D. H. (1997). Prey selection by juvenile walleye as influenced by prey
409 morphology and behavior. *Canadian Journal of Fisheries and Aquatic Sciences*, 54(11), 2618-
410 2626.
- 411 Estes, J. A., & Duggins, D. O. (1995). Sea otters and kelp forests in Alaska: generality and variation in a
412 community ecological paradigm. *Ecological Monographs*, 65(1), 75-100.
- 413 Gosling, S. D. (2001). From mice to men: what can we learn about personality from animal research?
414 *Psychological bulletin*, 127(1), 45.
- 415 Graham, M. H. (2004). Effects of local deforestation on the diversity and structure of southern California
416 giant kelp forest food webs. *Ecosystems*, 7(4), 341-357.
- 417 Hamilton, S. L., Caselle, J. E., Malone, D. P., & Carr, M. H. (2010). Incorporating biogeography into
418 evaluations of the Channel Islands marine reserve network. *Proceedings of the National Academy*
419 *of Sciences*, 107(43), 18272-18277.
- 420 Harrold, C., & Reed, D. C. (1985). Food availability, sea urchin grazing, and kelp forest community
421 structure. *Ecology*, 66(4), 1160-1169.
- 422 Hoyle, J. A., & Keast, A. (1987). The effect of prey morphology and size on handling time in a piscivore,
423 the largemouth bass (*Micropterus salmoides*). *Canadian Journal of Zoology*, 65(8), 1972-1977.
- 424 Huey, R. B., & Pianka, E. R. (1981). Ecological consequences of foraging mode. *Ecology*, 62(4), 991-
425 999.
- 426 Hughes, A. R., & Stachowicz, J. J. (2004). Genetic diversity enhances the resistance of a seagrass
427 ecosystem to disturbance. *Proceedings of the National Academy of Sciences of the United States of*
428 *America*, 101(24), 8998-9002.
- 429 Johnson, M. T., Lajeunesse, M. J., & Agrawal, A. A. (2006). Additive and interactive effects of plant
430 genotypic diversity on arthropod communities and plant fitness. *Ecology letters*, 9(1), 24-34.
- 431 Jones, K. A., & Godin, J.-G. J. (2010). Are fast explorers slow reactors? Linking personality type and
432 anti-predator behaviour. *Proceedings of the Royal Society of London B: Biological Sciences*,
433 277(1681), 625-632.
- 434 Keiser, C. N., Ingle, S. J., Toscano, B. J., Scharf, I., & Pruitt, J. N. (2017). Habitat complexity dampens
435 selection on prey activity level. *Ethology*.
- 436 Lafferty, K. D. (2004). Fishing for lobsters indirectly increases epidemics in sea urchins. *Ecological*
437 *Applications*, 14(5), 1566-1573.
- 438 Lichtenstein, J. L., Wright, C. M., McEwen, B., Pinter-Wollman, N., & Pruitt, J. N. (2017). The
439 multidimensional behavioural hypervolumes of two interacting species predict their space use and
440 survival. *Animal Behaviour*, 132, 129-136.
- 441 Lima, S. L., & Dill, L. M. (1990). Behavioral decisions made under the risk of predation: a review and
442 prospectus. *Canadian Journal of Zoology*, 68(4), 619-640.
- 443 MacDiarmid, A., Hickey, B., & Maller, R. (1991). Daily movement patterns of the spiny lobster *Jasus*
444 *edwardsii* (Hutton) on a shallow reef in northern New Zealand. *Journal of Experimental Marine*
445 *Biology and Ecology*, 147(2), 185-205.
- 446 McGhee, K. E., Pintor, L. M., Suhr, E. L., & Bell, A. M. (2012). Maternal exposure to predation risk
447 decreases offspring antipredator behaviour and survival in threespined stickleback. *Functional*
448 *Ecology*, 26(4), 932-940.
- 449 Miller, J. R., Ament, J. M., & Schmitz, O. J. (2014). Fear on the move: predator hunting mode predicts
450 variation in prey mortality and plasticity in prey spatial response. *Journal of Animal Ecology*,
451 83(1), 214-222.
- 452 Nannini, M. A., Parkos III, J., & Wahl, D. H. (2012). Do Behavioral Syndromes Affect Foraging Strategy
453 and Risk-Taking in a Juvenile Fish Predator? *Transactions of the American Fisheries Society*,
454 141(1), 26-33.
- 455 Pawson, D. L., & Pawson, D. J. (2013). Bathyal sea urchins of the Bahamas, with notes on covering
456 behavior in deep sea echinoids (Echinodermata: Echinoidea). *Deep Sea Research Part II: Topical*
457 *Studies in Oceanography*, 92, 207-213.

- Pearish, S., Hostert, L., & Bell, A. M. (2013). Behavioral type–environment correlations in the field: a study of three-spined stickleback. *Behavioral Ecology and Sociobiology*, 67(5), 765-774.
- Pearse, J. S. (2006). Ecological role of purple sea urchins. *Science*, 314(5801), 940-941.
- Perry, G. (2007). Movement patterns in lizards: measurement, modality, and behavioral correlates. *Lizard ecology: The evolutionary consequences of foraging mode*, 13-48.
- Post, D. M., Palkovacs, E. P., Schielke, E. G., & Dodson, S. I. (2008). Intraspecific variation in a predator affects community structure and cascading trophic interactions. *Ecology*, 89(7), 2019-2032.
- Preisser, E. L., & Bolnick, D. I. (2008). The many faces of fear: comparing the pathways and impacts of nonconsumptive predator effects on prey populations. *PloS one*, 3(6), e2465.
- Pruitt, J. N., Berdahl, A., Riehl, C., Pinter-Wollman, N., Moeller, H. V., Pringle, E. G., . . . Yeh, P. (2018). Social tipping points in animal societies. *Proceedings of the Royal Society of London B: Biological Sciences*, 285(1887), 20181282.
- Pruitt, J. N., Riechert, S. E., & Jones, T. C. (2008). Behavioural syndromes and their fitness consequences in a socially polymorphic spider, *Anelosimus studiosus*. *Animal Behaviour*, 76(3), 871-879.
- R Development Core Team. (2010). R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing.
- Reusch, T. B., Ehlers, A., Hämmerli, A., & Worm, B. (2005). Ecosystem recovery after climatic extremes enhanced by genotypic diversity. *Proceedings of the National Academy of Sciences of the United States of America*, 102(8), 2826-2831.
- Rudman, S. M., Rodriguez-Cabal, M. A., Stier, A., Sato, T., Heavyside, J., El-Sabaawi, R. W., & Crutsinger, G. M. (2015). Adaptive genetic variation mediates bottom-up and top-down control in an aquatic ecosystem. *Proceedings of the Royal Society of London B: Biological Sciences*, 282(1812), 20151234.
- Selkoe, K. A., Blenckner, T., Caldwell, M. R., Crowder, L. B., Erickson, A. L., Essington, T. E., . . . Hunsicker, M. E. (2015). Principles for managing marine ecosystems prone to tipping points. *Ecosystem Health and Sustainability*, 1(5), 1-18.
- Sih, A., Cote, J., Evans, M., Fogarty, S., & Pruitt, J. (2012). Ecological implications of behavioural syndromes. *Ecology letters*, 15(3), 278-289.
- Skelly, D. K. (1994). Activity level and the susceptibility of anuran larvae to predation.
- Stoffel, M. A., Nakagawa, S., & Schielzeth, H. (2017). rptR: repeatability estimation and variance decomposition by generalized linear mixed-effects models. *Methods in Ecology and Evolution*, 8(11), 1639-1644.
- Sweeney, K., Cusack, B., Armagost, F., O'Brien, T., Keiser, C. N., & Pruitt, J. N. (2013). Predator and prey activity levels jointly influence the outcome of long-term foraging bouts. *Behavioral Ecology*, 24(5), 1205-1210.
- Tegner, M., & Levin, L. (1983). Spiny lobsters and sea urchins: analysis of a predator-prey interaction. *Journal of Experimental Marine Biology and Ecology*, 73(2), 125-150.
- Werner, E. E., & Hall, D. J. (1974). Optimal foraging and the size selection of prey by the bluegill sunfish (*Lepomis macrochirus*). *Ecology*, 55(5), 1042-1052.
- Wilson, D. S., Coleman, K., Clark, A. B., & Biederman, L. (1993). Shy-bold continuum in pumpkinseed sunfish (*Lepomis gibbosus*): An ecological study of a psychological trait. *Journal of Comparative Psychology*, 107(3), 250.
- Wolf, M., & Weissing, F. J. (2012). Animal personalities: consequences for ecology and evolution. *Trends in Ecology & Evolution*, 27(8), 452-461.
- Zhao, C., Ji, N., Zhang, B., Sun, P., Feng, W., Wei, J., & Chang, Y. (2014). Effects of covering behavior and exposure to a predatory crab *Charybdis japonica* on survival and HSP70 expression of juvenile sea urchins *Strongylocentrotus intermedius*. *PloS one*, 9(5), e97840.

507 **6 | FIGURE LEGENDS**

508

509 **FIGURE 1.** Box plots depicting differences in average velocity between surviving and eaten *S.*

510 *purpuratus* in staged interactions with *P. interruptus*. Dots denote outliers, whiskers represent 10th and

511 90th percentiles, gray boxes indicate interquartile range: calculated median inclusive, and the central line

512 denotes the median velocity.

513

514 **FIGURE 2.** Synergistic effects of covering behavior and size (panel A, top) and activity and size (panel

515 B, bottom) on identity of *S. purpuratus* that survived (white circles) and were eaten (black circles) during

516 staged interactions with *P. interruptus*.

517

518

7 | FIGURES

Figure 1.

523 **Figure 2.**

524

525