The conspicuousness of the toxic *Heliconius* butterflies across time and habitat

Denise Dalbosco Dell’Aglia^{1,2*}

Jolyon Troscianko³

Martin Stevens³

W Owen McMillan²

Chris D Jiggins¹

Affiliations

1. Butterfly Genetics Group, Department of Zoology, University of Cambridge, Cambridge, United Kingdom

2. Smithsonian Tropical Research Institute, Panama City, Panama

3. Centre for Ecology and Conservation, College of Life and Environmental Sciences, University of Exeter, Penryn, United Kingdom

*** Corresponding author**

E-mail: denise.ddd3@gmail.com

Key words

aposematism; light environment; avian vision; butterfly vision; colour signal

Abstract

Forests are a mosaic of light spectra, and colour signal efficiency might change in different light environments. Local adaptation in *Heliconius* butterflies is linked to microhabitat use and the colourful wing colour patterns may also be adapted for signalling in different light environments. These butterflies exhibit conspicuous colours as a warning to predators that they are toxic and should be avoided, but also find and choose potential mates based on colour signals. The two selection pressures of predation and mate preference are therefore acting together. In this study we analysed the contrast of two *Heliconius* mimicry rings in their natural habitats under varying degrees of forest fragmentation and light conditions. We used digital image analyses and mapped the bird and butterfly vision colour space in order to examine whether warning colours have greater contrast and if they transmit a consistent signal across time of the day and habitat in a tropical forest. We tested conspicuousness using opponent colour channels against a natural green background. For avian vision, colours are generally very stable through time and habitat. For butterfly vision, there is some evidence that species are more contrasting in their own habitats, where conspicuousness is higher for red and yellow bands in the border and for white in the forest. Light environment affects *Heliconius* butterflies' warning signal transmission to a higher degree through their own vision, but to a lesser degree through avian predator vision. This work provides insight into the use of colour signals in sexual and natural selection in the light of ecological adaptation.

Introduction

The success of a signal is related to its effectiveness in a specific environment and how strongly it influences the behaviour of the receiver (Endler 1978). Forests are a mosaic of light colours, and the same colour pattern can have an altered appearance in different light environments (Endler 1993). If an individual shows high reflectance of a specific wavelength, but the environment lacks light in that part of the spectrum, the region of high reflection will be unimportant as a signal (Stevens et al. 2007). Ambient light spectra also vary from dawn to dusk, hence species that signal only at certain times and places are expected to evolve characteristics and predictable combinations of colours for particular environments (Endler 1993). Therefore, ambient light characteristics should be included together with the receiver visual system to understand the microhabitat choice and behaviour of animals.

Signals depend on the habitat where animals live in, since light conditions can alter colour perception by filtering wavelengths and altering visual backgrounds (Endler 1993; Lovell et al. 2005). Sensory drive explains the process of adaptation of signalling and sensory systems to the local environment (Endler 1992; Endler and Basolo 1998). Environment tuned spectral sensitivity is better known in aquatic habitats, such as in guppies (Endler 1980) and cichlid fish (Seehausen et al. 2008), as compared to terrestrial light environments. On land, colour depends on the reflection of the surroundings and has greater variability over time (Boughman 2002). Habitat signal transmission can favour diversification of mating signals through local adaptation, leading to reproductive isolation. Distinct *Anolis* lizards male dewlaps are found in different microhabitats (Fleishman et al. 1997). Male dewlap colours are more conspicuous in their own habitat than in other habitats, mainly because of the contrast against the background in the ultraviolet (UV) range (Leal and Fleishman 2002). Perception of colours in different light

conditions can also influence attacks by predators, for example among butterflies in an environment with high UV light, birds aimed at the butterfly wings, more specifically the marginal white eyespots that have UV reflectance, instead of the head (Olofsson et al. 2010).

Local adaptation in *Heliconius* butterflies commonly involves adaptation to specific microhabitat use (Estrada and Jiggins 2002; Elias et al. 2008; Jiggins 2008). Mimicry rings are groups of unpalatable species that share the same warning colour, and these tend to be found in different microhabitats such as forest or open areas. The *Heliconius* habitats are associated with the use of larval host-plants, adult food plants, sexual behaviour and gregarious roosting (Mallet and Gilbert 1995). Species that lay eggs on *Passiflora* species that occur in second growth tend to be seen in open areas, while species that lay eggs on canopy *Passiflora* vines are seen flying high in the forest. The choice of microhabitat also might be connected with light differences between those environments, such as the choice of using shady areas in communal roosting (Mallet and Gilbert 1995; Finkbeiner 2014).

Therefore, different light environments should create microhabitats where butterfly signals would be more efficient. Although mimicry rings differ in their microhabitat, the light environment has not been measured to verify whether colour patterns could be specifically adapted to particular light environments. The colourful wing colours of *Heliconius* butterflies may also be subject to evolution caused by sensory drive due to their potentially conflicting roles in predation and mate preference. Many species exhibit Müllerian mimicry (Müller 1879), in which two or more species share the same conspicuous colour as a warning to predators that they are toxic and should be avoided (Benson 1972). Also, these butterflies find and choose potential mates based on colour signals, which can lead to reproductive isolation (Jiggins et al. 2001; Sweeney et

al. 2003; Kronforst et al. 2006). Furthermore, communication between conspecifics might be based on UV signals, since *H. erato* females express the duplicate UV opsin gene, which allows a greater degree of discrimination of the UV-yellow wing patches (Briscoe et al. 2010; Bybee et al. 2012; McCulloch et al. 2016; Dell’Aglia et al. 2018).

This microhabitat structuring allows mimicry rings to remain distinct. This may be because there are sets of predators in different habitats, each of which perceive a different mimicry ring as the most abundant pattern (Joron and Mallet 1998). Although little is known of the specific predators that attack *Heliconius*, it seems likely that their aposematic signals are directed at several predators with different visual abilities and spectral sensitivities (Dell’Aglia et al. 2018). Ambient light together with predator sensitivity can interfere with the interpretation of the information perceived from colour signals.

Warning coloration should, therefore, be easy to detect and memorize even in heterogeneous environments and light conditions (Guilford and Dawkins 1991; Endler 1992). Warning signals are often dominated by red, yellow and orange, frequently contrasting with black, which are the main colours in *Heliconius*. The reason why these long-wavelength colours are widely represented in aposematic coloration is that they are highly conspicuous against natural backgrounds, are more stable across light conditions, allowing long distance discrimination and detectability, and influence memorability (Guilford and Dawkins 1991; Stevens and Ruxton 2012; Arenas et al. 2014; Dell’Aglia et al. 2016).

Perception of colour depends on several neurophysiological mechanisms, such as the presence of opponent colour channels. This chromatic mechanism involves comparisons of receptors outputs, in which opposite neural pathways are either activated or inhibited depending on the stimuli reaching the eye (Kelber et al. 2003; Renoult et al.

2015). This mechanism is useful especially regarding colour stability against spatial and temporal variation in illumination (Lovell et al. 2005; Renoult et al. 2015). For example birds, the major predator of aposematic butterflies, have tetrachromatic vision and seemingly have at least three opponent channels, as found in domestic chicks (Vorobyev et al. 1998; Osorio et al. 1999). Opponent channels have also been described for insects (Chittka et al. 1992; Chittka 1996) and butterflies (Kelber 1999), and have been hypothesized for *Heliconius* butterflies although more behavioural analyses are needed to confirm which opponent channels are actually used (Swihart 1971, 1972; Bybee et al. 2012). Butterflies in the genus *Papilio* have duplicate LW opsin genes to see in the red and green range (Kelber 1999; Briscoe 2008), while *Heliconius* has only one LW opsin to see red and green, and differences in sensitivity are associated with the presence of red filtering pigments in the ommatidia (Zaccardi et al. 2006; McCulloch et al. 2016). Thus, we expect avian predators and butterflies to rely on these high-contrast systems to process information under a changing light environment.

The aim of this study was to analyse *Heliconius* warning colouration under different light conditions in their natural habitats. In particular, to test conspicuousness of the wing colouration against a natural green background, encoded by opponent colour channels. Using digital image analyses, butterfly wings were photographed and mapped to UVS and VS avian predator vision and to *Heliconius erato* vision (Figure 1). Our predictions are that (1) signal contrast and conspicuousness for avian predators should have constancy, that should be stable throughout the day and in different light environments (Stevens and Ruxton 2012; Arenas et al. 2014). Warning signals might be honest indicators of prey unprofitability to predators, and if signals fluctuate through the day and between light environments, we would predict that this could delay learning by predators and be costly to the prey. Similarly for internal contrasts (i.e. contrast between

black and the coloured bands), therefore conspicuousness would not rely totally on background contrast but also on internal patterns which account for close-distance conspicuousness (Endler 1978; Aronsson and Gamberale-Stille 2009). (2) From a *Heliconius* butterfly perspective; we predict that signal contrast and conspicuousness should show habitat-specific maximum background contrast and higher colour differences in their own habitats (Table 1), which would facilitate detection and species identification. We therefore predict that selection for signal constancy will be much stronger in the avian visual system as compared to the butterfly visual system.

Figure 1. Co-mimics used in this study. (a) *Heliconius erato demophoon*. (b) *H. melpomene rosina*. (c) *H. cydno*. (d) *H. sapho*.

Table 1. Colour patches for each co-mimic pair studied and typical microhabitats and light conditions where these co-mimics occur. Microhabitats descriptions are based on Estrada and Jiggins (2002) and light conditions based on Endler (1993) and personal observations.

Co-mimics	Colour	Microhabitat	Light conditions
<i>H. erato demophoon</i> & <i>H. melpomene rosina</i>	red & yellow	forest border open area	partial shade no shade
<i>H. sapho</i> & <i>H. cydno</i>	white	closed forest canopy	full shade partial shade

Material and Methods

Study site and species

Fieldwork was performed during the dry season, along Pipeline Road, a tropical lowland rainforest in the Panama Canal Zone (Parque Nacional Soberanía, 9°7'33"N, 79°42'90"W). Pipeline Road makes a transect through the forest, creating a heterogeneous habitat with open sunny areas and close canopy exceeding 30 m in height. All specimens were collected in the area. Two pairs of co-mimics that live in sympatry were selected, *H. erato demophoon* (n = 8) and *H. melpomene rosina* (n = 8), *H. sapho* (n = 5) and *H. cydno* (n = 5), belonging to two different mimicry rings, red and yellow, and white, respectively (Figure 1 and Table 1).

Digital photography

The general approach and methodology for this work was based on previous work with colour stability using opponent signals (Lovell et al. 2005; Arenas et al. 2014). The spectral reflectance of mimetic pairs was investigated using digital photography. This provides a way to control for natural variation in luminance intensity (shadowing) that is not captured by spectrometry, and also allows non-invasive colour measurements easily applied in the field (Stevens et al. 2007; Troscianko and Stevens 2015). Therefore, through this method we could obtain colour measures under the sensitivity of all receiver photoreceptors (300-700 nm) in the actual viewing conditions of conspecifics and avian predators.

Fresh wings of each specimen were photographed following the same methods of image collection in Dell'Aglio et al. (2018). The camera was fitted to a tripod and pointed towards the ground (90°) at a height of approximately 80 cm. Each photo setup included two individuals, one of each species of the co-mimic pair, a 40% grey standard

(Spectralon® Labsphere) used for calibration and a leaf freshly collected to make background measures. The species used was *Guazuma ulmifolia* (Sterculiaceae), a small abundant shrub across all Pipeline Road, which facilitated the collection of fresh leaves.

Photos were taken under three different arboreal canopy conditions, forest border, closed forest, and open area, where those butterflies are usually seen (Table 1). All photos were taken under sunny to part-cloudy days, with three replicates in each habitat making sure that the amount of light was similar. In order to standardize the replicates, light measures were taken with a digital light meter (Digital Lux meter, Tondaj LX-1010B), which measures the total amount of LW (555 nm) per square meter (Lux) (Figure 2). Also photographs of the canopy were taken in order to measure vegetative cover, which was 81.5% (SE $\pm$ 0.2) for closed forest, 59.8 % (SE $\pm$ 3.4) for forest border, and 0% for open area. The aim was to analyze how colour signals are perceived throughout the morning when butterflies are most active. Therefore, photos were taken at dawn (7 am), morning (9 am) and noon (12 pm) during a short period of 15min as light conditions change rapidly. Open area photographs were taken only at 12 pm because of high sunlight incidence (Figure 2) and to represent a highly used environment by butterflies at this period of day.

Figure 2. Amount of light differs across light environment and time of day. Showing the average log Lux between the three replicates of each habitat ($\pm$ SE). Lux represents the amount of long-wave (555 nm) per square meter.

Image analysis and visual modelling

All images were processed and analyzed into the imaging software ImageJ (Rasband, 1997-2012). RAW human-visible and UV images were linearized and aligned following the methodology of Troscianko and Stevens (2015) and Arenas et al. (2014). Normally, photos would be normalized to the grey standard, which removes effects of light conditions (Arenas et al. 2014). Since our main interest was to measure how coloration changes in different light environments, the images were not normalized. Instead, an average grey standard value was obtained from all photographs. Photon catch values were obtained for each colour using the entire patch from linearized photos, and subsequently these values were multiplied by each photo exposure time and normalized with the average grey standard. With this methodology we were able to calculate how particular environment and time varies from the natural average light, and also assume

colour constancy, a neural mechanism that compensates for changes in illumination (Stevens et al. 2007; Arenas et al. 2014). We used the average photon catch results from the three habitats replicates. Predicted photon catch values were obtained using spectral sensitivity for each cone type of the blue tit (*Cyanistes caeruleus*) for the UV-sensitive vision (UVS) (Hart et al. 2000), peafowl (*Pavo cristatus*) for the violet-sensitive vision (VS) (Hart 2002) and *Heliconius erato* (Briscoe et al. 2010; McCulloch et al. 2016).

Background of many terrestrial habitats is dominated by greenish vegetation; therefore, a green leaf was chosen to make contrast calculations. Differences between light environment and time of the day were calculated using the contrast of warning colours against an average green leaf. Channel activation in avian vision was calculated using the Red-Green (RG), Blue-Yellow (BY) and Blue-UV opponent channel (Osorio et al. 1999; Lovell et al. 2005; Stevens et al. 2009). For the achromatic signal, we used avian double cones (DBL). Using a ratio-based approach suggested by Lovell et al. (2005), we calculated the opponent channel responses as follows:

$$RG = LW - MW / LW + MW$$

$$BY = SW - (LW + MW) / SW + (LW + MW)$$

$$Blue-UV = SW - UV / SW + UV$$

$$Achromatic = DBL$$

Opponent channels for *Heliconius* were based on existing bird opponent channels and on what has been proposed in earlier studies (Swihart 1971, 1972; Bybee et al. 2012). The *H. erato* compound eye has red filtering pigments that shift LW photoreceptor sensitivity from green to red and as the physiological mechanisms underlying these two LW photoreceptors are not known, sensitivities in Green (560 nm) and Red (600 nm) were used (McCulloch et al. 2016, 2017). To investigate differences between co-mimic

species, we calculated opponent channel activation based on the prediction that *Heliconius* mating system might use UV2-UV1 and RG contrasts for mate choice (Bybee et al. 2012; McCulloch et al. 2016, 2017). Channel activation in *Heliconius* vision was calculated using the opponent channels as follows:

$$RG = Red - Green / Red + Green$$

$$BY = Blue - (Red + Green) / Blue + (Red + Green)$$

$$Blue-UV2 = Blue - UV2 / Blue + UV2$$

$$UV2-UV1 = UV2 - UV1 / UV2 + UV1$$

To examine whether warning colours have greater contrast against green background we calculated the Weber Contrast (Whittle 1994), which takes into account the image value of the objects of interest as a fraction of background appearance using the formula:

$$C = (object - background) / background$$

Where *background* corresponds to the green leaf opponent channel values, and *object* corresponds to warning colour opponent channel values. This measure is suited to comparisons between small objects against larger backgrounds, such as butterflies against the green forest. For internal contrast, achromatic values of the warning colours were used against the black of each individual wing as background (Arenas et al. 2014). We plotted the mean absolute contrast of each colour signal as a function of time and light environment for the three vision models.

Statistical analyses

All statistical calculations were processed in the software R 3.5.3 (R Core Team 2019). Our approach was to model colour contrasts over the course of a day and under different habitats in term of both predator and butterfly vision. Normality tests showed

that contrast data were not normally distributed, therefore data were transformed to normality using square-root transformation and the transformed data were used in all statistical analyses. Raw data was plotted to illustrate the results. To test our predictions, general linear mixed models were performed using Satterthwaite approximations with random effects (packages *lme4* and *lmerTest*) and Tukey's post-hoc (package *multcomp*). The models were fitted accordingly to the predictions outlined above. Analyses were carried out using contrast values as the dependent variable, and fixed and random factors varied depending on the question. Factors were individuals, colour (red, yellow, white), habitat (border, forest, area), time (7am, 9am, 12pm), and bird vision (UVS, VS). For *Heliconius* vision, we also added side of the wing (dorsal, ventral) because this trait might be more important for butterflies than for their avian predators.

Results

Signal contrast and conspicuousness for avian predators

Red was generally the most contrasting colour against a green background in the RG opponent channel as compared to yellow ($z = -11.10$, $P < 0.001$, Table S1) and white ($z = -18.0$, $P < 0.001$, Table S1). In contrast, white had higher contrasts against a green background in the BY channel, as compared to red ($z = 22.88$, $P < 0.001$, Table S1) and yellow ($z = -31.47$, $P < 0.001$, Table S1) (Figure 3). Colours in open areas showed a higher contrast, such in the RG channel for red band ($t = 7.54$, $P < 0.001$, Table S2) with no difference between border and forest ($z = -0.31$, $P = 0.94$, Table S2) (Figure 3). In the Blue-UV opponent channel, UVS and VS birds could perceive red and yellow with less stability, and yellow showed higher contrast early in the morning than at noon (7 am: $z = 12.15$, $P < 0.001$; 9 am: $z = 14.24$, $P < 0.001$, Table S3).

Internal achromatic contrast was higher for yellow, compared to red ($z = 41.89$, $P < 0.001$, Table S1) and white ($z = 12.42$, $P < 0.001$, Table S1). Moreover, yellow has more contrast in the border, which is the preferred habitat of yellow band butterflies, than in the forest ($z = -3.42$, $P = 0.001$, Table S4) (Figure 3).

Signal contrast and conspicuousness for Heliconius conspecifics

In some cases, contrasts followed our prediction that species would be more contrasting in their own habitats (Figure 4). The yellow colour was more contrasting in the border in the UV2-UV1 channel, especially early hours such as 7 am ($t = -23.1$, $P < 0.001$, Table S5) and 9 am ($t = -13.3$, $P < 0.001$, Table S5). White was more contrasting in the forest than in the border at 7 am in the UV2-UV1 channel ($t = 2.32$, $P = 0.014$, Table S5) and also at 7 am in the Blue-UV2 channel ($t = 6.12$, $P < 0.001$, Table S5). Also, in the Blue-UV2 channel, while white colour contrast decreased during the day in the forest, it increased at 12 pm in the border ($z = -4.11$, $P < 0.001$, Table S5) (Figure 4). The red colour showed large differences in the RG channel between dorsal and ventral side, with dorsal side with the higher contrast ($t = -40.04$, $P < 0.001$, Table S6). Same results were found for red in the Blue-Yellow and Blue-UV2 opponent channel (Table S6) (Figure 4).

Figure 3. Colour conspicuousness for avian predators. Mean absolute contrast of colour signals ($\pm$ SE, standard error) in the bird vision systems analyzed (circles, UVS; triangles, VS) through habitats (red, border; green, forest; blue, open) and time (7am, 9am, 12pm). Vertical panels show the three colour signals (red, yellow and white), horizontal panels show opponent channels against green leaf (top, Red-Green; middle, Blue-Yellow and Blue-UV) and against the black of the wing (bottom, Achromatic). Note: Channels have different y-axis values. Error bars smaller than data points are not shown.

Figure 4. Colour conspicuousness for *Heliconius* conspecifics. Mean absolute contrast of colour signals against green leaf in *Heliconius* vision through habitats (red, border; green, forest; blue, open), time (7 am, 9 am, 12 pm) and side of the wing (circles, dorsal; triangles, ventral). Vertical panels show colour signals (red, yellow and white), horizontal panels show opponent channels against green leaf (top, Red-Green; middle, Blue-Yellow and Blue-UV2; bottom, UV2-UV1). Error bars: $\pm$ 1 standard error (SE), error bars smaller than data points are not shown.

Discussion

The bright and contrasting *Heliconius* wing patterns appear well adapted for signalling distastefulness to predators. However, their colour constancy and appearance in different light environments remains poorly studied. Here we have shown that colours are indeed very stable for avian predator vision, but somewhat less so for *Heliconius* vision. This is consistent with the idea that wing patterns are primarily selected for their role in signalling distastefulness to predators.

Signal stability and conspicuousness to avian predators

Generally, warning signals involve combinations of long-wavelength colours such as red, orange, and yellow which are highly conspicuous against natural backgrounds and stable under different natural conditions (Lovell et al. 2005; Stevens and Ruxton 2012). In our results, *Heliconius* red colouration has higher detectability against average green background in the RG output and these results are consistent regardless of habitat and time of the day. Previous work has investigated colour stability through opponent colour channels and also showed that red coloration is more contrasting and stable against green backgrounds over the course of a day and across light conditions to the bird visual system (Lovell et al. 2005; Arenas et al. 2014). Moreover, the *Heliconius* yellow colouration also is highly conspicuous against its internal black pattern in the achromatic output. Achromatic information is one of the main cues used for motion detection (Hämäläinen et al. 2015). Therefore, our results suggest that red and yellow signals work together and are likely effective in stimulating avian opponent channels in order to be conspicuous in all light environments.

There is evidence that red and yellow colouration serve as reliable warning signal to avian predators (Ham et al. 2006; Svádová et al. 2009; Arenas et al. 2015), but that this

is less true of white colouration. One explanation is that white is more variable across time and habitat, so provides a less reliable signal under varied light conditions (Stevens and Ruxton 2012; Arenas et al. 2014). As an example, field and aviary experiments with polymorphic yellow and white wood tiger moths, *Parasemia plantaginis*, showed that yellow males are avoided more than white males by predators, but white males have higher mating success (Nokelainen et al. 2012). Our results showed that white contrasts against green background were lower and rather variable for avian vision. The co-mimics *H. sapho* and *H. cydno* also contain iridescence blue that was not measured with this methodology. However, the lack of high contrast in white colouration might be balanced with the fact that polarized light might act as a signal, especially in forest habitats (Sweeney et al. 2003; Douglas et al. 2007; Pegram et al. 2015).

Highly conspicuous warning signals are expected to evolve to be stable in their appearance throughout the day and between light environments, in order to remain honest indicators of prey unpalatability (Blount et al. 2009; Cortesi and Cheney 2010; Stevens and Ruxton 2012; Arenas et al. 2015). If warning signals fluctuate through time and space this could alter bird foraging experiences and reduce the effectiveness of the aposematic signal. The final decision on whether or not to attack a prey results from a combination of information reaching the predator brain, and for greater efficiency, aposematic coloration needs to be easy to remember (Endler 1988). Our results support this prediction, as colours were generally stable through time and light environments in all opponent systems with only a few exceptions. Notably these occurred where contrasts were higher in open areas and in the early morning. This might also be favourable as the prey would be more conspicuous when they are most vulnerable to predation, since birds are more active and forage early in the morning (Buskirk et al. 1972; Poulin et al. 2001;

Steiger et al. 2009). In agreement with this, *Heliconius* predation and roost disturbance has been observed in the early morning (Mallet 1986; Finkbeiner 2014).

Habitat and time influence conspicuousness in Heliconius conspecifics

Butterflies belonging to the two mimetic rings studied here tend to be segregated between habitats, corresponding to areas where the photographs were taken, although there is considerable overlap (Estrada and Jiggins 2002). We showed that the colours were more unstable when seen through *Heliconius* vision as compared to avian vision and some colours tend to be more contrasting in their respective habitats.

Our results provide some evidence that co-mimic rings are more conspicuous in their own habitat as seen through *Heliconius* vision, reinforcing the idea that ecological adaptation leads to spatial segregation to where detection would be facilitated. Some colours had higher contrast against green backgrounds in their respective habitat, such as yellow in the border and white in the forest. Nonetheless, red showed the opposite trend and was generally more contrasting in the forest. Differences across light environments could affect mating preferences by altering search costs for a specific colour pattern, and perhaps changing the fitness of different colour patterns. Adaptation in different microhabitats within the forest might have an influence on how closely related species commonly differ in pattern, while convergence in pattern occurs between more distantly related species (Joron and Mallet 1998). Ecological adaptation is attributed to habitat preference and leads to assortative mating (Jiggins 2008). The two sister species studied here, *H. melpomene rosina* and *H. cydno*, are known to rarely hybridise in the wild, hence microhabitat segregation reduces potential mating encounters between these two species and reduces gene flow (Mallet et al. 1998; Merrill et al. 2013). Subtle environmental conditions could affect recognition in mating behaviour as seen in the jumping spider,

Habronattus pyrrithrix, which red males were more successful in approaching females in the sunlight (Taylor and McGraw 2013).

The activation of opponent channels was often higher in the early hours of the morning, at the time when the butterflies are more active and leave their roost or perches to forage (Mallet 1986; Finkbeiner et al. 2012). This was especially the case for *Heliconius* white and yellow wing colours in the UV channel, which might act in intraspecific communication (Briscoe et al. 2010; Bybee et al. 2012). There is evidence that distinct UV colour signals are being transmitted between co-mimics, which may reduce costs of mating confusion (Dell’Aglia et al. 2018). Similarly, in two species of newt, belly colour is distinct in the UV range and females often made mistakes choosing the wrong males in the absence of UV light (Secondi and Théry 2014).

In this context, the duplicate genes encoding two distinct visual pigments with sensitivity peaks in the UV range in *H. erato* females offer the potential for enhanced spectral discrimination in light environments and time of the day where UV is more prominent. A UV2-UV1 opponent channel was proposed by Bybee et al. (2012), who showed that this receptor combination would have lower error rates for discrimination between *Heliconius* and *Dryas* yellows. There is no direct evidence for such a mechanism yet, but the fact that males and females show differences in the expression of the two UV proteins suggests that UV2-UV1 contrasts could be an important opponent channel for a female specific behaviour, perhaps mate recognition or host plant finding (McCulloch et al. 2016, 2017). There is similarly no direct evidence for a Red-Green opponent channel, although the presence of red filters in their eyes means that this is a possibility (McCulloch et al. 2016). Differences between species in Red-Green channel activity for red colouration might have a role in mate recognition since *Heliconius* tend to be attracted to red (Merrill et al. 2011).

The sensory drive hypothesis describes evolutionary relationships among visual systems, conditions of the light environment and mating preferences (Endler and Basolo 1998). *Heliconius* mating preference is highly linked to colour and in *H. melpomene*, the gene responsible for red colour pattern is genetically linked to the preference for the same pattern (Jiggins et al. 2001; Naisbit et al. 2001; Merrill et al. 2011). Visual sensitivity data used here is only from *H. erato* whereas it might differ for other species in their visual systems and perhaps match with colour preference or habitat (Frentiu et al. 2007; Briscoe et al. 2010; McCulloch et al. 2017). In addition, mating behaviour might benefit from some habitats in maximizing conspicuousness, such as in tropical dwelling birds and wire-tailed manakins which visual contrast is increased during display by habitat choice (Endler and Théry 1996; Heindl and Winkler 2003). Nevertheless, our results suggest that selection for conspicuousness in the preferred habitat could explain in part the divergence in colour pattern in these species.

Conclusion

In conclusion, the transmission of *Heliconius* warning signals varies due to light environment to a much greater degree through their own visual system, but to a smaller degree through avian predator vision. Selection for signal detectability under different habitat conditions is a mechanism that is proposed to lead to evolution of signal diversity, as seen in species of *Anolis* lizards that occupy habitats that match their visual system and signal design (Leal and Fleishman 2002), in species of warblers which different cone opsin gene expression correlate with sexual selection and habitat use (Bloch 2015) and also colour patterns of guppies are more conspicuous to guppies at the times and places of courtship and relatively less conspicuous at times and places of predator risk (Endler 1991). *Heliconius* butterfly warning colours are highly contrasting against the forest

background and stable through time and habitat in terms of predator avoidance but also conspicuous to attract the attention of conspecifics. However, more extensive studies considering spectral sensitivities of different *Heliconius* species and their responses to environmental changes in their signal visibility are needed to confirm the conspicuousness to mates. Opponent channel colour contrasts can predict behaviour of perceivers, however, additional behavioural experiments on how light environment influences prey detectability, such with poison frogs (Rojas et al. 2014), are necessary to verify our results.

Funding

This study was supported by Smithsonian Tropical Research Institute, Cambridge Trust, and CAPES Brazil (9423/11-7) to D.D.D, by European Research Council (Speciation Genetics 339873) to C.D.J., and by Biotechnology and Biological Sciences Research Council David Phillips Research Fellowship (BB/G022887/1) to M.S.

Author contributions

D.D.D. collected all the data, analysed, and wrote the manuscript; J.T. developed digital image methodology and data analysis; W.O.M, M.S, and C.D.J conceived the study, edited, and wrote the manuscript. We have no conflict of interest to declare.

References

- Arenas, L. M., J. Troscianko, and M. Stevens. 2014. Color contrast and stability as key elements for effective warning signals. *Front. Ecol. Evol.* 2:1–12.
- Arenas, L. M., D. Walter, and M. Stevens. 2015. Signal honesty and predation risk among a closely related group of aposematic species. *Sci. Rep.* 5:11021.
- Aronsson, M., and G. Gamberale-Stille. 2009. Importance of internal pattern contrast and contrast against the background in aposematic signals. *Behav. Ecol.* 20:141.
- Benson, W. W. 1972. Natural Selection for Mullerian Mimicry in *Heliconius erato* in Costa Rica. *Science* 176:936–939.
- Bloch, N. I. 2015. Evolution of opsin expression in birds driven by sexual selection and habitat. *Proc. R. Soc. B* 282:20142321.
- Blount, J. D., M. P. Speed, G. D. Ruxton, and P. A. Stephens. 2009. Warning displays may function as honest signals of toxicity. *Proc. R. Soc. Lond. B* 276:871–877.
- Boughman, J. W. 2002. How sensory drive can promote speciation. *Trends Ecol. Evol.* 17:571–577.
- Briscoe, A. D. 2008. Reconstructing the ancestral butterfly eye: focus on the opsins. *J. Exp. Biol.* 211:1805–1813.
- Briscoe, A. D., S. M. Bybee, G. D. Bernard, Y. Furong, M. P. Sison-Mangus, R. D. Reed, A. D. Warren, J. Llorente-bousquets, and C. Chiao. 2010. Positive selection of a duplicated UV-sensitive visual pigment coincides with wing pigment evolution in *Heliconius* butterflies. *PNAS* 107:3628–3633.
- Buskirk, W. H., G. V. N. Powell, J. F. Wittenberger, R. E. Buskirk, and T. U. Powell. 1972. Interspecific Bird Flocks in Tropical Highland Panama. *Am. Ornithol. Union* 89:612–624.
- Bybee, S. M., F. Yuan, M. D. Ramstetter, J. Llorente-Bousquets, R. D. Reed, D. Osorio, and A. D. Briscoe. 2012. UV Photoreceptors and UV-Yellow Wing Pigments in *Heliconius* Butterflies Allow a Color Signal to Serve both Mimicry and Intraspecific Communication. *Am. Soc. Nat.* 179:38–51.
- Chittka, L. 1996. Optimal Sets of Color Receptors and Color Opponent Systems for Coding of Natural Objects in Insect Vision. *J. Theor. Biol.* 181:179–196.
- Chittka, L., W. Beier, H. Hertel, E. Steinmann, and R. Menzel. 1992. Opponent colour coding is a universal strategy to evaluate the photoreceptor inputs in Hymenoptera. *J. Comp. Physiol. A* 170:545–563.

528 Cortesi, F., and K. L. Cheney. 2010. Conspicuousness is correlated with toxicity in marine
529 opisthobranchs. *J. Evol. Biol.* 23:1509–1518.

530 Dell’Aglia, D. D., M. Stevens, and C. D. Jiggins. 2016. Avoidance of an aposematically
531 coloured butterfly by wild birds in a tropical forest. *Ecol. Entomol.*, doi:
532 10.1111/een.12335.

533 Dell’Aglia, D. D., J. Troscianko, W. O. McMillan, M. Stevens, and C. D. Jiggins. 2018.
534 The appearance of mimetic *Heliconius* butterflies to predators and conspecifics.
535 *Evolution* 72:2156–2166.

536 Douglas, J. M., T. W. Cronin, T. Chiou, and N. J. Dominy. 2007. Light habitats and the
537 role of polarized iridescence in the sensory ecology of neotropical nymphalid
538 butterflies (Lepidoptera: Nymphalidae). *J. Exp. Biol.* 210:788–799.

539 Elias, M., Z. Gompert, C. D. Jiggins, and K. Willmott. 2008. Mutualistic Interactions
540 Drive Ecological Niche Convergence in a Diverse Butterfly Community. *PLoS*
541 *Biol.* 6:e300.

542 Endler, J. A. 1978. A predator’s view of animal color patterns. Pp. 319 – 364 *in*
543 *Evolutionary Biology*.

544 Endler, J. A. 1988. Frequency-dependent predation, crypsis and aposematic coloration.
545 *Philos. Trans. R. Soc. B Biol. Sci.* 319:505–523.

546 Endler, J. A. 1980. Natural Selection on Color Patterns in *Poecilia reticulata*. *Evolution*
547 34:76–91.

548 Endler, J. A. 1992. Signals, Signal Conditions, and the Direction of Evolution. *Am. Nat.*
549 139:S125–S153.

550 Endler, J. A. 1993. The color of light in forests and its implications. *Ecol. Monogr.* 63:1–
551 27.

552 Endler, J. A. 1991. Variation in the appearance of guppy color patterns to guppies and
553 their predators under different visual conditions. *Vision Res.* 31:587–608.

554 Endler, J. A., and A. L. Basolo. 1998. Sensory ecology, receiver biases and sexual
555 selection. *Trends Ecol. Evol.* 13:415–420.

556 Endler, J. A., and M. Théry. 1996. Interacting Effects of Lek Placement, Display
557 Behavior, Ambient Light, and Color Patterns in Three Neotropical Forest-
558 Dwelling Birds. *Am. Nat.* 148:421–452.

559 Estrada, C., and C. D. Jiggins. 2002. Patterns of pollen feeding and habitat preference
560 among *Heliconius* species. *Ecol. Entomol.* 27:448–456.

561 Finkbeiner, S. D. 2014. Communal roosting in heliconius butterflies (nymphalidae): roost
562 recruitment, establishment, fidelity, and resource use trends based on age and sex.
563 J. Lepidopterists Soc. 68:10–16.

564 Finkbeiner, S. D., A. D. Briscoe, and R. D. Reed. 2012. The benefit of being a social
565 butterfly: communal roosting deters predation. Proc. R. Soc. B 279:2769–2776.

566 Fleishman, L. J., M. Bowman, D. Saunders, W. E. Miller, M. J. Rury, and E. R. Loew.
567 1997. The visual ecology of Puerto Rican anoline lizards: Habitat light and
568 spectral sensitivity. J. Comp. Physiol. A 181:446–460.

569 Frentiu, F. D., G. D. Bernard, M. P. Sison-Mangus, A. Van Zandt Brower, and A. D.
570 Briscoe. 2007. Gene duplication is an evolutionary mechanism for expanding
571 spectral diversity in the long-wavelength photopigments of butterflies. Mol. Biol.
572 Evol. 24:2016–2028.

573 Guilford, T., and M. S. Dawkins. 1991. Receiver psychology and the evolution of animal
574 signals. Anim. Behav. 42:1–14.

575 Ham, A. D., E. Ihalainen, L. Lindström, and J. Mappes. 2006. Does colour matter? The
576 importance of colour in avoidance learning, memorability and generalisation.
577 Behav. Ecol. Sociobiol. 60:482–491.

578 Hämäläinen, L., J. Valkonen, J. Mappes, and B. Rojas. 2015. Visual illusions in predator–
579 prey interactions: birds find moving patterned prey harder to catch. Anim. Cogn.
580 18:1059–1068.

581 Hart, N. S. 2002. Vision in the peafowl (Aves: Pavo cristatus). J. Exp. Biol. 205:3925–
582 3935.

583 Hart, N. S., J. C. Partridge, I. C. Cuthill, and A. T. D. Bennett. 2000. Visual pigments, oil
584 droplets, ocular media and cone photoreceptor distribution in two species of
585 passerine bird: the blue tit (Parus caeruleus L.) and the blackbird (Turdus merula
586 L.). J. Comp. Physiol. A 186:375–387.

587 Heindl, M., and H. Winkler. 2003. Interacting effects of ambient light and plumage color
588 patterns in displaying wire-tailed manakins (Aves, Pipridae). Behav. Ecol.
589 Sociobiol. 53:153–162.

590 Jiggins, C. D. 2008. Ecological Speciation in Mimetic Butterflies. BioScience 58:541–
591 548.

592 Jiggins, C. D., R. E. Naisbit, R. L. Coe, and J. Mallet. 2001. Reproductive isolation caused
593 by colour pattern mimicry. Nature 411:302–305.

594 Joron, M., and J. Mallet. 1998. Diversity in mimicry: paradox or paradigm? Trends Ecol.
595 Evol. 13:461–466.

596 Kelber, A. 1999. Ovipositing butterflies use a red receptor to see green. J. Exp. Biol.
597 202:2619–2630.

598 Kelber, A., M. Vorobyev, and D. Osorio. 2003. Animal colour vision – behavioural tests
599 and physiological concepts. Biol. Rev. 78:81–118.

600 Kronforst, M. R., L. G. Young, D. D. Kapan, C. McNeely, R. J. O’Neill, and L. E. Gilbert.
601 2006. Linkage of butterfly mate preference and wing color preference cue at the
602 genomic location of wingless. PNAS 103:6575–6580.

603 Leal, M., and L. J. Fleishman. 2002. Evidence for habitat partitioning based on adaptation
604 to environmental light in a pair of sympatric lizard species. Proc. R. Soc. Lond. B
605 269:351–359.

606 Lovell, P. G., D. J. Tolhurst, C. A. Párraga, R. Baddeley, U. Leonards, J. Troscianko, and
607 T. S. Troscianko. 2005. Stability of the color-opponent signals under changes of
608 illuminant in natural scenes. J. Opt. Soc. Am. 22:2060–2071.

609 Mallet, J. 1986. Gregarious roosting and home range in *Heliconius* butterflies. Natl.
610 Geogr. Res. 2:198–215.

611 Mallet, J., and L. E. Gilbert. 1995. Why are there so many mimicry rings? Correlations
612 between habitat, behaviour and mimicry in *Heliconius* butterflies. Biol. J. Linn.
613 Soc. 55:159–180.

614 Mallet, J., W. O. Mcmillan, and C. D. Jiggins. 1998. Estimating the Mating Behavior of
615 a Pair of Hybridizing *Heliconius* Species in the Wild. Evolution 52:503–510.

616 McCulloch, K. J., D. Osorio, and A. D. Briscoe. 2016. Sexual dimorphism in the
617 compound eye of *Heliconius erato*: a nymphalid butterfly with at least five
618 spectral classes of photoreceptor. J. Exp. Biol. 219:2377–2387.

619 McCulloch, K. J., F. Yuan, Y. Zhen, M. L. Aardema, G. Smith, J. Llorente-Bousquets, P.
620 Andolfatto, and A. D. Briscoe. 2017. Sexual Dimorphism and Retinal Mosaic
621 Diversification following the Evolution of a Violet Receptor in Butterflies. Mol.
622 Biol. Evol. 34:2271–2284.

623 Merrill, R. M., R. E. Naisbit, J. Mallet, and C. D. Jiggins. 2013. Ecological and genetic
624 factors influencing the transition between host-use strategies in sympatric
625 *Heliconius* butterflies. J. Evol. Biol. 26:1959–1967.

626 Merrill, R. M., B. Van Schooten, J. A. Scott, and C. D. Jiggins. 2011. Pervasive genetic
627 associations between traits causing reproductive isolation in *Heliconius*
628 butterflies. *Proc. R. Soc. B* 278:511–518.

629 Müller, F. 1879. *Ituna* and *Thyridia*; a remarkable case of mimicry in butterflies. *Trans.*
630 *Linn. Soc. Lond.* 1879:xx–xxix.

631 Naisbit, R. E., C. D. Jiggins, and J. Mallet. 2001. Disruptive sexual selection against
632 hybrids contributes to speciation between *Heliconius cydno* and *Heliconius*
633 *melpomene*. *Proc. R. Soc. B* 268:1849–1854.

634 Nokelainen, O., R. H. Hegna, J. H. Reudler, C. Lindstedt, and J. Mappes. 2012. Trade-
635 off between warning signal efficacy and mating success in the wood tiger moth.
636 *Proc. R. Soc. B* 279:257–265.

637 Olofsson, M., A. Vallin, S. Jakobsson, and C. Wiklund. 2010. Marginal Eyespots on
638 Butterfly Wings Deflect Bird Attacks Under Low Light Intensities with UV
639 Wavelengths. *PLoS ONE* 5:e10798.

640 Osorio, D., M. Vorobyev, and C. D. Jones. 1999. Colour vision of domestic chicks. *J.*
641 *Exp. Biol.* 202:2951–2959.

642 Pegram, K. V., H. A. Han, and R. L. Rutowski. 2015. Warning Signal Efficacy: Assessing
643 the Effects of Color, Iridescence, and Time of Day in the Field. *Ethology* 121:1–
644 13.

645 Poulin, B., G. Lefebvre, R. Ibáñez, C. Jaramillo, C. Hernández, and A. S. Rand. 2001.
646 Avian predation upon lizards and frogs in a neotropical forest understorey. *J.*
647 *Trop. Ecol.* 17:21–40.

648 R Core Team. 2019. R: A Language and Environment for Statistical Computing. R
649 Foundation for Statistical Computing, Vienna, Austria.

650 Renoult, J. P., A. Kelber, and H. M. Schaefer. 2015. Colour spaces in ecology and
651 evolutionary biology. *Biol. Rev.*, doi: 10.1111/brv.12230.

652 Rojas, B., P. Rautiala, and J. Mappes. 2014. Differential detectability of polymorphic
653 warning signals under varying light environments. *Behav. Processes* 109:164–
654 172.

655 Secondi, J., and M. Théry. 2014. An ultraviolet signal generates a conflict between sexual
656 selection and species recognition in a newt. *Behav. Ecol. Sociobiol.* 68:1049–
657 1058.

658 Seehausen, O., Y. Terai, I. S. Magalhaes, K. L. Carleton, H. D. J. Mrosso, R. Miyagi, I.
659 van der Sluijs, M. V. Schneider, M. E. Maan, H. Tachida, H. Imai, and N. Okada.
660 2008. Speciation through sensory drive in cichlid fish. *Nature* 455:620–627.

661 Steiger, S. S., J. P. Kelley, W. W. Cochran, and M. Wikelski. 2009. Low Metabolism and
662 Inactive Lifestyle of a Tropical Rain Forest Bird Investigated via Heart-Rate
663 Telemetry. *Physiol. Biochem. Zool.* 82:580–589.

664 Stevens, M., C. A. Párraga, I. C. Cuthill, J. C. Partridge, and T. S. Troscianko. 2007.
665 Using digital photography to study animal coloration. *Biol. J. Linn. Soc.* 90:211–
666 237.

667 Stevens, M., and G. D. Ruxton. 2012. Linking the evolution and form of warning
668 coloration in nature. *Proc. R. Soc. B* 279:417–426.

669 Stevens, M., M. C. Stoddard, and J. P. Higham. 2009. Studying Primate Color: Towards
670 Visual System-dependent Methods. *Int. J. Primatol.* 30:893–917.

671 Svádová, K., A. Exnerová, P. Štys, E. Landová, J. Valenta, A. Fučíková, and R. Socha.
672 2009. Role of different colours of aposematic insects in learning, memory and
673 generalization of naïve bird predators. *Anim. Behav.* 77:327–336.

674 Sweeney, A., C. D. Jiggins, and S. Johnsen. 2003. Polarized light as a butterfly mating
675 signal. *Nature* 423:31–32.

676 Swihart, C. A. 1971. Colour discrimination by the butterfly, *Heliconius charitonius* Linn.
677 *Anim. Behav.* 19:156–164.

678 Swihart, S. L. 1972. The neural basis of colour vision in the butterfly, *Heliconius erato*.
679 *J. Insect Physiol.* 18:1015–1025.

680 Taylor, L. a., and K. J. McGraw. 2013. Male ornamental coloration improves courtship
681 success in a jumping spider, but only in the sun. *Behav. Ecol.* 24:955–967.

682 Troscianko, J., and M. Stevens. 2015. Image Calibration and Analysis Toolbox – a free
683 software suite for objectively measuring reflectance, colour and pattern. *Methods*
684 *Ecol. Evol.* 6:1320–1331.

685 Vorobyev, M., D. Osorio, A. T. D. Bennett, N. J. Marshall, and I. C. Cuthill. 1998.
686 Tetrachromacy, oil droplets and bird plumage colours. *J. Comp. Physiol. A*
687 183:621–633.

688 Whittle, P. 1994. The psychophysics of contrast brightness. Pp. 35–110 in A. L. Gilchrist,
689 ed. *Lightness, Brightness, and Transparency*. Lawrence Erlbaum Associates,
690 Hillsdale, NJ.

Zaccardi, G., A. Kelber, M. P. Sison-mangus, and A. D. Briscoe. 2006. Color discrimination in the red range with only one long-wavelength sensitive opsin. J. Exp. Biol. 209:1944–1955.

Supplementary Information

Table S1. Colour contrast differences for avian vision per opponent channel. General linear mixed models results using Satterthwaite approximations with random effects and Tukey's post-hoc (y ~ Colour + (1 | Individuals) + (1 | Vision:Habitat:Time)).

Opponent channel	Colour	N	Linear mixed model			Tukey's post-hoc		
			df	t value	p-value	Contrasts	z value	p-value
Red-Green								
	Red (intercept)	1176	16.3	34.59	< 2E-16	White - Red	-18.00	< 2E-16
	White		1160	-18.00	< 2E-16	Yellow - Red	-11.10	< 2E-16
	Yellow		1160	-11.10	< 2E-16	Yellow - White	8.26	3.33E-16
Blue-Yellow								
	Red (intercept)	1176	17.4	29.01	4.44E-16	White - Red	22.88	<2E-16
	White		49.3	22.88	< 2E-16	Yellow - Red	-11.98	<2E-16
	Yellow		1140	-11.98	< 2E-16	Yellow - White	-31.47	<2E-16
Blue-UV								
	Red (intercept)	1176	14.5	16.84	6.44E-11	White - Red	-6.11	2.85E-09
	White		1160	-6.11	1.34E-09	Yellow - Red	6.81	< 1E-09
	Yellow		1160	6.80	1.57E-11	Yellow - White	12.08	< 1E-09
Achromatic								
	Red (intercept)	1176	36.6	16.27	< 2E-16	White - Red	4.69	5.42E-06
	White		28.4	4.69	6.25E-05	Yellow - Red	41.89	< 1E-07
	Yellow		1136	41.88	< 2E-16	Yellow - White	12.42	< 1E-07

Table S2. Habitat contrast differences for avian vision per colour for the Red-Green opponent channel. General linear mixed models results using Satterthwaite approximations with random effects and Tukey's post-hoc ($y \sim \text{Habitat} + (1 | \text{Individuals}) + (1 | \text{Vision:Time})$).

Colour	Habitat	N	Linear mixed model			Tukey's post-hoc		
			df	t value	p-value	Contrasts	z value	p-value
Red	Border (intercept)	448	445	33.05	<2E-16	Forest - Border	-0.31	0.947
	Forest		445	-0.31	0.755	Open - Border	7.55	<1E-05
	Open		445	7.54	2.54E-13	Open - Forest	7.77	<1E-05
Yellow	Border (intercept)	448	9.2	53.37	7.90E-13	Forest - Border	10.77	<2E-16
	Forest		425	10.77	<2E-16	Open - Border	21.46	<2E-16
	Open		429.2	21.46	<2E-16	Open - Forest	14.63	<2E-16
White	Border (intercept)	280	9.05	24.88	1.21E-09	Forest - Border	-3.14	0.005
	Forest		262.9	-3.14	0.002	Open - Border	16.84	< 1E-04
	Open		265.3	16.84	<2E-16	Open - Forest	18.83	< 1E-04

Table S3. Time contrast differences for avian vision per colour for the Blue-Yellow opponent channel. General linear mixed models results using Satterthwaite approximations with random effects and Tukey's post-hoc ($y \sim \text{Time} + (1 | \text{Individuals}) + (1 | \text{Vision:Habitat})$).

Colour	Time	N	Linear mixed model			Tukey's post-hoc		
			df	t value	p-value	Contrasts	z value	p-value
Red	12PM (intercept)	448	9.4	17.31	2.03E-08	7AM - 12PM	4.44	2.37E-05
	7AM		367.3	4.44	1.18E-05	9AM - 12PM	4.17	8.14E-05
	9AM		367.3	4.17	3.73E-05	9AM - 7AM	-0.25	0.964
Yellow	12PM (intercept)	448	5.7	14.04	1.15E-05	7AM - 12PM	12.15	<1E-04
	7AM		436.6	12.15	2.00E-16	9AM - 12PM	14.24	<1E-04
	9AM		436.6	14.24	2.00E-16	9AM - 7AM	2.07	0.0969
White	12PM (intercept)	280	5.1	25.46	1.41E-06	7AM - 12PM	-1.59	0.250
	7AM		250.1	-1.58	0.113	9AM - 12PM	-0.08	0.996
	9AM		250.1	-0.08	0.936	9AM - 7AM	1.46	0.306

Table S4. Habitat contrast differences for avian vision for the achromatic opponent channel per colour. General linear mixed models results using Satterthwaite approximations with random effects and Tukey's post-hoc ($y \sim \text{Habitat} + (1 \mid \text{Individuals}) + (1 \mid \text{Vision:Time})$).

Colour	Habitat	N	Linear mixed model			Tukey's post-hoc		
			df	t value	p-value	Contrasts	z value	p-value
Red	Border (intercept)	448	12.5	22.28	2.00E-11	Forest - Border	-3.50	0.001
	Forest	425		-3.51	0.001	Open - Border	-7.52	< 1E-04
	Open	429.8		-7.52	3.19E-13	Open - Forest	-5.30	< 1E-04
Yellow	Border (intercept)	448	14.4	24.45	3.89E-13	Forest - Border	-3.42	0.002
	Forest	424.8		-3.42	0.001	Open - Border	-5.56	< 1E-04
	Open	376.1		-5.56	5.07E-08	Open - Forest	-3.34	0.002
White	Border (intercept)	280	13.9	21.02	5.89E-12	Forest - Border	-2.41	0.041
	Forest	263		-2.41	0.017	Open - Border	-0.02	1.000
	Open	267.7		-0.02	0.981	Open - Forest	1.51	0.280

Table S5. Habitat contrast differences for *Heliconius* vision for yellow and white colours, per opponent channel and time. General linear mixed models results using Satterthwaite approximations with random effects and Tukey's post-hoc ($y \sim \text{Habitat} + (1 | \text{Individuals}) + (1 | \text{Side})$).

Colour	Channel	Time	Habitat	N	Linear mixed model			Tukey's post-hoc		
					df	t value	p-value	Contrasts	z value	p-value
Yellow	UV2-UV1	7AM	Border (intercept)	64	1	7.41	0.081			
			Forest		46	-23.1	<2e-16			
		9AM	Border (intercept)	64	1	6.56	0.090			
			Forest		61	-13.31	<2e-16			
		12PM	Border (intercept)	96	1	4.28	0.124	Forest - Border	2.20	0.071
			Forest		92	2.20	0.03	Open - Border	7.49	0.001
			Open		92	7.49	4.02E-11	Open - Forest	5.29	0.001
White	UV2-UV1	7AM	Border (intercept)	40	11.4	6.57	3.37E-05			
			Forest		29	2.62	0.014			
		Blue-UV2 7AM	Border (intercept)	40	13.3	4.52	0.001			
			Forest		29	6.12	1.14E-06			
		12AM	Border (intercept)	60	21.6	13.1	9.32E-12	Forest - Border	-4.11	1.24E-04
			Forest		48	-4.11	1.54E-04	Open - Border	-3.43	0.002
			Open		48	-3.43	0.001	Open - Forest	0.67	0.776

Table S6. Side of the wing contrast differences for *Heliconius* vision for the red colour, per opponent channel. General linear mixed models results using Satterthwaite approximations with random effects and Tukey's post-hoc ($y \sim \text{Side} + (1 \mid \text{Individuals}) + (1 \mid \text{Habitat:Time})$).

Opponent Channel	Side of the wing	N	Linear mixed model		
			df	t value	p-value
Red-Green	Dorsal (intercept)	224	6.2	9.78	5.59E-05
	Ventral		201	-40.04	< 2E-16
Blue-Yellow	Dorsal (intercept)	224	7.3	22.24	5.63E-08
	Ventral		201	-25.5	< 2E-16
Blue-UV2	Dorsal (intercept)	224	16.6	36.51	< 2E-16
	Ventral		201	-34.61	< 2E-16
UV2-UV1	Dorsal (intercept)	224	8	17.89	8.76E-08
	Ventral		216	2.84	0.0048