Title:

Introducing Rhythmic Sinusoidal Amplitude-Modulated Auditory Stimuli with Multiple Message Frequency Coding for Fatigue Reduction in Normal Subjects: An EEG Study

Abbreviated title:

Rhythmic Multi-Message SAM Stimuli Reduces Fatigue

Author names and affiliations:

Elham Shamsi,^{1,2} Ahmadreza Keihani,^{1,2} Zahra Shirzhiyan,^{1,2} Morteza Farahi,^{1,2} Amin Mahnam,³ Mohsen Reza Heydari,⁴ and Amir Homayoun Jafari^{1,2}

¹Department of Medical Physics and Biomedical Engineering, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran, ²Research Center for Biomedical Technologies and Robotics, Tehran University of Medical Sciences, Tehran, Iran, ³Department of Biomedical Engineering, Faculty of Engineering, University of Isfahan, Isfahan, Iran, ⁴Department of Neurology, Faculty of Medicine, Baqiyatallah University of Medical Sciences, Tehran, Iran

Corresponding author: Amir Homayoun Jafari, Department of Medical Physics and Biomedical Engineering, School of Medicine, Tehran University of Medical Sciences, Poursina Street, Ghods Street, Keshavarz Boulevard, Tehran, Iran. E-mail: h_jafari@tums.ac.ir.

Number of total pages: 29

Number of figures: 7

Number of tables: 4

Number of words for Abstract: 250 (including “Abstract” and all the key words)

Number of words for Significance Statement: 110 (including “Significance Statement”)

Number of words for Introduction: 650 (including “Introduction” and citations)

Number of words for Discussion: 1372 (including “Discussion” and citations)

Conflict of interest: The authors declare no competing financial interests.

Acknowledgements: This work was supported by a grant from the deputy of the research review board and the ethics community of Tehran University of Medical Sciences (Grant NO: 30863)

Abstract

Many of the brain-computer interface (BCI) systems depend on the user's voluntary eye movements. However, voluntary eye movement is impaired in people with some neurological disorders. Since their auditory system is intact, auditory paradigms are getting more patronage from researchers. However, lack of appropriate signal-to-noise ratio in auditory BCI necessitates using long signal processing windows to achieve acceptable classification accuracy at the expense of losing information transfer rate. Because users eagerly listen to their interesting stimuli, the corresponding classification accuracy can be enhanced without lengthening of the signal processing windows. In this study, six sinusoidal amplitude-modulated auditory stimuli with multiple message frequency coding has been proposed to evaluate two hypotheses: 1) these novel stimuli provide high classification accuracies (greater than 70%), 2) the novel rhythmic stimuli set reduces the subjects' fatigue compared to its simple counterpart. We recorded EEG from nineteen normal subjects (twelve female). Five-fold cross-validated naïve Bayes classifier classified EEG signals with respect to power spectral density at message frequencies, Pearson's correlation coefficient between the responses and stimuli envelopes, canonical correlation coefficient between the responses and stimuli envelopes. Our results show that each stimuli set elicited highly discriminative responses according to all the features. Moreover, compared to the simple stimuli set, listening to the rhythmic stimuli set caused significantly lower subjects' fatigue. Thus, it is worthwhile to test these novel stimuli in a BCI experiment to enhance the number of commands and reduce the subjects' fatigue.

Key words: rhythm; amplitude modulation; multiple message frequency coding; classification; fatigue

Significance Statement

Auditory BCI users eagerly listen to the stimuli they are interested in. Thus, response classification accuracy may be enhanced without the need for trial lengthening. Since humans enjoy listening to rhythmic sounds, this study was carried out for introducing novel rhythmic sinusoidal amplitude-modulated auditory stimuli with multiple message frequency coding. Our results show that each stimuli set evoked reliably discriminative responses according to all the features, and rhythmic stimuli set caused significantly lower fatigue in subjects. Thus, it is worthwhile to test these novel stimuli in a BCI study to increase the number of commands (by N^N permutations of just N message frequencies) and reduce the subjects' fatigue.

Introduction

Brain-computer interfaces (BCI) makes muscle-independent communication between brain and computer possible. Thus, translating the user's intention to an external command (e.g., wheelchair control) comes true (Wolpaw et al., 2002). BCI can improve the quality of life for people with motor disabilities. Electroencephalogram (EEG) has been considered a reliable modality for using in BCI studies due to its noninvasiveness, good temporal resolution, easy implementation, and low cost (Wang et al., 2004; Hoffmann et al., 2008; Nijboer et al., 2008).

Patients with some neurological disorders, such as late-stage amyotrophic lateral sclerosis (ALS) and minimally conscious state (MCS), cannot perform voluntary eye movements or fixate their gaze. Moreover, daily usage of tactile BCI is hard because most people do not have tactile stimulators at home (Kaufmann et al., 2013). Thus, there has been an increasing interest towards auditory BCI (aBCI) (Hill et al., 2004; Kanoh et al., 2008; Nijboer et al., 2008; Furdea et al., 2009; Klobassa et al., 2009; Kübler et al., 2009; Halder et al., 2010; Schreuder et al., 2010; Higashi et al., 2011; Höhne et al., 2011; Kim et al., 2011; Schreuder et al., 2011; Kim et al., 2012; Lopez-Gordo et al., 2012; Käthner et al., 2013; Nakamura et al., 2013; Simon et al., 2014; Kleih et al., 2015; Zhou et al., 2016; Heo et al., 2017; Kaongoen and Jo, 2017), which uses auditory selective attention to influence event-related potentials (ERPs) and auditory steady-state responses (ASSRs). ASSR is chiefly evoked by listening to amplitude-modulated (AM) tones, and its spectrum has peaks at message frequency (f_m) (Picton et al., 2003; Lopez et al., 2009; Tanaka et al., 2013; Tanaka et al., 2015).

In aBCI, lengthening the processing window enhances the classification accuracy, but it reduces the speed (Lopez-Gordo et al., 2012). However, because the users eagerly listen to their interesting stimuli, the classification accuracy is enhanced (Zhou et al., 2016; Heo et al., 2017). Moreover, rhythmic stimulation modulates the intrinsic neural oscillatory characteristics (Herrmann et al., 2016). Rhythmic sinusoidal AM tones elicited EEG (Heo et

al., 2017; Shamsi et al., 2017) and MEG (Kuriki et al., 2013), but each of those stimuli had just one message frequency. Further, responses were not classified in (Kuriki et al., 2013) and subjects' fatigue were not evaluated in (Kuriki et al., 2013; Heo et al., 2017).

To our knowledge, there is not any research on AM sequences with multiple message frequencies. In this paper, six novel stimuli with multiple message frequency coding were introduced to test our hypotheses: 1) the resulting ASSRs are highly discriminative, and 2) listening to the novel rhythmic set reduces the subjects' fatigue compared to the simple set.

Materials and Methods

Subjects. Nineteen healthy (twelve female) volunteers took part in this study. They all participated in our previous study (Shamsi et al., 2017), too. Their age was in the range of 22-29 years (25.26 ± 2.05). All of them were right-handed according to Edinburgh Handedness Inventory (Oldfield, 1971) (0.75 ± 0.26). Participants reported no musical expertise. The instructions were explained to them. Subjects signed written informed consent form before conducting the experiments. All the procedures were approved by the ethics committee and the deputy of research review board of Tehran University of Medical Sciences.

Stimuli. In order to maintain consistency with other ASSR studies, double-sideband transmitted-carrier amplitude modulation with a modulation depth of 1 was used to generate the stimuli (to get more details, see (Lopez et al., 2009; Kuriki et al., 2013; Tanaka et al., 2013; Heo et al., 2017)):

$$s(t) = \sin(2\pi f_c t) (1 + \sin(2\pi f_m t)) \quad (1)$$

Where $s(t)$ stands for the stimulus signal. In addition, f_c and f_m are carrier and modulation (i.e., message) frequency, respectively. Two sets of stimuli were designed, each of which contained

three stimuli. All the stimuli had a duration of 180 s. Each stimulus comprised three f_m s. The stimuli sets is schematically represented in **Figure 1**. Using multiple message frequencies in each stimulus enriches its spectral content, because each f_m can elicit its corresponding peak in ASSR spectrum. In this way, different orderings and permutations of f_m s make it possible to produce various commands in aBCI. That is to say, using this coding, only N message frequencies can generate N^N permutations, which means N^N stimuli and N^N commands, whereas N^N message frequencies in single-message sinusoidal AM tones are required for generating the same number of stimuli and commands. This is important because there is limitation for message frequency selection in the sense that strong ASSRs were elicited by message frequencies in the range of [30-50] Hz (Picton et al., 1987), so using multiple message frequency coding facilitates the construction of stimuli corresponding to possible commands.

It is noteworthy that in this paper, whenever only a single carrier was present in the stimuli, those stimuli are called “simple”, while the stimuli containing more than one carrier are referred to as “rhythmic”. In other words, “rhythm” was generated using multiple carriers. For both sets of stimuli, f_m s were chosen to be among the (30, 35, 40) Hz. This is because consistent and robust ASSRs were elicited by message frequencies in the range of [30-50] Hz (Picton et al., 1987). Carrier frequencies were selected among the musical notes to be interesting for the subjects to listen to them. In this way, f_c s were members of the (262, 392, 494) Hz corresponding to “do”, “sol”, and “si” musical notes, respectively. For the rhythmic stimuli, the presence of each carrier was set to 0.5 s according to the best tempo sensitivity time interval (Drake and Botte, 1993). Frequency details are displayed in **Table 1**. Each stimuli set contained ascending, descending, and one of the possible zigzagging codings of message/carrier frequency. In other words, within each set, we constructed only three permutations (out of 27

B

Figure 1. Schematic representation of the stimuli sets. **A**, simple set. **B**, rhythmic set.

possible permutations). For instance, a pattern with 30-30-35 Hz (two identical f_{ms} at first and second portions of the triple pattern) or any similar permutation was not constructed. The reason is that we wanted to see the distinguishability that all of 30, 35 and 40 Hz within the proposed coding can provide in the corresponding ASSRs. Therefore, the presence of all three f_{ms} in the proposed coding was required in this study. All the stimuli were generated in MATLAB R2016b (MathWorks Inc., Natick, MA, USA). Sampling frequency for all the stimuli was 4410 Hz.

Table 1. Message and carrier frequency details in the stimuli sets

		Frequency (Hz)					
		f_{m1}	f_{m2}	f_{m3}	f_{c1}	f_{c2}	f_{c3}
Stimuli name							
Simple set	SA	30	35	40	262		-
	SB	40	35	30	494	-	-
	SC	40	30	35	392	-	-
Rhythmic set	RA	30	35	40	494	262	392
	RB	40	35	30	262	392	494
	RC	40	30	35	494	392	262

Task. In order to assess the subjects' level of some psychological factors (i.e., depression, stress, anxiety) through the week before the experiment, they were asked to fill DASS questionnaire (Lovibond and Lovibond, 1995) preceding the stimuli presentation. Also, the subjects filled questionnaire of current motivation (QCM) (Rheinberg et al., 2001; Vollmeyer and Rheinberg, 2006) before the beginning of the experiment. In this way, it was possible to measure their motivation and interest for participation, sense of challenge about the task, and anxiety they feel about the task. Then, the participants were requested to remain eyes-closed, still, and listen to the stimuli. For preventing from effects of the stimuli presentation order on the reported fatigue, the stimuli were presented in a random order for each participant. In other words, stimuli presentation order differed between the subjects. After listening to each stimulus, participants reported the amount of stimulus-induced fatigue that they experienced, as an integer number from 0 (minimum fatigue) to 10 (maximum fatigue) according to the visual analog scale (VAS) and they were given a short break of 60-120 seconds before presentation of the next stimulus. This procedure was performed for every stimulus. There were

Figure 2. Stimuli presentation procedure. “Stim” is the abbreviation of stimulus. The stimuli were presented in a random order. After listening to each stimulus, which was 180 s long, the subjects reported the level of stimulus-induced fatigue that they experienced, as an integer number from 0 (minimum fatigue) to 10 (maximum fatigue) according to VAS. Then, they were given a short break of 60-120 s before presentation of the next stimulus.

two reasons for this separate presentation: 1) we wanted to ensure that the stimuli in each set (i.e., simple, rhythmic) elicit sufficient inherently distinguishable responses in the brain, 2) we wanted to measure the amount of fatigue that each stimulus caused to each subject, so we had to present the stimuli separately (i.e., one by one). The stimuli presentation procedure is displayed in **Figure 2**. It is worth mentioning that all the previously mentioned psychological data were later used to explore whether there is a relationship between those factors and the fatigue level that subjects reported.

Experiment apparatus and recording. Insert earphones ER-3A (Etymotic Research, Elk Grove Village, IL) presented the stimuli to the subjects. For each stimulus, the volume was set according to equal loudness level contours at the standard ISO 226:2003.

Electrode placement was performed according to 10-20 international system. Active g.LADYbird electrodes were placed on Fz, Cz, T7 and T8. The reason for selecting Fz was that it is shown that Fz has the highest average amplitude of responses in subjects to whom a music is pleasant (Kayashima et al., 2017). Three other channels were consistently used in a number of previous ASSR studies (Lopez et al., 2009; Higashi et al., 2011; Kim et al., 2011; Heo et al., 2017; Shamsi et al., 2017). According to (Heo et al., 2017; Shamsi et al., 2017), right earlobe and Fpz were considered as the reference and ground, respectively. EEG was recorded by g.USBamp (g.tec Medical Engineering GmbH, Austria) at

a sampling frequency of 4800 Hz. Online filters consisted of a bandpass with a bandwidth of [0.5-2000] Hz, and a notch with a center frequency of 50 Hz.

Signal analysis. Firstly, in order to detect ASSR, we explored that whether the amplitude spectrum at f_m was larger than the mean+3×standard deviation (SD) of the amplitude spectrum at frequencies in the range of (f_m-1 to f_m-5) and (f_m+1 to f_m+5) (Tanaka et al., 2015). Then, prominent features (i.e., power spectral density (PSD), Pearson's correlation coefficient (PCC), canonical correlation coefficient (CCC)) were extracted. Feature extraction and classification were performed across every 20-s segment of the EEGs to be consistent with the relevant literature and practical applications (Kim et al., 2011; Heo et al., 2017). All the analyses were carried out in MATLAB R2016b (MathWorks Inc., Natick, MA, USA) on a laptop, which had Intel® Core™ i7-2670QM CPU @ 2.20 GHz as its processor.

Power spectral density. Keeping in mind that PSD is a robust feature for analyzing ASSR, it was computed (using amplitude spectrum at f_m and its adjacent frequencies in the range of f_m-5 - f_m-1 and f_m+1 - f_m+5) according to the literature (Tanaka et al., 2013; Tanaka et al., 2015), as follows:

$$PSD(f_m) = \frac{\sum |X(f_m-1:f_m+1)|^2}{\sum (|X(f_m-5:f_m-1)|^2 + |X(f_m+1:f_m+5)|^2)} \quad (2)$$

Where $|X(f_m)|$ is the amplitude spectrum of the brain response at frequency of f_m .

Pearson correlation coefficient. To investigate the amount of correlation between each stimulus and its corresponding ASSR, Pearson's correlation coefficient was used, which, through this paper, will be referred to as "PCC". As previously mentioned, the spectrum of this response has a peak at the modulation frequency (f_m) (Tanaka et al., 2013; Tanaka et al., 2015), which is exactly the same as the fundamental frequency of the stimulus envelope. Thus, PCC

was calculated for investigating the amount of correlation between each stimulus` envelope (i.e., X) and the ASSR (i.e., Y) that stimulus elicited. It was calculated as follows:

$$\rho_{X,Y} = \frac{\text{cov}(X,Y)}{\sigma_X \sigma_Y} \quad (3)$$

Where $\text{cov}(X,Y)$, σ_X , and σ_Y are the covariance of the two signals, the standard deviation of X and the standard deviation of Y, respectively. $\rho_{X,Y}$ has a value within the interval of [-1, 1]. Obviously, if there is not any linear relationship between X and Y, it will be zero.

Canonical correlation coefficient. This method seeks for a pair of linear combinations for two signals in such a way that the correlation between two canonical signals being maximized. In this way, pairs having linear combinations with the most linear correlation are chosen in a way that the previously identified pairs are orthogonal to them. If the EEG is represented by X and the stimulus envelope is considered to be Y, their projection vectors are denoted by $\tilde{x} = w_x^T X$ and $\tilde{y} = w_y^T Y$, respectively. Solving the equation below, w_x and w_y can be obtained:

$$\max_{w_x, w_y} \rho = \frac{E(\tilde{x}\tilde{y}^T)}{\sqrt{E(\tilde{x}\tilde{x}^T)E(\tilde{y}\tilde{y}^T)}} \quad (4)$$

Where, ρ is called the canonical correlation coefficient.

Classification. There were two cases for the classification, one for the responses to simple stimuli set, and the other for those of the rhythmic stimuli set. That is to say, in this study, two three-class classification problems existed. Classification was conducted by means of five-fold cross-validated naïve Bayes classifier. The chosen classifier utilizes the total probability theorem and the Bayes theorem to estimate the posterior probability (i.e., the probability that the features of an observation belong to a particular class) for each class. Then, for each

observation, corresponding posterior probabilities are compared to each other and the most will be selected as the outcome of the classification. Naïve Bayes classifier performs classification on the assumption that features of each class have statistical independence, whereas sometimes this is not the case. This classifier, however, works well in practice (Hastie et al., 2009). Posterior probability was calculated as follows:

$$\hat{P}(k|X_1, \dots, X_P) = \frac{\pi(k) \prod_{j=1}^P P(X_j|k)}{\sum_{k=1}^M \pi(k) \prod_{j=1}^P P(X_j|k)} \quad (5)$$

Where, k is the class index, $X_1, \dots, X_P$ are the features for each observation, and $\pi(k)$ is the empirical prior probability of class k . It is worth mentioning that the hyperparameters of the naïve Bayes classifier for each training fold was determined by Bayesian optimization.

In order to evaluate the amount of classification performance, classification accuracy and Cohen's kappa value were computed. Classification accuracy was defined to be the number of correctly classified observations divided by the number of classified observations. According to (Billinger et al., 2012), Cohen's kappa value was calculated as follows:

$$p_e = \frac{\sum_{i=1}^M C_{i,:} C_{:,i}}{N^2} \quad (6)$$

$$\kappa = \frac{\text{Acc} - p_e}{1 - p_e} \quad (7)$$

Where, p_e is the chance level, $C_{i,:}$ is the i -th row of confusion matrix, $C_{:,i}$ is the i -th column of confusion matrix, M represents the number of classes, and N is the total number of classified observations. Besides, κ and Acc are the Cohen's kappa value and the classification accuracy, respectively.

Experimental Design and Statistical Analysis. In order to examine the effect of stimuli type (i.e., simple, rhythmic) and feature (i.e., PSD, PCC, CCC) on the classification performances, we used the classification performance measures (i.e., accuracy, Cohen`s kappa value) for all the subjects (12 female, 7 male) as the dependent variable. Features and stimuli types were the within-subjects factors. All these were carried out via within-subjects repeated measures

Figure 3. Signal recording and analyses procedure.

ANOVA. Mauchly`s test checked whether the sphericity assumption held. Further, Greenhouse-Geisser approximation corrected the degrees of freedom. We selected Tukey`s honest significant difference to perform *post hoc* comparisons.

In addition, Wilcoxon signed rank test was used to see whether the level of stimuli-induced fatigue corresponding to the two sets of stimuli differ significantly within the subjects. In this

statistical design, subjects' fatigue was the dependent variable and type of the stimuli sets (i.e., simple and rhythmic) were within-subjects factors.

To investigate whether there is a relationship between psychological factors, which were evaluated via the questionnaires, and the reported fatigue, Spearman's correlation test was performed. All the analyses were conducted in MATLAB R2016b (MathWorks Inc., Natick, MA, USA).

Moreover, signal recording and analyses procedure is illustrated in **Figure 3**.

Results

Response detection

For each stimulus, amplitude spectrum of its corresponding EEG was computed. We checked whether the amplitude spectrum at f_m was larger than the mean + 3 × standard deviation (SD) of the amplitude spectrum at frequencies in the range of ($f_m - 1$ to $f_m - 5$) and ($f_m + 1$ to $f_m + 5$) (Tanaka et al., 2015) to make sure that ASSR was appeared. For simple and rhythmic stimuli set, amplitude spectrum corresponding to one stimulus is denoted in **Figure 4**. as a representative, because all the ASSRs satisfied the aforementioned condition (Tanaka et al., 2015). Consequently, these newly designed auditory stimuli with multiple message frequency coding elicited robust ASSRs.

Behavioral results

After listening to each stimulus, participants reported the amount of fatigue that they experienced by listening to that stimulus. All the subjects reported their level of fatigue as an integer number in the range from 0 (minimum) to 10 (maximum) according VAS. In comparison to the simple stimuli, the rhythmic stimuli caused significantly lower fatigue ($p = 0.005$, Wilcoxon signed rank,). This confirms our second hypothesis (fatigue reduction using

Figure 4. Amplitude spectrum of the responses to **A**, simple set. **B**, rhythmic set.

the proposed novel rhythmic stimuli set). Boxplot representation of fatigue due to the stimuli in each set is illustrated in **Figure 5**. In addition, there was not any significant correlation between psychological factors and fatigue caused by listening to each stimulus. This is an indication of the fact that the subjects truly reported the fatigues that were chiefly caused by listening to the stimuli, regardless of their psychological factors. For each stimulus, scatter plot of the data that yielded strongest Spearman's correlation coefficient, along with the best monotone curve fitted to the data are denoted on **Figure 6**. In each case, the best-fit curve was obtained through nonlinear least squares method. Details of all the Spearman tests (Spearman's correlation coefficients and corresponding p -values) are illustrated **Table 2**.

Figure 5. Boxplot representation of fatigue caused by the stimuli (fatigue levels were integers from 0 (minimum fatigue) to 10 (maximum fatigue) according to VAS). Rhythmic set significantly reduced the subjects' fatigue ($p = 0.005$, Wilcoxon signed rank).

Figure 6. Scatter plots of the data that yielded strongest Spearman's correlation coefficient between the psychological factors and fatigue caused by each stimulus (each plot's name), along with the best monotone curve fitted to the data. There was not any significant correlation between psychological factors and fatigue caused by listening to each stimulus. In each plot, r represents Spearman's correlation coefficient, while P is the p obtained in the Spearman's correlation test.

Classification performance

To investigate whether the designed stimuli yield intrinsically discriminative responses, we performed classification (for features and classifier parameters used, see Materials and Methods). For both sets of stimuli, high classification accuracy and Cohen's kappa value (up to a maximum of 100% and 1, respectively) were obtained. There was no significant difference between the responses to the simple and rhythmic stimuli sets in terms of classification performance ($F(1,15) = 4.06$, $p = 0.062$, repeated measures ANOVA). Furthermore, there was not any significant difference between PSD, PCC, and CCC features in terms of classification performance ($F(2,30) = 1.21$, $p = 0.307$, repeated measures ANOVA). These indicate that the responses to the stimuli in each set are sufficiently discriminative. Further, the results show that all the extracted features were discriminant measures for the responses to each set. Group

307 **Table 2. Spearman`s correlation test (r: Spearman`s correlation coefficient, P: *p* obtained via the test).**

308 **There was not any significant correlation between the psychological factors and stimuli-caused fatigue**

	Challenge	Interest	Success Probability	Anxiety (QCM)	Depression	Anxiety (DASS)	Stress
Fatigue Caused by SA	r = -0.088 P = 0.721	r = -0.055 P = 0.823	r = 0.099 P = 0.685	r = 0.086 P = 0.726	r = -0.122 P = 0.619	r = -0.098 P = 0.691	r = -0.217 P = 0.372
Fatigue Caused by SB	r = 0.049 P = 0.843	r = 0.185 P = 0.449	r = 0.086 P = 0.726	r = 0.114 P = 0.643	r = 0.025 P = 0.920	r = 0.263 P = 0.276	r = 0.015 P = 0.952
Fatigue Caused by SC	r = 0.149 P = 0.543	r = 0.314 P = 0.191	r = 0.109 P = 0.658	r = 0.084 P = 0.736	r = -0.127 P = 0.603	r = 0.176 P = 0.470	r = -0.101 P = 0.680
Fatigue Caused by RA	r = -0.074 P = 0.765	r = -0.160 P = 0.512	r = -0.055 P = 0.823	r = 0.015 P = 0.951	r = 0.071 P = 0.772	r = 0.268 P = 0.267	r = -0.006 P = 0.981
Fatigue Caused by RB	r = -0.012 P = 0.962	r = -0.190 P = 0.435	r = -0.047 P = 0.848	r = 0.170 P = 0.488	r = 0.371 P = 0.118	r = 0.454 P = 0.051	r = 0.308 P = 0.200
Fatigue Caused by RC	r = 0.176 P = 0.472	r = -0.070 P = 0.776	r = -0.298 P = 0.216	r = 0.049 P = 0.844	r = 0.247 P = 0.308	r = 0.438 P = 0.061	r = 0.413 P = 0.079

309

310 **Table 3. Between-subjects classification performance for different stimuli sets and features**

	PSD		PCC		CCC	
	Simple	Rhythmic	Simple	Rhythmic	Simple	Rhythmic
Accuracy (%)	92.98	88.89	93.57	81.29	94.15	83.04
Cohen`s kappa value	0.90	0.83	0.90	0.72	0.91	0.75

311

312

313

314 **Figure 7. Top,** Group means of the classification accuracy for the three features (PSD, PCC, CCC) as a function
 315 of stimuli type (illustrated as Simple, Rhythmic). **Bottom,** Group means of the Cohen's kappa value for the three
 316 features (PSD, PCC, CCC) as a function of stimuli type (illustrated as Simple, Rhythmic).

317

318 **Table 4. A comparison between the results of this paper and those of some relevant ASSR studies**

Study	Subjects	Number of stimuli / Type	Segment (seconds)	Average classification accuracy (%)	Fatigue measurement
(Kim et al., 2011)	6 normal	2 / single-message AM tones	20	76.25	Not included
(Nakamura et al., 2013)	8 normal	2 / single-message AM speech sentence	9	78.6	Not included
(Kaongoen and Jo, 2017)	10 normal	2 / simple single-message AM tones	21	82.90	Not included
(Heo et al., 2017)	6 normal	2 / simple single-message AM tones	20	74	Not included
(Heo et al., 2017)	6 normal	2 / single-message AM natural sound carrier	20	87.67	Not included
(Heo et al., 2017)	6 normal	2 / single-message AM musical carrier	20	89.67	Not included
(Shamsi et al., 2017)	19 normal	3 / simple single-message sinusoidal AM tone	20	82.46	Median: 4
(Shamsi et al., 2017)	19 normal	3 / rhythmic single-message sinusoidal AM sequence	20	80.70	Median: 2
This work	19 normal	3 / simple multiple-message sinusoidal AM tone	20	PSD: 91.23 PCC: 97.66 CCC: 93.57	Median: 5
This work	19 normal	3 / rhythmic multiple-message sinusoidal AM sequence	20	PSD: 82.46 PCC: 89.48 CCC: 83.63	Median: 3

means of the classification accuracy and Cohen's kappa value for different conditions (i.e., features) as a function of the stimuli type (i.e., simple, rhythmic) are illustrated in **Figure 7**. The results show that all the average classification accuracies are well above 70%, which is the minimum acceptable classification accuracy in BCI systems. Thus, the responses to the stimuli in each set are highly discriminative. In other words, our first hypothesis was confirmed. To investigate whether there is generalizability in terms of response discrimination, we performed between-subjects classification for the responses to each set. Classification accuracy and Cohen's kappa value for between-subjects classification according to each feature are listed in **Table 3**.

Discussion

According to the facts that humans enjoy listening to rhythmic sounds (Zhou et al., 2016; Heo et al., 2017) and rhythmic stimulation influences the intrinsic neural oscillatory characteristics (Herrmann et al., 2016), it seems that utilizing rhythmic auditory stimuli in the experiments that aim to evoke and examine auditory responses in the brain reduces the subjects' fatigue. Thus, this study was carried out to test our two hypotheses 1) the ASSRs to the novel stimuli with multiple message frequency coding are highly discriminative, and 2) listening to the novel rhythmic stimuli set with multiple message frequency coding reduces the subjects' fatigue. All these were conducted to determine whether the stimuli introduced in this paper have enough feasibility (in terms of classification performance and subjects' fatigue) to be used in an aBCI. In some previous studies, rhythmic sinusoidal amplitude-modulated tones were used to elicit EEG (Heo et al., 2017; Shamsi et al., 2017) and MEG (Kuriki et al., 2013), but all of them utilized single message frequency. Further, response classification and subjects' fatigue evaluation were not conducted in (Kuriki et al., 2013). Although user acceptance evaluated in (Heo et al., 2017), user fatigue was not taken into account. Thus, both the stimuli sets designed in this study were novel in the sense of having multiple message frequency coding. In addition,

response classification and fatigue evaluation were carried out for these stimuli for the very first time. For a better insight into the novelty of this work, a comparison between the results of this paper and those of some relevant ASSR studies, which performed response classification, are illustrated in **Table 4**.

Robust peak in ASSR spectrum at message frequencies (corresponding to the envelope of the stimuli) is consistent with previous findings (Lopez et al., 2009; Kim et al., 2011; Nozaradan et al., 2012; Kuriki et al., 2013; Kaongoen and Jo, 2017) and confirms the notion that human's auditory system acts like an envelope detector, which can be an amplitude demodulator (Miyazaki et al., 2013). In addition, amplitudes of the responses to the rhythmic set were lower than those of the simple set. This may be due to the more complex structure of the rhythmic set, compared to that of the simple set. There is relevant supporting evidence that more complex stimuli elicited less amplitude, when compared to stimuli with simpler structure (Nakamura et al., 2013; Shamsi et al., 2017).

Moreover, the rhythmic stimuli set resulted in less fatigue in the subjects, compared to that of the simple stimuli set. This is in agreement with the findings in our previous study on the comparison between the fatigue levels that simple and rhythmic single-message sinusoidal AM stimuli can cause (Shamsi et al., 2017) and confirms our second hypothesis. In addition, the insignificant and infinitesimal correlation between the fatigue and the psychological factors can ensure us that the subjects truly reported the fatigue that was chiefly caused by listening to the stimuli, regardless of their psychological factors.

We were able to perform highly accurate, precise and reliable classification on within- and between-subjects responses without any artifact rejection. This shows that there was adequate inherent discrimination even at the raw signal level for the responses to each stimuli set. It can be seen from the within-subjects classification performance results that: 1) stimuli with multiple message frequencies generate highly distinguishing ASSRs, so they have the potential

be utilized in aBCI to increase the number of available commands, and therefore the information transfer rate, by means of multiple permutations of just a few message frequencies, which is accessible via the coding presented in this paper. In other words, fewer message frequencies (N in the proposed multiple message coding, compared to N^N in single-message SAM tones) can generate N^N commands in aBCI, 2) the rhythmic stimuli elicit discriminative responses, which are as distinct as that of the simple stimuli, 3) all the features (PSD, PCC, and CCC) are discriminant measures for the classification of the ASSR to the stimuli with multiple message frequencies. Also, high amounts of between-subjects classification performance indicates that the ASSRs to the stimuli in each set were reliably distinct and generalizable. Furthermore, all the average classification accuracies were far above 70%, which is sufficient for a BCI system. In other words, our first hypothesis was confirmed. Thus, the stimuli designed in this paper have the adequate potential to be corresponding to several different commands and generate distinct responses in BCI systems.

The average classification performances obtained in this study outperformed previous studies, which utilized single-message AM tones (Lopez et al., 2009; Kim et al., 2011; Heo et al., 2017; Kaongoen and Jo, 2017; Shamsi et al., 2017) and single-message AM sentences (Nakamura et al., 2013). Particularly, the average classification performances obtained for our simple set was higher than those of a research, which used single-message AM natural sound carriers (Heo et al., 2017). However, the average classification performances for our rhythmic set was a bit lower than those of a study, which made use of single-message AM instrumental music carriers (Heo et al., 2017). It is worth mentioning that in the current study, each stimulus was presented separately, while the stimuli in most of the compared studies were played simultaneously, which may decrease their classification performance. In other words, for each subject of the current study, each stimulus played, the fatigue reported by the subject was written down, and another stimulus was presented, and so on. There were two reasons for this:

1) we wanted to ensure that whether the stimuli in each introduced set evoke adequate inherently distinguishable responses in the brain, 2) we wanted to measure the amount of fatigue that each stimulus caused to each subject, so we had to present the stimuli separately (i.e., one by one). Although simultaneous presentation of the stimuli is required in BCI paradigms, this is not the case in our study, which is not a BCI paradigm. This study is a preliminary step that investigated the feasibility of utilizing the proposed stimuli in a BCI paradigm. For this purpose, the amount of inherent distinguishability between the responses in each set, along with subjects' fatigue were measured through the separate presentation. Therefore, our purpose required this kind of stimuli presentation. However, in simultaneous presentation of the stimuli, f_c coding in the rhythmic set will help the users to focus on and discriminate between the stimuli. This implies that the classification performance of the responses in the simultaneous stimuli presentation would not be too different from our results, which are obtained via separate stimuli presentation.

The results showed that stimuli in each set have sufficient inherent discrimination to the extent that it is worthwhile to use these novel auditory stimuli with multiple message frequency coding in a BCI experiment. If we are asked to choose one of our proposed stimuli sets to be utilized in BCI studies, the choice will be the "rhythmic set". The reason is listening to the rhythmic set reduced the subjects' fatigue and the brain responses to the rhythmic set were classified via a common classifier, with a high performance close to the simple set, so this set will be able to increase the number of possible commands by permutation of the message frequencies of its stimuli.

Sinusoidal amplitude-modulated tones are helpful in studies concerning encoding of envelope and periodicity in human's auditory system. Moreover, they can be used in ASSR-based BCI systems. Therefore, exploring sinusoidal AM tone-evoked ASSR is chiefly important. In this paper, each stimuli set contained ascending, descending and one of the

possible zigzagging codings of message/carrier frequency. For future work, it is suggested to explore ASSR to other possible zigzagging permutations of message/carrier frequency, and make a comparison between the responses to stimuli with different coding types (ascending, descending and zigzagging) and frequency effects. Also, testing auditory stimuli constructed with other modulations (e.g., frequency modulation (FM), pulse width modulation (PWM), etc.) would be valuable. Further, conducting the experiment performed in this paper on completely locked-in state syndrome (CLIS) patients is proposed for future work to see whether they are useful for those individuals. In this study, we aimed at exploring the responses in common domains (e.g., time and frequency). However, nonlinear and/or time-frequency analyses can be performed and compared in future studies.

References

- Billinger M, Daly I, Kaiser V, Jin J, Allison BZ, Müller-Putz GR, Brunner C (2012) Is it significant? Guidelines for reporting BCI performance. In: Towards Practical Brain-Computer Interfaces, pp 333-354: Springer.
- Drake C, Botte M-C (1993) Tempo sensitivity in auditory sequences: Evidence for a multiple-look model. *Attention, Perception, & Psychophysics* 54:277-286.
- Furdea A, Halder S, Krusienski D, Bross D, Nijboer F, Birbaumer N, Kübler A (2009) An auditory oddball (P300) spelling system for brain-computer interfaces. *Psychophysiology* 46:617-625.
- Halder S, Rea M, Andreoni R, Nijboer F, Hammer E, Kleih S, Birbaumer N, Kübler A (2010) An auditory oddball brain-computer interface for binary choices. *Clinical Neurophysiology* 121:516-523.
- Hastie T, Tibshirani R, Friedman J (2009) *The Elements of Statistical Learning*, 2nd Edition: Springer-Verlag New York.

- Heo J, Baek HJ, Hong S, Chang MH, Lee JS, Park KS (2017) Music and natural sounds in an auditory steady-state response based brain-computer interface to increase user acceptance. *Computers in Biology and Medicine* 84:45-52.
- Herrmann CS, Murray MM, Ionta S, Hutt A, Lefebvre J (2016) Shaping Intrinsic Neural Oscillations with Periodic Stimulation. *The Journal of Neuroscience* 36:5328-5337.
- Higashi H, Rutkowski TM, Washizawa Y, Cichocki A, Tanaka T (2011) EEG auditory steady state responses classification for the novel BCI. In: 2011 Annual International Conference of the IEEE Engineering in Medicine and Biology Society, pp 4576-4579.
- Hill NJ, Lal TN, Bierig K, Birbaumer N, Schölkopf B (2004) An auditory paradigm for brain-computer interfaces. In: *Proceedings of the 17th International Conference on Neural Information Processing Systems*, pp 569-576. Vancouver, British Columbia, Canada: MIT Press.
- Hoffmann U, Vesin J-M, Ebrahimi T, Diserens K (2008) An efficient P300-based brain-computer interface for disabled subjects. *Journal of Neuroscience Methods* 167:115-125.
- Höhne J, Schreuder M, Blankertz B, Tangermann M (2011) A Novel 9-Class Auditory ERP Paradigm Driving a Predictive Text Entry System. *Frontiers in Neuroscience* 5:99.
- Kanoh Si, Miyamoto K-i, Yoshinobu T (2008) A brain-computer interface (BCI) system based on auditory stream segregation. In: *Engineering in Medicine and Biology Society, 2008. EMBS 2008. 30th Annual International Conference of the IEEE*, pp 642-645: IEEE.
- Kaongoen N, Jo S (2017) A novel hybrid auditory BCI paradigm combining ASSR and P300. *Journal of Neuroscience Methods* 279:44-51.

468 Käthner I, Ruf CA, Pasqualotto E, Braun C, Birbaumer N, Halder S (2013) A portable auditory
469 P300 brain–computer interface with directional cues. *Clinical Neurophysiology*
470 124:327-338.

471 Kaufmann T, Holz EM, Kübler A (2013) Comparison of tactile, auditory, and visual modality
472 for brain-computer interface use: a case study with a patient in the locked-in state.
473 *Frontiers in Neuroscience* 7:129.

474 Kayashima Y, Yamamuro K, Makinodan M, Nakanishi Y, Wanaka A, Kishimoto T (2017)
475 Effects of Canon chord progression on brain activity and motivation are dependent on
476 subjective feelings, not the chord progression per se. *Neuropsychiatric Disease and*
477 *Treatment* 13:1499-1508.

478 Kim D-W, Lee J-C, Park Y-M, Kim I-Y, Im C-H (2012) Auditory brain-computer interfaces
479 (BCIs) and their practical applications. *Biomedical Engineering Letters* 2:13-17.

480 Kim D-W, Hwang H-J, Lim J-H, Lee Y-H, Jung K-Y, Im C-H (2011) Classification of selective
481 attention to auditory stimuli: toward vision-free brain–computer interfacing. *Journal of*
482 *neuroscience methods* 197:180-185.

483 Kleih SC, Herweg A, Kaufmann T, Staiger-Sälzer P, Gerstner N, Kübler A (2015) The WIN-
484 speller: a new intuitive auditory brain-computer interface spelling application. *Frontiers*
485 *in Neuroscience* 9:346.

486 Klobassa DS, Vaughan T, Brunner P, Schwartz N, Wolpaw J, Neuper C, Sellers E (2009)
487 Toward a high-throughput auditory P300-based brain–computer interface. *Clinical*
488 *Neurophysiology* 120:1252-1261.

489 Kübler A, Furdea A, Halder S, Hammer EM, Nijboer F, Kotchoubey B (2009) A brain–
490 computer interface controlled auditory event-related potential (P300) spelling system
491 for locked-in patients. *Annals of the New York Academy of Sciences* 1157:90-100.

492 Kuriki S, Kobayashi Y, Kobayashi T, Tanaka K, Uchikawa Y (2013) Steady-state MEG
493 responses elicited by a sequence of amplitude-modulated short tones of different carrier
494 frequencies. *Hearing Research* 296:25-35.

495 Lopez-Gordo MA, Pelayo F, Prieto A, Fernandez E (2012) An Auditory Brain-Computer
496 Interface with Accuracy Prediction. *International Journal of Neural Systems*
497 22:1250009.

498 Lopez M-A, Pomares H, Pelayo F, Urquiza J, Perez J (2009) Evidences of cognitive effects
499 over auditory steady-state responses by means of artificial neural networks and its use
500 in brain-computer interfaces. *Neurocomputing* 72:3617-3623.

501 Lovibond SH, Lovibond PF (1995) Manual for the depression anxiety stress scales. Sydney,
502 N.S.W.: Psychology Foundation of Australia.

503 Miyazaki T, Thompson J, Fujioka T, Ross B (2013) Sound envelope encoding in the auditory
504 cortex revealed by neuromagnetic responses in the theta to gamma frequency bands.
505 *Brain Research* 1506:64-75.

506 Nakamura T, Namba H, Matsumoto T (2013) Classification of auditory steady-state responses
507 to speech data. In: 2013 6th International IEEE/EMBS Conference on Neural
508 Engineering (NER), pp 1025-1028.

509 Nijboer F, Furdea A, Gunst I, Mellinger J, McFarland DJ, Birbaumer N, Kübler A (2008) An
510 auditory brain-computer interface (BCI). *Journal of Neuroscience Methods* 167:43-50.

511 Nozaradan S, Peretz I, Mouraux A (2012) Selective neuronal entrainment to the beat and meter
512 embedded in a musical rhythm. *Journal of Neuroscience* 32:17572-17581.

513 Oldfield RC (1971) The assessment and analysis of handedness: the Edinburgh inventory.
514 *Neuropsychologia* 9:97-113.

515 Picton TW, John MS, Dimitrijevic A, Purcell D (2003) Human auditory steady-state responses:
516 respuestas auditivas de estado estable en humanos. International Journal of Audiology
517 42:177-219.

518 Picton TW, Skinner CR, Champagne SC, Kellett AJC, Maiste AC (1987) Potentials evoked by
519 the sinusoidal modulation of the amplitude or frequency of a tone. The Journal of the
520 Acoustical Society of America 82:165-178.

521 Rheinberg F, Vollmeyer R, Burns BD (2001) FAM: Ein Fragebogen zur Erfassung aktueller
522 Motivation in Lern-und Leistungssituationen (Langversion, 2001). Diagnostica 2:57-
523 66.

524 Schreuder M, Blankertz B, Tangermann M (2010) A new auditory multi-class brain-computer
525 interface paradigm: spatial hearing as an informative cue. PloS one 5:e9813.

526 Schreuder M, Rost T, Tangermann M (2011) Listen, You are Writing! Speeding up Online
527 Spelling with a Dynamic Auditory BCI. Frontiers in Neuroscience 5:112.

528 Shamsi E, Shirzhiyan Z, Keihani A, Farahi M, Mahnam A, Heydari MR, Jafari AH (2017)
529 Classification of the EEG Evoked by Auditory Stimuli with a Periodic Carrier
530 Frequency Coding in Order to Be Used in BCI Systems. Frontiers in Biomedical
531 Technologies 3:41-48.

532 Simon N, Käthner I, Ruf CA, Pasqualotto E, Kübler A, Halder S (2014) An auditory multiclass
533 brain-computer interface with natural stimuli: Usability evaluation with healthy
534 participants and a motor impaired end user. Frontiers in Human Neuroscience 8:1039.

535 Tanaka F, Matsubara A, Nishifuji S (2015) Evidence of stochastic resonance of auditory
536 steady-state response in electroencephalogram for brain machine interface. In:
537 Consumer Electronics (GCCE), 2015 IEEE 4th Global Conference on, pp 195-199:
538 IEEE.

- Tanaka K, Kuriki S, Nemoto I, Uchikawa Y (2013) Auditory Steady-State Responses in Magnetoencephalogram and Electroencephalogram: Phenomena, Mechanisms, and Applications. *Advanced Biomedical Engineering* 2:55-62.
- Vollmeyer R, Rheinberg F (2006) Motivational effects on self-regulated learning with different tasks. *Educational Psychology Review* 18:239-253.
- Wang T, Deng J, He B (2004) Classifying EEG-based motor imagery tasks by means of time–frequency synthesized spatial patterns. *Clinical Neurophysiology* 115:2744-2753.
- Wolpaw JR, Birbaumer N, McFarland DJ, Pfurtscheller G, Vaughan TM (2002) Brain–computer interfaces for communication and control. *Clinical Neurophysiology* 113:767-791.
- Zhou S, Allison BZ, Kübler A, Cichocki A, Wang X, Jin J (2016) Effects of Background Music on Objective and Subjective Performance Measures in an Auditory BCI. *Frontiers in Computational Neuroscience* 10:105.