Species-specific recognition of Sulfolobales mediated by UV-inducible pili and S-layer glycosylation patterns

Marleen van Wolferen¹, Asif Shajahan², Kristina Heinrich¹, Susanne Brenzinger³, Ian M. Black², Alexander Wagner¹, Ariane Briegel³, Parastoo Azadi², and Sonja-Verena Albers^{1*}

¹Molecular Biology of Archaea, Institute of Biology II - Microbiology, University of Freiburg, Schaezenlestrasse 1, 79104, Freiburg, Germany

²Complex Carbohydrate Research Center, The University of Georgia, 315 Riverbend Road, Athens, GA, 30602, USA.

³Institute of Biology, Leiden University, Sylviusweg 72, 2333 BE, Leiden, The Netherlands

*Correspondence to S.V. Albers, e-mail: sonja.albers@biologie.unifreiburg.de, tel.: +49(0)761 203 2630

Keywords: type IV pili, archaea, Sulfolobus, DNA exchange, glycosylation

Abstract

The UV-inducible pili system of Sulfolobales (Ups) mediates the formation of species-specific cellular aggregates. Within these aggregates, cells exchange DNA in order to repair DNA double strand breaks via homologous recombination. Substitution of the *S. acidocaldarius* pilin subunits UpsA and UpsB with their homologs from *Sulfolobus tokodaii* showed that these subunits facilitate species-specific aggregation. A region of low conservation within the UpsA homologs is primarily important for this specificity. Aggregation assays in the presence of different sugars showed the importance of *N*-glycosylation in the recognition process. In addition, the *N*-glycan decorating the S-layer of *S. tokodaii* is different from the one of *S. acidocaldarius*. Therefore, each *Sulfolobus* species seems to have developed a unique UpsA binding pocket and unique *N*-glycan composition to ensure aggregation and consequently also DNA exchange with cells from only the same species, which is essential for DNA repair by homologous recombination.

Importance

Type IV pili can be found on the cell surface of many archaea and bacteria where they play important roles in different processes. The Ups-pili from the crenarchaeal Sulfolobales species are essential in establishing species-specific mating partners, ensuring genome stability. With this work, we show that different *Sulfolobus* species have species-specific regions in their Ups-pilin subunits, which allow them to interact only with cells from the same species. Additionally, different *Sulfolobus* species all have unique S-layer *N*-glycosylation patterns. We propose that the unique features of each species allow the recognition of specific mating partners. This knowledge for the first time gives insights into the molecular basis of archaeal self-recognition.

42

43

Introduction

Type IV pili (T4P) are cell surface appendages that can be found on the cell-surfaces of many bacteria and archaea (1, 2). They have been implicated in motility, secretion, DNA transformation, adhesion to surfaces and the formation of intercellular associations (3, 4). In bacteria, many examples of T4P with cellular-binding properties have been described. The major pilin subunit PilE from *Neisseria* T4P, was shown to bind endothelial cells and hemagglutinate erythrocytes whereas the *Neisseria* minor pilin PilV is essential for adherence to host cells (5–10). Additionally, major pilin PilA from *Myxococcus xanthus* binds to self-produced exopolysaccharides, subsequent retraction of the T4P allows gliding motility and fruiting body formation (11, 12). T4P also form intercellular connections that are essential for conjugational exchange of DNA. For instance, PAPI-1 encoded T4P bring *Pseudomonas aeruginosa* cells in close proximity by binding to lipopolysaccharides of the recipient cells and thereby promote exchange of PAPI-1 DNA (13, 14).

In archaea, several gene clusters have been found to encode T4P-like structures (4, 15–19). The best-characterized archaeal T4P-like structure is the archaellum, which is essential for swimming motility (4, 19–21). However, little is known about the role and mode of action of archaeal non-archaellum T4P in attachment to biotic or abiotic surfaces. T4P from the thermophilic crenarchaeon *Sulfolobus acidocaldarius* (Aap: archaeal adhesive pili) and the euryarchaea *Haloferax volcanii* and *Methanococcus maripaludis* were shown to be involved in attachment to surfaces (22–27). However, their exact mode of binding has not been studied. Next to Aap-pili, Ups-pili can be found in Sulfolobales (UV inducible pili of Sulfolobales) (28–31). These T4P assemble upon treatment of the cells with UV-stress and other DNA double strand break inducing agents. They are crucial in cellular self-interactions thereby mediating the formation of species-specific cellular aggregates (32, 33). Within these

aggregates, cells are able to exchange chromosomal DNA using the Ced-system (Crenarchaeal exchange of DNA), suggesting a community based DNA repair system via homologous recombination (32, 34). Interestingly, the Ced-system was found to function independently of the Ups-pili, even though both systems are essential for DNA transport (35).

The *ups*-operon encodes two pilin subunits with a class III signal peptide: UpsA and UpsB (29). Deletion mutants of either *upsA* or *upsB* still form pili (though less and smaller), but do not aggregate after UV induction. The pilins are therefore both suggested to be major subunits forming mixed Ups-pili (31, 32). While the importance of Ups-pili in cellular recognition is known, the underlying molecular mechanism of the species-specific cellular aggregation of *Sulfolobus* species has not been determined.

In this study, we investigated the role of Ups-pili in species-specific aggregation on a molecular level. To this end, *in vivo* chimera mutants were constructed in which we exchanged (parts of) the genes encoding the pilin subunits UpsA and UpsB of *S. acidocaldarius* and *S. tokodaii*. By using these strains in aggregation assays and fluorescence in situ hybridization (FISH) experiments, we were able to assign a specific region of UpsA to be required for species-specific cell aggregation of archaeal cells. Furthermore, aggregation assays in the presence of different sugars suggested a role of *N*-glycosylation in cellular recognition. Glycan analysis on the thus far unstudied *S. tokodaii* S-layer showed a different *N*-glycan composition compared to that of other *Sulfolobus* species. Based on these experiments, we propose that a specific region of UpsA forms a binding site to bind species-specific *N*-glycan chains of S-layer components, thereby allowing species-specific cell aggregation and subsequent DNA exchange.

Results

The role of pilin subunits in species-specificity

To study the role of the Ups-pilin subunits (UpsA and UpsB) in species-specific recognition of *Sulfolobus* cells, a *S. acidocaldarius* strain was constructed in which the genomic region from the start codon of *upsA* until the stop codon of *upsB* was exchanged with the orthologous region from *S. tokodaii* (resulting in strain MW135; Figure 1A, Table 1). Upon UV-induction *S. acidocaldarius* MW135 was still found to produce Ups-pili (Figure S3), however, interestingly, it showed little to no cellular aggregation (Figure 1B). In order to test if this *S. acidocaldarius* Ups-hybrid strain was able to recognize and therefore aggregate with *S. tokodaii* cells, fluorescence in situ hybridisation (FISH) with species-specific probes was performed on mixed *S. acidocaldarius*/*S. tokodaii* strains after UV-induction. A positive control with a mixture of background strain *S. acidocaldarius* MW501 ($\Delta flaI/\Delta aapF$, a strain that does not produce archaella or Aap-pili; Table 1) and *S. tokodaii*, confirmed previously observed species-specific aggregation (Figure 1C, first panel). The negative control in which a *S. acidocaldarius* $\Delta upsAB$ strain was mixed with *S. tokodaii*, revealed, as expected, no aggregation of the *S. acidocaldarius* $\Delta upsAB$ strain and normal aggregation of *S. tokodaii* (Figure 1C, second panel). Interestingly, cells from *S. acidocaldarius* MW135 interacted with *S. tokodaii* cells and thereby formed mixed species aggregates (Figure 1C, third panel).

To find putative species-specific regions in the pilin subunits involved in species-specific recognition, alignments were made using UpsA and UpsB amino acid sequences, from several Sulfolobales (Figure S1). Additionally, the relationship between UpsA and UpsB homologs was studied by creating a phylogenetic tree (Figure S2, Supplementary results and methods). A region with low conservation was revealed in UpsA (Figure S1, amino acid 84-98 for *S. acidocaldarius*, red box). To test whether this region plays a role in

cell-cell recognition, a strain was constructed in which only the region of low conservation in *S. acidocaldarius* UpsA (amino acid 84-98) was exchanged with the corresponding part from *S. tokodaii* UpsA (amino acid 80-101) (MW137 Figure 1A, Table 1). Similar to what was observed for the *S. acidocaldarius* mutant in which *upsA* and *upsB* were exchanged completely (MW135), *S. acidocaldarius* MW137 showed pili formation (Figure S3) but showed little to no aggregation with itself (Figure 1B). Instead, it was found to aggregate with *S. tokodaii* (Figure 1C, fourth panel). This observation strongly suggests that the non-conserved region (exchanged in MW137) defines the species-specificity during cellular aggregation.

The role of glycosylation in species-specificity

The fact that *Sulfolobus* Ups wild-type strains are able to form mating pairs with Ups-deletion strains (32), suggests that factors, other than Ups-pili, play a role in species-specific recognition. The surface proteins of Sulfolobales are heavily glycosylated (36, 37). We therefore suggested that Ups-pili might recognize glycosylated proteins and thereby initiate cellular interactions. To confirm this hypothesis, UV induced aggregation assays were performed in the presence of monosaccharides that are also part of the *S. acidocaldarius* *N*-glycan chain (Glc₁Man₂GlcNAc₂QuiS, containing glucose, mannose, *N*-acetylglucosamine and the *Sulfolobus*-specific sulfoquinovose residues) (37) (Figure 2). The addition of *N*-acetylglucosamine or glucose did not result in altered cellular aggregation (Figure 2A and B); however, in the presence of mannose, cell aggregates were significantly smaller (Figure 2B). This suggests that the mannose molecules partially saturate the binding sites of the Ups-pili and thereby inhibit interactions between pili and the glycan chains on the S-layer of the host cell resulting in reduced aggregation.

Defining the glycosylation pattern of *S. tokodaii* S-layer proteins

Our hypothesis that S-layer glycosylation is important for species-specificity suggests that different *Sulfolobus* species have different glycosylation patterns. So far, the glycan structure of *S. tokodaii* is unknown. To analyze the glycan structures on the S-layer of *S. tokodaii*, *N*-glycans were released from isolated S-layer by hydrazinolysis. Using MALDI/TOF-MS profiling, one main *N*-glycan species and two other low abundant species could be identified in both positive (Figure S4A) and negative ion mode (Figure S4B). The structures of *N*-glycans were proposed based on mass-to-charge ratio of each *N*-glycan ions observed (Figure S4, Table 2) as well as its MS/MS fragmentation pattern (Figure 3). The three *N*-glycan species could be identified as QuiS₁Hex₄HexNAc₂, QuiS₁Hex₃HexNAc₂, and QuiS₁Hex₄HexNAc₁, respectively (Table 2). To determine the linkages between the sugars in the deduced *N*-glycan species, linkage analysis (38) was performed on the permethylated *N*-glycans released from S-layer proteins. The various types of linkages observed on each monosaccharide and their relative abundances on the *N*-glycans are shown in the Figure S5. The most plausible position of this linkage in the glycan chain can be observed on the right-side column in Figure S5. Based on this linkage information, MSⁿ determination of glycan branching (Figure 3), and the glycan masses (Figure S4), the *N*-glycan glycoforms and their isomers were deduced (Figure S6). Figure 4 schematically shows the most prominent glycan structures from *S. acidocaldarius* (37), *S. solfataricus* (39) and *S. tokodaii* (this study). In agreement with our hypothesis, the core of these structures are similar, whereas the outer saccharides differ. A typical sulfated glycan is present in all three *Sulfolobus* glycan structures. Using LC-MS profiling on the tryptic digest of SlaA and SlaB, several different glycopeptides could indeed be observed (Supplementary results and methods, Figure S7 and Figure S8 respectively).

Determination of the binding site in UpsA

We know that a *S. acidocaldarius* mutant in which both Ups-pilin subunits are deleted, does not aggregate upon UV-stress (31). Here we could successfully complement this phenotype by expressing the *upsAB* genes from a maltose inducible plasmid (Figure 5, $\Delta upsAB + upsAB$). Using site directed-mutagenesis on this plasmid, we moreover created point mutations within the above-described region of interest of UpsA (black squares in Figure S1): D85A, N87A, N94A and Y96A. All mutants still produced pili upon UV induction (Figure S8). Interestingly, when expressing UpsA in which the poorly conserved residues D85 or Y96 were mutated to alanine, respectively, UV induced aggregation was significantly reduced. On the other hand, mutation of the conserved N87 or N94 showed wild-type aggregation (Figure 5). These results suggest that the region of low conservation within UpsA is specifically adapted to the glycan structure of the same species in order to ensure species-specific aggregation only.

Discussion

Both bacterial and archaeal T4P have shown to be essential for surface-adherence. Given the fact that bacterial T4P are strongly related to pathogenicity, their mode of binding has primarily been studied for pathogenic bacteria such as *P. aeruginosa*, *Vibrio cholerae*, *Myxococcus*, *Neisseria* and Enteropathogenic *E. coli* species. However, also non-pathogenic bacteria and archaea encode several T4P involved in adhesion, which are studied in far less detail. The Crenarchaeal Sulfolobales encode three types of T4P: archaella, involved in swimming motility (20); Aap-pili, involved in attachment to diverse surfaces (24, 25); and Ups-pili, mediating intraspecies cellular aggregation and DNA exchange (29, 31, 32, 35). During this study, we have examined the role that Ups-pili play in the formation of *Sulfolobus* mating partners. In particular, we focused on the role that pilin subunit UpsA plays in cell-recognition.

The Ups-pilus is formed by two pilin subunits UpsA and UpsB, which are both thought to be major pilin subunits that build up mixed pili structures (31). We revealed that UpsA is involved in species-specific cellular interactions and we were able to alter this specificity by exchanging (parts of) the pilin subunit with that of another species (Figure 1). The binding of bacterial T4P to other cells is often based on pilin-sugar interactions. Surface exposed glycans can be found on cells from all domains of life where they display an enormous range of different structures that are often highly specific to certain species (40). Glycans are therefore perfect anchors to bind specific host- or partner cells. In *Saccharomyces cerevisiae*, surface exposed lectins can bind to surface exposed sugars in a calcium-dependent manner, thereby forming cellular aggregates, a process which is called flocculation (41). This behavior can be inhibited by saturating the binding of the lectins through addition of loose sugars to the medium (42) (Figure 2). In similar experiments with *S. acidocaldarius* we found that mannose has an inhibiting effect on UV-induced cellular aggregation. Since two outer mannose residues are present in the *S. acidocaldarius* *N*-glycan tree, binding of Ups-pili to this side of the glycans tree is probable. When analyzing the *N*-glycans of *S. tokodaii*, we could indeed find differences in this part of the *N*-glycan structure when compared to that of *S. acidocaldarius* (37) and *S. solfataricus*. (39) (Figure 4). As observed for Eukarya (43), the core or the glycan structure is similar in all three species, whereas the outer residues differ. Our results thereby suggest that UpsA contains a specific binding pocket that is able to bind specific sugar moieties of the *N*-glycans presented on the S-layer of distinct *Sulfolobus* species (Figure 6).

Among the euryarchaeal *Haloferax* species, glycosylation was found to be essential for cell fusion (44), emphasizing the importance of glycosylation in Archaeal cellular recognition in general. It is unclear if pili or other types of lectin molecules are involved in cellular

interactions that initiate *Haloferax* fusion events. Similar to our findings, different *Haloferax* species are also known to be differentially glycosylated (45) leading to semi-specific cell-cell recognition (44). Cell fusion between different *Haloferax* species could also be observed but with far lower efficiency (46). In addition, under different environmental conditions, *Haloferax* glycosylation patterns change, leading to more or less favorable *N*-glycans for mating (47). One could envision that low frequency interactions between different *Sulfolobus* species also occur and might occasionally lead to horizontal gene transfer (34), thereby affecting speciation. In a single hot spring in Kamchatka (Russia) two different groups of *Sulfolobus islandicus* strains were found to be present. Despite their coexistence, it was postulated that *S. islandicus* species mainly exchange DNA within these groups (48). It is likely that *N*-glycan patterns and Ups-pili between the species are different serving as a barrier to gene transfer. This behavior might be seen as the two groups diverging into different species.

The Ced-system that is involved in DNA transfer among Sulfolobales can also be found in several crenarchaea that do not encode Ups-pili (34, 35), it is therefore likely these species have developed a different mechanism to initiate cellular interactions. Given the importance of glycosylation in cell-cell interactions in both euryarchaeal *Haloferax* and crenarchaeal *Sulfolobus* species, glycosylation most likely also plays a role in these interactions.

This study has given molecular insights in the cellular recognition mechanism of the previously described crenarchaeal Ups-system (29, 31, 32). Our current model suggests that upon DNA damage, Ups-pili are formed; the UpsA pilin subunits contain a species-specific glycan binding pocket in pilin subunit UpsA that can only bind glycans presented on cells from the same species (Figure 6). This system allows the formation of species-specific cellular connections prior to DNA-exchange via the Ced-system (35). In that way, only DNA

from the same species is exchanged and used for DNA repair via efficient homologous recombination. This proposed cellular recognition mechanism in Sulfolobales restricts exchange of genomic DNA to cells from the same species, thereby playing an important in role genome integrity and the maintenance of species. Co-evolution of *N*-glycosylation and pilin subunit UpsA might play an important role in speciation.

Experimental procedures

Culture conditions.

Sulfolobus acidocaldarius strains and derived mutants (Table 2) were grown aerobically at 75 °C in basic Brock medium (49), supplemented with 0.1% NZ amine, 0.2% dextrin and 20 µg/ml uracil and adjusted to pH 3.5 with sulfuric acid. For solid media the medium was supplemented with 1.5% gelrite. Plates were incubated for 5-6 days at 76°C. *E. coli* competent cells DH5α and ER1821 (NEB) used for respectively cloning and methylation of plasmid DNA were grown in LB medium (10 g/l tryptone; 5 g/l yeast extract; 10 g/l NaCl) at 37°C supplemented with the appropriate antibiotics. Growth of cells was monitored by optical density measurements at 600 nm.

Deleting, exchanging and complementation of genes in *S. acidocaldarius*.

To construct deletion and pilin exchange mutants; up- and downstream flanking regions of the genes of interest (approximately 600 bp) were amplified with primers listed in Table S1. Overlap PCR was performed to connect the up- and downstream fragments. To replace (parts of) *upsA* and *upsB* from *S. acidocaldarius* with their homologues from *Sulfolobus tokodaii*, synthetic DNA was ordered (GenScript) consisting out of *S. acidocaldarius upsAB* flanking regions and (parts of) *S. tokodaii upsAB* genes (Table S1). The PCR product and synthetic DNA fragments were subsequently cloned into pSVA406, resulting in the plasmids listed in Table 1. The plasmids were methylated in *E. coli* ER1821 containing pM.EsaBC4I (NEB)

(50) and transformed into *S. acidocaldarius* MW501 ($\Delta fla/\Delta aap$) (Table 1) (51). This background strain lacks Aap-pili and archaella, allowing easy EM analysis. Integrants were selected on plates lacking uracil and grown in 24-well plates for 2 days in the same medium. Subsequently cultures were plated and grown for 5 days on second selection plates containing uracil and 100 μ g/ml 5-FOA to select for clones in which the plasmid looped out by homologous recombination. Obtained colonies were tested by PCR for successful deletion/replacement of the genes. Correctness of strains was confirmed by DNA sequencing. Strains that were made during this study are listed in Table 1.

For complementation of a $\Delta upsAB$ mutant (MW143), the DNA region comprising *upsA* and *B* was amplified using primers listed in Table S1 and cloned into pSVA1450 under control of a maltose inducible promoter resulting in plasmid pSVA1855 (Table S1). This plasmid was subsequently used to as a template to introduce point mutations into *upsA* (D85A, N87A, N94A and Y96A) (Table 1) using two overlapping primers per mutation (Table S1). Resulting plasmids were then transformed via electroporation into MW143 as described previously (51). Cultures were grown without the addition of uracil. Expression of (mutated) UpsA and B was induced by addition of 0.2% maltose.

UV treatment, aggregation assays.

UV light treatment was performed as described in (29); 10 ml culture (OD_{600} 0.2-0.3) was treated with a UV dose of 75 J/m² (254 nm, Spectroline, UV crosslinker) in a plastic petri dish. For FISH experiments *S. acidocaldarius* and *S. tokodaii* were first mixed in equal amounts. For complementation the $\Delta upsAB$ strain, expression of UpsAB (derivatives) was additionally induced with 0.2% maltose. Afterwards cultures were put back at 76°C for 3 h. Samples taken at different time points were analyzed with phase contrast microscopy. To quantify aggregated cells after induction with UV, 5 μ l of cell culture (diluted to OD 0.2) was

spotted on a microscope slide covered with a thin layer of 2% agarose in Brock minimal medium. Cells were visualized with phase contrast microscopy (Zeiss, Axio Observer.Z1). Free and aggregated cells (≥ 3) were counted for at least three fields per strain using ImageJ cell counter. Percentages of cells in aggregates were subsequently calculated.

Fluorescence *in situ* hybridization (FISH).

For FISH experiments, 10 μ l of a mixed UV induced (described above) culture was spotted and dried on a glass slide. To fix the cells, 10 μ l of 37% formaldehyde was spotted on top of the cells and incubated for 20 min at room temperature. Afterwards formaldehyde was removed and the cells were washed for 10 min with a drop of 1x PBS. Glass slides were subsequently dried at room temperature. Cells were permeabilized by incubating the slides 3 min in 50, 80 and 96% ethanol, respectively. After drying the slides, 10 μ l of hybridization buffer (900 mM NaCl, 20mM Tris HCl pH 8.0, 10% formamid) mixed with 50 ng/ μ l FISH probes (for *S. acidocaldarius* and *S. tokodaii*, Table S1) was spotted on the cells. Slides were incubated in the dark at 46 °C for 1.5 h for hybridization. Subsequently the cells were washed by incubating the slides for 10 min in wash buffer (450 mM NaCl, 20 mM Tris HCl pH 8.0) at 48°C. Slides were then dipped in ice cold water and dried. For microscopy, 1x PBS was spotted on the cells and a coverslip was added. Cells were examined using fluorescence microscopy (Zeiss, Axio Observer.Z1).

S-layer isolation

A cell pellet from a 50 ml *S. tokodaii* str. 7 culture with an OD₆₀₀ of about 0.6 was resuspended and incubated whilst shaking (500 rpm) for 60 min at 37°C in 30 ml of buffer (10 mM NaCl, 20 mM MgSO₄, 0.5% sodium lauroylsarcosine, pH 5.5). Samples were centrifuged for 45 min in an Avanti J-26 XP centrifuge (Beckman Coulter) at 21,000 g (rotor JA-25.50), yielding a brownish tan pellet. The pellet was resuspended and incubated for 30

min at 37°C in 1 ml of buffer A. Subsequent centrifugation for 20 min (tabletop centrifuge at maximum speed), yielded in a translucent tan pellet containing S-layer proteins. Purified S-layer proteins were washed a few times with water and then stored in water at 4°C.

Electron microscopy analysis

Ups-pili on *S. acidocaldarius* cells were visualized with TEM. Cells were negatively stained with 2% uranyl acetate on carbon-coated copper grids. Transmission electron microscopy images were recorded using the Talos L120C (Thermo Scientific™) microscope equipped with a 4k × 4k Ceta CMOS camera. Acceleration voltage was set to 120kV and magnification to 2.27Å/ pixel.

Acknowledgements

AS, IMD and PA were supported in part by the National Institutes of Health grants 1S10OD018530 and P41GM10349010 to the Complex Carbohydrate Research Center. SB was supported by a postdoctoral fellowship from the German Academy of Sciences Leopoldina. We thank Małgorzata Ajon (University of Groningen) for technical support.

References

1. Craig L, Forest KT, Maier B. 2019. Type IV pili: dynamics, biophysics and functional consequences. *Nat Rev Microbiol*.
2. Giltner CL, Nguyen Y, Burrows LL. 2012. Type IV pilin proteins: versatile molecular modules. *Microbiol Mol Biol Rev* 76:740–772.
3. Maier B, Wong GCL. 2015. How Bacteria Use Type IV Pili Machinery on Surfaces. *Trends Microbiol* 23:775–788.
4. Denise R, Abby SS, Rocha EP. 2019. Diversification of the type IV filament superfamily into machines for adhesion, secretion, DNA transformation and motility. *bioRxiv* 576694.
5. Coureuil M, Join-Lambert O, Lécuyer H, Bourdoulous S, Marullo S, Nassif X. 2012. Mechanism of meningeal invasion by *Neisseria meningitidis*. *Virulence* 3:164–172.
6. Bernard SC, Simpson N, Join-Lambert O, Federici C, Laran-Chich M-P, Maïssa N, Bouzinba-Ségard H, Morand PC, Chretien F, Taouji S, Chevet E, Janel S, Lafont F, Coureuil M, Segura A, Niedergang F, Marullo S, Couraud P-O, Nassif X, Bourdoulous S. 2014. Pathogenic *Neisseria meningitidis* utilizes CD147 for vascular colonization. *Nat Med* 20:725–731.
7. Kolappan S, Coureuil M, Yu X, Nassif X, Egelman EH, Craig L. 2016. Structure of the *Neisseria meningitidis* Type IV pilus. *Nat Commun* 7:13015.
8. Hung M-C, Christodoulides M. 2013. The biology of *Neisseria* adhesins. *Biology (Basel)* 2:1054–109.
9. Scheuerpflug I, Rudel T, Ryll R, Pandit J, Meyer TF. 1999. Roles of PilC and PilE proteins in pilus-mediated adherence of *Neisseria gonorrhoeae* and *Neisseria meningitidis* to human erythrocytes and endothelial and epithelial cells. *Infect Immun* 67:834–843.
10. Winther-Larsen HC, Hegge FT, Wolfgang M, Hayes SF, van Putten JP, Koomey M. 2001. *Neisseria gonorrhoeae* PilV, a type IV pilus-associated protein essential to human epithelial cell adherence. *Proc Natl Acad Sci U S A* 98:15276–15281.
11. Hu W, Yang Z, Lux R, Zhao M, Wang J, He X, Shi W. 2012. Direct visualization of the interaction between pilin and exopolysaccharides of *Myxococcus xanthus* with eGFP-fused PilA protein. *FEMS Microbiol Lett* 326:23–30.
12. Li Y, Sun H, Ma X, Lu A, Lux R, Zusman D, Shi W. 2003. Extracellular polysaccharides mediate pilus retraction during social motility of *Myxococcus xanthus*. *Proc Natl Acad Sci U S A* 100:5443–5448.
13. Carter MQ, Chen J, Lory S. 2010. The *Pseudomonas aeruginosa* pathogenicity island PAPI-1 is transferred via a novel type IV pilus. *J Bacteriol* 192:3249–3258.
14. Hong TP, Carter MQ, Struffi P, Casonato S, Hao Y, Lam JS, Lory S, Jousson O. 2017. Conjugative type IVb pilus recognizes lipopolysaccharide of recipient cells to initiate PAPI-1 pathogenicity island transfer in *Pseudomonas aeruginosa*. *BMC Microbiol* 17:31.
15. Albers S V, Pohlschroder M. 2009. Diversity of archaeal type IV pilin-like structures.

- Extremophiles 2009/04/07. 13:403–410.
16. Ng SY, Zolghadr B, Driessen AJ, Albers S V, Jarrell KF. 2008. Cell surface structures of archaea. J Bacteriol 2008/07/16. 190:6039–6047.
17. Szabo Z, Stahl AO, Albers S V, Kissinger JC, Driessen AJ, Pohlschroder M. 2007. Identification of diverse archaeal proteins with class III signal peptides cleaved by distinct archaeal prepilin peptidases. J Bacteriol 2006/11/23. 189:772–778.
18. Makarova KS, Koonin E V, Albers S-V. 2016. Diversity and Evolution of Type IV pili Systems in Archaea. Front Microbiol 7:667.
19. Chaudhury P, Quax TEF, Albers S-V. 2018. Versatile cell surface structures of archaea. Mol Microbiol 107:298–311.
20. Jarrell KF, Albers S-V. 2012. The archaellum: an old motility structure with a new name. Trends Microbiol 20:307–312.
21. Albers S-V, Jarrell KF. 2018. The Archaellum: An Update on the Unique Archaeal Motility Structure. Trends Microbiol 26:351–362.
22. Tripepi M, Imam S, Pohlschröder M. 2010. Haloferax volcanii flagella are required for motility but are not involved in PibD-dependent surface adhesion. J Bacteriol 192:3093–102.
23. Esquivel R, Xu R, Pohlschroder M. 2013. Novel, archaeal adhesion pilins with a conserved N-terminus. J Bacteriol.
24. Henche A-L, Ghosh A, Yu X, Jeske T, Egelman E, Albers S-V. 2012. Structure and function of the adhesive type IV pilus of *Sulfolobus acidocaldarius*. Env Microbiol 14:3188–3202.
25. Zolghadr B, Klingl A, Koerdts A, Driessen AJ, Rachel R, Albers S V. 2010. Appendage-mediated surface adherence of *Sulfolobus solfataricus*. J Bacteriol 2009/10/27. 192:104–110.
26. Bardy SL, Eichler J, Jarrell KF. 2003. Archaeal signal peptides--a comparative survey at the genome level. Protein Sci 2003/08/22. 12:1833–1843.
27. Jarrell KF, Stark M, Nair DB, Chong JPJJ. 2011. Flagella and pili are both necessary for efficient attachment of *Methanococcus maripaludis* to surfaces. FEMS Microbiol Lett 319:44–50.
28. Fröls S, Gordon PMK, Panlilio MA, Duggin IG, Bell SD, Sensen CW, Schleper C. 2007. Response of the hyperthermophilic archaeon *Sulfolobus solfataricus* to UV damage. J Bacteriol 2007/10/02. 189:8708–8718.
29. Fröls S, Ajon M, Wagner M, Teichmann D, Zolghadr B, Folea M, Boekema EJ, Driessen AJ, Schleper C, Albers S V. 2008. UV-inducible cellular aggregation of the hyperthermophilic archaeon *Sulfolobus solfataricus* is mediated by pili formation. Mol Microbiol 2008/11/08. 70:938–952.
30. Götz D, Paytubi S, Munro S, Lundgren M, Bernander R, White MF. 2007. Responses of hyperthermophilic crenarchaea to UV irradiation. Genome Biol 8:R220.
31. van Wolferen M, Ajon M, Driessen AJM, Albers S-V. 2013. Molecular analysis of the UV-inducible pili operon from *Sulfolobus acidocaldarius*. Microbiologyopen 2:928–937.

- 406 32. Ajon M, Fröls S, van Wolferen M, Stoecker K, Teichmann D, Driessen AJ, Grogan
407 DW, Albers S V, Schleper C. 2011. UV-inducible DNA exchange in
408 hyperthermophilic archaea mediated by type IV pili. *Mol Microbiol* 2011/10/18.
409 82:807–817.
- 410 33. Allers T. 2011. Swapping genes to survive - a new role for archaeal type IV pili. *Mol*
411 *Microbiol* 82:789–791.
- 412 34. Wagner A, Whitaker RJ, Krause DJ, Heilers J-H, Van Wolferen M, Van Der Does C,
413 Albers S-V. 2017. Mechanisms of gene flow in archaea. *Nat Rev Microbiol* 15.
- 414 35. van Wolferen M, Wagner A, van der Does C, Albers S-V. 2016. The archaeal Ced
415 system imports DNA. *Proc Natl Acad Sci U S A* 113:2496–2501.
- 416 36. Albers S-V, Meyer BH. 2011. The archaeal cell envelope. *Nat Rev Microbiol* 9:414–
417 426.
- 418 37. Peyfoon E, Meyer B, Hitchen PG, Panico M, Morris HR, Haslam SM, Albers S V, Dell
419 A. 2010. The S-layer glycoprotein of the crenarchaeote *Sulfolobus acidocaldarius* is
420 glycosylated at multiple sites with chitobiose-linked N-glycans. *Archaea* 2010/10/12.
421 2010.
- 422 38. Shajahan A, Heiss C, Ishihara M, Azadi P. 2017. Glycomic and glycoproteomic
423 analysis of glycoproteins—a tutorial. *Anal Bioanal Chem* 409:4483–4505.
- 424 39. Palmieri G, Balestrieri M, Peter-Katalinić J, Pohlentz G, Rossi M, Fiume I, Pocsfalvi
425 G. 2013. Surface-exposed glycoproteins of hyperthermophilic *Sulfolobus solfataricus*
426 P2 Show a common N-glycosylation profile. *J Prot Res* 12:2779–2790.
- 427 40. National research council. 2012. Transforming glycoscience: a roadmap for the future.
428 National Academies Press (US).
- 429 41. Soares E V. 2011. Flocculation in *Saccharomyces cerevisiae*: a review. *J Appl*
430 *Microbiol* 110:1–18.
- 431 42. Masy CL, Henquinet A, Mestdagh MM. 1992. Flocculation of *Saccharomyces*
432 *cerevisiae*: inhibition by sugars. *Canad J Microbiol* 38:1298–306.
- 433 43. Varki A, Lowe JB. 2009. Biological Roles of GlycansEssentials of Glycobiology.
- 434 44. Shalev Y, Turgeman-Grott I, Tamir A, Eichler J, Gophna U. 2017. Cell Surface
435 Glycosylation Is Required for Efficient Mating of *Haloferax volcanii*. *Front Microbiol*
436 8:1253.
- 437 45. Kaminski L, Naparstek S, Kandiba L, Cohen-Rosenzweig C, Arbiv A, Konrad Z,
438 Eichler J. 2013. Add salt, add sugar: N-glycosylation in *Haloferax volcanii* : Figure 1.
439 *Biochem Soc Trans* 41:432–435.
- 440 46. Naor A, Lapierre P, Mevarech M, Papke RT, Gophna U. 2012. Low Species Barriers in
441 Halophilic Archaea and the Formation of Recombinant Hybrids. *Curr Biol* 22:1444–
442 1448.
- 443 47. Guan Z, Naparstek S, Calo D, Eichler J. 2012. Protein glycosylation as an adaptive
444 response in Archaea: growth at different salt concentrations leads to alterations in
445 *Haloferax volcanii* S-layer glycoprotein N-glycosylation. *Environ Microbiol* 14:743–
446 753.
- 447 48. Cadillo-Quiroz H, Didelot X, Held NL, Herrera A, Darling A, Reno ML, Krause DJ,

- Whitaker RJ. 2012. Patterns of Gene Flow Define Species of Thermophilic Archaea. PLoS Biol 10:e1001265.
49. Brock TD, Brock KM, Belly RT, Weiss RL. 1972. *Sulfolobus*: a new genus of sulfur-oxidizing bacteria living at low pH and high temperature. Arch Mikrobiol 1972/01/01. 84:54–68.
50. Kurosawa N, Grogan DW. 2005. Homologous recombination of exogenous DNA with the *Sulfolobus acidocaldarius* genome: properties and uses. FEMS Microbiol Lett 253:141–149.
51. Wagner M, van Wolferen M, Wagner A, Lassak K, Meyer BH, Reimann J, Albers S-V. 2012. Versatile Genetic Tool Box for the Crenarchaeote *Sulfolobus acidocaldarius*. Front Microbiol 3:214.
52. Suzuki T, Iwasaki T, Uzawa T, Hara K, Nemoto N, Kon T, Ueki T, Yamagishi A, Oshima T. 2002. *Sulfolobus tokodaii* sp. nov. (f. *Sulfolobus* sp. strain 7), a new member of the genus *Sulfolobus* isolated from Beppu Hot Springs, Japan. Extremophiles 6:39–44.
53. van Wolferen M, Ajon M, Driessen AJM, Albers S-V. 2013. Molecular analysis of the UV-inducible pili operon from *Sulfolobus acidocaldarius*. Microbiologyopen 2:928–37.
54. Meyer BH, Zolghadr B, Peyfoon E, Pabst M, Panico M, Morris HR, Haslam SM, Messner P, Schaffer C, Dell A, Albers S V. 2011. Sulfoquinovose synthase - an important enzyme in the N-glycosylation pathway of *Sulfolobus acidocaldarius*. Mol Microbiol 2011/11/09. 82:1150–1163.

Figure legends

Figure 1: *S. acidocaldarius upsAB* mutants and their aggregation behavior. (A) (Parts of) *upsA* and *B* from *S. acidocaldarius* or MW501 ($\Delta flaI / \Delta aapF$) (green) were replaced with the same regions from *S. tokodaii* (red), resulting in MW135 (exchange from start codon of *upsA* until stop codon of *upsB*) and MW137 (*Saci upsA* aa 84-98::*ST upsA* aa 80-101) (see also Figure S1). (B) Quantitative analysis of UV-induced cellular aggregation of mutants shown in A. Percentage of cells in aggregates 3h after induction with or without 75 J/m² UV (dark or light grey, respectively). (C) Aggregation behavior of mixtures of *S. tokodaii* (red) with different *S. acidocaldarius* mutants (green) after treatment with UV-light (UV). Untreated cells were used as a control. Mutants used for this experiment were: MW501 (wt *upsAB*), MW143 ($\Delta upsAB$), MW135 and MW137. FISH labeled cells were visualized with fluorescence microscopy. Scale bar 10 μ m.

Figure 2: UV-induced aggregation of *S. acidocaldarius* MW001 upon addition of 20 mM mannose, glucose or *N*-acetylglucosamine. (A) Percentage of cells in aggregates. (B) Average sizes of formed aggregates. Light grey bars represent non-induced cells and dark grey bars represent cells induced with 75 J/m² UV.

Figure 3: HCD MS² spectra of heptasaccharide (*m/z* - 1651.7, Figure S4a) released from the S-layer proteins from *S. tokodaii* by hydrazinolysis.

Figure 4: Structure of the glycan trees present on the S-layer of *S. tokodaii* in comparison to those from *S. acidocaldarius* (37) and *S. solfataricus* (39).

Figure 5: UV induced cellular aggregation of *S. acidocaldarius* $\Delta upsAB$ complementation strains. A *S. acidocaldarius* $\Delta upsAB$ mutant (MW143) was complemented with maltose inducible plasmids carrying *upsAB* or *upsAB* with a D85A, N87, N94A or Y96A mutation in

495 UpsA (see also Figure S1). Percentage of cells in aggregates 3h after induction with or
496 without 75 J/m² UV (dark or light grey, respectively).

497 **Figure 6:** Proposed model species-specific interactions between Ups-pili and *N*-glycosylated
498 S-layer of Sulfolobales. Ups-pili of *S. acidocaldarius* (green) only form interactions with the
499 *N*-glycan of the same species and not with that of other species *S. tokodaii* (red).

500

Table 1: Strains used during this study

Strain	Background strain	Genotype	Source/ reference
<i>S. tokodaii</i> 7			Suzuki et al. 2002
	<i>S. acidocaldarius</i>		
MW001	DSM639	Δ pyrEF (91–412 bp)	Wagner et al. 2012
	<i>S. acidocaldarius</i>	<i>Saci upsAB::ST upsAB</i> ,	
MW135	MW501	Δ flaI (Δ bp 1-672), Δ aapF	This study
	<i>S. acidocaldarius</i>	<i>Saci upsA</i> (aa 84-98):: <i>ST</i>	
MW137	MW501	<i>upsA</i> (aa 80-101), Δ flaI	
		(Δ bp 1-672), Δ aapF	This study
	<i>S. acidocaldarius</i>	Δ upsAB, Δ flaI (Δ bp 1-	
MW143	MW501	672), Δ aapF	This study
	<i>S. acidocaldarius</i>		
MW501	MW001	Δ flaI (Δ bp 1-672), Δ aapF	
			(53)
	<i>S. acidocaldarius</i>		
Δ Agl3	MW001	Δ agl3 (Δ saci0423)	Meyer et al. 2011

Table 2. List of *N*-linked glycans released from S-layer glycoprotein from *S. tokodaii* detected by MALDI/TOF- MS. Abbreviations: QuiS: sulfoquinovose, Hex: hexose, HexNAc *N*-acetyl hexosamine.

Positive ion mode

Permethylated mass (m/z) ¹	Text description of structures	Percentage of glycans
1406	QuiS ₁ Hex ₄ HexNAc ₁	3.92
1447	QuiS ₁ Hex ₃ HexNAc ₂	7.09
1651	QuiS ₁ Hex ₄ HexNAc ₂	88.99

¹All masses (mass+ 2Na - H) are single-charged.

²Calculated from the area units of detected *N*-linked glycans.

Negative ion mode

Permethylated mass (m/z) ¹	Text description of structures	Percentage of glycans
1360	QuiS ₁ Hex ₄ HexNAc ₁	1.27
1401	QuiS ₁ Hex ₃ HexNAc ₂	1.82
1605	QuiS ₁ Hex ₄ HexNAc ₂	96.91

¹All masses (mass - H) are single-charged.

²Calculated from the area units of detected *N*-linked glycans.

510 **Supplementary material:**

511 **Supplementary results and methods:** describing the phylogenetic analysis of UpsA and
512 UpsB from different Sulfolobales and the *N*-glycan analysis of the glycosylated S-layer of *S.*
513 *tokodaii* S-layer proteins SlaA and SlaB.

514 **Table S1:** Plasmids and primers used during this study.

515 **Figure S1:** Alignments of UpsA from different Sulfolobales. The class III cleavage site,
516 cleaved by PibD is depicted by a red line. The red box in UpsA indicates the less conserved
517 region. Shown are UpsA amino acid sequences from *Sulfolobus acidocaldarius* DSM 639
518 (*Saci*), *Sulfolobus tokodaii* Str. 7 (*ST*), *Sulfolobus solfataricus* P2 (*Sso*), *Stygiolobus azoricus*
519 (*Staz*), *Metallosphaera cuprina* Ar-4 (*Mcup*), *Metallosphaera hakonensis* DSM 7519 (*Mhak*),
520 *Metallosphaera sedula* DSM5348 (*Msed*), and *Metallosphaera yellowstonensis* MK1 (*Myel*).

521 **Figure S2:** Maximum-likelihood phylogenetic tree of UpsA and UpsB homologs from
522 different archaeal species (*Sulfolobus acidocaldarius* strains: DSM 639, N8, Ron12/I,
523 SUSAZ; *Sulfolobus solfataricus* strains: P2, 98/2, P1; *Sulfolobus islandicus* strains: REY15A,
524 HVE10/4, M.16.4; *Sulfolobus tokodaii* str. 7; *Stygiolobus azoricus*; *Metallosphaera sedula*
525 DSM5348; *Metallosphaera cuprina* Ar-4; *Metallosphaera hakonensis* DSM 7519;
526 *Metallosphaera yellowstonensis* MK1). Branch numbers represent bootstrap values above
527 80% (100 replicates).

528 **Figure S3:** Transmission electron micrographs of UV induced *S. acidocaldarius* mutants
529 (upper panel) and expression strains (lower panel). Upper panel: Ups-pili of *S. acidocaldarius*
530 MW501 ($\Delta flaI$ / $\Delta aapF$), MW135 (exchange *S. acidocaldarius upsAB* genes with those of *S.*
531 *tokodaii* from start codon of *upsA* until stop codon of *upsB*) and MW137 (*Saci upsA* aa 84-
532 98::*ST upsA* aa 80-101). Lower panel: UV induced *S. acidocaldarius* expression strains.

Wildtype and mutated *upsAB* genes (black squares in Figure S1) were expressed in a $\Delta upsAB/\Delta flaI/\Delta aapF$ strain (MW143). The following maltose-inducible expression plasmids were used: pSVA1855 expressing wildtype *upsAB*; pSVA1860, expressing *upsAB* with a D85A mutation in *upsA*; pSVA1860, expressing *upsAB* with a D85A mutation in *upsA*. pSVA1861, expressing *upsAB* with a N87A mutation in *upsA*; pSVA1862, expressing *upsAB* with a N94A mutation in *upsA*; pSVA1863, expressing *upsAB* with a Y96A mutation in *upsA*. Scale bar 100 nm.

Figure S4: (A) MALDI MS spectra of *N*-glycans released from the S-layer protein from *S. tokodaii* by hydrazinolysis observed (positive ion mode). (B) MALDI MS spectra of *N*-glycans released from the S-layer protein from *S. tokodaii* by hydrazinolysis observed (negative ion mode). Structures are assigned based on MS/MS analysis. *During hydrazinolysis a fraction of glycans gets derivatized by hydrazine reagent.

Figure S5: Glycosyl linkages of monosaccharides of *N*-glycans from the S-Layer proteins of *S. tokodaii* were determined by GC-MS analysis using the PMAA (Partially Methylated Alditol Acetate) method.

Figure S6: Different possible glycoforms of *N*-glycans identified on the S-layer proteins from *S. tokodaii*. Multiple isomers of each glycoforms were also observed. The structure, branching and linkage of *N*-glycans were characterized by MSⁿ fragmentation by ESI-MSⁿ and linkage analysis by GC-MS.

Figure S7: *N*-linked glycosylation sites identified from SlaA of *S. tokodaii* by LC-MS/MS analysis (Tryptic digestion and semi-specific cleavage search using Byonic software).

(A) HCD MS² spectra of glycopeptide ¹¹⁷⁵IYYN[SuphoQuinovose₁Hex₄HexNAc₂] ATSGR¹¹⁸³ from SlaA. (B) HCD MS² spectra of glycopeptide

556 ¹¹⁸⁸NVYGQVVLN[SuphoQuinovose₁Hex₄HexNAc₂]AS GN¹²⁰⁰ from SlaA.

557 (C) HCD MS² spectra of glycopeptide

558 ¹²²²AVLPN[SuphoQuinovose₁Hex₄HexNAc₂]NTLTTL TFNK¹²³⁶ from SlaA.

559 (D) HCD MS² spectra of glycopeptide

560 ¹²⁹⁸IIPAN[SuphoQuinovose₁Hex₄HexNAc₂]ITPIR¹³⁰⁷ from SlaA.

561 (E) HCD MS² spectra of glycopeptide

562 ¹³⁶²EGVN[SuphoQuinovose₁Hex₄HexNAc₂]ASVTSPV VYYSYQAV VAK¹³⁸³ from SlaA.

563 (F) HCD MS² spectra of glycopeptide ¹⁴²¹AVGPAISEYPVNLVFTN[SuphoQuinovose₁Hex₄

564 HexNAc₂]VT VEK¹⁴⁴² from SlaA.

565 **Figure S8:** N-linked glycosylation sites identified from SlaB of *S. tokodaii* by LC-MS/MS

566 analysis (Tryptic digestion and semi-specific cleavage search using Byonic software).

567 (A) HCD MS² spectra of glycopeptide

568 ²⁰⁰GN[SuphoQuinovose₁Hex₄HexNAc₂]QTISLTLK²⁰⁹ from SlaB.

569 (B) HCD MS² spectra of glycopeptide

570 ³⁴⁷EIETVN[SuphoQuinovose₁Hex₄HexNAc₂]QTVYTL MNEIK³⁶³ from SlaB.

571 (C) HCD MS² spectra of glycopeptide

572 ³⁶⁴SLN[SuphoQuinovose₁Hex₄HexNAc₂]ASISQLSTTL SSTTTEITTL NDIK³⁹² from

573 SlaB.

574

575

576

A

B

C

S. acidocaldarius MW501
+ *S. tokodaii*

S. acidocaldarius Δ *upsAB*
+ *S. tokodaii*

S. acidocaldarius MW135
+ *S. tokodaii*

S. acidocaldarius MW137
+ *S. tokodaii*

A **B**

S. acidocaldarius

S. solfataricus

S. tokodaii