Bioactive compounds from *Chrysosporium multifidum*, a fungus isolated from *Hermetia illucens* gut microbiota

Yesenia Correa¹, Billy Cabanillas^{1*}, Valérie Jullian², Daniela Álvarez¹, Denis Castillo¹,
Cédric Dufloer², Beatriz Bustamante³, Elisa Roncal¹, Edgar Neyra¹, Patricia Sheen¹,
Michel Sauvain^{1,2}

¹Laboratorios de Investigación y Desarrollo, Facultad de Ciencias y Filosofía,
Universidad Peruana Cayetano Heredia, Lima, Peru

²UMR 152 PharmaDev, Université de Toulouse, IRD, UPS, France

³Clinical Mycology Laboratory, Instituto de Medicina Tropical Alexander von
Humboldt, Universidad Peruana Cayetano Heredia, Lima, Peru.

* Corresponding autor

E-mail: bjcabanillas@hotmail.com

Abstract

The gut microbiota of insects contains a wide range of organisms that protect them against the attack of pathogens by releasing various types of bioactive compounds. In the present study, we report the isolation and identification of the fungus *Chrysosporium multifidum* as a component of the microbiota from the larval gut of *Hermetia illucens*. Extract from the broth culture of *C. multifidum* showed moderate activity on a strain of methicillin-resistant *Staphylococcus aureus* (MRSA). The bioguided isolation of the extract resulted in the characterization of six α -pyrone derivatives (**1-6**) and one diketopiperazines (**7**), among them 5,6-dihydro-4-methoxy-6-(1-oxopentyl)-2H-pyran-2-one (**4**) showed the best activity ($IC_{50} = 11.4 \pm 0.7 \mu\text{g/ml}$ and $MIC = 62.5 \mu\text{g/ml}$).

Introduction

Hermetia illucens, also known as the black soldier fly (BSF), is native species of Americas. In the last decade, interest in this insect has increased because its larvae can reduce various types of organic waste [1]. BSF larvae have the ability to extract efficiently the protein and lipid content of the wastes they feed on. The larvae and their derivatives appear as a promising alternative in sustainable production for the animal feed industry and biodiesel production. Several research groups and companies around the world are conducting studies in order to optimize their production on a large scale [2]. BSF larvae can feed on different types of organic waste unaffected by bacteria entering their gut [3, 4]. The control of larvae over these pathogens would take place through the release of bioactive substances released into the larva [5]. Thus, compounds generated by BSF larvae could be an alternative to drugs used to treat bacteria harmful to humans, especially those that have shown increased antibiotic resistance in recent years [6, 7]. The gut

microbiota arises as an important source of antibacterial compounds due to the existing evidence on the different biological activities found in the fungi associated with insects [8-13]. Recently, a study showed the diversity of fungi isolated from the gut of BSF larvae. Among them, *Trichosporon asahii* showed to be active on sensitive strains of *Candida glabrata* and *Candida lusitanae* [14]. Other antibacterial substances such as peptides [15, 16] and lipids [17] have also been isolated from BSF, but information is still limited. The purpose of this work is to describe the isolation and identification of a fungus with antimicrobial activity from the gut of BSF larvae, as well as the identification of isolated compounds from the *in vitro* culture of the fungus.

Material and methods

Larvae rearing

The BSF larvae used in this experiment were obtained from the breeding colony established at the Universidad Peruana Cayetano Heredia (Lima, Peru) maintained at 28 ± 1°C and 70% of relative humidity. Specimens were fed with fresh unsterilized chicken guano for 11 days. After this time larvae exhibited lengths between 1.5 - 2 cm.

Extraction of gut from larvae and isolation of active fungi

Samples were collected in triplicate; each collection corresponded to larvae obtained from a different breeding cycle. The larvae were washed with EtOH 70% and sterilized during 15 min with UV light. Each larva was dissected transversally and the gut was extracted. A sample of 0.5 cm was taken from the middle gut to isolate the fungi. The gut samples were homogenized in 200 µl of saline solution 0.89%. The resulting solutions were

diluted (10^{-1} - 10^{-4}) and 10 μ l were seeded in plates containing potato dextrose agar (PDA) (BD-Difco®) and Sabouraud agar (SBA) (BD-Difco®) both supplemented with chloramphenicol (100 mg/L) and gentamicin (50 mg/). To confirm the absence of external contamination two controls were used, a first control using the saline solution used in homogenization, and a second control resulting from the swabbing of the larvae surface after disinfection and before dissection. Plates with sample dilutions were incubated for 21 days at $30 \pm 1^{\circ}\text{C}$. Strains with different morphology were separated repeatedly and constantly in the same media to obtain pure colonies. The activity of fungi colonies to *Staphylococcus aureus* subsp. *aureus* ATCC 43300 and *Salmonella enterica* subsp. *enterica* var. Typhimurium ATCC 13311 were determinate by a previously described method [18], with small modifications. From all the 25 isolates obtained, one of them showed the best activity being submitted to DNA analysis to establish its identity at the specie level.

DNA extraction from the active fungus

A sample of 100 mg of mycelium was transferred into a 2 ml tube and 700 μ l of extraction buffer consisting of 0.1 M Tris-HCl (pH=8) , 20 mM EDTA (pH=8), 1.4 M NaCl, 0.2% (v/v) 2-mercaptoethanol and 2% (w/v) CTAB was added with mixed acid-washed 150-212 μ m glass beads. The mixture was placed inside a Quiagen Tissue Lyser II for 30 seconds, and aliquot of 15 μ l RNAsa A (20 mg/ml) was added and then mixed at 55°C and 850 rpm for 30 min. After, 700 μ l of chloroform:isoamyl alcohol (24:1) was added, and the mixture was centrifuged at 14,000 rpm for 10 min at room temperature. The supernatant was mixed with 50 μ l of 10% (w/v) CTAB and 600 μ l of chloroform and then centrifuged at 14,000 rpm for 10 min. The new supernatant was transferred to a clean 1.5

ml tube an equal volume of ice-cold was added, leaved at -20°C for one night, and then centrifuged at 14,000 rpm for 20 min. The pellet was washed with 1 ml of 80% ethanol (4°C) and centrifuged at 14,000 rpm for 10 min, the operation was repeated and the resulting pellet was left to dry at room temperature for 3 hours. The quantification of DNA was carried out by Nanodrop. Finally, an electrophoresis was performed to verify the integrity of the DNA in 1.5% agarose gel. The extracted DNA was stored at -20°C until use.

PCR amplification and sequencing of the active fungus DNA

A fungal DNA amplification was performed by conventional PCR, two zones were chosen for the amplification, the first one amplified an area covering ITS1-2 rRNA with the universal primers ITS1 (TCCGTAGGTGAACCTGCGGG) and ITS4 (TCCTCCGCTTATTGATGGC); while the second covered D1/D2 domains of large subunit (LSU) ribosomal DNA(rDNA) with the universal primers NL1 (GCATCAATAAGCGGGAGGAAAG) and NL4 (GGTCCGTGTTTCAAGGGG. A master mix solution was prepared containing 25 µl of KOD (Hot start Master Mix- Sigma Aldrich), 1.5 µl of each primer and 18.5 µl of NFW. The DNA was included in 3.5 µl at a concentration of 20 ng/µl. Cycling conditions were as follow: initial denaturation at 94°C for 5 min, followed by 35 denaturation cycles at 94°C for 30 seconds, hybridization at 55.7°C for 30 seconds (for ITS1-2 zone amplification) or 58.1°C for 30 seconds (for D1/D2 region amplification), and extension at 72°C for 60 seconds, then a final extension was performed at 72°C for 7 min. The PCR product was verified by performing an 1.5 % agarose gel electrophoresis. The sequencing process was achieved by Macrogen USA, with an automated system based on Sanger's methodology. The sequences of each

amplified zone were analysed using Sequencer 5.4.6 Software (Gen Codes Corporation). Subsequently, Nucleotide BLAST tool (NCBI) was used to obtain the species with the highest homology between the sequences.

Preparation of the *C. multifidum* broth extract

A culture of *C. multifidum* (1×10^5 spores/ml) was inoculated into 50 ml of Sabouraud broth (BD-Difco®) and incubated at 30°C and 150 rpm for 2 days. The resulting culture was divided into two parts and transferred to a flask containing 500 ml of dextrose broth and incubated for 3 days at 30°C and 150 rpm. This operation was repeated until obtain 10 L of culture. The broth was separated by vacuum filtration and extracted with ethyl acetate (v/v, 1:1). The organic layers were collected and the solvent removed in a rotavapor resulting in 1.5 g of crude extract.

Compound isolation

The crude extract (1.5 g) was chromatographed on silica gel by MPLC with a gradient of CH_2Cl_2 –MeOH (v/v 0:1 to 1:0 v/v) to give 6 fractions (CM1-CM16). Fraction 5 showed the best activity on the bioautography test [19]. It was rechromatographed using silica gel and a gradient of petroleum ether-ethyl acetate (v/v, 90:10 to 80:20) giving as result 8 fractions (CM5.1 – CM5.8), fraction CM5.3 was identify as **2** (6.4 mg), fraction CM 5.5 as **4** (1.7 mg) and fraction CM 5.8 as **6** (8 mg). Fraction CM5.7 was filtered on a Sephadex LH-20 column using CH_2Cl_2 as eluent to yield **1** (4.2 mg). Compounds **3** (16.3 mg) and **5** (4.9 mg) were obtained from the purification of fraction CM9 on silica gel using a gradient of petroleum ether-ethyl acetate (v/v, 80:20 a 60:40). A fractionation of CM10

with a solvent system of MeOH-CH₂Cl₂ (v/v, 80:20 to 100:0) resulted in the isolation of **7** (5 mg).

Antibacterial activity assays

The half-maximal inhibitory concentration (IC₅₀) was determinate using a microdilution method [20, 21]. Bacteria strains were cultured in 7 ml of Mueller Hinton Broth (MHB) incubated at 37°C for 24 hours. The tested compounds were dissolved in DMSO, final concentration of solvent was less 1%. Dilutions were prepared in 96 well plates mixing prepared DMSO solutions with MHB medium to a final volume of 50 µl. Then, a bacteria culture aliquot of 50 µl was inoculated to the dilutions. Final concentrations of compounds in each well ranged from 500 to 0.98 µg/ml and bacteria density was 5 x 10⁵ CFU/ml. After 24 h of incubation at 37°C the optical density (OD) was read at 595 nm. Tetracycline was used as a positive control in a range of 0.3-0.025 µg/ml. A Probit analysis was performed to determine the IC₅₀ of the compounds. The minimum inhibitory concentration (MIC) is determined by observing the first concentration that did not present turbidity or bacterial growth.

Results and discussions

Isolation and identification of the active fungus

The culture of solutions prepared from *H. illucens* gut yielded to the isolation of 25 cultivable fungal strains with different morphotypes. The active fungus was identified as *Arthroderma multifidum* with 100% identity and coverage with the AB861747.1 and AB359438.1. This specie is known as the sexual stage (teleomorph) of *Chrysosporium*

multifidum (anamorph) [22, 23]. No teleomorph stage was show in the prepared cultures, in contrast it presented abundant pyriform microconidia and hyaline septate hyphae (Fig 1) belonging to its anamorph (*Chrysosporium multifidum*), so we finally named it as *C. multifidum* (GenBank accession numbers: MK982149 and MK982181). The use of this stage allowed us to determine the antimicrobial activity of its culture supernatant with different experimental methods.

Fig 1. *Chrysosporium multifidum* isolated from *H. illucens* gut after 7 days of incubation at 30°C. Macroscopic view (A). Microscopic view of pyriform microconidia and hyaline septate hyphae (B).

This specie is a saprotroph commonly found in the soil. It has not been studied as an endosymbiont however it is seen as opportunistic and in other cases as a pathogen. Its isolation is directly related to the food consumed, since there are reports that indicate the isolation of the genus *Chrysosporium* from chicken guano samples [24]. This fact increases the probability that they are organisms assimilated along with food (chicken guano diet) and that they would be selected within the fly's biological system for their convenience (for having enzymes or even beneficial antimicrobial substances) in exchange for providing an environment with enough nutrients for their development [25]. It is known that even some of these selected fungi can survive in glandular cavities or cuticular invaginations called micangias where they can develop and reproduce being favourably transported to new hosts by the insects [26, 27].

Compound isolation from *C. multifidum* broth extract

The bioguided isolation of ethyl acetate extract prepared from the *C. multifidum* broth resulted in the isolation of: 4-methoxy-2H-pyran-2-one (1) [28], 4-methoxy-6-pentyl-2H-

pyran-2-one (**2**) [29], 6-(1-hydroxypentyl)-4-methoxy-pyran-2-one (**3**) [29, 30], 6-
 [(7S,8R)-8-propyloxiran-1-yl]-4-methoxy-pyran-2-one (**4**) [31], pestalotin (**5**) [32, 33],
 5,6-dihydro-4-methoxy-6-(pentanoyloxy)-2H-pyran-2-one (**6**) [30, 33] y cyclo-(L-Pro-L-
 Phe) (**7**) [34]. All the compounds (Fig 2) were identified by comparison of their
 spectroscopic data (HRMS and ¹H and ¹³C NMR) with literature, as well as careful
 examination of their 2D NMR spectra (COSY, HSQC, HMBC). Optical rotations were
 also coherent with those published, except for (**4**), for which we found an optical rotation
 close to zero indicating the possible isolation of a racemic mixture (found [α]_D²⁰ -4.3,
 c=0.16, MeOH/CH₂Cl₂ 9/1 ; published [α]_D²⁵ -98.7, c=0.6, MeOH) [31].

Fig 2. Structure compounds of 1-7 isolated from *C. multifidum* broth extract

This study is the first published chemical characterization of *Chrysosporium multifidum*.
 The literature describes several chemical prospecting works carried out on species of the
Chrysosporium genus, which led to the discovery of groups of compounds such as:
 adenopectines [35], nucleosides [36], zearalenone derivatives, benzoquinones [38],
 naphthaquinones [39], anthraquinones [40], benzolactones [41], naphthopyrones [42],
 naphthalenes [43], phenyl-2(1H)-pyridinones [44], alkylphenols [45],
 bisdechlorogeodins [46], sterols [47, 48], and caryophyllenes [49]. However, there is no
 prior record of any α -pyrones derivatives, so this would be the first report of these
 compounds within the genus. Compound **8** is also reported here for the first time within
 the *Chrysosporium* genus; but this has also been reported from cultures of other fungi and
 bacteria [50, 51].

Biological analyses

The α -pyrone **4** showed to be the more active in the bioautography test. Then antimicrobial activity was quantified, results are shown in Table 1. The values of IC₅₀ (11.4 $\pm$ 0.7 μ g/ml) and MIC (62.5 μ g/ml) on the methicillin-resistant *Staphylococcus aureus* strain indicate only moderate activity compared to control. The known compounds in *Chrysosporium* genus have displayed biological activities as antitumour [44], antifungal [36, 37, 48, 49] and cytotoxic [41], only naphthaquinon-type compounds isolated from *C. queenslandicum* [39] have been shown to be active on gram-positive bacteria *Micrococcus luteus* and *Bacillus subtilis* with MIC values close to those obtained in this work. On the other hand, both natural and synthetic α -pyrones have shown antimicrobial and antifungal activity on a variety of species [52, 53]. Substitutions in positions 4 and 6 of the pyrone ring would be related to this activity. Subsequent trials should be carried out to test the activity of all derivatives of isolated α -pyrones on other groups of bacteria including gram-negative ones.

Table 1. Antimicrobial activity of compound **4** against MRSA strain

Conclusion

A total of 25 fungi colonies were isolated from the gut of *Hermetia illucens* larvae fed with chicken guano. These colonies were tested on methicillin-resistant *Staphylococcus aureus* (MRSA) ATCC 43300 and *Salmonella* Typhimurium ATCC 13311. One colony showed the best activity on the MRSA strain, this specimen was subsequently identified as *Chrysosporium multifidum*. A broth culture of the fungus was prepared and seven compounds were isolated using chromatographic methods and bioassay by bioautography. The active compound against MRSA was identified as the α -pyrone **4** with a MIC of 62.5 μ g/ml. These first results in the exploration of the microbiota of *H.*

illucens open a path to understand the interaction of this fungus with other microorganisms that allow the larva to control the pathogenic microbes introduced through its food made of contaminated organic waste.

Author Contributions

Conceptualization: Michel Sauvain, Yesenia Correa, Denis Castillo.

Investigation: Yesenia Correa, Billy Cabanillas, Valérie Jullian, Daniela Álvarez, Denis Castillo, Cédric Dufloer, Beatriz Bustamante, , Michel Sauvain.

Funding acquisition: Michel Sauvain.

Methodology: Elisa Roncal, Edgar Neyra, Patricia Sheen, Cédric Dufloer

Supervision: Billy Cabanillas, Michel Sauvain, Yesenia Correa, Valérie Jullian.

Validation: Billy Cabanillas, Michel Sauvain, Yesenia Correa, Valérie Jullian.

Writing - original draft: Billy Cabanillas, Yessenia Correa, Michel Sauvain, Daniela Alvarez.

Writing - review & editing: Billy Cabanillas, Yessenia Correa, Michel Sauvain, Daniela Alvarez

References

1. Wang YS, Shelomi M. Review of Black Soldier Fly (*Hermetia illucens*) as animal feed and human food. Foods. 2017; 6(10):91. doi: [10.3390/foods6100091](https://doi.org/10.3390/foods6100091).

2. Müller A, Wolf D, Gutzeit HO. The black soldier fly, *Hermetia illucens* – a promising source for sustainable production of proteins, lipids and bioactive substances. Z Naturforsch C. 2017; 72(9-10):351-363. doi: [10.1515/znc-2017-0030](https://doi.org/10.1515/znc-2017-0030).
3. Jeon H, Park S, Choi J, Jeong G, Lee SB, Choi Y et al. The intestinal bacterial community in the food waste-reducing larvae of *Hermetia illucens*. Curr Microbiol. 2011; 62(5):1390-1399. doi: [10.1007/s00284-011-9874-8](https://doi.org/10.1007/s00284-011-9874-8).
4. Bruno D, Bonelli M, De Filippis F, Di Lelio I, Tettamanti G, Casartelli M, et al. The intestinal microbiota of *Hermetia illucens* larvae is affected by diet and shows a diverse composition in the different midgut regions. Appl Environ Microbiol. 2019; 85 (2): e01864-18. doi: [10.1128/AEM.01864-18](https://doi.org/10.1128/AEM.01864-18).
5. Choi WH, Yun JH, Chu JP, Chu KB. Antibacterial effect of extracts of *Hermetia illucens* (Diptera: Stratiomyidae) larvae against Gram-negative bacteria. Entomol. Res. 2012; 42(5): 219–226. doi: [10.1111/j.1748-5967.2012.00465.x](https://doi.org/10.1111/j.1748-5967.2012.00465.x).
6. Tenover, FC. Mechanisms of antimicrobial resistance in bacteria. Am J Infect Control 2006; 34(5): S3-10. doi: [10.1016/j.ajic.2006.05.219](https://doi.org/10.1016/j.ajic.2006.05.219)
7. Mathur P, Singh S. Multidrug resistance in bacteria: a serious patient safety challenge for India. J Lab Physicians. 2011; 5(1):5-10. doi: [10.4103/0974-2727.115898](https://doi.org/10.4103/0974-2727.115898).
8. Nilanonta C, Isaka M, Kittakoop P, Palittapongarnpim P, Kamchonwongpaisan S, Pittayakhajonwut D, et al.. Antimycobacterial and antiplasmodial cyclodepsipeptides from the insect pathogenic fungus *Paecilomyces tenuipes* BCC 161. Planta Med 2000; 66(8): 756-758. doi: [10.1055/s-2000-9776](https://doi.org/10.1055/s-2000-9776).
9. Gebhardt K, Schimana J, Krastel P, Dettner K, Rheinheimer J, Zeeck A, et al. Endophenazines A–D, newphenazine antibiotics from the arthropod associated

endosymbiont *Streptomyces anulatus*. I. Taxonomy, fermentation, isolation and biological activities. J. Antibiot. 2002; 55(9):794–800. doi: [10.7164/antibiotics.55.794](https://doi.org/10.7164/antibiotics.55.794).

10. Schlörke O, Krastel P, Müller I, Usón I, Dettner K, Zeeck A. Structure and biosynthesis of ceto-niacytone A, a cytotoxic aminocarba sugar produced by an endosymbiotic *Actinomyces*. J. Antibiot. 2002; 55(7):635–642. doi: [10.7164/antibiotics.55.635](https://doi.org/10.7164/antibiotics.55.635).

11. Isaka M, Rugseree N, Maithip P, Kongsaree P, Prabpai S, Thebtaranonth Y. Hirsutellones A–E, antimycobacterial alkaloids from the insect pathogenic fungus *Hirsutella nivea* BCC 2594. *Tetrahedron* 2005; 61(23):5577–5583. doi: [10.1016/j.tet.2005.03.099](https://doi.org/10.1016/j.tet.2005.03.099)

12. Oh, DC, Scott JJ, Currie CR, Clardy J. Mycangimycin, A polyene peroxide from a mutualist *Streptomyces* ssp. *Org. Lett.* 2009; 11(3):633–636. doi: [10.1021/ol802709x](https://doi.org/10.1021/ol802709x).

13. Carr G, Derbyshire ER, Caldera E, Currie CR, Clardy J. Antibiotic and antimalarial quinones from fungus-growing ant-associated *Pseudonocardia* sp. *J. Nat. Prod.* 2012 ; 75(10):1806–1809. doi: [10.1021/np300380t](https://doi.org/10.1021/np300380t).

14. Varotto Boccazzi I, Ottoboni M, Martin E, Comandatore F, Vallone L, Spranghars T, et al. A survey of the mycobiota associated with larvae of the black soldier fly (*Hermetia illucens*) reared for feed production. *PLoS One* 2017; 12(8):e0182533. doi: [10.1371/journal.pone.0182533](https://doi.org/10.1371/journal.pone.0182533).

15. Park SI, Kim JW, Yoe SM. Purification and characterization of a novel antibacterial peptide from black soldier fly (*Hermetia illucens*) larvae. *Dev Comp Immunol.* 2015; 52(1):98–106. doi: [10.1016/j.dci.2015.04.018](https://doi.org/10.1016/j.dci.2015.04.018).

16. Elhag O, Zhou D, Song Q, Soomro AA, Cai M, Zheng L, Yu Z, Zhang J. Screening, expression, purification and functional characterization of novel antimicrobial peptide genes from *Hermetia illucens* (L.). PloS one 2017, 12(1), e0169582. doi: [10.1371/journal.pone.0169582](https://doi.org/10.1371/journal.pone.0169582).
17. Won HC, Jiang. M. Evaluation of antibacterial activity of hexanedioic acid isolated from *Hermetia illucens* larvae. J Appl Biomed. 2014; 12(3):179-189. doi: [10.1016/j.jab.2014.01.003](https://doi.org/10.1016/j.jab.2014.01.003).
18. Pereira E, Santos A, Reis F, Tavares RM, Baptista P, Lino-Neto T, et al. A new effective assay to detect antimicrobial activity of filamentous fungi. Microbiol Res 2013; 168(1), 1-5. doi: [10.1016/j.micres.2012.06.008](https://doi.org/10.1016/j.micres.2012.06.008).
19. Rahalison L, Hamburger M, Hostettmann K, Monod M, Frenk E. A bioautographic agar overlay method for the detection of antifungal compounds from higher plants. Phytochem Analysis 1991; 2:199-203. doi: [10.1002/pca.2800020503](https://doi.org/10.1002/pca.2800020503).
20. Wiegand I, Hilpert K, Hancock RE. Agar and broth dilution methods to determine the minimal inhibitory concentration (MIC) of antimicrobial substances. Nat Protoc. 2008; 3(2):163. doi: [10.1038/nprot.2007.521](https://doi.org/10.1038/nprot.2007.521).
21. Zgoda JR, Porter JR. A Convenient Microdilution method for screening natural products against bacteria and fungi. Pharm Biol. 2001; 39(3):221-5. doi: [10.1076/phbi.39.3.221.5934](https://doi.org/10.1076/phbi.39.3.221.5934).
22. Chabasse D, Guiguen C, Couatarmanac'h A, Launay H, Reece V, De Bièvre C. Contribution à la connaissance de la flore fongique kératinophile isolée des petits Mammifères sauvages et du lapin de garenne en France-Discussion sur les espèces fongiques rencontrées. Ann Parasit Hum Comp. 1987; 62(4):357-368. doi: [10.1051/parasite/1987624357](https://doi.org/10.1051/parasite/1987624357).

23. Metin B, Heitman J. Sexual reproduction in dermatophytes. *Mycopathologia*. 2017; 182(1-2):45-55. doi: [10.1007/s11046-016-0072-x](https://doi.org/10.1007/s11046-016-0072-x).
24. Allimuthu V. Implication of fungal growth in poultry management and biogas production. PhD Thesis, Periyar University, 2015. Available from: <http://hdl.handle.net/10603/151954>.
25. Moubasher AH, Abdel-Sater MA, Soliman Z. Yeasts and filamentous fungi inhabiting guts of three insect species in Assiut, Egypt. *Mycosphere* 2017; 8(9):1297-316. doi: 10.5943/mycosphere/8/9/4.
26. Beaver R, Wilding N, Collins N, Hammond P, Webber J. Insect-fungus relationships in the bark and ambrosia beetles. *Insect-fungus interactions*. 1989:121-43. doi: [10.1016/B978-0-12-751800-8.50011-2](https://doi.org/10.1016/B978-0-12-751800-8.50011-2).
27. Stone W, Nebeker T, Monroe W. Ultrastructure of the mesonotal mycangium of *Xylosandrus mutilatus* (Blandford), an exotic ambrosia beetle (Coleoptera: Curculionidae: Scolytinae) by light, scanning, and transmission electron microscopy. *Microsc Microanal* 2005; 11(S02):172-173. doi: [10.1017/S143192760550015](https://doi.org/10.1017/S143192760550015).
28. Rao KB, Reddy GCS. A new reaction of patulin. *J. Nat. Prod.* 1989, 52(6):1376–1378. doi: [10.1021/np50066a039](https://doi.org/10.1021/np50066a039).
29. Evidente A, Zonno MC, Andolfi A, Troise C, Cimmino A, Vurro M. Phytotoxic α -pyrones produced by *Pestalotiopsis guepinii*, the causal agent of hazelnut twig blight. *The Journal of antibiotics* 2012; 65, 203-206. doi: [10.1038/ja.2011.134](https://doi.org/10.1038/ja.2011.134).
30. Strunz GM, Heissne, CJ, Kakushima M, Stillwell MA. Metabolites of an unidentified Fungus: a new 5,6-dihydro-2-pyrone related to pestalotin. *Can J Chem.* 1974, 52(5): 825-826. doi: [10.1139/v74-128](https://doi.org/10.1139/v74-128).

31. Yang XL, Huang L, Li HY., Yang DF, Li ZZ. Two new compounds from the plant endophytic fungus *Pestalotiopsis versicolor*. J Asian Nat Prod Res. 2014; 17(4): 333-337. doi: [10.1080/10286020.2014.961918](https://doi.org/10.1080/10286020.2014.961918).
32. Ellestad GA, McGahren WJ, Kunstmann MP. Structure of a new fungal lactone, LL-P880.alpha., from an unidentified *Penicillium* species. J Org Chem. 1972; 37(12):2045-2047. doi: [10.1021/jo00977a044](https://doi.org/10.1021/jo00977a044).
33. Kimura Y, Susuki A, Tamura S. ¹³C-NMR Spectra of pestalotin and its analogues. Agr Biol Chem, 1980; 44(2): 451-452. doi: [10.1080/00021369.1980.10863966](https://doi.org/10.1080/00021369.1980.10863966).
34. Sansinenea E, Salazar F, Jiménez J, Mendoza A, Ortiz A. Diketopiperazines derivatives isolated from *Bacillus thuringiensis* and *Bacillus endophyticus*, establishment of their configuration by X-ray and their synthesis. Tetrahedron Lett. 2016, 57(24), 2604–2607. doi: [10.1016/j.tetlet.2016.04.117](https://doi.org/10.1016/j.tetlet.2016.04.117).
35. Hayakawa Y, Adachi H, Kim JW, Shin-ya K, Seto H. Adenopeptin, a new apoptosis inducer in transformed cells from *Chrysosporium* sp. Tetrahedron 1998; 54(52):15871-15878. doi: [10.1016/S0040-4020\(98\)00996-X](https://doi.org/10.1016/S0040-4020(98)00996-X).
36. Yamashita M, Kawai Y, Uchida I, Komori T, Kohsaka M, Imanaka H, et al. Chryscandin, a novel peptidyl nucleoside antibiotic. II. Structure determination and synthesis. J Antibiot. 1984; 37(11):1284-1293. doi: [10.7164/antibiotics.37.1284](https://doi.org/10.7164/antibiotics.37.1284).
37. Hoshino Y, Ivanova VB, Yazawa K, Ando A, Mikami Y, Zaki SM, et al. Queenslondon, a new antifungal compound produced by *Chrysosporium queenslandicum*: production, isolation and structure elucidation. J Antibiot. 2002; 55(5):516-519. doi: [10.7164/antibiotics.55.516](https://doi.org/10.7164/antibiotics.55.516).

38. Fredenhagen A, Petersen F, Tintelnot-Blomley M, Rösel J, Mett H, Hug P. Semicochliodinol A and B: inhibitors of HIV-1 protease and EGF-R protein tyrosine kinase related to asterriquinones produced by the fungus *Chrysosporium merdarium*. J. Antibiot. 1997; 50(5):395-401. doi: [10.7164/antibiotics.50.395](https://doi.org/10.7164/antibiotics.50.395).
39. Ivanova VB, Hoshino Y, Yazawa K, Ando A, Mikami Y, Zaki SM, et al. Isolation and structure elucidation of two new antibacterial compounds produced by *Chrysosporium queenslandicum*. J Antibiot. 2002; 55(10):914-918. doi: [10.7164/antibiotics.55.914](https://doi.org/10.7164/antibiotics.55.914).
40. Slater GP, Haskins RH, Hogge LR. Metabolites from a *Chrysosporium* species. Can J Microbiol. 1971; 17(12): 1576-1579. doi: [10.1139/m71-252](https://doi.org/10.1139/m71-252).
41. Jeon, JE, Julianti E, Oh H, Park W, Oh DC, Oh KB, et al. Stereochemistry of hydroxy-bearing benzolactones: isolation and structural determination of chrysoarticulins A–C from a marine-derived fungus *Chrysosporium articulatum*. Tetrahedron Lett. 2013; 54(24):3111–3115. doi: [10.1016/j.tetlet.2013.04.006](https://doi.org/10.1016/j.tetlet.2013.04.006).
42. Ogawa H, Hasumi K, Sakai K, Murakawa S, Endo A. Pannorin, a new 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor produced by *Chrysosporium pannorum*. J Antibiot 1991; 44(7):762-767. doi: [10.7164/antibiotics.44.762](https://doi.org/10.7164/antibiotics.44.762).
43. Tsipouras A, Goetz MA, Hensens OD, Liesch JM, Ostlind DA, Williamson JM, et al. Sporandol: a novel antiparasitic binaphthalene from *Chrysosporium meridarium*. Bioorg Med Chem Lett. 1997; 7(10):1279–1282. doi: [10.1016/S0960-894X\(97\)00226-6](https://doi.org/10.1016/S0960-894X(97)00226-6).
44. Hirano N, Kohno J, Tsunoda S, Nishio M, Kishi N, Okuda T, et al. TMC-69, a new Antitumor antibiotic with Cdc25A inhibitory activity, produced by *Chrysosporium* sp. TCI068. J Antibiot 2001, 54(5): 421-427. doi: [10.7164/antibiotics.54.421](https://doi.org/10.7164/antibiotics.54.421).

- 399 45. Sekhar Rao KC, Divaka S, Karanth NG, Sattur AP. 14-(2',3',5'-
400 trihydroxyphenyl)tetradecan-2-ol, a novel acetylcholinesterase inhibitor from
401 *Chrysosporium* sp. J. Antibiot. 2001; 54(10):848-849. doi:
402 [10.7164/antibiotics.54.848](https://doi.org/10.7164/antibiotics.54.848).
- 403 46. Tanaka Y, Matsuzaki K, Zhong CL, Yoshida H, Kawakubo T, Masuma R, et al.
404 Dechlorogeodin and its new dihydro derivatives, fungal metabolites with
405 herbicidal activity. J Antibiot. 1996; 49(10):1056-9. doi:
406 [10.7164/antibiotics.49.1056](https://doi.org/10.7164/antibiotics.49.1056).
- 407 47. Van der Pyl D, Cans P, Debernard JJ, Herman F, Lelievre Y, Tahraoui L, et al.
408 RPR113228, a novel farnesyl protein transferase inhibitor produced by
409 *Chrysosporium lobatum*. J Antibiot. 1995; 48(7):736-737. doi:
410 [10.7164/antibiotics.48.736](https://doi.org/10.7164/antibiotics.48.736).
- 411 48. Yang SW, Buevich A, Chan TM, Terracciano J, Chen G, Loebenberg D, et al. A
412 new antifungal sterol sulfate, Sch 601324, from *Chrysosporium* sp. J Antibiot.
413 2003; 56(4):419-422. doi: [10.7164/antibiotics.56.419](https://doi.org/10.7164/antibiotics.56.419).
- 414 49. Yang SW, Chan TM, Terracciano J, Boehm E, Patel R, Chen G, et al.
415 Caryophyllenes from a fungal culture of *Chrysosporium pilosum*. J Nat Prod.
416 2009; 72(3):484-487. doi: [10.1021/np8006414](https://doi.org/10.1021/np8006414).
- 417 50. Chen YS. Studies on the metabolic products of *Rosellinia necatrix*. I. Isolation
418 and characterization of several physiologically active neutral substances. Bull.
419 Agr. Chem. Soc. Japan 1960, 24 (4):372-381. doi:
420 [10.1080/03758397.1960.10857680](https://doi.org/10.1080/03758397.1960.10857680).
- 421 51. Takeda Y, Fujita T, Shingu T, Ogimi C. Studies on the bacterial gall of *Myrica*
422 *rubra*: isolation of a new [7.0]-Metacyclophan from the galland dl-β-phenyllactic

- 423 acid from the culture of gall-forming bacteria. Chem Pharm Bull. 1987; 35:2569–
 424 73. doi: [10.1248/cpb.35.2569](https://doi.org/10.1248/cpb.35.2569).
- 425 52. Fairlamb IJ, Marrison LR, Dickinson JM, Lu FJ, Schmidt JP. 2-pyrones
 426 possessing antimicrobial and cytotoxic activities. Bioorg Med Chem. 2004;
 427 12(15):4285-4299. doi: [10.1016/j.bmc.2004.01.051](https://doi.org/10.1016/j.bmc.2004.01.051).
- 428 53. Bhat ZS, Rather MA, Maqbool M, Lah HU, Yousuf SK, Ahmad Z. α -pyrones:
 429 Small molecules with versatile structural diversity reflected in multiple
 430 pharmacological activities-an update. Biomed Pharmacother. 2017; 91:265-277.
 431 doi: [10.1016/j.biopha.2017.04.012](https://doi.org/10.1016/j.biopha.2017.04.012).

432

433

434

Figure 1

Figure 2

TABLE 1

Compound	IC₅₀ (μg/mL)	MIC (μg/mL)
4	11.4 ± 0.7	62.5
Tetracycline	0.1 ± 0.02	0.4