

Neanderthal-Denisovan ancestors interbred with a distantly-related hominin

Alan R. Rogers,^{1*} Nathan S. Harris,¹ Alan A. Achenbach,¹

¹Department of Anthropology, University of Utah, Salt Lake City, USA

*To whom correspondence should be addressed; E-mail: rogers@anthro.utah.edu

One-sentence summary We document a middle Pleistocene expansion out of Africa and the earliest known admixture between human populations.

Previous research has shown that modern Eurasians interbred with their Neanderthal and Denisovan predecessors. We show here that hundreds of thousands of years earlier, the ancestors of Neanderthals and Denisovans interbred with their own Eurasian predecessors—members of a “superarchaic” population that separated from other humans about 2 mya. The superarchaic population was large, with an effective size between 10 and 46 thousand individuals. We confirm previous findings that: (1) Denisovans also interbred with superarchaics, (2) Neanderthals and Denisovans separated early in the middle Pleistocene, (3) their ancestors endured a bottleneck of population size, and (4) the Neanderthal population was large at first but then declined in size. We provide qualified support for the view that (5) Neanderthals interbred with the ancestors of modern humans.

Introduction

We used genetic data to study the history of human populations during the middle Pleistocene. Early in this period, the ancestors of modern humans separated from those of Neanderthals and Denisovans. Somewhat later, Neanderthals and Denisovans separated from each other. The paleontology and archeology of this period also record important changes, as large-brained hominins appear in Europe and Asia, and Acheulean tools appear in Europe [1, 2]. We studied this period using genetic data from modern Africans and Europeans, and from the two archaic populations, Neanderthals and Denisovans.

Fig. 1 illustrates our notation. Upper-case letters refer to populations, and combinations such as XY refer to the population ancestral to X and Y . X represents an African population (the




Figure 1: A population network including four episodes of gene flow, with an embedded gene genealogy. Upper case letters (*X*, *Y*, *N*, *D*, and *S*) represent populations (Africa, Europe, Neanderthal, Denisovan, and superarchaic). Greek letters label episodes of admixture. *d* and *xyn* illustrate two nucleotide site patterns, in which 0 and 1 represent the ancestral and derived alleles. A mutation on the red branch would generate site pattern *d*. One on the blue branch would generate *xyn*. For simplicity, this figure refers to Neanderthals with a single letter. Elsewhere, we use two letters to distinguish between the Altai and Vindija Neanderthals.




Figure 2: Observed site pattern frequencies. Horizontal axis shows the relative frequency of each site pattern in random samples consisting of a single haploid genome from each of *X*, *Y*, *V*, *A*, and *D*, representing Africa, Europe, Vindija Neanderthal, Altai Neanderthal, Denisovan, and superarchaic. Horizontal lines (which look like dots) are 95% confidence intervals estimated by a moving-blocks bootstrap [4]. Data: Simons Genome Diversity Project [5] and Max Planck Institute for Evolutionary Anthropology [6, 7, 8].

Yorubans), *Y* a European population, *N* Neanderthals, and *D* Denisovans. *S* is an unsampled “superarchaic” population that is distantly related to other humans. Lower-case letters at the bottom of Fig. 1 label “nucleotide site patterns.” A nucleotide site exhibits site pattern *xyn* if random nucleotides from populations *X*, *Y*, and *N* carry the derived allele, but those sampled from other populations are ancestral. Site pattern probabilities can be calculated from models of population history, and their frequencies can be estimated from data. Our Legofit [3] software estimates parameters by fitting models to these relative frequencies.

The current data include two high-coverage Neanderthal genomes: one from the Altai Mountains of Siberia [6] and the other from Vindija Cave in Croatia [8]. Rather than assigning the two Neanderthal fossils to separate populations, our model assumes that they inhabited the same population at different times. This implies that our estimates of Neanderthal population




Figure 3: Residuals from models α and $\alpha\beta\gamma\delta$. Key: red asterisks, real data; blue circles, bootstrap replicates.

size will refer to the Neanderthal metapopulation rather than to any individual subpopulation.

The Altai and Vindija Neanderthals appear in site pattern labels as “a” and “v”. Thus, *av* is the site pattern in which the derived allele appears only in nucleotides sampled from the two Neanderthal genomes. Fig. 2 shows the site pattern frequencies studied here. In contrast to our previous analysis [9], the current analysis includes singleton site patterns, *x*, *y*, *v*, *a*, and *d*, as advocated by Mafessoni and Prüfer [10].

Greek letters in Fig. 1 label episodes of admixture. We label models by concatenating greek letters, to indicate the episodes of admixture they include. For example, model “ $\alpha\beta$ ” includes only episodes α and β . Our model does not include gene flow from Denisovans into moderns, because there is no evidence of such gene flow into Europeans. Two years ago we studied a model that included only one episode of admixture: α , which refers to gene flow from Neanderthals into Europeans [9]. The left panel of Fig. 3 shows the residuals from this model, using the new data. Several are far from zero, suggesting that something is missing from the model [11].

Recent literature suggests some of what might be missing. There is evidence for admixture into Denisovans from a “superarchaic” population, which was distantly related to other humans

Table 1: Bootstrap estimate of predictive error (bepe) values and bootstrap model average (booma) weights

Model	bepe	weight
α	1.14×10^{-6}	0
$\alpha\delta$	0.81×10^{-6}	0
$\alpha\gamma$	0.62×10^{-6}	0
$\alpha\gamma\delta$	0.39×10^{-6}	0
$\alpha\beta$	0.17×10^{-6}	0
$\alpha\beta\gamma$	0.17×10^{-6}	0
$\alpha\beta\delta$	0.14×10^{-6}	0.04
$\alpha\beta\gamma\delta$	0.11×10^{-6}	0.96

[12, 13, 6, 8, 14] and also for admixture from early moderns into Neanderthals [14]. These episodes of admixture appear as β and γ in Fig. 1. Adding β and/or γ to the model improved the fit, yet none of the resulting models were satisfactory. For example, model $\alpha\beta\gamma$ implied (implausibly) that superarchaics separated from other hominins eight million years ago.

To understand what might still be missing, consider what we know about the early middle Pleistocene, around 600 kya. At this time, large-brained hominins appear in Europe along with Acheulean stone tools [1, 2]. They were probably African immigrants, because similar fossils and tools occur earlier in Africa. According to one hypothesis, these early Europeans were Neanderthal ancestors [15, 16]. Somewhat earlier—perhaps 750 kya [17, table S12.2]—the “neandersovan” ancestors of Neanderthals and Denisovans separated from the lineage leading to modern humans. Neandersovans may have separated from an African population and then expanded into Eurasia. If so, they would not have been expanding into an empty continent, for Eurasia had been inhabited since 1.85 mya [18]. Neandersovan immigrants would have met the indigenous superarchaic population of Eurasia. This suggests a fourth episode of admixture—from superarchaics into neandersovans—which appears as δ in Fig. 1.

Results

We considered eight models, all of which include α , and including all combinations of β , γ , and/or δ . In choosing among complex models, it is important to avoid overfitting. Conventional methods such as AIC [19] are not available, because we don’t have access to the full likelihood function. Instead, we use the *bootstrap estimate of predictive error* (bepe) [20, 21, 3]. The best model is the one with the lowest value of bepe. When no model is clearly superior, it is better to average across several than to choose just one. For this purpose, we used *bootstrap model averaging* (booma) [22, 3]. The booma weight of the i th model is the fraction of data sets (including the real data and 50 bootstrap replicates) in which that model “wins,” i.e. has the lowest value of bepe. The bepe values and booma weights of all models are in table 1.

The best model is $\alpha\beta\gamma\delta$, which includes all four episodes of admixture. It has smaller residuals (Fig. 3, right), the lowest bepe value, and the largest booma weight. One other model— $\alpha\beta\delta$ —has a positive booma weight, but all others have weight zero. To understand what this means, recall that bootstrap replicates approximate repeated sampling from the process that generated the data. The models with zero weight lose in *all* replicates, implying that their disadvantage is large compared with variation in repeated sampling. On this basis, we can reject these models. Neither of the two remaining models can be rejected. These results provide strong support for two episodes of admixture (β and δ) and qualified support for a third (γ). Not only does this support previously-reported episodes of gene flow, it also provides evidence of another, in which superarchaics contributed genes to neandersovans. Model-averaged parameter estimates, which use the weights in table 1, are listed in Supporting Online Material (SOM) table S1 and graphed in Fig. 4.

The superarchaic separation time, T_{XYNDs} , has a point estimate of 2.4 mya. This estimate may be biased upward, because our molecular clock assumes a fairly low mutation rate of 0.38×10^{-9} per nucleotide site per year. Other authors prefer slightly higher rates [23]. Although the mutation rate is apparently insensitive to generation time among the great apes, it is sensitive to the age of male puberty. If the average age of puberty during the past two million years were half-way between those of modern humans and chimpanzees, the yearly mutation rate would be close to 0.45×10^{-9} [24, Fig. 2B], and our estimate of T_{XYNDs} would drop to 2.0 mya—just at the origin of the genus *Homo*. Under this clock, the 95% confidence interval is 1.9–2.5 mya.

The lower end of this interval hardly differs from the 1.85 mya date of the earliest Eurasian archaeological remains at Dmanisi [18]. It is possible that superarchaics separated from an African population 1.9 mya, expanded into Eurasia, and left those remains at Dmanisi. If so, then superarchaics descend from the earliest human dispersal into Eurasia. On the other hand, some authors prefer a higher mutation rate of 0.5×10^{-9} per year [6]. Under this clock, the lower end of our confidence interval would be 1.7 mya. Thus, our results are also consistent with the view that superarchaics entered Eurasia after the earliest remains at Dmanisi.

Parameter N_S is the effective size of the superarchaic population. This parameter can be estimated because there are two sources of superarchaic DNA in our sample (β and δ), and this implies that coalescence time within the superarchaic population affects site pattern frequencies. Although this parameter has a broad confidence interval, even the low end implies a fairly large population of about 10,000. This does not require large numbers of superarchaic humans, because effective size can be inflated by geographic population structure [25]. Our large estimate may mean that neandersovans and Denisovans received gene flow from two different superarchaic populations.

Parameter T_{ND} is the separation time of Neanderthals and Denisovans. Our point estimate—731 kya—is remarkably old. Furthermore, the neandersovan population that preceded this split was remarkably small: $N_{ND} \approx 700$. This supports our previous results, which indicated an early separation of Neanderthals and Denisovans and a bottleneck among their ancestors [9].

Because our analysis includes two Neanderthal genomes, we can estimate the effective size of the Neanderthal population in two separate epochs. The early epoch extends from $T_{AV} =$




Figure 4: Model-averaged parameter estimates with 95% confidence intervals estimated by moving-blocks bootstrap [4]. Key: m_α , fraction of Y introgressed from N ; m_β , fraction of D introgressed from S ; m_γ , fraction of N introgressed from XY ; m_δ , fraction of ND introgressed from S ; T_{XYND} , superarchaic separation time; T_{XY} , separation time of X and Y ; T_{ND} , separation time of N and D ; T_{AV} , end of early epoch of Neanderthal history; T_A , age of Altai Neanderthal fossil; T_V , age of Vindija Neanderthal fossil; T_D , age of Denisovan fossil; N_S , size of superarchaic population; N_{XYND} , size of populations $XYND$ and $XYND$; N_{XY} , size of population XY ; N_{ND} , size of population ND ; N_{AV} , size of early Neanderthal population; N_N , size of late Neanderthal population. Parameters that exist in only one model are not averaged.

446 kya to $T_{ND} = 731$ kya, and within this epoch the effective size was large: $N_{AV} \approx 12,000$. It was smaller during the later epoch: $N_N \approx 3500$. These results support previous findings that the Neanderthal population was large at first but then declined in size [6, 8].

Discussion

Early in the middle Pleistocene—about 600 kya—large-brained hominins appear in the fossil record of Europe along with Acheulean stone tools. There is disagreement about how these early Europeans should be interpreted. Some see them as the common ancestors of modern humans and Neanderthals [26], others as an evolutionary dead end, later replaced by immigrants from Africa [27, 28], and others as early representatives of the Neanderthal lineage [15, 16]. Our estimates are most consistent with the last of these views. They imply that by 600 kya Neanderthals were already a distinct lineage, separate not only from the modern lineage but also from Denisovans.

These results resolve a discrepancy involving human fossils from Sima de los Huesos (SH). Those fossils had been dated to at least 350 kya and perhaps 400–500 kya [29]. Genetic evidence showed that they were from a population ancestral to Neanderthals and therefore more recent than the separation of Neanderthals and Denisovans [30]. However, genetic evidence also indicated that this split occurred about 381 kya [6, table S12.2]. This was hard to reconcile with the estimated age of the SH fossils. To make matters worse, improved dating methods later showed that the SH fossils are even older—about 600 ky, and much older than the molecular date of the Neanderthal-Denisovan split [31]. Our estimates resolve this conflict, because they push the date of the split back well beyond the age of the SH fossils.

Our estimate of the Neanderthal-Denisovan separation time conflicts with 381 kya estimate discussed above [6, 10]. This discrepancy results in part from differing calibrations of the molecular clock. Under our clock, the 381 ky date becomes 502 ky [9], but this is still far from our own 731 ky estimate. The remaining discrepancy may reflect differences in our models of history. Misspecified models often generate biased parameter estimates.

Our new results on Neanderthal population size differ from those we published in 2017 [9]. At that time, we argued that the Neanderthal population was substantially larger than others had estimated. Our new estimates are more in line with those published by others [6, 8]. The difference does not result from our new and more elaborate model, because we get similar results from model α , which (like our 2017 model) allows only one episode of gene flow (SOM table S2). Instead, it was including the Vindija Neanderthal genome that made the difference. Without this genome, we still get a large estimate ($N_N \approx 12,000$), even using model $\alpha\beta\gamma\delta$ (SOM table S3). This implies that the Neanderthals who contributed DNA to modern Europeans were more similar to the Vindija Neanderthal than to the Altai Neanderthal, as others have also shown [8].

Our results revise the date at which superarchaics separated from other humans. One previous estimate put this date between 0.9 and 1.4 mya [6, p. 47], which implied that superarchaics

arrived well after the initial human dispersal into Eurasia around 1.9 mya. This required a complex series of population movements between Africa and Eurasia [32, pp. 66-71]. Our new estimates do not refute this reconstruction, but they do allow a simpler one, which involves only three expansions of humans from Africa into Eurasia: an expansion of early *Homo* at about 1.9 mya, an expansion of neandersovans at about 700 kya, and an expansion of modern humans at about 50 kya.

It seems likely that superarchaics descend from the initial human settlement of Eurasia. As discussed above, the large effective size of the superarchaic population hints that it comprised at least two deeply-divided subpopulations, of which one mixed with neandersovans and another with Denisovans. We suggest that around 700 kya, neandersovans expanded from Africa into Eurasia, endured a bottleneck of population size, interbred with indigenous Eurasians, largely replaced them, and separated into eastern and western subpopulations—Denisovans and Neanderthals. These same events unfolded once again around 50 kya as modern humans expanded out of Africa and into Eurasia, largely replacing the Neanderthals and Denisovans.

References

- [1] R. G. Klein, *Journal of World Prehistory* **9**, 167 (1995).
- [2] R. G. Klein, *The Human Career: Human Biological and Cultural Origins* (University of Chicago Press, Chicago, 2009), third edn.
- [3] A. R. Rogers, *bioRxiv* **613067** (2019).
- [4] R. Y. Liu, K. Singh, *Exploring the “Limits” of the Bootstrap*, R. LePage, L. Billard, eds. (Wiley, New York, 1992), pp. 225–248.
- [5] S. Mallick, *et al.*, *Nature* **538**, 201 (2016).
- [6] K. Prüfer, *et al.*, *Nature* **505**, 43 (2014).
- [7] M. Meyer, *et al.*, *Science* **338**, 222 (2012).
- [8] K. Prüfer, *et al.*, *Science* **358**, 655 (2017).
- [9] A. R. Rogers, R. J. Bohlender, C. D. Huff, *Proceedings of the National Academy of Sciences, USA* **114**, 9859 (2017).
- [10] F. Mafessoni, K. Prüfer, *Proceedings of the National Academy of Sciences, USA* **114**, E10256 (2017).
- [11] A. R. Rogers, R. J. Bohlender, C. D. Huff, *Proceedings of the National Academy of Sciences, USA* **114**, E10258 (2017).

- [12] P. J. Waddell, J. Ramos, X. Tan, *ArXiv* **1112.6424** (2011).
- [13] P. J. Waddell, *ArXiv* **1312.7749** (2013).
- [14] M. Kuhlwilm, *et al.*, *Nature* **530**, 429 (2016).
- [15] J.-J. Hublin, *Proceedings of the National Academy of Sciences, USA* **106**, 16022 (2009).
- [16] J.-J. Hublin, *Neandertals and Modern Humans in Western Asia*, T. Akazawa, K. Aoki, O. Bar-Yosef, eds. (Kluwer, 1998), pp. 295–310.
- [17] H. Li, S. Mallick, D. Reich, Population size changes and split times, Supplementary Information 12 of Prüfer *et al.* [6] (2014).
- [18] R. Ferring, *et al.*, *Proceedings of the National Academy of Sciences, USA* **108**, 10432 (2011).
- [19] H. Akaike, *IEEE Transactions on Automatic Control* **19**, 716 (1974).
- [20] B. Efron, *Journal of the American Statistical Association* **78**, 316 (1983).
- [21] B. Efron, R. J. Tibshirani, *An Introduction to the Bootstrap* (Chapman and Hall, New York, 1993).
- [22] S. T. Buckland, K. P. Burnham, N. H. Augustin, *Biometrics* **53**, 603 (1997).
- [23] A. Kong, *et al.*, *Nature* **488**, 471 (2012).
- [24] G. Amster, G. Sella, *Proceedings of the National Academy of Sciences* **113**, 1588 (2016).
- [25] M. Nei, N. Takahata, *Journal of Molecular Evolution* **37**, 240 (1993).
- [26] G. P. Rightmire, *Evolutionary Anthropology* **6**, 218 (1998).
- [27] R. Foley, M. M. Lahr, *Cambridge Archaeological Journal* **7**, 3 (1997).
- [28] P. Endicott, S. Y. W. Ho, C. Stringer, *Journal of Human Evolution* **59**, 87 (2010).
- [29] J. L. Bischoff, *et al.*, *Journal of Archaeological Science* **30**, 275 (2003).
- [30] M. Meyer, *et al.*, *Nature* **531**, 504 (2016).
- [31] J. L. Bischoff, *et al.*, *Journal of Archaeological Science* **34**, 763 (2007).
- [32] D. Reich, *Who We Are and How We Got Here: Ancient DNA and the New Science of the Human Past* (Pantheon Books, New York, 2018).
- [33] Chimpanzee Sequencing and Analysis Consortium, *Nature* **437**, 69 (2005).

- [34] A. Scally, *et al.*, *Nature* **483**, 169 (2012).
- [35] D. Gordon, *et al.*, *Science* **352**, aae0344 (2016).
- [36] H. Li, *Bioinformatics* **27**, 2987 (2011).
- [37] K. R. Veeramah, M. F. Hammer, *Nature Reviews Genetics* **15**, 149 (2014).
- [38] K. Price, R. M. Storn, J. A. Lampinen, *Differential Evolution: A Practical Approach to Global Optimization* (Springer Science and Business Media, 2006).

Acknowledgments

We thank Ryan Bohlender, Elizabeth Cashdan, Fabrizio Mafessoni, Jon Seger, and Timothy Webster for comments. **Funding.** This work was supported by NSF BCS 1638840 (ARR), NSF GRF 1747505 (AAA), and the Center for High Performance Computing at the University of Utah (ARR). **Author contributions.** ARR was responsible for research design, software implementation, data analysis, and writing. NSH and AAA developed and implemented the quality control pipeline for genome sequence data sets. **Competing interests.** None. **Data and materials availability.** Pipelines and intermediate data files are available at `osf.io/vrwna`.

List of supplementary materials

Materials and Methods
 Figs. S1 to S3
 Tables S1 to S3