1 Chromosome level draft genomes of the fall
2 armyworm, *Spodoptera frugiperda* (Lepidoptera:
3 Noctuidae), an alien invasive pest in China

4 Huan Liu^{1,10*}, Tianming Lan^{1,11*}, Dongming Fang^{1,11*}, Furong Gui^{2,9*},
5 Hongli Wang^{1,10}, Wei Guo³, Xiaofang Cheng⁴, Yue Chang⁵, Shuqi
6 He², Lihua Lyu⁶, Sunil Kumar Sahu^{1,10}, Le Cheng⁷, Haimeng Li¹, Ping
7 Liu⁴, Guangyi Fan⁵, Tongxian Liu⁸, Ruoshi Hao⁹, Haorong Lu^{1,10}, Bin
8 Chen⁹, Shusheng Zhu², Zhihui Lu⁹, Fangneng Huang¹², Wei Dong^{1,10},
9 Yang Dong², Le Kang³, Huanming Yang^{1,10}, Jun Sheng^{2,9#}, Youyong
10 Zhu^{2#}, Xin Liu^{1,10#}.

11 1. BGI-Shenzhen, Beishan Industrial Zone, Yantian District, Shenzhen 518083,
12 China

13 2. Yunnan Agricultural University, Kunming 650201, China

14 3. State Key Laboratory of Integrated Management of Pest Insects and Rodents,
15 Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China

16 4. MGI, BGI-Shenzhen, Shenzhen 518083, China

17 5. BGI-Qingdao, BGI-Shenzhen, Qingdao 266555, China

18 6. Plant protection research institute , Guangdong Academy of Agricultural
19 Sciences, Guangzhou 510640, China

20 7. BGI-Yunnan, No. 389 Haiyuan Road, High-tech Development Zone, Kunming,
21 Yunnan 650106, China

22 8. Plant Medical College, Qingdao Agricultural University. Qingdao 266109,
23 China

24 9. Yunnan Plateau Characteristic Agriculture Industry Research Institute ,
25 Kunming 650201, China

10. State Key Laboratory of Agricultural Genomics, BGI-Shenzhen, Shenzhen
518083, China,

11. China National GeneBank, Jinsha Road, Dapeng New District, Shenzhen
518120, China,

12. Department of Entomology, Louisiana State University AgCenter

*These authors contributed equally to this work.

#Correspondence: Xin Liu(liuxin@genomics.cn), Youyong Zhu
(yyzhu@ynau.edu.cn), or Jun Sheng(shengjunpuer@163.com)

Abstract:

The fall armyworm (FAW), *Spodoptera frugiperda* (J.E. Smith) is a severely destructive pest native to the Americas, but has now become an alien invasive pest in China, and causes significant economic loss. Therefore, in order to make effective management strategies, it is highly essential to understand genomic architecture and its genetic background. In this study, we assembled two chromosome scale genomes of the fall armyworm, representing one male and one female individual procured from Yunnan province of China. The genome sizes were identified as 542.42 Mb with N50 of 14.16 Mb, and 530.77 Mb with N50 of 14.89 Mb for the male and female FAW, respectively. We predicted about 22,201 genes in the male genome. We found the expansion of cytochrome P450 and glutathione s-transferase gene families, which are functionally related to the intensified detoxification and pesticides tolerance. Further population analyses of corn strain (C strain) and rice strain (R strain) revealed that the Chinese fall armyworm was most likely invaded from Africa. These strain information, genome features and possible invasion source described in this study will be extremely important for making effective strategies to manage the fall armyworms.

Key Words: Fall armyworm, *Spodoptera frugiperda*, Chromosome-level genome, Insect, Pest

1 Introduction

It has been more than 100 years since the Fall armyworm (FAW), *Spodoptera frugiperda* (J.E. Smith) was reported to damage maize and other crops in the USA¹. It is a severely destructive agricultural pest native to Americas which survives the whole year in the tropical and subtropical area from far south

Argentina, Chile and La Pampa to far north Florida, Texas, Mexico and the Caribbean²⁻⁵. It cannot survive severe winters because of the lack of diapause. However, FAW has a remarkable capacity of long-distance migration, with which the FAW can fly over 100 km per night⁶. Each spring, it can migrate over 2000 km from the overwintering areas to reinvade more northern regions, even up to Canada^{4,7,8}. Recently, FAW spread out from its native region and invaded into Africa in 2016 with the report in São Tomé, Bénin, Togo and Nigeria⁹. This invasion rapidly became widespread in the whole sub-Saharan Africa till October 2017^{10,11}. Following this trend, FAW soon invaded into many Asian countries, including India, Yemen, Thailand, Myanmar and Sri Lanka in 2018¹²⁻¹⁴. Early R *et al* (2018)⁵ forecasted that China is one of the most vulnerable countries of being invaded by the FAW according to the information on frequent commercial trade and passenger transportation between Africa and China. Half a year later after this forecast, in January 2019, International Plant Protection Convention (IPPC) Contact Point for China spotted FAW for the first time in Puer and Dehong city, Yunnan Province, China (<https://www.ippc.int/en/news/first-detection-of-fall-armyworm-in-china/>). This invasive pest has rapidly invaded many provinces in China by June 2019. Now, the FAW has been detected in large parts of the world (Figure 1).

Figure 1 The distribution of the FAW all over the world¹⁵.

There are several hosts of FAW, which mainly includes 186 plant species from 42 families¹⁶. Although FAW is a highly polyphagous pest, graminaceous plants are their preferred hosts, such as maize, rice, sorghum and other major agricultural species. Especially, maize is most likely to be attacked than other plant species⁹. Maize is very important food security of many countries in Asia, Africa, and Latin America (<https://maize.org/projects-cimmyt-and-iita-2/>), and now are facing severe threats from the infestation of the FAW. The production of maize infested by the FAW can suffer yield loss of 40% to 72%, and in some plots 100% total has been reported¹⁷⁻¹⁹. The yield loss of maize can reach 8.3 to 20.6m tons per year in just 12 African countries without any control methods for FAW,

according to the study conducted by Centre for Agriculture and Biosciences International (CABI)²⁰. For Brazil alone, cost to control the FAW on maize is more than 600 million dollars per year²¹. The economic and the yield losses by FAW is a major concern worldwide.

The FAW is not a new species to science; it has been an herbivorous pest for many years. However, the mostly common used approach to mitigate the damage to crops is still the broad-spectrum insecticides²². The use of insecticides highly relies on the knowledge of farmers, but many farmers even do not know the name of the pest²³. Lack of scientific guidance leads to the inappropriate use of pesticides. Besides, the effective management of FAW may require five sprays of pesticides per maize cycle²¹, and many smallholder farmers cannot afford the expensive costs for it, resulting in the use of low-quality products or low dose sprays²⁴. Moreover, the FAW always hides inside the stem of maize, this makes the insecticide much less effective. Although many pesticides are less harmful to the environment and humans, all these factors can lead to the sublethal effects, which possibly help the FAW to evolve resistance against the pesticides²⁵⁻²⁷. An effective compensating management for insecticides is the use of *Bacillus thuringiensis* (Bt) toxins produced by the bacterium. Bt plants have been proved fatal to many insect pests, including the FAW²⁸⁻³⁰. The Bt toxin provides much longer protection than insecticides and less harmful to the environment and humans. Although some research reported the resistance of FAW to Bt maize^{29,30}, multiple genes or new gene with more or new Bt toxin expressed are still thought to have a good performance for resisting the FAW^{29,30}. Biological control, including the introduction of natural enemies and using companion cropping system^{19,21,25,31,32}, is also an effective way to resist the FAW.

The FAW at least consist of two morphological identical but genetically distinct subpopulations, the corn strain (C strain) and rice strain (R strain)³³⁻³⁵. The two strains have their own preferable host plants. The C strain is preferentially associated with maize and other large grasses, but the C strain prefers rice and large grasses^{36,37}. The C strain is subdivided into two subgroups, the FL-type and the TX-type^{38,39}. The TX-type is distributed in most of the Americas, but the FL-type is only limited to Florida and the Caribbean^{3,35,40,41}. Each strain has its strain specific physiological traits, leading to some strain-specific response to biological and chemical agents^{36,37,42}. The C strain larvae are more tolerant than the R strain to the methyl parathion, cypermethrin, cypermethrin and δ -endotoxin from transgenic Bt plant⁴³. Therefore, the origin of FAW needs to be considered to make effective strategies to manage the FAW. The *Cytochrome c oxidase subunit I* (COI) and the *Triosephosphate isomerase* (Tpi) gene are also selected for identifying

the subtype of the FAW, but these markers cannot always give the right identifications⁴⁴.

The management of the FAW needs more detailed genetic information to help people know more about the FAW, to find more new genes to develop more effective Bt plants, and to more accurately identify the different strains for precision spraying of pesticides. Although several genomes of the FAW has been sequenced and assembled⁴⁵⁻⁴⁷, the assemblies were fragmented. Moreover, two genomes were assembled using the Sf21⁴⁵ and Sf9⁴⁷ cell line, respectively, the resource was unique that cannot well provide a comprehensive reference. In this study, we assembled two FAW genomes to the chromosome level using two samples (one male, one female) collected from Yunnan Province, China. We analyzed the subtype of the FAW invaded Yunnan province, and also discussed the possible resource of the invaded FAW in China. We are also screening expanded gene families to seek some key genes with the function of polyphagia and tolerance to insecticides. China is the second largest corn producer after the USA; therefore, it is urgent to select a series of methods to control the FAW. This study provided key information to help make strategies to manage the FAW in China.

2 Materials and Methods

2.1 Samples and treatments

We collected seven FAW samples, including four adults, two fifth-instar larvae and two sixth-instar larvae (Table 1). The four adult FAWs were collected from Yunnan Province, China, and the larvae were collected from Guangdong Province, China. One male and one female adult individual were used for genome sequencing and assembly (Figure 2). Two other adult FAWs were used to capture the conformation of chromosomes to perform the chromosome level genome assembly. One fifth-instar larva and one sixth-instar larva were subjected for the transcriptomic studies. One sixth-instar larva was used for whole genome sequencing with 5K and 300bp insert size. All samples were intestinal and ovarian, and thoroughly cleaned before performing DNA or RNA isolation.

Table 1 Samples used in this study

Sample Identifier	Species	Location	Sex	Instars
SFynMstLFR	<i>Spodoptera frugiperda</i>	Kunming, Yunnan	Male	Adult
SFynFMstLFR	<i>Spodoptera frugiperda</i>	Kunming, Yunnan	Female	Adult
SFgdRNA 1	<i>Spodoptera frugiperda</i>	Guangzhou, Guangdong	Unknown	Fifth-instar
SFgdRNA 2	<i>Spodoptera frugiperda</i>	Guangzhou, Guangdong	Unknown	Sixth-instar
SFgdWGS	<i>Spodoptera frugiperda</i>	Guangzhou, Guangdong	Unknown	Sixth-instar

161

162

163 Figure 2 The two adult fall armyworm (FAWs) used for genome assembly. The left FAW is
164 female, the right FAW is the male individual.

165 2.2 DNA isolation, library preparation, sequencing and genome assembly.

166 The high molecular weight DNA was extracted using the separated muscle tissue
167 by following the protocol recommended by Cheng et al (2018)⁴⁸. We then used
168 the single tube long fragment read (stLFR) technology⁴⁹ to preparing the co-
169 barcoding DNA libraries with the MGIEasy stLFR Library Prep Kit (lot number:
170 1000005622), and the libraries were then loaded on the sequencer for sequencing
171 according to the protocol of MGISEQ-2000⁵⁰. To testify the accuracy of genome
172 assembly, we constructed a DNA library with a 5kb insert size and sequenced by
173 BGISEQ-500 sequencer. To finally ligate the scaffolds to chromosomes, Hi-C
174 technology⁵¹ was used to capture the conformation of chromosomes using
175 another two adult individuals. The primary genome was assembled using the
176 supernova (v2.1.1)⁵² software with the parameters `--maxreads=700M`. We filled
177 gaps by use of GapCloser⁵³ and GapCloser stLFR (unpublished method) with the
178 default parameters. Finally, we performed the chromosome concatenation using
179 the Hi-C generated data by 3d-DNA pipeline⁵⁴.

180 2.3 Comparative genomics analysis

Identification of orthology and paralogy groups of *Spodoptera frugiperda* genes and other considered genomes were done using OrthoMCL⁵⁵ methods on the all-versus-all BLASTP alignment (e-value, <1e-5). We constructed gene families for nine species including *Bombyx mori*, *Danaus plexippus*, *Drosophila melanogaster*, *Heliconius melpomene*, *Helicoverpa armigera*, *Manduca sexta*, *Plutella xylostella*, *Spodoptera litura* and *Spodoptera frugiperda*. The phylogenetic tree, including these nine species, was constructed using the combined set of all the single copy genes.

To search for homology, we compared protein-coding genes of *Spodoptera frugiperda* to that of other species using BLASTP with an E-value threshold of 1e-5. Base on the Whole-genome BLASTP and the genome annotation results, we detected the syntenic blocks using MCscan⁵⁶. A region with at least five syntenic genes and no more than 15 gapped genes was defined as a syntenic block.

2.4 RNA isolation, transcriptome libraries preparation and sequencing

The RNA extraction kit (RNeasy Mini Kit, Qiagen) was used for the total RNA isolation. We performed the RNase-free agarose gel to check the contamination, and then the RNA integrity and purity were measured by Agilent 2100 Bioanalyzer system (Agilent, United States) and NanoDrop Spectrophotometer (THERMO, United States), respectively. The extracted RNA was fragmented into 200-400 bp and reverse transcribed to cDNA for library preparation. The libraries were prepared to follow the manufacturer's instructions for the BGISEQ-500 sequencing platform. Pair-end 100 sequencing was performed on the BGISEQ-500 sequencer using the processed libraries.

2.4 Bioinformatics analysis for transcriptome data

Raw data were firstly processed using the Trimmomatic to filter the reads with adaptors and reads with the proportion of Ns and low-quality bases larger than 10% by SOAPfilter. Bridger software (v20141201) was used to *de novo* assemble the transcriptome, the redundancies were then removed by TGICL. The contigs were concatenated into scaffolds and further assembled to unigenes by clustering and removing redundancy.

FPKM was calculated to estimate the expression level of unigenes. In the study, the reads were mapped against the unigene library using Bowtie, and then unique mapped reads were selected for estimating the expression level by combining eXpress. Finally, DEG unigenes were selected with differential expression level with the parameter of FDR ≤ 0.01 and Fold Change ≥ 4.

2.5 Identifying the strains and the possible source of FAW invasion in China

We used the *Tpi* gene as a DNA maker to identify the strain of the FAW invaded into China. We identified the *Tpi* gene fragments from four FAWs, including two from Yunnan province (sequences were retrieved from the whole genome sequencing data) and two from Guangdong province (sequences were retrieved from the RNA-seq data). Eight sites (TpiE4-129, TpiE4-144, TpiE4-165, TpiE4-168, TpiE4-180, TpiE4-183, TpiE4-192, TpiE4-198) in the fourth exon of the *Tpi* gene were used for determining the strains and a possible source of the invaded FAW⁵⁷. The fourth intron of the *Tpi* gene⁵⁷ was used for constructing the phylogenetic tree using the PhyML (v3.0)⁵⁸ software with the Maximum-Likelihood methods to assist in identifying the strains. Only two *Tpi* fourth introns were used for phylogenetic analysis because we cannot retrieve the introns from RNAseq data. 112 sequences of the fourth intron of the *Tpi* gene with strain information were downloaded from NCBI (Table S1). The sequence alignments were performed using the Clustal W⁵⁹.

3 Results and discussion

3.1 Chromosome level genome assembly for two FAWs

Only one individual was used for genome sequencing for both the male and female genome assembly, which was less than the number used for genome assembly by Gouin A *et al* (2017)⁴⁶. This maximally decreased the heterozygosity level. A total of 2μg and 1.6μg total genomic DNA with the average band size larger than 20 kb were isolated for stLFR libraries preparation from the male and female FAW, respectively. Two stLFR libraries with more than 30 million barcodes were constructed for running on the BGISEQ-500 sequencer. 110.93 Gb and 91.41 Gb high-quality reads were generated for the male and female individual, respectively (Table S2). The primary assembly sizes were 542.42 Mb and 530.77 Mb for the male and female individual. The scaffold N50 and N90 for the male individual were 507.12 Kb and 6.43 Kb, and that for the female were 528.27 Kb and 5.11 Kb, respectively (Table 2). After getting the confirmation from Hi-C sequencing, we finally concatenated the scaffolds to 31 chromosomes with the scaffold N50 of 14.16 Mb and 14.89 Mb for the male and female individual, respectively. Genomes sizes assembled in this study are in the range of the lepidoptera genome sizes at 246M to 809M⁶⁰, but are larger than previously assembled FAW genomes⁴⁵⁻⁴⁷, probably due to the use of the new stLFR technology with the super high coverage and reads quality. This is the first time to assemble the genome of the FAW to the chromosome level, which will be in no doubt to accelerate the biological studies and making the effective strategy of pest managements.

Table2 Summary of the two *Spodoptera frugiperda* genome assemblies

Methods	Statistics	SFynMstLFR				SFynFMstLFR			
		Scaffold Length (bp)	Number	Contig Length (bp)	Number	Scaffold Length (bp)	Number	Contig Length (bp)	Number
stLFR	N50	507,121	226	91,970	1,220	528,269	231	83,007	1,299
	Total length (bp)	542,424,128		505,703,627		530,766,122		496,217,290	
	Ratio of Ns (%)	6.77				6.51			
	GC content (%)	36.52				36.57			
HiC	N50	14,162,803	16	91,970	1,220	14,883,732	13	124,992	813
	Total length (bp)	543,659,128		505,703,627		531,931,622		496,217,290	
	Ratio of Ns (%)	6.98				6.71			
	GC content (%)	36.52				36.58			
Chromosome Level		461,198,141	31			435,876,255	31		
Chromosome Level (%)		84.83				81.94			

254

255 3.2 Evaluation of the assembly

256 The GC content for the genomes of the male and female individual was found to
257 be 35.52% and 36.57% (Table 2), respectively, which showed a similar level with
258 most closely related Lepidoptera species ranging from 31.6% to 37.7%⁶¹. We used
259 the Benchmarking Universal Single-Copy Orthologs (BUSCO: version 2.0)⁶² and
260 Core Eukaryotic Genes (CEGMA)⁶³ to evaluated the completeness of the two
261 assemblies. In BUSCO analysis, genomes of the male and female samples
262 covered 95% and 94.5% complete BUSCO genes (Table S3). In the CEGMA
263 analysis, 83.47% and 85.48% complete core eukaryotic genes were found for the
264 two genomes (Table S4). This is better than all FAW genomes that has been
265 published (PRJNA380964; PRJNA257248; PRJEB13110; and PRJNA344686).
266 Besides, we also mapped the sequencing data generated from libraries of Hi-C,
267 MatePair5K, WGS, and the RNA-seq to the assembled male genome. The
268 mapping rates were all higher than 90% (Table 3), and the insert size were also
269 consistent with the libraries, except for the MatePair5K, probably because the
270 large insert size cannot ensure that the one pair reads they mapped to the same
271 scaffold. EST sequences of the FAW were downloaded from NCBI and the
272 transcripts were assembled without reference. We mapped these EST sequences
273 and transcripts to the male reference genome we assembled, and the results
274 showed that more than 90% EST sequences and more than 80% transcripts we
275 assembled could be found on the assembled male genome (Table S5). However,
276 it is noteworthy that the transcript from SFgdRNA 2 has a lower mapping rate
277 than that of SFgdRNA 1. We inferred that this resulted due to the genetic
278 differences between the C strain and the R strain, because the SFgdRNA 2

sample was identified belong to the R strain. Overall, all the above results well testified the completeness of the two genomes.

Table 3 Mapping reads against the assembled male genome using raw reads generated by different libraries

Type	HIC	MatePair	stLFR	RNA-seq	WGS
Total Mapped Reads	93.58%	93.68%	95.60%	98.98%	90.71%
Perfect Match	38.81%	38.54%	43.67%	52.14%	27.76%
Unique Match	76.39%	77.15%	83.23%	76.99%	74.77%
Total Unmapped Reads	6.42%	6.32%	4.40%	1.02%	9.29%
Total FullMapped Reads	32.48%	30.06%	71.45%	52.46%	49.40%

3.3 Annotation

We firstly used Repeat Modeler (v1.0.11), LTR finder (v1.0.5) and repeatscount (v1.0.5) methods to identify *de novo* repeat motifs by modeling *ab initio*, and these repeat motifs were added into the RepBase⁶⁴ library as known repeat elements. We then performed the RepeatMasker⁶⁵ to mask the assembly, using the combined RepBase library. Usually, Repeat elements take a substantial part of the genome and contribute as important events to genome evolution^{61,66,67}. In this study, by the combination of *de novo* and homology-based searching, 153 Mbp repeat elements were finally identified for the male FAW, and accounting for 28.24% the FAW genomes.

Gene prediction was carried out by both the homology-based and *de novo* methods using repeat masked genomes. For the *de novo* prediction, we used Augustus, glimmerHMM and SNAP (Table S6). For the homology-based approaches, *Bombyx mori*, *Danaus plexippus*, *Drosophila melanogaster* and *Spodoptera litura* genomes were used for homology alignments using the TblastN. Moreover, transcripts that were predicted with RNA-seq, Gene sets were then merged to form a non-redundant gene set with GLEAN; then all annotated genes were checked and filtered manually. A total of 22201 genes were finally obtained for the male samples (Table S6).

In the final gene set we identified, we found 94.2% compete for BUSCO genes and 95.16% CEGMA genes, which were all better than the published FAW genome (Table S7, Table S8). Of these identified genes, 93.48% was confirmed that have functions (Table S9), which was facilitated the further exploration of the functions. Besides, we also found 60 miRNAs, 840 tRNAs and 197 rRNAs by using the homology prediction method.

3.4 The transcriptome analysis of the larvae

After filtering, we finally obtained 58Gb clean data with 341,526,489 cleaned reads. These reads were assembled into 72,604 contigs with the N50 of 2077bp. These contigs were further assembled into scaffolds, and the scaffolds were further assembled to 51,495 unigenes by clustering and removing redundancy. The contig number in our study are significantly higher than that in the study of Kakumani *et al*⁶⁸. This maybe result from that we only used a single method for assembly. We also calculated the expression abundance for unigenes between the fifth-instar and sixth-instar larvae. The result showed 2,648 differentially expressed genes (DEGs). We further performed the clustering analysis to cluster genes with identical or similar expressed behaviors. Remarkable expression difference was found between the fifth-instar larvae and the sixth-instar larvae (Figure S1). This difference was in consistent with the different strains of the two larvae (we described in 3.6). However, if the different instar contributes to the differential expression, it was further confirmed by more detailed analysis.

3.5 Comparison to other published lepidopteran genomes

To further explore the detailed relationship between the FAW and its other lepidopteran relatives. We constructed a phylogenetic tree of nine genomes using 2,001 single copy genes downloaded from NCBI and insectbase (Table S10). The result showed that the *S. frugiperda* which we sequenced actually clustered with its most related species *S. litura*, which is in accordance with the study by Cheng *et al*⁶⁹(Figure 3).

Figure 3 The phylogenetic relationships among nine lepidopteran genomes.

Through the gene family analysis, a total of 12,516 gene families were found in the *S. frugiperda* genome, including 20,012 genes. Of these gene families, 324 are specific to the *S. frugiperda* compared to the other seven species (Table S11). Then, we analyzed the 34 functional gene families of insects, finding some expanded gene families, including cytochrome p450, glutathione s-transferase, and hydrolase (Table 4). The cytochrome p450 gene family is closely related with

intensified detoxification⁶⁹, the genes in this family of the *S. frugiperda* is 200, more than that of *S. litura*, which indicated that the *S. frugiperda* was more polyphagous than *S. litura*. This is also consistent with the habits of *S. frugiperda*. The expanded glutathione s-transferase gene family was proved that could enhance the insecticides tolerance of the *S. litura*⁶⁹. In this study, we found more genes for *S. frugiperda*, which indicated that the *S. frugiperda* was probably easier to gain resistance to pesticides. These gene families are a valuable genetic source to develop more effective pesticides or other methods to manage the FAW.

Table 4 Identified genes in gene families of 9 insects

	DME	PXY	MSE	BMO	HME	DPL	HAR	SLI	SFR
ABC transporter	55	76	40	40	35	38	36	86	66
acetylcholine receptor	18	27	20	27	24	28	22	22	27
Acetylcholinesterase	27	39	62	51	32	27	82	87	46
alkaline phosphatase	13	10	11	10	8	8	8	11	12
aminopeptidase	45	53	49	37	35	39	15	19	24
carboxylesterase	30	55	90	70	50	54	100	110	84
Chitinase	45	27	28	24	26	32	23	26	46
chloride channel	16	25	20	23	18	19	19	16	24
CTL	2	4	3	2	2	2	3	4	4
cytochrome p450	104	108	144	96	130	99	121	132	200
DNA methyltransferase	1	1	1	1	1	1	1	1	1
ecdysone receptor	1	1	3	3	3	2	3	1	3
GABA	15	16	32	9	12	15	16	14	10
glutamate-gated chloride channel	10	14	12	13	10	10	12	12	14
Glutathione s-transferase	50	28	42	25	23	25	46	47	60
glycosyltransferase	65	25	23	19	24	22	19	21	31
G protein	23	42	36	31	26	29	29	29	30
gustatory receptor	54	69	45	76	73	68	197	238	220
heat shock protein	45	38	54	35	29	46	40	48	45
hydrolase	213	250	243	209	181	204	267	310	396
immunoglobulin	65	51	70	76	56	69	71	67	80
odorant-binding protein	62	65	70	51	87	63	77	61	70
odorant receptor	67	82	79	43	73	64	87	75	75
Painless	1	1	1	1	1	1	1	1	1
pheromone	43	22	29	15	19	14	16	14	19
protease inhibitor	75	33	60	38	26	36	46	39	51
Ryanodine receptor	6	12	3	7	4	8	6	3	6
sensory neuron membrane protein	3	5	4	5	4	7	7	3	10
serpin	35	23	30	18	19	24	21	18	26
sirtuin	6	4	3	5	5	5	7	7	7

sodium channel	38	30	33	24	15	27	30	25	36
sugar transporter	17	51	42	36	40	37	41	40	59
superoxide dismutase	17	18	12	12	9	11	12	13	36
Vitellogenin receptor	8	14	15	11	9	12	16	14	15

Note: gene families come from <http://www.insect-genome.com/genefamily/gene-family.php>

344

345 3.6 FAW in China includes both the C and R strain, possibly invaded from Africa

346 There are three strain-specific sites (E4165, E4168 and E4183) in the fourth exon of
347 the *Tpi* gene that can identify the C strain from the R strain for Western
348 Hemisphere populations⁵⁷. Especially, the E4183 is an effective diagnostic marker
349 for *Tpi* gene to define the C or R strain. In this study, two Yunnan samples and
350 one Guangdong sample were identified as the C strain and the other one
351 Guangdong sample was identified as the R strain (Figure 4). The phylogenetic
352 tree showed that the two Yunnan samples were clustered in the clade that
353 consisted of all C strain individuals, which strengthened the results inferred
354 using the strain-specific sites in the fourth exon (Figure 5). This result at least
355 showed that the FAW invaded into China included both the strains. However,
356 the detailed population genetic structure and the frequencies of the two strains in
357 the Chinese population need more information from the population-level
358 studies.

359

360

Figure 4 The identification of C and R strains.

To further confirm the possible source of the Chinese FAW, we compared the haplotype consisted of all eight polymorphisms in the fourth exon of the *Tpi* gene as shown in the Figure 4. We found that all the four individuals hold identical haplotype with the African population, including three *Tpi*Ca1 and one *Tpi*Ra1. Although the haplotype of C strain was shared by the African and the Western Hemisphere populations⁵⁷, the *Tpi*Ra1 has not been detected in any Western Hemisphere populations⁵⁷ showing the uniqueness to African populations. The finding of the *Tpi*Ra1 haplotype in the Guangdong population indicated that there were at least parts of the FAW populations in China that was invaded from Africa, probably through the frequent commercial trade and passenger transportation between Africa and China⁵. However, we cannot confirm other sources because of the small sample size we used here. The strain information and possible invasion source found in this study will be extremely important for making effective strategies to manage the FAWs in China.

Figure 5 The phylogenetic tree to identify the FAW strains of collected from Yunnan, China. The red branched presented the C strain, the red triangles present the two Yunnan samples.

4 Conclusions

In summary, we assembled two chromosome scale genomes of the fall armyworm, representing one male and one female individual procured from Yunnan province of China. The genome sizes were identified as 542.42 Mb with N50 of 14.16 Mb, and 530.77 Mb with N50 of 14.89 Mb for the male and female FAW, respectively. . The completeness of the two genomes are better than all

previously published FAW genomes which is evident by the BUSCO and CEGMA analysis. A total of 22,201 genes were predicted in the male genome, and 12,516 gene families were found in the *S. frugiperda* genome, including 20,012 genes. Of these gene families, we found expansion of cytochrome p450 and glutathione s-transferase gene families, which were closely related to the function of intensified detoxification and pesticides tolerance. We finally identified both the R strain and C strain individuals in the Chinese population, showed that the Chinese FAW was most likely invaded from Africa. The strain information and possible invasion source found in this study will be extremely important for making the effective strategies to manage the FAWs in China.

Acknowledgements

This study was supported by the Guangdong Provincial Key Laboratory of core collection of corp genetic resources research and application (NO.2011A091000047), Guangdong Provincial Key Laboratory of Genome Read and Write (grant no. 2017B030301011) and Key Laboratory of Genomics ,Ministry of Agriculture.

Data availability

The Raw sequencing data and the two chromosome level genome assemblies have been deposited to the CNSA (CNGB Nucleotide Sequence Archive) with accession CNP0000513 (<https://db.cngb.org/cnsa/>).

Author Contributions

Huanming Yang, Le Kang, Jun Sheng, Youyong Zhu, Yang Dong, Xin Liu and Huan Liu designed the research. Dongming Fang, Tianming Lan and Yue Chang, Hongli Wang, Fangneng Huang, Wei Dong and Guangyi Fan performed the data analysis. Xiaofang Chen, Haorong Lu, Ping Liu, Tongxian Liu, Rushi Hao, Bin Chen, Shusheng Zhu, Zhihui Lu and Haimeng Li performed the DNA and RNA extraction and the library preparation. Huan Liu, Tianming Lan, Yang Dong, Wei Guo, Shuqi He, Le Chen and Lihua Lyu collected the samples. Tianming Lan, Dongming Fang, Hongli Wang, Sunil Kumar Sahu and Furong Gui wrote and revised the manuscript. All the authors read and revised the final version of the manuscript.

Competing interests

The authors declare no competing interests.

- 1 Hinds, W. E. & Dew, J. A. The grass worm or fall army worm. *Alabama Agricultural and Experiment Station Bulletin* **186**, 25 (1915).

420 2 Pair, S. D., Raulston, J. R., Westbrook, J. K., Wolf, W. W. & Adams, S. D. Fall
421 Armyworm (Lepidoptera Noctuidae) Outbreak Originating in the Lower Rio
422 Grande Valley, 1989. *Florida Entomologist* **74**, 14 (1991).

423 3 Nagoshi, R. N., Meagher, R. L. & Hay-Roe, M. Inferring the annual migration
424 patterns of fall armyworm (Lepidoptera: Noctuidae) in the United States from
425 mitochondrial haplotypes. *Ecology and evolution* **2**, 1458-1467, doi:10.1002/ece3.268
426 (2012).

427 4 Mitchell, E. R. *et al.* Seasonal Periodicity of Fall Armyworm, (Lepidoptera
428 Noctuidae) in the Caribbean Basin and Northward to Canada. *Journal of*
429 *entomological science* **26**, 12 (1991).

430 5 Early, R., González-Moreno, P., Murphy, S. T. & Day, R. Forecasting the global
431 extent of invasion of the cereal pest *Spodoptera frugiperda*, the fall armyworm.
432 *NeoBiota* **40**, 25-50, doi:10.3897/neobiota.40.28165 (2018).

433 6 Johnson, S. J. Migration and the life history strategy of the fall armyworm,
434 *Spodoptera frugiperda* in the Western Hemisphere. *Insect Sci. Appli* **8**, 7 (1987).

435 7 du Plessis, H., van den Berg, J., Ota, N. & Kriticos, D. J. *Spodoptera frugiperda*
436 (Fall Armyworm). *CSIRO-InSTePP Pest Geography* (2018).

437 8 Westbrook, J. K., Nagoshi, R. N., Meagher, R. L., Fleischer, S. J. & Jairam, S.
438 Modeling seasonal migration of fall armyworm moths. *International journal of*
439 *biometeorology* **60**, 255-267, doi:10.1007/s00484-015-1022-x (2016).

440 9 Goergen, G., Kumar, P. L., Sankung, S. B., Togola, A. & Tamo, M. First Report of
441 Outbreaks of the Fall Armyworm *Spodoptera frugiperda* (J E Smith)
442 (Lepidoptera, Noctuidae), a New Alien Invasive Pest in West and Central Africa.
443 *PloS one* **11**, e0165632, doi:10.1371/journal.pone.0165632 (2016).

444 10 FAO. Briefing Note on FAO Actions on Fall Armyworm in Africa
445 <http://www.fao.org/3/a-bt415e.pdf> (2017).

446 11 Cock, M. J. W., Beseh, P. K., Buddie, A. G., Cafa, G. & Crozier, J. Molecular
447 methods to detect *Spodoptera frugiperda* in Ghana, and implications for
448 monitoring the spread of invasive species in developing countries. *Scientific*
449 *reports* **7**, 4103, doi:10.1038/s41598-017-04238-y (2017).

450 12 Sharanabasappa *et al.* First report of fall armyworm, *Spodoptera frugiperda* (J E
451 smith) (Lepidoptera: Noctuidae), an alien invasive pest on maize in India. *Pest*
452 *Management in Horticultural Ecosystems* **24**, 7 (2018).

453 13 Ganiger, P. C. *et al.* Occurrence of the new invasive pest, fall armyworm,
454 *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae), in the maize fields
455 of Karnataka, India. *Current Science* **115**, 3 (2018).

456 14 Portal, C. F. A. Available online: <https://www.cabi.org/isc/fallarmyworm>
457 (accessed on 6 June 2019).

458 15 CABI. Invasive Species Compendium. <https://www.cabi.org/isc/fallarmyworm>.

459 16 Casmuz, A. *et al.* Revisión de los hospederos del gusano cogollero del maíz,
460 *Spodoptera frugiperda* (Lepidoptera: Noctuidae). *Revista de la Sociedad*
461 *Entomológica Argentina* **69**, 23 (2010).

- 462 17 Murúa, G., Molina-Ochoa, J. & Coviella, C. Population dynamics of the fall
463 armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae) and its parasitoids
464 in northwestern Argentina. *Florida Entomologist* **89**, 8 (2006).
- 465 18 Silva-Aguayo, G. *et al.* Bioactivity of Boldo (*Peumus boldus* Molina) (Laurales:
466 Monimiaceae) on *Spodoptera frugiperda* (J.E. Smith) and *Helicoverpa zea* (Boddie)
467 (Lepidoptera: Noctuidae). *Southwestern Entomologist* **35**, 215-231,
468 doi:10.3958/059.035.0301 (2010).
- 469 19 Wyckhuys, K. A. G. & O'Neil, R. J. Population dynamics of *Spodoptera*
470 *frugiperda* Smith (Lepidoptera: Noctuidae) and associated arthropod natural
471 enemies in Honduran subsistence maize. *Crop Protection* **25**, 1180-1190,
472 doi:10.1016/j.cropro.2006.03.003 (2006).
- 473 20 Day, R. *et al.* Fall Armyworm: Impacts and Implications for Africa. *Outlooks on*
474 *Pest Management* **28**, 196-201, doi:10.1564/v28_oct_02 (2017).
- 475 21 Sisay, B. *et al.* First report of the fall armyworm, *Spodoptera frugiperda*
476 (Lepidoptera: Noctuidae), natural enemies from Africa. *Journal of Applied*
477 *Entomology* **142**, 800-804, doi:10.1111/jen.12534 (2018).
- 478 22 Gordy, J. W., Leonard, B. R., Blouin, D., Davis, J. A. & Stout, M. J. Comparative
479 Effectiveness of Potential Elicitors of Plant Resistance against *Spodoptera*
480 *frugiperda* (J. E. Smith) (Lepidoptera: Noctuidae) in Four Crop Plants. *PloS one*
481 **10**, e0136689, doi:10.1371/journal.pone.0136689 (2015).
- 482 23 Midega, C. A. O. *et al.* Farmers' perceptions of cotton pests and their
483 management in western Kenya. *Crop Protection* **42**, 193-201,
484 doi:10.1016/j.cropro.2012.07.010 (2012).
- 485 24 Dal Pogetto, M. H. F. A. *et al.* Corn Yield with Reduction of Insecticidal Sprayings
486 Against Fall Armyworm *Spodoptera frugiperda* (Lepidoptera Noctuidae). *Journal*
487 *of Agronomy* **11**, 5, doi:10.3923/ja.2012.17.21 (2012).
- 488 25 Fotso Kuate, A. *et al.* *Spodoptera frugiperda* Smith (Lepidoptera: Noctuidae) in
489 Cameroon: Case study on its distribution, damage, pesticide use, genetic
490 differentiation and host plants. *PloS one* **14**, e0215749,
491 doi:10.1371/journal.pone.0215749 (2019).
- 492 26 Yu, S. J., Nguyen, S. N. & Abo-Elghar, G. E. Biochemical characteristics of
493 insecticide resistance in the fall armyworm, *Spodoptera frugiperda* (J.E. Smith).
494 *Pesticide Biochemistry and Physiology* **77**, 1-11, doi:10.1016/s0048-3575(03)00079-8
495 (2003).
- 496 27 de Bon, H. *et al.* Pesticide risks from fruit and vegetable pest management by
497 small farmers in sub-Saharan Africa. A review. *Agronomy for Sustainable*
498 *Development* **34**, 723-736, doi:10.1007/s13593-014-0216-7 (2014).
- 499 28 Sousa, F. F. *et al.* Life-History Traits of *Spodoptera frugiperda* Populations
500 Exposed to Low-Dose Bt Maize. *PloS one* **11**, e0156608,
501 doi:10.1371/journal.pone.0156608 (2016).
- 502 29 Bernardi, D. *et al.* Cross-Resistance between Cry1 Proteins in Fall Armyworm
503 (*Spodoptera frugiperda*) May Affect the Durability of Current Pyramided Bt

504 Maize Hybrids in Brazil. *PloS one* **10**, e0140130, doi:10.1371/journal.pone.0140130
505 (2015).

506 30 Aguirre, L. A. *et al.* Evaluation of Foliar Damage by Spodoptera
507 frugiperda (Lepidoptera: Noctuidae) to Genetically Modified Corn (Poales:
508 Poaceae) in Mexico. *Florida Entomologist* **99**, 276-280, doi:10.1653/024.099.0218
509 (2016).

510 31 Midega, C. A. O., Pittchar, J. O., Pickett, J. A., Hailu, G. W. & Khan, Z. R. A
511 climate-adapted push-pull system effectively controls fall armyworm,
512 Spodoptera frugiperda (J E Smith), in maize in East Africa. *Crop Protection* **105**,
513 10-15, doi:10.1016/j.cropro.2017.11.003 (2018).

514 32 Kenis, M. *et al.* Telenomus Remus, a Candidate Parasitoid for the Biological
515 Control of Spodoptera Frugiperda in Africa, is already Present on the Continent.
516 *Insects* **10**, doi:10.3390/insects10040092 (2019).

517 33 Pashley, D. P. Host-associated Genetic Differentiation in Fall Armyworm
518 (Lepidoptera: Noctuidae): A Sibling Species Complex? *Annals of the Entomological*
519 *Society of America* **79**, 7 (1986).

520 34 Juárez, M. L. *et al.* Population structure of Spodoptera frugiperda maize and rice
521 host forms in South America: are they host strains? *Entomologia Experimentalis et*
522 *Applicata* **152**, 182-199, doi:10.1111/eea.12215 (2014).

523 35 Nagoshi, R. N. *et al.* Analysis of strain distribution, migratory potential, and
524 invasion history of fall armyworm populations in northern Sub-Saharan Africa.
525 *Scientific reports* **8**, 3710, doi:10.1038/s41598-018-21954-1 (2018).

526 36 Nagoshi, R. N. & Meagher, R. L. Behavior and distribution of the two fall
527 armyworm host strains in Florida. *Florida Entomologist* **87**, 10,
528 doi:[http://dx.doi.org/10.1653/0015-4040\(2004\)087\[0440:BADOTT\]2.0.CO;2](http://dx.doi.org/10.1653/0015-4040(2004)087[0440:BADOTT]2.0.CO;2) (2004).

529 37 Meagher, R. L. & Nagoshi, R. N. Differential Feeding of Fall Armyworm
530 (Lepidoptera: Noctuidae) Host Strains on Meridic and Natural Diets. *Annals of*
531 *the Entomological Society of America* **105**, 462-470, doi:10.1603/an11158 (2012).

532 38 Nagoshi, R. N., Silvie, P. & Meagher, R. L. Comparison of Haplotype Frequencies
533 Differentiate Fall Armyworm (Lepidoptera Noctuidae) Corn-Strain Populations
534 from Florida and Brazil. *Molecular Entomology* **100**, 8, doi:10.1603/0022-
535 0493(2007)100[954:cohfdf]2.0.co;2 (2007).

536 39 Nagoshi, R. N. *et al.* Haplotype Profile Comparisons Between Spodoptera
537 frugiperda (Lepidoptera: Noctuidae) Populations From Mexico With Those From
538 Puerto Rico, South America, and the United States and Their Implications to
539 Migratory Behavior. *Journal of economic entomology* **108**, 135-144,
540 doi:10.1093/jee/tou044 (2015).

541 40 Nagoshi, R. N. *et al.* Using Haplotypes to Monitor the Migration of Fall
542 Armyworm (Lepidoptera Noctuidae) Corn-Strain Populations from Texas and
543 Florida. *Journal of economic entomology* **101**, 8, doi:10.1603/0022-
544 0493(2008)101[742:uhtmtm]2.0.co;2 (2008).

545 41 Nagoshi, R. N. *et al.* Fall armyworm migration across the Lesser Antilles and the
546 potential for genetic exchanges between North and South American populations.
547 *PloS one* **12**, e0171743, doi:10.1371/journal.pone.0171743 (2017).

548 42 Unbehend, M. *et al.* Geographic variation in sexual attraction of *Spodoptera*
549 *frugiperda* corn- and rice-strain males to pheromone lures. *PloS one* **9**, e89255,
550 doi:10.1371/journal.pone.0089255 (2014).

551 43 Adamczyk Jr, J. J., Holloway, J. W., Leonard, B. R. & Graves, J. B. Susceptibility of
552 Fall Armyworm Collected from Different Plant Hosts to Selected Insecticides and
553 Transgenic Bt Cotton. *Journal of Cotton Science* **1**, 8 (1997).

554 44 Nagoshi, R. N. Evidence that a major subpopulation of fall armyworm found in
555 the Western Hemisphere is rare or absent in Africa, which may limit the range of
556 crops at risk of infestation. *PloS one* **14**, e0208966,
557 doi:10.1371/journal.pone.0208966 (2019).

558 45 Kakumani, P. K., Malhotra, P., Mukherjee, S. K. & Bhatnagar, R. K. A draft
559 genome assembly of the army worm, *Spodoptera frugiperda*. *Genomics* **104**, 134-
560 143, doi:10.1016/j.ygeno.2014.06.005 (2014).

561 46 Gouin, A. *et al.* Two genomes of highly polyphagous lepidopteran pests
562 (*Spodoptera frugiperda*, Noctuidae) with different host-plant ranges. *Scientific*
563 *reports* **7**, 11816, doi:10.1038/s41598-017-10461-4 (2017).

564 47 Nandakumar, S., Ma, H. & Khan, A. S. Whole-Genome Sequence of the
565 *Spodoptera frugiperda* Sf9 Insect Cell Line. *Genome Announcements* **5**:e00829-17,
566 doi:https://doi.org/10.1128/genomeA.00829-17 (2017).

567 48 Cheng, X. *et al.* A simple bead-based method for generating cost-effective co-
568 barcoded sequence reads. 11 October 2018, *PROTOCOL (Version 1) available at*
569 *Protocol Exchange* [+https://doi.org/10.1038/protex.2018.116+].

570 49 Wang, O. *et al.* Efficient and unique cobarcoding of second-generation
571 sequencing reads from long DNA molecules enabling cost-effective and accurate
572 sequencing, haplotyping, and de novo assembly. *Genome research* **29**, 798-808,
573 doi:10.1101/gr.245126.118 (2019).

574 50 Huang, J. *et al.* A reference human genome dataset of the BGISEQ-500 sequencer.
575 *GigaScience* **6**, 1-9, doi:10.1093/gigascience/gix024 (2017).

576 51 Lieberman-Aiden, E. *et al.* Comprehensive mapping of long-range interactions
577 reveals folding principles of the human genome. *Science* **326**, 289-293,
578 doi:10.1126/science.1181369 (2009).

579 52 Weisenfeld, N. I., Kumar, V., Shah, P., Church, D. M. & Jaffe, D. B. Direct
580 determination of diploid genome sequences. *Genome research* **27**, 757-767,
581 doi:10.1101/gr.214874.116 (2017).

582 53 Luo, R. *et al.* SOAPdenovo2: an empirically improved memory-efficient short-
583 read de novo assembler. *GigaScience* **1**, 18, doi:10.1186/2047-217X-1-18 (2012).

584 54 Dudchenko, O. *et al.* De novo assembly of the *Aedes aegypti* genome using Hi-C
585 yields chromosome-length scaffolds. *Science* **356**, 92-95,
586 doi:10.1126/science.aal3327 (2017).

587 55 Li, L., Stoeckert, C. J., Jr. & Roos, D. S. OrthoMCL: identification of ortholog
588 groups for eukaryotic genomes. *Genome research* **13**, 2178-2189,
589 doi:10.1101/gr.1224503 (2003).

590 56 Wang, Y. *et al.* MCSanX: a toolkit for detection and evolutionary analysis of
591 gene synteny and collinearity. *Nucleic acids research* **40**, e49,
592 doi:10.1093/nar/gkr1293 (2012).

593 57 Nagoshi, R. N., Goergen, G., Plessis, H. D., van den Berg, J. & Meagher, R., Jr.
594 Genetic comparisons of fall armyworm populations from 11 countries spanning
595 sub-Saharan Africa provide insights into strain composition and migratory
596 behaviors. *Scientific reports* **9**, 8311, doi:10.1038/s41598-019-44744-9 (2019).

597 58 Guindon, S. *et al.* New algorithms and methods to estimate maximum-likelihood
598 phylogenies: assessing the performance of PhyML 3.0. *Systematic biology* **59**, 307-
599 321, doi:10.1093/sysbio/syq010 (2010).

600 59 Chenna, R. *et al.* Multiple sequence alignment with the Clustal series of
601 programs. *Nucleic acids research* **31**, 3497-3500, doi:10.1093/nar/gkg500 (2003).

602 60 Triant, D. A., Cinel, S. D. & Kawahara, A. Y. Lepidoptera genomes: current
603 knowledge, gaps and future directions. *Current opinion in insect science* **25**, 99-105,
604 doi:10.1016/j.cois.2017.12.004 (2018).

605 61 Nowell, R. W. *et al.* A high-coverage draft genome of the mycalesine butterfly
606 *Bicyclus anynana*. *GigaScience* **6**, 1-7, doi:10.1093/gigascience/gix035 (2017).

607 62 Simao, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E.
608 BUSCO: assessing genome assembly and annotation completeness with
609 single-copy orthologs. *Bioinformatics* **31**, 3210-3212,
610 doi:10.1093/bioinformatics/btv351 (2015).

611 63 Parra, G., Bradnam, K. & Korf, I. CEGMA: a pipeline to accurately annotate core
612 genes in eukaryotic genomes. *Bioinformatics* **23**, 1061-1067,
613 doi:10.1093/bioinformatics/btm071 (2007).

614 64 Bao, W., Kojima, K. K. & Kohany, O. Repbase Update, a database of repetitive
615 elements in eukaryotic genomes. *Mobile DNA* **6**, 11, doi:10.1186/s13100-015-0041-9
616 (2015).

617 65 Saha, S., Bridges, S., Magbanua, Z. V. & Peterson, D. G. Empirical comparison of
618 ab initio repeat finding programs. *Nucleic acids research* **36**, 2284-2294,
619 doi:10.1093/nar/gkn064 (2008).

620 66 Lorenzi, H. *et al.* Genome wide survey, discovery and evolution of repetitive
621 elements in three *Entamoeba* species. *BMC genomics* **9**, 595, doi:10.1186/1471-
622 2164-9-595 (2008).

623 67 Thybert, D. *et al.* Repeat associated mechanisms of genome evolution and
624 function revealed by the *Mus caroli* and *Mus pahari* genomes. *Genome research* **28**,
625 448-459, doi:10.1101/gr.234096.117 (2018).

626 68 Kakumani, P. K. *et al.* De novo transcriptome assembly and analysis of Sf21 cells
627 using illumina paired end sequencing. *Biology direct* **10**, 44, doi:10.1186/s13062-
628 015-0072-7 (2015).

629 69 Cheng, T. *et al.* Genomic adaptation to polyphagy and insecticides in a major East
630 Asian noctuid pest. *Nature ecology & evolution* **1**, 1747-1756, doi:10.1038/s41559-
631 017-0314-4 (2017).

632