Title: Attraction of *Triatoma infestans* (Klug) to adhesive yeast-baited trap under laboratory conditions

Cardozo Miriam¹, Fiad Federico G.², Crocco Liliana B. ¹, Gorla David E.³

Institutional affiliations:

1. Universidad Nacional de Córdoba. Facultad de Ciencias Exactas, Físicas y Naturales. Cátedra de Introducción a la Biología. Córdoba, Argentina. Instituto de Investigaciones Biológicas y Tecnológicas (IIBYT-CONICET)- Avda. Vélez Sarsfield 299, piso 5. X5000JJC, Córdoba, Argentina.
2. Universidad Nacional de Córdoba. Facultad de Ciencias Exactas, Físicas y Naturales. Cátedra de Introducción a la Biología. Córdoba, Argentina. Avda. Vélez Sarsfield 299, piso 5. X5000JJC, Córdoba, Argentina.
3. Universidad Nacional de Córdoba. Grupo de Ecología y Control de Vectores. Instituto de Diversidad y Ecología Animal (IDEA-CONICET). Rondeau 798 (Jardín Zoológico) X5000AVP, Córdoba, Argentina.

Corresponding author

E-mail address: cardozo.miri@gmail.com (MC)

ORCID: 0000-0002-4339-974X

Postal address: Instituto de Investigaciones Biológicas y Tecnológicas, Facultad de Ciencias Exactas, Físicas y Naturales, Universidad Nacional de Córdoba. Avda. Vélez Sarsfield 299, piso 5. X5000JJC, Córdoba, Argentina.

Telephone: 54+351+4332100 ext. 248 - Fax: 54+351+4332100

Abstract

Existing methods to detect domestic triatomines have low sensitivity. As early house infestation detection is epidemiologically important, the exploration of better methods is required. Hence, we measured the attractiveness of a yeast-baited trap to adults and nymphs of *Triatoma infestans*, under laboratory conditions.

The assays were conducted in an experimental arena, with an experimental and a control traps placed at opposite sides and one refuge in the center area. Insects were released and the number of triatomines in the yeast and control traps were counted, after 3, 6 and 24 hours of the beginning of the experiment. We use generalized linear models within a multimodel inference approach to model the number of insects in the trap, using insect age classes, time after assay initiation and date of the experiment as predictors.

Our results show that the attraction to CO₂ depends upon the life stage of the insects. During the 24 hours of experiment a constant number of adults were attracted to the yeast trap, while nymphs show attraction only up to the first three hours after the initiation of CO₂ liberation. Undoubtedly, the orientation response to chemical cues deserves further studies to be fully understood.

Key words: Triatominae, Chagas Disease, yeasts, carbon dioxide

Introduction

Chagas disease is considered one of the most important endemic diseases in Latin America, affecting approximately 5–6 million individuals. The disease is caused by *Trypanosoma cruzi* (Trypanosomatidae), which not only infects humans but also more than 100 species of domestic and sylvatic mammals and can be transmitted by over 150 species of triatomines (Triatominae, Reduviidae) (WHO 2015).

Triatoma infestans, characterized by its high adaptive capacity to domestic environments, is the vector with the greatest epidemiological importance in the Southern Cone countries of South America (Rabinovich 1972; Lent and Wygodzinsky 1979).

The maximum geographical expansion of *T. infestans* distribution occurred between 1970 and 1980 with an estimated occupation area of 6.28 million km², including Argentina, Bolivia, Brazil, Chile, Paraguay, Peru and Uruguay. The Southern Cone Initiative, coordinated by the Pan American Health Organization to control the transmission of Chagas disease in Latin America from 1991 interrupted the vector transmission of *T. cruzi* in Chile, Uruguay and Brazil through insecticide-based vector control, health education and house improvement programs (Dias et al. 2002). The Initiative produced a significant reduction of the distribution area of *T. infestans* to less than 1 million km² (Schofield et al. 2006). Nevertheless, in arid Gran Chaco areas of Argentina, Paraguay, and Bolivia, reinfestations of human dwellings continue to occur in several provinces or departments (Ceballos et al. 2011; Bustamante-Gómez et al. 2016).

Although this species has long been found almost exclusively in domiciliary and peridomestic environments, a significant increase in the number of wild population found in sylvatic environments was reported, mainly in the Inter-Andean Valleys of Bolivia, in the Gran Chaco of Argentina, Bolivia and Paraguay (Noireau et al.1997; Rolón et al. 2011) and in a Metropolitan region from Chile (Bacigalupo et al. 2010). Recent studies also evidenced the presence of gene flow between sylvatic and intra-peridomestic *T. infestans* populations in Argentina (Piccinali et al. 2011), suggesting that sylvatic populations may be involved in the reinfestation observed in some places.

The prevention of Chagas disease depends on the elimination of the domestic colonies of triatomines. Insecticide residual spraying is very effective, but re-infestation of treated dwellings is frequent. Early detection and elimination of triatomine reinfestation is critical for long-term control; however, current methods used for vector-detection have low sensitivity. A number of alternatives have been evaluated (Abad-Franch et al. 2011), either for the detection of domestic triatomine species (like *T. infestans*), or other secondary vector species that frequently invade domestic and peridomestic structures (Cavallo et al. 2016; Cecere et al. 2016; Giraldez et al. 2016). A number of methods for triatomine detection have been tested in different ecotopes (sylvatic, domestic and peridomestic environments). Current method for routine entomological surveillance used by vector control programs in Latin America is the fixed effort active search, sometimes using a flushing out agent. Although widely adopted, it depends heavily on operator experience and has low sensitivity when vector abundance is low (Gürtler et al.1999).

Passive bio-sensors such as the Gómez-Nuñez box or Maria sensors (Gómez-Nuñez 1965; Wisnivesky-Colli et al. 1987) were tested with poor results because of methodological and operational limitations and low sensitivity to detect colonization (Pinto Dias et al. 2005). The dissection of microhabitats in which triatomines feed and shelter, such as tree holes, palm crowns, bromeliads, rock piles, burrows and bird nests, has been effective to capture sylvatic specimens, but it requires important sampling efforts, human resources and sometimes generates negative impact on the environment. Light traps have the disadvantage of attracting only hungry adults, although it has been possible to capture some species that are otherwise difficult to collect (Noireau and Dujardin 2001; Vazquez-Prokopec et al. 2004). Traps with animal bait, such as the Noireau adhesive trap, which uses a mouse as bait, were successfully used in sylvatic ecotopes (Abad-Franch et al. 2000; Gürgel-Gonçalves et al. 2003; Noireau et al. 1999). However, it is expensive due to host maintenance and some authors reported that its efficiency depends upon the triatomine species studied and the biotic region of study (Reyes-Novelo et al. 2012). As none of the explored methods show reasonable sensitivity, there is a need to develop a sensitive detection method for entomological vigilance.

Two types of non-live baited traps were also evaluated based on semiochemicals (Rojas de Arias et al. 2012) and yeast. The host orientation behavior of triatomines is controlled by physical and chemical signals, including olfactory clues such as carbon dioxide. The carbon dioxide released by the *Saccharomyces cerevisiae* cultures, is a chemical signal indicative of a food source for hematophagous insects and therefore it can evoke both behavioral

responses: activation and attraction of the triatomines to the source (Lazzari et al. 2013; Guerenstein and Lazzari 2009).

Several authors have demonstrated the effectiveness of yeast traps to attract and capture *T. infestans* under both laboratory (Guerenstein et al. 1995; Barrozo and Lazzari 2004, 2006) and natural conditions (Lorenzo et al. 1998, 1999; Bacigalupo et al. 2006). Other studies have also demonstrated that yeast traps are a useful tool for the detection of potential new sylvatic habitats of *T. infestans* as well as other triatomine species and they can be a suitable alternative for their control (Bacigalupo et al. 2006; Botto-Mahan et al. 2002).

Previous studies using yeast traps for triatomine detection evaluated the device during a fixed time, and generally using nymph triatomines.

Within the exploration efforts to find a method that improves the detection sensitivity of triatomines, we report a study that measured the temporal variation in attractiveness of an adhesive yeast trap for adults and nymphs of *T. infestans* under laboratory conditions.

Materials and methods

Experimental setting

The study was designed as an experiment involving the release of insects in an arena containing a baited trap (with yeast) and a non-baited trap (without yeast, control trap).

The experimental arena measured 100 x 80 x 80 cm, with kraft paper ground and non-climbable walls. Control and yeast traps were placed at opposite sides of the arena and folded paper as an artificial shelter (15 x 10 cm) was placed in the center.

During the assay, 8 insects were released in the area center and only after the insects had hidden in the folded shelter the traps were placed in the arena. After 3, 6 and 24 hours of the beginning of the experiment, the number of triatomines captured in the yeast trap, those adhered to the control, the loose ones in the arena and those that remained hidden in the refuge were counted. Each time the number of bugs was recorded in the traps, the replacement of the attached bugs was made, making sure that there were always 8 individuals in the experimental box.

Some tested insects were used up to 2 times with a difference of at least ten days between tests to ensure independence.

We performed seventeen series of assays for adults of *T. infestans* and thirteen series of assays for 4th and 5th nymph instars. During the test, the triatomines were able to move freely throughout the experimental arena. Assays were conducted at room temperature (approximately 25°C ± 2°C), in darkness starting at 10.00 am and finishing 24 hours later. The position of the control and the experimental traps was changed randomly in the successive trials, to compensate for possible external asymmetries.

Traps

The experimental adhesive yeast traps (Figure 1) consisted of a plastic container of 500 cm³, with a perforated cover containing a solution of 5 g of dry yeast LEVEX[®] + 10 g of sucrose + 100 ml of water. For the purpose of this work, we used the same concentration employed by Bacigalupo et al. (2006), as this concentration proved to be effective for the capture of wild *T. infestans* colonies in field studies.

The control trap had the same design of the yeast trap although it only contained a solution of sucrose (10 g of sucrose + 100 ml of water). The plastic container of both the yeast trap and the control trap was surrounded by a rectangle of corrugated plastic, covered with adhesive paper tape Doble A®. The corrugated plastic increases the adhesive surface and offers a refuge site. This yeast trap design is simple and cheap, easy to transport due to its low weight and volume and it can be used either horizontally or vertically so that it can be put in a great variety of sylvatic and peridomestic ecotopes.

Fig 1 Front and side view of the yeast trap prototype used to test the attractiveness of the yeast culture for *Triatoma infestans*

Insects

The triatomines used in the assays were 180 adults and 103 nymphs (83 fifth nymphal instar and 20 fourth nymphal instar) of *T. infestans*. The insects came from colonies reared during many generations (>25) in the Centro de Referencia de Vectores from the Servicio Nacional de Chagas (CeReVe-SNCh), located in Santa María de Punilla (Córdoba, Argentina).

All the triatomines were fed on chicken and kept under a natural illumination cycle under room temperature conditions ($26^{\circ}\text{C} \pm 5^{\circ}\text{C}$ and at $40 \pm 60\%$ relative humidity). The insects were starved for approximately 15 to 30 days before the experiments.

Data analysis

Although the results of this type of experiment has been analyzed comparing proportions of trapped insects through ANOVA tests to detect contrast signification (e.g. Guerenstein et al. 1995), we adopted the effect estimation approach (Cumming 2012) instead of significance test. To evaluate the attractiveness of the yeast solution we used generalized linear models (GLMs) with Poisson error distribution. We consider alternative hypothesis that would explain the variation in the number of bugs captured in the yeast trap (response variable) as a function of predictor variables including the insect stages (adults or nymphs), the time intervals at which the response variable was recorded (3, 6 and 24 hours after the beginning of the assay) and the date of the experiment. The latter was considered as a temporal control to detect possible asymmetries between the assays. The candidate models considered the individual effects of each predictor on the response variable as well as joint models evaluating the additive effects and the interactions of the possible combinations. The GLMs were fitted through maximum likelihood and their relative performance were measured with Akaike's information criterion (AICc), using the function `aictab` of the package `AICmodavg` (Mazerolle 2017) in the R software version 3.4.3 (R Core Team 2017). We assessed the significance of the effects by comparing size, unconditional standard error and 95% confidence interval. The effect-size estimates for each factor were averaged, using the `modavg` function from the

package AICmodavg in the R software version 3.4.3 (Mazerolle 2017), for all coefficients included in models that showed a difference in AIC values ≤ 3 with the model that showed the lowest AIC.

Results

In every replicate an average of 7.58 (95% CI [7.43;7.73]) insects, either nymph instar or adults, moved out of the central refuge. On average, 3.53 (95% CI [2.97;4.09]) adults and 2.71 (95% CI [1.97;3.46]) nymph instar that moved out of the refuge were attracted by the yeast trap, and only 5% of adults and 7% of nymph instars remained in the refuge after 24 hs.

Considering that the 8 insects released in each assay were able to choose among 4 sites (yeast trap, control trap, refuge and out of refuge), 2 is the expected number of insects to be counted at each time interval in any of these sites under the hypothesis of random selection. The median number of adults captured in the yeast trap was always > 2 (Fig. 2A), whereas in the nymph group, the median number of captured insects in the yeast trap was 2 after the first 3 hours (Fig. 2B).

Fig 2 Number of *T. infestans* adults (A) and nymph instars (B) captured in the adhesive yeast trap 3, 6 and 24 hours after the beginning of the assay. The dotted line indicates the expected number of insects under the hypothesis of random selection of sites by the triatomines.

From the nine candidate models of the multimodel inference approach, three of them described equally well the results ($\Delta AICc \leq 3.0$) (Table 1). Two of the models included the interaction between life stage factor and the time as predictors. The presence of interaction indicates that the time effect depends on the life stage of the triatomine, meaning that attraction to the carbon dioxide

source at 3, 6 and 24 hours after the beginning of the assay, differ between adults and nymphs of *T. infestans*.

Table 1 Model set. Exhaustive list of all GLMs considered in this study to explain variation of the number of insects recorded in the yeast trap (Y).

Model structure	$\Delta AICc$
Y (life stage + time + life stage * time)	0.000
Y (life stage + time + life stage * time + date)	1.377
Y (life stage)	2.715
Y (life stage + date)	4.495
Y (life stage + time)	4.620
Y (life stage + time + date)	6.365
Y (date)	6.825
Y (time)	6.903
Y (time + date)	8.767

$\Delta AICc$, represents the difference in the value of the Akaike's information criterion (AIC) with respect to the AIC value of the best candidate model

+, additive effect of the factor

*, interaction effect of the factors

After detecting the life stage x time interaction, the response variable was modelled for nymph instars and adults separately using GLMs with Poisson link, considering the effects of date, time and their additive effects. Models for the nymph stage showed no effect of date (coefficient estimate = 0.01, 95%CI [0;0.01]), but a negative effect of time (coefficient estimate= -0.04, 95%CI [-0.07; -0.01]). Models for the adult stage showed no effect of date or time, and

an interception estimate of 1.17 (95%CI [0.93;1.41]), meaning that number of adults remained constant during the experimental periods between 2.5 and 4.1.

Discussion

Our results confirm the effectiveness of CO₂ liberated by a small yeast culture to attract *T. infestans*. The detection of natural host odour blends or a single constituent (CO₂) tends to increase the triatomine locomotor activity and trigger both behavioral responses of activation and attraction (Guerenstein and Lazzari 2009). For the first time we report the comparative attraction during 24 hours of *T. infestans* nymph instars and adults to the CO₂ liberated by a small yeast culture. Our study shows that *T. infestans* adults and nymphs had higher locomotion activity after the experiment started than insects in similar experiments performed using other triatomine species (*T. dimidiata*, *T. pallidipennis*, *T. brasiliensis*, *T. sordida*, *T. pseudomaculata*), where more than 50% of the nymphs remained in the central refuge (Pires et al. 2000; Pimenta et al. 2007), compared to 5% of adults and 7% of nymphs in the present study. The proportion of nymphs captured in the yeast trap (34%, out of the total used in the assays) was similar to that obtained for *T. infestans* in a previous study (44%, Guerenstein et al. 1995).

Our study shows that the attractiveness to the CO₂ liberated by a small yeast culture depends upon the age classes of the insects. On average, adults were more attracted to the yeast trap than nymphs. During the 24 hours of experimental period, a constant number of adults (2.5 - 4.1) was attracted and captured in the yeast trap. However, our results showed a different behavior of nymphs, that presented attraction to CO₂ only the first three hours of the assay, and then declining significantly over time.

Even though it has been demonstrated that the attractiveness and orientation towards CO₂ by *T. infestans* is limited to a temporal window at the beginning of the night (Barrozo and Lazzari 2004), we observed that if a source of CO₂ is offered, starved adults of *T. infestans* can respond to the chemical stimulus long before the beginning of the scotophase, under laboratory conditions. This behavior might be based on the possibility that the sensitivity to one specific odour is not “switched-off” outside the temporal window associated with the search of food, as Bodin et al. (2008) suggested.

The results presented in this work add new questions about the mechanisms that modulate the CO₂ attraction and the possible influence of vital stage of *T. infestans*.

Several studies evidence that the orientation response to stimuli is more complex than believed. This process is influenced mainly by internal factors, such as circadian clocks (Barrozo and Lazzari 2004), as well as the triatomine’s physiological state concerning its nutritional status, the proximity to oviposition and the moult cycle (Bodin et al. 2008,2009). Undoubtedly, the orientation response to stimuli and chemosensitivity deserves further studies to be fully understood. Not only to comprehend the biological basis but also because it is fundamental for the improvement of detection techniques or development of new detection tools.

The interpretation of the results of this study should take into account that the nymph’s age and the reproductive status of females were not measured. Further laboratory studies should be carried out to confirm and understand the mechanisms determining the relationship between attraction to CO₂ and the life stage. The adhesive yeast-baited trap used in this work should be tested under

field conditions to determine its sensitivity and easy-of-use before it could be recommended for the use of routine activities of triatomine surveillance.

References

Abad-Franch F, Noireau F, Paucar, A, Aguilar H M, Carpio C, Racines VJ (2000) The use of live-bait traps for the study of sylvatic *Rhodnius* populations (Hemiptera: Reduviidae) in palm trees. *Papers in Entomology*. Museum, University of Nebraska State at Digital Commons.

Abad-Franch F, Vega MX, Rolón MS, Santos WS, Rojas de Arias A (2011) Community Participation in Chagas Disease Vector Surveillance: Systematic Review. *PLoS NTD* 5(6): e1207 doi:10.1371/journal.pntd.0001207

Bacigalupo A, Torres-Pérez F, Segovia V, García A, Correa JP, Moreno L, Arroyo P, Cattán PE (2010) Sylvatic foci of the Chagas disease vector *Triatoma infestans* in Chile: description of a new focus and challenges for control programs. *Mem Inst Oswaldo Cruz* 105(5):633–41

Bacigalupo AB, Segura JA, García A, Hidalgo J, Galuppo SG, Cattán PE (2006) Primer hallazgo de vectores de la enfermedad de Chagas asociados a matorrales silvestres en la Región Metropolitana, Chile. *Rev Med Chil* 134:1230-1236

Barrozo RB, Lazzari CR (2004) Orientation behavior of the blood-sucking bug *Triatoma infestans* to short chain fatty acids: synergistic effect of L-lactic acid and carbon dioxide. *Chem Senses* 29:833-841

Barrozo RB, Lazzari CR (2006) Orientation response of hematophagous bugs to CO₂: the effect of the temporal structure of the stimulus. *J Comp Physiol* 192:827–831

Bodin A, Barrozo RB, Couton L, Lazzari C (2008) Temporal modulation and adaptive control of the behavioural response to odours in *Rhodnius prolixus*. J Insect Physiol 54:1343–1348

Bodin A, Vinauger C, Lazzari CR (2009) Behavioral and physiological state dependency of host seeking in the bloodsucking insect *Rhodnius prolixus*. J Exp Biol 212, 2386-2393

Botto-Mahan C, Cattán PE, Canals M (2002) Field tests of carbon dioxide and conspecifics as baits for *Mepraia spinolai*, wild vector of Chagas disease. Acta Trop 82 377–380.

Bustamante-Gómez M, Gonçalves Diotaiuti L, Gorla DE (2016) Distribution of pyrethroid resistant populations of *Triatoma infestans* in the Southern Cone of South America. PLoS NTD 10(3).e0004561. doi.org/10.1371/journal.pntd.0004561

Cavallo MJ, Amelotti I, Gorla DE (2016) Invasion of rural houses by wild Triatominae in the arid Chaco. J Vector Ecol 41(1):97 -102

Ceballos LA, Piccinali RV, Marcet PL, Vazquez-Prokopec GM, Cardinal MV, Schachter-Broide J, Dujardin JP, Dotson EM, Kitron U, Gürtler RE (2011) Hidden sylvatic foci of the main vector of Chagas disease *Triatoma infestans*: Threats to the vector elimination campaign? PLoS NTD 5(10): e1365. doi.org/10.1371/journal.pntd.0001365

Cecere MC, Leporace M, Fernández MP, Zárate JE, Moreno C, Gürtler Re, Cardinal MV (2016) Host- feeding sources and infection with *Trypanosoma cruzi* of *Triatoma infestans* and *Triatoma eratyrusiformis* (Hemiptera: Reduviidae) from the Calchaqui Valleys in Northwestern Argentina. J Med Entomol 53(3):666-673

Cecere MC, Vazquez-Prokopec GM, Gürtler RE, Kitron U (2006) Reinfestation sources for Chagas disease vector, *Triatoma infestans*, Argentina. *Emerg Infect Dis* 12(7):1096-1102

Cumming G (2012) Understanding the new statistics. Effect size, confidence intervals and meta-analysis. New York: Routledge.

Dias JCP, Silveira AC, Schofield CJ (2002) The impact of Chagas disease control in Latin America - A Review. *Mem Inst Oswaldo Cruz* 97(5):603-612

Giraldez EL, López Ureta MP, Nepote M (2011) Acercamiento de triatomíneos silvestres hacia la vivienda rural e intervenciones sobre el medio ambiente sin planificación adecuada: potencial riesgo en la transmisión de la enfermedad de Chagas. Resúmenes V Congreso Iberoamericano sobre desarrollo y ambiente (VCISDA)

Gómez-Núñez JC (1965) Desarrollo de un nuevo método para evaluar la infestación intradomiciliaria por *Rhodnius prolixus*. *Acta Cient Venez* 16:26-31

Guerenstein PG, Lazzari CR (2009) Host-seeking: How triatomíneos acquire and make use of information to find blood. *Acta Trop* 110(2-3):148-58

Guerenstein PG, Lorenzo MG, Núñez JA, Lazzari CR (1995) Baker's yeast, an attractant for baiting traps for Chagas' disease vectors. *Experientia* 51: 834–837

Gürgel-Gonçalves R, Palma AR, Menezes MN, Leite RN, Cuba CA (2003) Sampling *Rhodnius neglectus* (Triatominae) in *Mauritia flexuosa* palm trees (Arecaceae): A field study in the Brazilian Savanna. *Med Vet Entomol* 17:347-9

Gürtler RE, Cecere MC, Canale DM, Castañera R, Chuit R, Cohen JE (1999) Monitoring house reinfestation by vectors of Chagas disease: a comparative trial of detection methods during a four-year follow-up. *Acta Trop* 72:213–234

Gürtler RE, Petersen RM, Cecere MC, Schweigmann NJ, Chuit R, Gualtieri JM, Wisnivesky-Colli C (1994) Chagas disease in northwest Argentina: risk of domestic reinfestation by *Triatoma infestans* after a single community-wide application of deltamethrin. *Trans R Soc Trop Med Hyg* 88:27-30

Lazzari CR, Pereira MH, Lorenzo MG (2013) Behavioral biology of Chagas disease vectors. *Mem Inst Oswaldo Cruz* 108(1):34-47

Lent H, Wygodzinsky P (1979) Revision of the Triatominae (Hemiptera, Reduviidae), and their significance as vectors of Chagas disease. *Bull Am Mus Nat Hist* 163:123-520

Lorenzo, M.G. (1998). *Triatoma infestans* can be captured under natural climatic conditions using yeast-baited traps. *Acta Trop* 30;70(3):277-84

Lorenzo MG, Lazzari CR (1998) Activity pattern in relation to refuge exploitation and feeding in *Triatoma infestans* (Hemiptera:Reduviidae). *Acta Trop* 70:163–170

Lorenzo MG, Manrique G, Pires HHR, de Brito Sánchez MG, Diotaiuti L, Lazzari CR (1999) Yeast culture volatiles as attractants for *Rhodnius prolixus*: electroantennogram responses and captures in yeast-baited traps. *Acta Trop* 72(1):119-124

Mazerolle MJ (2017) AICcmodavg: Model selection and multimodel inference based on (Q)AIC(c). R package version 2.1-1. <https://cran.r-project.org/package=AICcmodavg>.

McGwire K, Segura EL, Scavuzzo M, Gómez AA, Lamfri M (2006) Spatial pattern of reinfestation by *Triatoma infestans* in Chancaní, Argentina. *J Vector Ecol* 31(1):17-28

Noireau F, Dujardin JP (2001) Flight and nutritional status of sylvatic *Triatoma sordida* and *Triatoma guasayana*. Mem Inst Oswaldo Cruz 96(3):385-389

Noireau F, Flores R and Vargas (1999) Trapping sylvatic Triatominae (Reduviidae) in hollow trees. Trans Roy Soc Trop Med Hyg 93:13-4

Noireau F, Flores R, Gutierrez T, Dujardin JP (1997) Detection of sylvatic dark morphs of *Triatoma infestans* in the Bolivian Chaco. Mem Inst Oswaldo Cruz 92:583-4

Piccinalli RV, Marcet PL, Ceballos LA, Kitron U, Gürtler RE, Dotson EM (2011) Genetic variability, phylogenetic relationships and gene flow in *Triatoma infestans* dark morphs from the Argentinean Chaco. Infect Genet Evol 11(5):895-903 doi: 10.1016/j.meegid.2011.02.013.

Pimenta FE, Diotaiuti L, Lustosa-Lima AC, Lorenzo MG (2007) Evaluation of cultures of *Saccharomyces cerevisiae* as baits for *Triatoma dimidata* and *Triatoma pallidipennis*. Mem Inst Oswaldo Cruz 102(2): 229-231

Pinto-Dias JC, Vieira EP, Tadashi H, Azeredo BVM (2005) About the use of “Maria” bio-sensors in the activities of epidemiological surveillance against Chagas’ disease in the north of Minas Gerais State. Rev Soc Bras Med Trop 38(5):377-382

Pires HRH, Lazzari CR, Diotaiuti I, Lorenzo MG (2000) Performance of yeast-baited traps with *Triatoma sordida*, *Triatoma brasiliensis*, *Triatoma pseudomaculata* and *Panstrongylus megistus* in laboratory assays. PJP 7(6): 384-388

R Core Team (2017) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>

Rabinovich JE (1972) Vital statistics of Triatominae (Hemiptera: Reduviidae) under laboratory conditions. I. *Triatoma infestans* Klug. J Med Entomol 9:351-370

Reyes-Novelo E, Salazar-Fleites C, Chi-Poot M, Chan-Espinoza D, Pérez-Canul S, Ruiz-Piña H (2012) Eficacia de trampas cebadas con ratón en la colecta de *Triatoma dimidiata* (Latreille, 1811) (Hemiptera: Reduviidae) silvestre en el norte de Yucatán. Conference: Entomología Mexicana at Mexico. January 2012. Volume 11

Rojas de Arias A, Abad-Franch F, Acosta N, López E, González N, Zerba, E, Tarelli G, Masuh H (2012) Post-Control Surveillance of *Triatoma infestans* and *Triatoma sordida* with Chemically-Baited Sticky Traps. PLoS NTD 6(9): e1822. doi:10.1371/journal.pntd.0001822

Rolón M, Vega MC, Román F, Gómez A, Rojas de Arias A (2011) First report of colonies of sylvatic *Triatoma infestans* (Hemiptera: Reduviidae) in the Paraguayan Chaco, using a trained dog. PLoS NTD 5(5):e1026. doi:10.1371/journal.pntd.0001026

Schofield CJ, Jannin J, Salvatella R (2006) The future of Chagas disease control. Trends Parasitol 12:583–588.

Vazquez-Prokopec, G.M., Ceballos, L.A., Kitron, U. and Gürtler, R.E. (2004). Active dispersal of natural populations of *Triatoma infestans* (Hemiptera: Reduviidae) in Rural Northwestern Argentina. Mem Inst Oswaldo Cruz 96: 385-389

Wisnivesky-Colli C, Paulone I, Pérez A, Chuit R, Gualtieri J, Smith A, Segura EL (1987) A new tool for continuous detection of the presence of triatomine bugs, vectors of Chagas disease, in rural households. Med 47(1):45-50.

World Health Organization (2015) Chagas disease in Latin America: an epidemiological update based on 2010 estimates. Weekly epidemiological record. No. 6, 90, 33–44.

Acknowledgement

The authors thank Emma Bosco and Elisa Barbero for their valuable contribution with the design and development of the adhesive yeast trap. We also thank Raul Stariolo from the Centro de Referencia de Vectores from the Servicio Nacional de Chagas (CeReVe-SNCh, Santa María de Punilla Córdoba, Argentina) for the provision of the insects for experimentation. MC and DEG are supported by the National Research Council of Argentina (CONICET).

Funding

This work was supported by a grant given by the National University of Córdoba to LBC.

Conflict of Interest

The authors declare that they have no conflict of interest.