Sun exposure drives Antarctic cryptoendolithic community structure and composition

Claudia Coleine¹, Jason E. Stajich², Laura Zucconi¹, Silvano Onofri¹, Laura Selbmann^{1,3*}

¹*Department of Ecological and Biological Sciences (DEB), University of Tuscia, Viterbo, Italy*

²*Department of Microbiology and Plant Pathology and Institute of Integrative Genome Biology, University of California, Riverside, CA, USA*

³*Italian National Antarctic Museum (MNA), Mycological Section, Genoa, Italy*

*Corresponding author: Laura Selbmann (selbmann@unitus.it)

Abstract

The harsh environmental conditions of the ice-free regions of Continental Antarctica are considered one of the closest Martian analogues on Earth. There, rocks play a pivotal role as substratum for life and endolithism represents a primary habitat for microorganisms when external environmental conditions become incompatible with active life on rock surfaces. Due to the thermal inertia of rock, the internal airspace of lithic substratum is where microbiota find a protected and buffered microenvironment, allowing life to spread throughout these regions with extreme temperatures and low water availability. The high degree of adaptation and specialization of the endolithic communities makes them highly resistant but scarcely resilient to any external perturbation and thus, any shifts in microbial community composition may serve as early-alarm systems of environmental perturbation, including climate change.

Previous research concluded that altitude and distance from sea do not play as driving factors in shaping microbial abundance and diversity, while sun exposure was hypothesized as significant parameter influencing endolithic settlement and development. This study aims to explore our hypothesis that changes in sun exposure translate to shifts in community composition and abundances of main biological compartments (fungi, algae and bacteria) in the Antarctic cryptoendolithic communities. We performed a preliminary molecular survey, based on DGGE and qPCR techniques, of 48 rocks with varying sun exposure, collected in Victoria Land along an altitudinal transect from 834 to 3100 m a.s.l.

Our findings demonstrate that differences in sun radiation between north and south exposure influence temperature of rocks surface, availability of water and metabolic activity and also have significant impact on community composition and microbial abundance.

Keywords: Antarctica, Cryptoendolithic communities, Sun exposure, DGGE, qPCR, Sampling

Introduction

The rate of warming due to increased levels of greenhouse gases in the atmosphere is amplified with elevation and at high latitudes due to the polar amplification phenomenon. Polar amplification predicts that as global mean temperature climbs, the greatest warming will occur at the Polar regions (Bekryaev et al., 2010). The impact of climate change is, therefore, particularly intense at the Poles and in mountain environments, nowadays known as the Third Pole (Yao et al., 2012; Yang et al., 2014). As consequence of warming, range-restricted species, particularly polar and mountain top species, have already shown severe contractions and have been the first groups in which entire species have gone extinct due to recent climate change (Parmesan, 2006; Descamps et al., 2017; Bhatta et al., 2018).

The Arctic regions are melting faster than the Antarctic and, if the heating trend continues, studies forecast an ice-free North Pole in summer by mid-century. Strong evidence of warming in Antarctica, is also documented; researchers from the British Antarctic Survey report a warming trend up to 2.5 °C since the 1940s in the Antarctic Peninsula and Maritime Antarctica, the most rapid changes in mean air temperatures on Earth (e.g. Turner et al. 2005, 2007). Previous research reported an apparent contrast between strong warming of the Antarctic Peninsula and slight cooling of the Antarctic continental interior; there are now evidences that significant warming extends well beyond the Antarctic Peninsula and covers most of West Antarctica with a warming exceeding 0.1 °C per decade over the past 50 years (Steig et al., 2009).

Progressions of this warming trend will influence Antarctica's biodiversity by the introduction of allochthonous, competitive species and the consequent extinction of highly specialized and less competitive autochthonous ones (Farrell et al., 2011; Olech and Chwedorzewska, 2011; Selbmann et al., 2012), which will have impacts on the ecosystem functions of glaciers, freshwater systems and atmosphere. Thus, it is urgent to develop a strong base knowledge for Antarctic terrestrial ecosystems and use this to identify ecosystem changes (NAS, 2011).

Endolithism is a specialized colonization by microbes to enable dwelling inside airspaces of rocks. This lifestyle represents adaptation at the edge inhabitable conditions. Airspaces within rocks offer to microbiota a protected and buffered microenvironment, allowing life to expand into different extreme conditions, i.e., hot and cold deserts or geothermal environments (Friedmann and Ocampo, 1976; Friedmann, 1982; Bell, 1993; Walker et al., 2005). Rocks are the prevailing substratum for life in the ice-free areas of Antarctica, supporting the highest standing biomass in the Antarctic ice-free desert and mountain tops emerging from the Polar Plateau (Cowan and Tow, 2004; Cary et al., 2010; Cowan et al., 2014; Selbmann et al., 2017). Endolithic microbial life represents the predominant recorded life-form in these areas (Nienow and Friedmann, 1993). The harsh conditions are considered one of the closest analogues to Mars on Earth (Quintal et al., 2018).

Different from soil microbial communities of these areas, endolithic microbes develop as very tiny and stable communities thanks to the stable and concrete nature of rocks. Various typologies have been observed for the microbial composition and the most complex and widespread are the lichen-dominated communities (Nienow and Friedmann, 1993). These self-supporting microbial ecosystems are composed of algae, mainly lichenized fungi, bacteria and cyanobacteria, many of which are endemic species to the regions (Nienow and Friedmann, 1993; Selbmann et al., 2005, 2008; Egidi et al., 2014). The high degree of adaptation and specialization in exploiting such ultimate niches makes these communities very susceptible to physical and climatic alteration (Selbmann et al., 2017) and any shift in microbial communities composition may serve as early-alarm system of environmental perturbation. Based on a substantial sampling of different typologies (volcanic and sedimentary) of colonised rocks in the Victoria Land, Antarctica, sandstone was determined to be the most suitable substratum for microbial endoliths, allowing them to spread and persist under stronger environmental pressure (Zucconi et al., 2016; Selbmann et al., 2017). To get clues to the future effects of climate change on these unique ecosystems, the response of the communities to increasing environmental pressure, due to altitude (from sea level to 3600 m a.s.l.) and sea distance (up to 100 km) was recently

investigated. The results suggested that these two parameters alone do not play as driving factors in shaping the community diversity and composition, and highlighted the needs to consider additional environmental parameters to elucidate how, in the long run, future environmental changes will impact these unique communities (Coleine et al., 2018a).

With this in mind, a new sampling campaign (Dec.2015- Jan. 2016) of sandstones was performed in the Victoria Land (Antarctica) in the frame of the Italian National Program for Antarctic Researches (PNRA), along an altitudinal gradient from 834 to 3100 m a.s.l., adding sun-exposure as new parameter to investigate its influence on endolithic settlement and development (Friedmann and Weed, 1987). Rocks with varying sun exposure (north and south faced surfaces) were collected in all the localities surveyed. Based on the prior experiences (Selbmann et al., 2017; Coleine et al., 2018a), it is suggested that sampling strategy and environmental parameters considered may have important consequences on a proper and exhaustive biodiversity description; thus, before proceeding with metabarcoding and metagenomics experiments to develop a detailed picture of microbial diversity and functionality, we have selected a suite of more rapid and cheaper complementary approaches. We have used Denaturing Gradient Gel Electrophoresis (DGGE) and quantitative PCR, to test hypothesis that changes in sun exposure impacts community composition and abundances of primary biological assemblages.

Materials and methods

Study area

Eight localities were visited in Victoria Land (Continental Antarctica) during the XXXI Italian Antarctic Campaign (Dec. 2015 - Jan. 2016). North and south exposed sandstone samples were collected along a latitudinal transect from 74°10'44.0''S 162°30'53.0''E (Mt. New Zealand, Northern Victoria Land) to 77°54'43.6''S 161°34'39.3''E (Knobhead, Southern Victoria Land), ranging from 834 (Battleship Promontory, Southern Victoria Land) to 3100 m a.s.l. (Mt. New Zealand) (Table 1). All rock samples were excised aseptically and collected in triplicate, transported at -20 °C at the Tuscia University (Viterbo, Italy) and stored at Mycological Section of the Italian Antarctic National Museum (MNA) until downstream analysis.

125

126

127 **Table 1.** Table lists sampling details of eight visited localities in Victoria Land, Antarctica: sun exposure,

Locality	Sample	Sun	Altitude	Temperature	Humidity	Coordinates
----------	--------	-----	----------	-------------	----------	-------------

128 altitude, air temperature (measured when sampling), relative humidity and geographic coordinates.

Battleship Promontory	1	North	834	-4.4	22.9	76°54'04.0''S 160°54'36.6''E
Battleship Promontory	2	North	834	-4.4	22.9	76°54'04.0''S 160°54'36.6''E
Battleship Promontory	3	North	834	-4.4	22.9	76°54'04.0''S 160°54'36.6''E
Battleship Promontory	1	South	834	-4.4	22.9	76°54'04.0''S 160°54'36.6''E
Battleship Promontory	5	South	834	-4.4	22.9	76°54'04.0''S 160°54'36.6''E
Battleship Promontory	6	South	834	-4.4	22.9	76°54'04.0''S 160°54'36.6''E
Pudding Butte	2	North	1573	-8.5	32.4	75°51'30.2''S 159°58'25.7''E
Pudding Butte	3	North	1573	-8.5	32.4	75°51'30.2''S 159°58'25.7''E
Pudding Butte	4	North	1573	-8.5	32.4	75°51'30.2''S 159°58'25.7''E
Pudding Butte	1	South	1588	-8.5	32.4	75°51'33.0''S 159°58'26.6''E
Pudding Butte	2	South	1588	-8.5	32.4	75°51'33.0''S 159°58'26.6''E
Pudding Butte	3	South	1588	-8.5	32.4	75°51'33.0''S 159°58'26.6''E
Siegfried Peak	2	North	1620	-9.3	52.8	77°34'43.3''S 161°47'11.7''E
Siegfried Peak	3	North	1620	-9.3	52.8	77°34'43.3''S 161°47'11.7''E
Siegfried Peak	5	North	1620	-9.3	52.8	77°34'43.3''S 161°47'11.7''E
Siegfried Peak	3	South	1620	-6.8	54.9	77°34'39.9''S 161°47'17.4''E
Siegfried Peak	5	South	1620	-6.8	54.9	77°34'39.9''S 161°47'17.4''E
Siegfried Peak	6	South	1620	-6.8	54.9	77°34'39.9''S 161°47'17.4''E
Linnaeus Terrace	3	North	1649	-9.6	58.6	77°36'01.3''S 161°05'00.5''E
Linnaeus Terrace	4	North	1649	-9.6	58.6	77°36'01.3''S 161°05'00.5''E
Linnaeus Terrace	6	North	1649	-9.6	58.6	77°36'01.3''S 161°05'00.5''E
Linnaeus Terrace	2	South	1761	-12.6	68.2	77°37'09.9''S 161°11'50.8''E
Linnaeus Terrace	3	South	1761	-12.6	68.2	77°37'09.9''S 161°11'50.8''E
Linnaeus Terrace	4	South	1761	-12.6	68.2	77°37'09.9''S 161°11'50.8''E
Finger Mt.	2	North	1720	-6.4	35.1	77°45'0.9"S 160°44'44.5" E
Finger Mt.	3	North	1720	-6.4	35.1	77°45'0.9"S 160°44'44.5" E
Finger Mt.	6	North	1720	-6.4	35.1	77°45'0.91"S 160°44'42.9"
Finger Mt.	2	South	1720	-6.4	35.1	77°45'10''S 160°44'40''E
Finger Mt.	4	South	1720	-6.4	35.1	77°45'10"S 160°44'44.39.7" E
Finger Mt.	5	South	1720	-6.4	35.1	77°45'10.1"S 160°44'45.1" E
University Valley	2	North	2090	-14.3	18	77°52'28.6''S 160°44'22.6''E
University Valley	3	North	2090	-14.3	18	77°52'28.6"S 160°44'22.6" E
University Valley	4	North	2090	-14.3	18	77°52'29"S 160°44'22.3" E
University Valley	3	South	2200	-11.2	39.1	77°52'21.5''S 160°45'19.2''E
University Valley	5	South	2200	-11.2	39.1	77°52'21.5''S 160°45'19.2''E
University Valley	6	South	2200	-11.2	39.1	77°52'21.5''S 160°45'19.2''E
Knobhead	1	North	2150	-12.5	50	77°54'37.8''S 161°34'48.8''E
Knobhead	2	North	2150	-12.5	50	77°54'37.8''S 161°34'48.8''E
Knobhead	3	North	2150	-12.5	50	77°54'37.8''S 161°34'48.8''E
Knobhead	1	South	2150	-8.9	38.9	77°54'43.6''S 161°34'39.3''E
Knobhead	2	South	2150	-8.9	38.9	77°54'43.6''S 161°34'39.3''E

Knobhead	3	South	2150	-8.9	38.9	77°54'43.6''S 161°34'39.3''E
Mt. New Zealand	1	North	3100	-17.2	47.6	74°10'44.0''S 162°30'53.0''E
Mt. New Zealand	2	North	3100	-17.2	47.6	74°10'44.0''S 162°30'53.0''E
Mt. New Zealand	3	North	3100	-17.2	47.6	74°10'44.0''S 162°30'53.0''E
Mt. New Zealand	1	South	3100	-17.2	47.6	74°10'44.0''S 162°30'53.0''E
Mt. New Zealand	2	South	3100	-17.2	47.6	74°10'44.0''S 162°30'53.0''E
Mt. New Zealand	3	South	3100	-17.2	47.6	74°10'44.0''S 162°30'53.0''E

129

130 *Environmental DNA extraction and Denaturing Gel Gradient Electrophoresis*

131 Environmental DNA was extracted from 0.3 g of crushed rocks using NucleoSpin® Plant II Kit (Macherey-
132 Nagel, GmbH & Co. KG, Duren, Germany) according to the manufacturer's instructions and quantified by
133 Quant-iT dsDNA HS assay kit (Invitrogen molecular probes- Eugene, Oregon, USA). Microbial diversity
134 was screened by Denaturing Gel Gradient Electrophoresis (DGGE). A semi-nested PCR was performed
135 using primers with a GC-clamp (Muyzer et al., 1993); fungal and algal ITS rRNA was amplified from
136 environmental DNA with primers ITS1F-GC/ITS2 (Gardes and Bruns, 1993; White et al., 1990), following
137 the protocol reported in Selbmann et al. (2017), while 341F-GC/518R primers (Muyzer et al., 1993) were
138 utilized for 16S rRNA amplification (Valášková and Baldrian, 2009) (Table 2).

139 Amplicons were purified using NucleoSpin® Gel and PCR Clean-up (Macherey-Nagel, GmbH & Co. KG,
140 Düren, Germany). One-hundred ng of final DNA concentration were loaded into each well for DGGE and
141 runs were performed on DGGE Electrophoresis System (C.B.S. Scientific, Del Mar, California, USA). Gels
142 at 7.5% polyacrylamide (37.5:1 acrylamide:bisacrylamide) were mixed in a gradient maker (Hoefer, USA),
143 using two different concentrations: for fungi and algae (0% and 70%) and for bacteria (0% and 60%). The
144 electrophoresis run was performed for 5 h for fungi and algae and 3.5 h for bacteria in 1× TAE Buffer at a
145 constant temperature of 60 °C and 200 V.

146 Bands were visualized by staining for 40 min with GelRed solution (Biotium Inc, CA, USA) (1.34 g NaCl,
147 66.7 pi GelRed and 200 mL dw) and then visualized with an UV transilluminator (Chemidoc, Bio-Rad).
148 Scanned gels were analyzed with TotalLab Quant Software (Cleaver Scientific Ltd; United Kingdom): bands
149 were assigned and matched automatically and then checked manually. Profile similarity was calculated by
150 determining Dice's similarity coefficient, considering presence/absence of bands, for the total number of lane
151 patterns from the DGGE gels. The similarity coefficients calculated were then used to generate the
152 dendrograms utilizing the clustering method Unweighted Pair Group Method using Arithmetic mean
153 (UPGMA).

154 Triplicate samples were processed and analyzed independently.

155

156

157 *NMDS ordination plots*

158 Multivariate statistical analyses were performed to determine the effects of sun exposure on microbial
159 diversity composition using PAST software (PAleontological STatistics, ver. 2.17) The effect of this abiotic
160 parameter was tested displaying changes in communities' composition with Non-Metric Multidimensional
161 Scaling (NMDS) based both on abundance data (bands intensity), calculating Bray-Curtis distance index and
162 presence-absence data, using Jaccard index (Clarke, 1993). Means of abundance data were square-root
163 transformed and analyses were carried out with 999 permutations as described in Coleine et al. (2018b).
164 NMDS were plotted using the combined occurrence and abundance data of the three replicates from each
165 site. Permutational multivariate analyses of variance (PERMANOVA, $p < 0.05$) based on the Euclidean
166 distance were utilized to establish differences in the two differently sun-exposed rock-inhabiting
167 communities.

Quantitative PCR

Total fungal and bacterial abundances were measured for all samples by quantitative PCR (qPCR) using NS91F (5'-GTCCCTGCCCTTTGTACACAC-3') and ITS51R (5'-ACCTTGTTACGACTTTTACTTCCTC-3') for total fungi and Eub338 (5'-ACTCCTACGGGAGGCAGCAG-3') and Eub518 (5'-ATTACCGCGGCTGCTGG-3') for total bacteria (Fierer et al., 2005). To determine relative gene-copies abundances, standard curves were generated using a 10-fold serial dilution of a plasmid containing a copy of *Cryomyces antarcticus* ITS rRNA gene for fungi and *Escherichia coli* 16S rRNA gene for bacteria. The amplicons were generated in 100 µl reactions, containing 50 µl of 2x PCR BioMix™ (Bioline, London, UK), 5 pmol of both forward and reverse primers, 43 µl of DEPC water and 5 µl template of genomic DNA. For fungi, amplification of ITS region was carried out as follows: 95 °C for 3 min and the 35 cycles of 95 °C 40 s, 55 °C 30 s, and final extension at 72 °C 40 s; while bacterial 16S region was amplified as follows: 95 °C for 3 min and the 35 cycles of 95 °C 40 s, 53 °C 30 s, and final extension at 72 °C 40 s. PCR products were then purified using the NucleoSpin® PCR Clean-up kit (MACHEREY-NAGEL, GmbH & Co. KG), quantified with Qubit dsDNA HS Assay kit and cloned using the pGEM®-T Easy Vector Systems (Promega, Madison, Wisconsin, US). Plasmids were isolated using the NucleoSpin Plasmid kit (Macherey-Nagel, GmbH & Co. KG).

Five standards were utilized for qPCR in series from 10⁷ to 10³ copies.

The 25 µl qPCR reactions contained 12.5 µl iQ™ SYBR® Green Supermix (Bio-Rad, Hercules, California, US), 1 µl of each forward and reverse primers, 0.3 ng of environmental DNA or standard and 9.5 µl nuclease-free water. The reactions were carried out on Quantitative real-time BioRad CFX96™ PCR detection system (Bio-Rad, Hercules, California, US). Primers and amplification protocols are mentioned before. Melting curves were generated to confirm that the amplified products were of the appropriate size. Fungal and bacterial gene copy numbers were generated using a regression equation for each assay relating the cycle threshold (C_t) value to the known number of copies in the standards.

Each assay included no-template controls (NTC). All qPCR reactions were run in triplicate.

Means and standard deviations were calculated and statistical analysis were performed using one-way analysis of variance (Anova) and pairwise multiple comparison procedures, carried out using the statistical software SigmaStat 2.0 (Jandel, USA). Significant differences were calculated by Tukey test (p<0.05).

The fungal-to bacterial ratio was calculated from log-transformed abundance values.

Table 2. PCR primers used for DGGE in the present study.

Amplified group	Primer	Sequence 5 ¹ -3 ¹
Fungi	ITS1F-GC ^a / ITS2	GCACGGGGGGCTTGGTCATTTAG/AG GCTGCGTTCTTCATCGATGC
Bacteria	341F-GC ^b /518R	GCACGGGGGGCCTACGGGAGGC/AG ATTACCGCGGCTGCTGG
CGCCCCGCCGCGCCCCGCGCCCGTCCCGCCGCCCGCCCCG-3 ¹		
^b 5'-CGCCCCGCCGCGCGGGCGGGGCGGGGGCACGGG-3 ¹		

Results

DGGE profiles

DNA was efficiently extracted from almost all rock samples, with few exceptions (i.e. Linnaeus Terrace south) where DNA extraction was not quantifiable, but PCR-DGGE worked out for all samples. Because results were similar when analysing fungi and algae, only data based on fungi are reported. Profile similarity based only on band presence/absence, was calculated by the Dice coefficient and UPGMA was used to create dendrograms describing pattern similarities (Figs. 1, 2). Overall, the banding patterns of the replicates showed a high degree of similarity (data not shown), which was also supported by the dendrograms, generating DGGE patterns that grouped together as most similar to each other. A coherent grouping according to locations and to sun exposure was generated in the clustering based on fungal, and, even more clearly, on bacterial profiles, with few scattered exceptions. In particular, in the clustering generated on fungal profiles, samples were split according to the localities for University Valley, Pudding Butte, Battleship Promontory, Mt. New Zealand and Finger Mt. (Fig. 1). The grouping was also evident in the clustering based on DGGE bacterial profiles (Fig. 2); as for fungi, the splitting was obtained in the communities from Pudding Butte (south), Battleship Promontory (south) and Finger Mt. (south), but also for Linnaeus Terrace and Knobhead for both sun expositions.

NMDS ordination plots

To investigate the similarity of the fungal and bacterial communities' composition amongst different sun exposures, a Non-metric Multi-Dimensional Scaling (NMDS) analysis was computed. NMDS ordination plots were generated both with the only presence-absence matrix using the Jaccard index and with the combined frequency of occurrence using the Bray-Curtis index. Because both approaches produced similar results, we showed results based on abundance only. When stress values are <0.1 , the NMDS plot is considered to be an acceptable representation of the original data; in fact, a stress value below 0.1 indicates a reliable ordination of data, without a real probability of misinterpretation (Clarke, 1993). In this analysis, the stress value was 0.09 for fungi and 0.07 for bacteria, fitting with the ideal ordination. NMDS plots generated from the DGGE profiles of amplified fungal ITS and bacterial 16S rDNA revealed that only small changes occurred among samples collected in the same sun-exposed rock surface ($p>0.05$) and did not exhibit any changes endolithic communities by sampled localities (data not shown). On the contrary, a major change (1-way NPMANOVA, $p<0.05$) occurred between north and south sun-exposed communities, showing a strong structuring of fungal and bacterial communities according to the sun exposure (Fig. 3).

Abundance of fungal and bacterial communities

Both ITS and 16S rRNA gene copy numbers varied mostly significantly ($p < 0.05$) along the sampled sites, ranging from 6.1×10^4 (Knobhead south sun exposure, Dry Valleys, Southern Victoria Land) to 9.8×10^6 fungal copies (Battleship Promontory south sun exposure, Dry Valleys); conversely, the differently north and south exposed rock surfaces of Linnaeus Terrace (Dry Valleys) showed the highest (7.7×10^5) and the lowest (1.3×10^3) bacterial gene-copies, respectively (Fig. 4 and Table 1S). Overall, both fungal and bacterial abundances varied between the two sun-exposures, except in the case of Finger Mt. and University Valley in Dry Valleys, where abundance of the two major biological compartments was similar. Furthermore, microbial abundance was generally higher in north-exposed rocks, with only few exceptions: fungi were most abundant in south sun-exposed samples at Battleship Promontory, while bacteria were larger in number at Mt. New Zealand south (Fig.4).

The fungal-to bacterial ratio (F/B) (based on log-copy numbers) showed a slightly higher dominance of fungi in these ecosystems in both sun-exposed sampled localities and were significantly different in all samples ($p < 0.05$). In north-exposed rocks F/B was 1.24 ± 0.2 (mean $\pm$ SD), while in south-exposed surface it varied between 1.37 ± 0.31 (data not shown).

Discussions

The Antarctic cryptoendolithic communities host among the most resistant microorganisms on Earth, being able to survive under environmental conditions once accounted as incompatible with life (Horowitz et al., 1972).

Recent studies have improved knowledge of the microbial diversity and composition of these ecosystems (Wei et al., 2016; Archer et al., 2017) and provided important insights to clarify the complex relationship between environmental parameters (i.e. altitude and distance from sea) and biodiversity (Selbmann et al., 2017; Coleine et al., 2018a, b). The lack of consistent microbial diversity patterns along altitudinal and sea distance gradients has suggested that new hypotheses about which can act as driving factor shaping biodiversity must put forward (Selbmann et al., 2017; Coleine et al., 2018a).

Towards this end, the aim of this study was to test the potential effects on the structure and composition of the bacterial, fungal and algal assemblages of a new parameter, namely sun exposure, as significant variable influencing endolithic settlement and development, using a preliminary molecular screening approach on a considerable amount of rock samples collected along the Victoria Land (Continental Antarctica).

We found that DGGE-based similarity assessment visualized as dendrograms indicated a clear grouping of samples according mostly by sun exposure, highlighting an evident relationship between microbial composition and this environmental parameter. This trend is consistent in all biological compartments examined. Particularly, a clear clustering was observed for DGGE profiles obtained from samples collected in Pudding Butte, Battleship Promontory, Mt. New Zealand and Finger Mt. in fungi and algae (Fig. 2, data not shown); as for eukaryotes, in bacteria, clear groups were observed in samples belonging to Battleship Promontory, Finger Mt., Pudding Butte, Knobhead and Linnaeus Terrace.

Considering all the data, higher similarities in fingerprinting profiles were observed within samples collected in the same site, with very few scattered exceptions, likely due to variability among samples.

These findings highlighted the effectiveness of sampling strategy applied for the Antarctic Expedition 2015-16 and did not reflect what was observed by Selbmann et al. (2017). In that study, based on a previous sampling (PNRA, Antarctic Expedition 2010-11), the largest Antarctic sampling to date of rocks hosting lithic communities, including different rock typologies (i.e. sandstone, granite, dolerite) from 46 different localities in Victoria Land, was investigated. Results indicated a remarkable local variability found even in rocks from the same site, maybe due to the size of sampled area (about 100 m²) and to variability of rock typologies collected.

DGGE band profiles were analysed by statistical analysis and, consistently to clustering analysis, samples were mainly grouped by sun exposure along the three analysed biological groups. The observed shift in community composition correlated to sun exposure was also confirmed by Non-Metric Multidimensional scaling (NMDS) analysis, which organizes data into 2-D spatial graphs by reducing dimensionality. An apparent gradient in abundance and community composition in response to sun exposure was previously observed in Coleine et al. (2018b), where authors investigated biodiversity and composition of functional groups of fungi in Antarctic endolithic communities and reported the absence of any correlation with altitude and sea distance, while a remarkable variability was observed considering the sun exposure parameter.

It was previously hypothesized that the distribution of endoliths reflects the degree of insulation on the rock surfaces; in northern exposed rocks, environmental conditions are more favourable than southern exposed

faces, and cryptoendolithic colonization is more often observed and favoured (Friedmann, 1977; Friedmann and Weed, 1987). The capability to maintain biological activity may depend on sufficient insulation of the rock to allow an efficient photosynthetic process; moreover, warmer temperatures will allow metabolic activity and more water due to snow melt (McKay and Friedmann, 1985; Deegenaars and Watson, 1998). Our findings support the hypothesis that sun radiation not only affects temperature of rocks surface, availability of water and metabolic activity, but even the biodiversity.

Community structure was also shaped by north and south exposure when fungal and bacterial abundances were estimated with a qPCR assay. In the most localities, fungal (ranging from 6.1×10^4 to 9.8×10^6 gene copies) and bacterial (7.7×10^5 to 1.3×10^3 gene copies) abundances changed significantly according to sun exposure.

We also calculated the fungal:bacterial (F/B) ratio, a metric to assess environmental impacts and the functional implications of microbial communities (Raeymaekers, 2000; Fierer et al., 2005). The F/B dominance was significantly affected by sun exposure and showed a significant dominance of fungi in these microbiomes. Fungi predominate in the south-exposed rocks (with the only exception of Finger Mt. in McMurdo Dry Valleys) where conditions are much more extreme; indeed, we found a greater fungal dominance (F/B, 1.37) respect than north-exposed sites (F/B, 1.24).

We propose that further studies are needed to elucidate the relationship between fungi and bacteria, and to reveal their functions in these ecosystems. Nevertheless, this result is not surprisingly: extreme environments (i.e. the ice-free desert of Victoria Land in Continental Antarctica) are not an only prerogative of archaea and bacteria. Among eukaryotes, fungi (alone or in symbiosis with cyanobacteria or algae forming lichens) are the most versatile and ecologically successful phylogenetic lineage; they evolved to survive and proliferate in the extremes (Magan, 2007), even in habitats normally precluded to the most (Selbmann et al., 2013). Fungi are, for instance, generally more resistant to desiccation than bacteria with hyphae that may cross air-filled soil pores to access nutrients and water (Gordon et al., 2008; de Vries et al., 2012), showing a remarkable ability to survive in stress conditions (e.g. low water availability) of Antarctic ice-free areas (Onofri et al., 2004).

The most remarkable example for stress resistance is, in fact, given by the fungus *Cryomyces antarcticus*, a cryptoendolithic black fungus isolated from the McMurdo Dry Valleys in Antarctica, chosen as the best eukaryotic test organisms for astrobiological investigations for its stunning resistance to temperature cycles ($-20/+20$ °C), high temperature ($+90$ °C) and saline concentration (up to 25% NaCl) (Onofri et al., 2008; 2012; 2015). It was found to resist to ionizing radiation up to 55.81 kGy (Selbmann et al., 2018), while the bacterium *Deinococcus radiodurans*, widely considered the extremophile par excellence and gold-medalist of radiation resistance (Battista et al., 1997; Venkateswaran et al., 2000), survives up to 20 kGy of gamma radiation.

In conclusion, even though DGGE-based approach does not show individual taxa responses and gives limited insight into taxonomy, it is still currently considered a powerful, rapid and costly-effective method for determining shapes in microbial community composition (Zheng et al., 2013; Kovalski Mitter et al., 2018). This technique gave us the advantage to rapidly and inexpensively analyze a great number of rock samples and three biological domains (fungi, algae and bacteria), providing a clear trend of how sun exposure influences cryptoendolithic community biodiversity and composition.

This study expands existing knowledge on the relationship between environmental parameters and Antarctic endolithic biodiversity and examines endolithic community structure. The results inform planning of future Antarctic campaigns to maximize identification of active endolithic communities.

With the advent of -omics approaches such as metabarcoding, metagenomics and metabolomics, we may explore the composition deeply focusing on a targeted selection of rock samples. These samplings will

353 characterize taxa identity and abundance in rocks hosting endolithic communities, examine the complex
 354 community dynamics, their stress-adaptation strategies, and potential functions across an environmental
 355 variation gradient that is influenced by sun exposure.
 356 A detailed picture on how these communities respond to increasing environmental pressures will also give
 357 clues for predicting and monitoring the effects of global change on these unique border ecosystems and their
 358 vulnerable biodiversity.

ACKNOWLEDGMENTS

L.S., C.C. and L.Z. wish to thank the Italian National Program for Antarctic Researches (PNRA) for funding sampling campaigns and researches activities in Italy in the frame of Projects 2009/A1.11, 2013/AZ-17, 2015/AZ1.02 and AMunDsEN PNRA_00006. The Italian Antarctic National Museum (MNA) is acknowledged for financial support to the Mycological Section on the MNA for preserving rock Antarctic samples analysed in this study and stored in the Culture Collection of Fungi from Extreme Environments (CCFEE), University of Tuscia, Italy.

Figure captions

Figure 1. Dendrogram obtained by Unweighted Pair Group Method using Arithmetic average (UPGMA) based on the DGGE profiles of the fungal component of the endolithic communities.

The relationships among samples are based on similarity, evaluated by the Dice coefficient.

Triplicate samples were analysed.

Figure 2. Dendrogram obtained by Unweighted Pair Group Method using Arithmetic average (UPGMA) based on the DGGE profiles of the bacterial component of the endolithic communities.

The relationships among samples are based on similarity, evaluated by the Dice coefficient.

Triplicate samples were analysed.

Figure 3. Non-Metric Multidimensional Scaling (N-MDS) ordination plots for fungal (A) and bacterial endolithic communities (B), calculating the Bray–Curtis index, based on square-root transformed abundance data.

Stress value (A): 0.09.

Stress value (B): 0.07

Figure 4. Abundances of total fungi and bacteria in endolithic communities, as estimated using the qPCR assays. Error bars are the standard errors. Significant differences are calculated by Tukey test with p value<0.05 and indicated by different letters.

384 **Tables 1S.** Fungal and Bacterial gene-copies abundances. Mean and deviation standard are listed in for each
385 site.

Sites	rRNA gene numbers		386
	Fungi	Bacteria	387
Battleship Promontory North sun exposure	3,682,594±876,867	31,142±1,007	
Battleship Promontory South sun exposure	9,771,455±1,338,254	6,444±68	
Pudding Butte North sun exposure	8,584,316±503,423	29,294±443	
Pudding Butte South sun exposure	234,884±57,037	17,665±607	
Siegfried Peak North sun exposure	2,097,544±697,632	406,064±49,436	
Siegfried Peak South sun exposure	688,525±145,531	270,656±38,024	
Linnaeus Terrace North sun exposure	775,910±121,907	768,939±166,909	
Linnaeus Terrace South sun exposure	434,961±110,021	1,341±105	
Finger Mt. North sun exposure	1,569,445±810,611	445,405±133,202	
Finger Mt. South sun exposure	1,506,548±92,730	466,154±111,781	
University Valley North sun exposure	1,179,191±516,881	248,738±114,353	
University Valley South sun exposure	632,387±204,927	248,642±445	
Knobhead North sun exposure	753,157±211,738	421,314±43,770	
Knobhead South sun exposure	60,691±39,833	1,536±637	
Mt. New Zealand North sun exposure	5,249,232±113,652	35,924±358	
Mt. New Zealand South sun exposure	801,119±118,617	51,578±3,629	

References

- Archer SD, de los Ríos A, Lee KC, Niederberger TS, Cary SC, Coyne, KJ, *et al.* (2017) Endolithic microbial diversity in sandstone and granite from the McMurdo Dry Valleys, Antarctica Polar Biol 40(5):997-1006. [10.1007/s00300-016-2024-9](https://doi.org/10.1007/s00300-016-2024-9)
- Battista JR (1997) Against all odds: the survival strategies of *Deinococcus radiodurans*. Ann Rev Microbiol 51(1):203-224. [10.1146/annurev.micro.51.1.203](https://doi.org/10.1146/annurev.micro.51.1.203)
- Bekryaev RV, Polyakov IV, Alexeev VA (2010) Role of polar amplification in long-term surface air temperature variations and modern Arctic warming. J Climate 23(14):3888-3906. [10.1175/2010JCLI3297.1](https://doi.org/10.1175/2010JCLI3297.1)
- Bell RA. (1993) Cryptoendolithic algae of hot semiarid lands and deserts. J Phycol 29:133-139. [10.1111/j.00223646.1993.00133.x](https://doi.org/10.1111/j.00223646.1993.00133.x)
- Bhatta KP, Grytnes JA, Vetaas OR (2018) Downhill shift of alpine plant assemblages under contemporary climate and land use changes. Ecosphere 9(1):e02084. [10.1002/ecs2.2084](https://doi.org/10.1002/ecs2.2084)
- Cary SC, McDonald IR, Barrett JE, Cowan, DA (2010) On the rocks: the microbiology of Antarctic dry valley soils. Nat Rev Microbiol 8:129-138. [10.1038/nrmicro2281](https://doi.org/10.1038/nrmicro2281)
- Clarke KR (1993) Non-parametric multivariate analyses of changes in community structure. Aust J Ecol 18:117-143. [10.1111/j.1442-9993.1993.tb00438.x](https://doi.org/10.1111/j.1442-9993.1993.tb00438.x)
- Coleine C, Stajich JE, Zucconi L, Onofri S, Pombubpa N, Egidì E *et al.* (2018a) Antarctic cryptoendolithic fungal communities are highly adapted and dominated by Lecanoromycetes and Dothideomycetes. Front Microbiol 9. [10.3389/fmicb.2018.01392](https://doi.org/10.3389/fmicb.2018.01392)
- Coleine C, Zucconi L, Onofri S, Pombubpa N, Stajich J, Selbmann L (2018b) Sun exposure shapes functional grouping of fungi cryptoendolithic Antarctic communities. Life 8(2). [10.3390/life8020019](https://doi.org/10.3390/life8020019)
- Cowan D, Tow L (2004) Endangered Antarctic environments. Annu Rev Microbiol 58:649-690. [10.1146/annurev.micro.57.030502.090811](https://doi.org/10.1146/annurev.micro.57.030502.090811)
- Cowan DA, Makhalanyane TP, Dennis PG, Hopkins DW (2014) Microbial ecology and biogeochemistry of continental Antarctic soils. Front Microbiol 5:154. [10.3389/fmicb.2014.00154](https://doi.org/10.3389/fmicb.2014.00154)
- Deegenaars ML, Watson K (1998) Heat shock response in psychrophilic and psychrotrophic yeast from Antarctica. Extremophiles 2(1):41-50.
- Descamps S, Aars J, Fuglei E, Kovacs KM, Lydersen C, Pavlova O *et al.* (2017) Climate change impacts on wildlife in a High Arctic Archipelago–Svalbard, Norway. Global Change Biol 23(2):490-502. [10.1111/gcb.13381](https://doi.org/10.1111/gcb.13381)
- De Vries FT, Liiri ME, Bjørnlund L, Bowker MA, Christensen S, Setälä HM, Bardgett RD (2012) Land use alters the resistance and resilience of soil food webs to drought. Nat Clim Change 2(4):276. [10.1038/nclimate1368](https://doi.org/10.1038/nclimate1368)
- Egidì E, De Hoog GS, Isola D, Onofri S, Quaedvlieg W, De Vries, *et al.* (2014) Phylogeny and taxonomy of meristematic rock-inhabiting black fungi in the dothideomycetes based on multi-locus phylogenies. Fungal Divers 65:127-165. [10.1007/s13225-013-0277-y](https://doi.org/10.1007/s13225-013-0277-y)
- Farrell RL, Arenz BE, Duncan SM, Held BW, Jurgens JA, Blanchette RA (2011) Introduced and indigenous fungi of the Ross Island historic huts and pristine areas of Antarctica. Polar Biol 34:1669-1677.
- Fierer N, Jackson JA, Vilgalys R, Jackson RB (2005) Assessment of soil microbial community structure by use of

426 taxon-specific quantitative PCR assays. *Appl Environ Microbiol* 71(7):4117-4120. [10.1128/AEM.71.7.4117-4120.2005](https://doi.org/10.1128/AEM.71.7.4117-4120.2005)

427 Friedmann EI, Ocampo R (1976) Endolithic blue-green algae in dry valleys-primary producers in Antarctic desert
428 ecosystem. *Science* 193, 1247-1249 [10.1126/science.193.4259.1247](https://doi.org/10.1126/science.193.4259.1247)

429 Friedmann EI (1977) Microorganisms in Antarctic desert rocks from dry valleys and Dufek Massif. *Antarctic J US*
430 12(4): 26-29.

431 Friedmann EI (1982) Endolithic microorganisms in the Antarctic cold desert. *Science* 215:1045-1053

432 Friedmann EI, Weed R (1987) Microbial trace-fossil formation, biogenous, and abiotic weathering in the Antarctic cold
433 desert. *Science* 236(4802):703-705. [10.1126/science.215.4536.1045](https://doi.org/10.1126/science.215.4536.1045)

434 Gardes M, Bruns TD (1993) ITS primers with enhanced specificity for basidiomycetes—application to the identification
435 of mycorrhizae and rusts. *Mol Ecol* 2(2):113-118. [10.1111/j.1365-294X.1993.tb00005.x](https://doi.org/10.1111/j.1365-294X.1993.tb00005.x)

436 Gordon H, Haygarth PM, Bardgett RD (2008) Drying and rewetting effects on soil microbial community composition
437 and nutrient leaching. *Soil Biol Biochem* 40(2):302-311. [10.1016/j.soilbio.2007.08.008](https://doi.org/10.1016/j.soilbio.2007.08.008)

438 Horowitz NH, Cameron RE, Hubbard JS (1972) Microbiology of the dry valleys of Antarctica. *Science* 176(4032), 242-
439 245. [10.1126/science.176.4032.242](https://doi.org/10.1126/science.176.4032.242)

440 Kovalski-Mitter E, de Freitas R, Germida JJ (2018) Microbial communities associated with barley growing in an oil
441 sands reclamation area in Alberta, Canada. *Can J Microbiol* 64(12):1004-1019. [10.1139/cjm-2018-0324](https://doi.org/10.1139/cjm-2018-0324)

442 Magan N (2007) Fungi in extreme environments. *The Mycota* 4:85-103. [10.1016/j.funeco.2012.04.003](https://doi.org/10.1016/j.funeco.2012.04.003)

443 McKay CP, Friedmann EI (1985) The cryptoendolithic microbial environment in the Antarctic cold desert: temperature
444 variations in nature. *Polar Biol* 4(1):19-25. [10.1007/BF00286813](https://doi.org/10.1007/BF00286813)

445 Muyzer G, de Waal EC, Uitterlinden AG (1993) Profiling of complex microbial populations by denaturing gradient gel
446 electrophoresis analysis of polymerase chain reaction amplified genes coding for 16S rRNA. *Appl Environ Microbiol*
447 59: 695-700.

448 NAS (2011) Future science opportunities in Antarctica and the Southern Ocean (report in brief). National Academy of
449 Sciences <http://www.dels.nas.edu/prb>.

450 Nienow JA, Friedmann EI (1993) Terrestrial lithophytic (rock) communities. In: Friedmann EI (ed) *Antarctic*
451 *microbiology*. Wiley-Liss, New York, 343-412.

452 Olech M, Chwedorzewska KJ (2011) Short note. The first appearance and establishment of an alien vascular plant in
453 natural habitats on the forefield of a retreating glacier in Antarctica. *Antarct Sci* 23:153-154
454 [10.1017/S0954102010000982](https://doi.org/10.1017/S0954102010000982)

455 Onofri S, Selbmann L, Zucconi L, Pagano S (2004) Antarctic microfungi as models for exobiology. *Planet Space Sci*
456 52(1-3):229-237. [10.1016/j.pss.2003.08.019](https://doi.org/10.1016/j.pss.2003.08.019)

457 Onofri S, Barreca D, Selbmann L, Isola D, Rabbow E, Horneck G *et al.* (2008) Resistance of Antarctic black fungi and
458 cryptoendolithic communities to simulated space and Mars conditions. *Stud Mycol* 61:99-109. [10.3114/sim.2008.61.10](https://doi.org/10.3114/sim.2008.61.10)

459 Onofri S, de la Torre R, de Vera JP, Ott S, Zucconi L, Selbmann L (2012) Survival of rock-colonizing organisms after
460 1.5 years in outer space. *Astrobiology* 12:508-516. [10.1089/ast.2011.0736](https://doi.org/10.1089/ast.2011.0736)

461 Onofri S, de Vera JP, Zucconi L, Selbmann L, Scalzi G, Venkateswaran KJ *et al.* (2015) Survival of Antarctic
462 Cryptoendolithic Fungi in Simulated Martian Conditions on Board the International Space Station. *Astrobiology*

- 463 15:1052-1059. [10.1089/ast.2015.1324](https://doi.org/10.1089/ast.2015.1324).
- 464 Parmesan C (2006) Ecological and evolutionary responses to recent climate change. *Annu Rev Ecol.Evol Syst* 37:637-
465 669. [10.1146/annurev.ecolsys.37.091305.110100](https://doi.org/10.1146/annurev.ecolsys.37.091305.110100)
- 466 Quintal H, Head J, Palumbo A, Dickson J (2018) McMurdo Dry Valleys: Exploring Antarctica as a Mars Analogue. In
467 Lunar and Planetary Science Conference, 49.
- 468
- 469 Raeymaekers L (2000) Basic principles of quantitative PCR. *Molecular Biotechnol* 15(2):115-122.
470 [10.1385/MB:15:2:115](https://doi.org/10.1385/MB:15:2:115)
- 471 Selbmann L, de Hoog GS, Mazzaglia A, Friedmann EI, Onofri S *et al.* (2005) Fungi at the edge of life: cryptoendolithic
472 black fungi from Antarctic deserts. *Stud Mycol* 51:1-32.
- 473 Selbmann L, de Hoog GS, Zucconi L, Isola D, Ruissi S, van den Ende AG *et al.* (2008) Drought meets acid: three new
474 genera in a dothidealean clade of extremotolerant fungi. *Stud Mycol* 61:1-20. [10.3114/sim.2008.61.01](https://doi.org/10.3114/sim.2008.61.01)
- 475 Selbmann L, Isola D, Fenice M, Zucconi L, Sterflinger K, Onofri S (2012) Potential extinction of Antarctic endemic
476 fungal species as a consequence of Global Warming. *Science Total Environ* 438:127-134.
477 [10.1016/j.scitotenv.2012.08.027](https://doi.org/10.1016/j.scitotenv.2012.08.027)
- 478 Selbmann L, Egidio E, Isola D, Onofri S, Zucconi L, de Hoog GS *et al.* (2013) Biodiversity, evolution and adaptation of
479 fungi in extreme environments. *Plant Biosyst* 147(1):237-246. [10.1080/11263504.2012.753134](https://doi.org/10.1080/11263504.2012.753134)
- 480 Selbmann L, Onofri S, Coleine C, Buzzini P, Canini F, Zucconi L (2017) Effect of environmental parameters on
481 biodiversity of the fungal component in the lithic Antarctic communities. *Extremophiles* 21:1069-1080.
482 [10.1007/s00792-017-0967-6](https://doi.org/10.1007/s00792-017-0967-6).
- 483 Selbmann L, Pacelli C, Zucconi L, Dadachova E, Moeller R, de Vera JP, Onofri S (2018) Resistance of an Antarctic
484 cryptoendolithic black fungus to radiation gives new insights of astrobiological relevance. *Fungal Biol* 122(6):546-554.
485 [10.1016/j.funbio.2017.10.012](https://doi.org/10.1016/j.funbio.2017.10.012)
- 486 Steig EJ, Schneider DP, Rutherford SD, Mann ME, Comiso JC, Shindell DT (2009) Warming of the Antarctic ice-sheet
487 surface since the 1957 International Geophysical Year. *Nature* 457(7228):459. [10.1038/nature07669](https://doi.org/10.1038/nature07669)
- 488 Turner J, Colwell SR, Marshall GJ, Lachlan Cope TA, Carleton AM, Jones PD *et al.* (2005) Antarctic climate change
489 during the last 50 years. *Int J Climatol* 25:279-294. [10.1002/joc.1130](https://doi.org/10.1002/joc.1130)
- 490 Turner J, Overland JE, Walsh JE (2007) An Arctic and Antarctic perspective on recent climate change. *Int J Climatol*
491 27:277-293. [10.1002/joc.1406](https://doi.org/10.1002/joc.1406)
- 492 Valášková V, Baldrian P (2009) Denaturing gradient gel electrophoresis as a fingerprinting method for the analysis of
493 soil microbial communities. *Plant Soil Environ* 55(10):413-423. [10.17221/132/2009-PSE](https://doi.org/10.17221/132/2009-PSE)
- 494 Venkateswaran A, McFarlan SC, Ghosal D, Minton KW, Vasilenko A, Makarova K, *et al.* (2000) Physiologic
495 determinants of radiation resistance in *Deinococcus radiodurans*. *Applied Environ Microbiol* 66(6):2620-2626.
- 496 Walker JJ, Spear JR, Pace NR (2005) Geobiology of a microbial endolithic community in the Yellowstone geothermal
497 environment. *Nature* 434:1011-1014. [10.1038/nature03447](https://doi.org/10.1038/nature03447)
- 498 Wei ST, Lacap-Bugler DC, Lau MC, Caruso T, Rao S, de los Rios A, *et al.* (2016) Taxonomic and functional diversity
499 of soil and hypolithic microbial communities in Miers Valley, McMurdo Dry Valleys, Antarctica *Front Microbiol*
500 7:1642. [10.3389/fmicb.2016.01642](https://doi.org/10.3389/fmicb.2016.01642)
- 501 White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for

502 phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (eds) PCR Protocols: a Guide to Methods and
503 Applications, 315-322. Academic Press, New York. [10.1016/B978-0-12-372180-8.50042-1](https://doi.org/10.1016/B978-0-12-372180-8.50042-1)

504 Yao T, Thompson LG, Mosbrugger V, Zhang F, Ma Y, Luo T *et al.* (2012) Third pole environment (TPE).
505 Environmental Development 3:52-64. [10.1016/j.envdev.2012.04.002](https://doi.org/10.1016/j.envdev.2012.04.002)

506 Yang K, Wu H, Qin J, Lin C, Tang W, Chen Y (2014) Recent climate changes over the Tibetan Plateau and their
507 impacts on energy and water cycle: A review. Glob Planet Change 112:79-91. [10.1016/j.gloplacha.2013.12.001](https://doi.org/10.1016/j.gloplacha.2013.12.001)

508 Zheng J, Liang R, Zhang L, Wu C, Zhou R, Liao X (2013) Characterization of microbial communities in strong
509 aromatic liquor fermentation pit muds of different ages assessed by combined DGGE and PLFA analyses. Food Res Int
510 54(1):660-666. [10.1016/j.foodres.2013.07.058](https://doi.org/10.1016/j.foodres.2013.07.058)

511 Zucconi L, Onofri S, Cecchini C, Isola D, Ripa C, Fenice M, *et al.* (2016) Mapping the lithic colonization at the
512 boundaries of life in Northern Victoria Land, Antarctica. Polar Biol 39(1):91-102. [10.1007/s00300-014-1624-5](https://doi.org/10.1007/s00300-014-1624-5)

A**FUNGAL COMMUNITIES** **B****BACTERIAL COMMUNITIES**

A

FUNGAL GENE COPIES NUMBER

B

BACTERIAL GENE COPIES NUMBER