Circadian control of intrinsic heart rate via a sinus node clock and the pacemaker channel

Yanwen Wang BSc, PhD¹; Servé Olieslagers BSc²; Anne Berit Johnsen BSc, PhD³;

Svetlana Mastitskaya BSc, PhD⁴; Haibo Ni MSc⁵; Yu Zhang BSc, PhD¹;

Nicholas Black BM BCh (Oxon)¹; Cali Anderson BSc¹; Charlotte Cox MRes¹;

Annalisa Bucchi BSc, PhD⁶; Sven Wegner MD, PhD⁷; Beatriz Bano-Otalora BSc, PhD⁷;

Cheryl Petit MRes⁷; Eleanor Gill MRes¹; Sunil Jit Logantha BPharm, MSc, PhD¹;

Nick Ashton BSc, PhD¹; George Hart DM, FRCP¹; Henggui Zhang BSc, PhD⁵;

Elizabeth Cartwright BSc, PhD¹; Ulrik Wisloff BSc, PhD³; Paula Da Costa Martins BSc, PhD²;

Dario DiFrancesco BSc, PhD⁶; Halina Dobrzynski BSc, PhD¹;

Hugh D. Piggins BSc, PhD⁷; Mark R. Boyett BSc, PhD, FSB, FRCP*^{†1}; Alicia D'Souza BSc, PhD^{†1}

[†]Joint senior authors

¹Division of Cardiovascular Sciences, University of Manchester, UK. ²Faculty of Health, Medicine and Life Science, Maastricht University, Netherlands. ³Department of Circulation and Medical Imaging, Norwegian University of Science and Technology, Norway. ⁴Neuroscience, Physiology and Pharmacology, University College London, UK. ⁵School of Physics and Astronomy, University of Manchester, UK. ⁶Department of Biosciences, University of Milan, Italy. ⁷Division of Diabetes, Endocrinology and Gastroenterology, University of Manchester

***Corresponding author:** Professor Mark R. Boyett; Division of Cardiovascular Sciences, University of Manchester, 46 Grafton Street, Manchester M13 9NT, UK
Tel: +44 161 275 1192 Fax: +44 161 275 1183 Email: mark.boyett@manchester.ac.uk

ABSTRACT

In the human, there is a circadian rhythm in the resting heart rate and it is higher during the day in preparation for physical activity. Conversely, slow heart rhythms (bradyarrhythmias) occur primarily at night. Although the lower heart rate at night is widely assumed to be neural in origin (the result of high vagal tone), the objective of the study was to test whether there is an *intrinsic* change in heart rate driven by a local circadian clock. In the mouse, there was a circadian rhythm in the heart rate *in vivo* in the conscious telemetrized animal, but there was also a circadian rhythm in the intrinsic heart rate in denervated preparations: the Langendorff-perfused heart and isolated sinus node. In the sinus node, experiments (qPCR and bioluminescence recordings in mice with a *Per1* luciferase reporter) revealed functioning canonical clock genes, e.g. *Bmal1* and *Per1*. We identified a circadian rhythm in the expression of key ion channels, notably the pacemaker channel *Hcn4* (mRNA and protein) and the corresponding ionic current (funny current, measured by whole cell patch clamp in isolated sinus node cells). Block of funny current in the isolated sinus node abolished the circadian rhythm in the intrinsic heart rate. Incapacitating the local clock (by cardiac-specific knockout of *Bmal1*) abolished the normal circadian rhythm of *Hcn4*, funny current and the intrinsic heart rate. Chromatin immunoprecipitation demonstrated that *Hcn4* is a transcriptional target of BMAL1 establishing a pathway by which the local clock can regulate heart rate. In conclusion, there is a circadian rhythm in the intrinsic heart rate as a result of a local circadian clock in the sinus node that drives rhythmic expression of *Hcn4*. The data reveal a novel regulator of heart rate and mechanistic insight into the occurrence of bradyarrhythmias at night.

Living things including humans are attuned to the 24 h day-night cycle and many biological processes exhibit an intrinsic ~24 h (i.e. circadian) rhythm. In the human, the resting heart rate (in the absence of physical activity) shows a circadian rhythm and is higher during the day when we are awake^{1,2}. The heart is therefore primed, anticipating the increase in demand during the awake period. Conversely, bradyarrhythmias primarily occur at night^{1,3}. Previously, the circadian rhythm in heart rate *in vivo* has been attributed to the autonomic nervous system: to high sympathetic nerve activity accompanying physical activity during the awake period and high vagal tone during the sleep period⁴. This is partly based on heart rate variability as a surrogate measure of autonomic tone⁵⁻⁷; however, we have shown that heart rate variability cannot be used in any simple way as a measure of autonomic tone⁸. Therefore, the mechanism underlying the circadian rhythm in heart rate is still unknown. We asked the question whether there is a circadian rhythm in the *intrinsic* heart rate set by the pacemaker of the heart, the sinus node.

In a cohort of nine mice, ECG telemetry *in vivo* showed a circadian rhythm in mean heart rate and other electrophysiological parameters (PR interval, QRS duration and QT interval) during a 12 h light:12 h dark lighting regime, and circadian rhythms in these parameters were sustained when the mice were placed in constant dark conditions (Figure 1). A circadian rhythm in heart rate, PR interval, QRS duration and QT interval are also observed in the human^{7,9-11}. Mice are nocturnal and as expected they rested more and were less active from Zeitgeber time (ZT) 0 to ZT 12 (lights-on) and were more active from ZT 12 to ZT 0 (lights-off); this activity pattern continued in constant darkness (Figure 1)^{12,13}. The heart rate was highest at ~ZT 13 and it varied by 76 ± 4 beats/min (as determined by the least-squares best fit sine wave) over the course of 24 h (Figure 1; Table S1 in the Supplementary Information).

Circadian rhythm in intrinsic sinus node pacemaking and ion channel expression

Experiments on the isolated, denervated, Langendorff-perfused heart (Figure 2A) dissected at projected ZT 0 or ZT 12 demonstrated that there was a circadian rhythm in *intrinsic* sinus node pacemaking: the intrinsic heart rate was still higher at ZT 12 than ZT 0 – by 107 beats/min (Table S1). In the Langendorff-perfused heart, a suitable pacing protocol (Figure S1 in the Supplementary Information) showed the corrected sinus node recovery time (cSNRT), a commonly used indicator of sinus node function, to be significantly different at ZT 0 and ZT 12 (Figure 2B). It is concluded

that there is a circadian rhythm in the intrinsic heart rate. What is responsible for the intrinsic circadian rhythm? Molecular circadian clock machinery is ultimately responsible for all circadian rhythms. The master clock is in the suprachiasmatic nucleus of the brain, but there are peripheral clocks in other organs¹⁴. To determine if there is circadian clock machinery in the sinus node, qPCR was used – qPCR showed that mRNA for a wide range of molecules known to be involved in the circadian clock is present in the sinus node and the expression varied in the expected manner from ZT 0 to ZT 12 (Figure 2C). Two key transcripts, *Bmal1* and *Clock*, were measured in the sinus node at four time points (BMAL1 and CLOCK form a heterodimer); Figure 2D,E shows that they varied in a circadian manner and were at a maximum at ~ZT 0 (Table S1). The presence of an intrinsic circadian clock in the sinus node was confirmed by measuring the bioluminescence in the isolated sinus node from a transgenic mouse (*Per1::LUC*) carrying a luciferase reporter gene reporting the activity of *Per1*, a key circadian clock component (Figure 2F). *Per1*-driven bioluminescence fluctuated in the expected circadian manner and this periodicity was lost in *Cry1*^{-/-} *Cry2*^{-/-} mice lacking the *Cryptochrome* genes and consequently lacking a functional clock¹⁵ (Figure 2F). Pacemaking is the result of the concerted action of ion channels and Ca²⁺-handling proteins and it is likely that the local circadian clock in the sinus node controls the heart rate by controlling their expression. Expression of mRNA for many of these molecules (and also some key regulatory transcription factors) was measured by qPCR and some transcripts, for example the pacemaking ion channel *Hcn4*, demonstrated significant daily rhythms (Figure 2G; Table S2).

Circadian rhythm in HCN4 and I_f

Of the potential mechanisms controlling the intrinsic heart rate, we principally focused on *Hcn4*, because this is known to play a central role in pacemaking in the sinus node¹⁶. *Hcn4* mRNA was measured at six time points and was at a maximum at ~ZT 0 (Figure 3A; Table S1). Expression of HCN4 protein in the sinus node was measured using western blot at ZT 0 and ZT 12 – the western blot is shown in Figure 3B and the mean HCN4 band intensities are shown in Figure 3C. HCN4 protein expression was higher at ZT 12 than ZT 0 (Figure 3C). Expression of HCN4 protein in the sinus node was also measured using immunohistochemistry (Figure 3D,E). Figure 3D shows examples of immunolabelling of HCN4 in tissue sections through the sinus node from mice culled at ZT 0 and ZT 12); consistent with the western blot, the labelling was brighter, indicating higher

107 expression, at ZT 12. This is confirmed by the mean data in Figure 3E. In summary, the data show
 108 a daily rhythm in HCN4 protein, but it is out of phase with the circadian rhythm in mRNA. Western
 109 blot and immunohistochemistry were used to measure HCN4 at four time points and this showed
 110 that HCN4 protein peaked at between ZT 6 and 12 (Figure S2; Table S1) – HCN4 protein,
 111 therefore, lags behind mRNA - this is expected. If there are changes in HCN4 protein, there should
 112 be changes in the corresponding ionic current, I_f . Figure 4A shows patch clamp recordings of I_f
 113 from isolated sinus node cells – the cells were isolated at ZT 0 and ZT 10 and the recordings were
 114 made at ZT 2 and ZT 12 (the earliest time points at which recordings could be made). The current
 115 density was approximately double at ZT12 than at ZT2 and this is confirmed by the mean current-
 116 voltage relationships for I_f at the two time points in Figure 4B. Once cells were isolated at ZT 0 and
 117 ZT 10, recordings from different cells could continue to be made for ~3 h. In the case of cells
 118 isolated at ZT 0 we observed the current density to increase the later the recording, but in the case
 119 of cells isolated at ZT 10 we observed the current density to decrease the later the recording. In
 120 Figure 4C we have plotted the density of I_f in a total of 367 sinus node cells against the time of the
 121 recording (this includes cells which were isolated at ZT 3, 6, 9, 12 and 15). Figure 4C shows a
 122 projected daily rhythm in the density of I_f – clearly the circadian clock does not stop ticking on
 123 isolation of single sinus node cells. I_f density reached a maximum at ~ZT 10, approximately at the
 124 same time as the peak in HCN4 protein as assessed by western blot and immunohistochemistry
 125 (Figure S2; Table S1). Current density determines the rate of change of membrane potential and
 126 is, therefore, the physiologically important variable. It is calculated by dividing the current amplitude
 127 by the cell capacitance, C_m , and these variables are shown in Figure 4D,E. As anticipated there is
 128 a projected daily rhythm in I_f amplitude, which peaks at the same time as I_f density, but the
 129 fractional change in amplitude is less than the fractional change of density (Figure 4D,E). This is
 130 because there is a circadian rhythm in C_m (Figure 4E); in other words, a circadian rhythm in C_m
 131 contributes to the circadian rhythm in I_f density. C_m is determined by cell size and other
 132 experiments confirmed that there is a circadian rhythm in cell size (Figure S3). A circadian rhythm
 133 in cell size may be surprising. However, a circadian rhythm in cell size has been reported in other
 134 tissues: in mouse liver, cell size at ZT 12 is less than at ZT 2¹⁷; axonal volume shows a circadian
 135 rhythm in s-LNV clock neurons in *Drosophila*¹⁸; glial cells and neurons in the housefly's visual

system show rhythmic size changes¹⁹.

Role of I_f and other ionic mechanisms in circadian rhythm in intrinsic heart rate

To test whether the changes in I_f density could account for the daily rhythm in the intrinsic heart rate, the observed changes in I_f density (Figure 4C) were incorporated into a biophysical model of the spontaneous action potential of the mouse sinus node (Figure 4F). The model predicted a circadian rhythm in heart rate of 101 ± 11 beats/min with a maximum heart rate at ~ZT 10 (Figure 4G; Table S1), roughly consistent with experimental data for the intrinsic heart rate (Figure 2A). This suggests that HCN4 and I_f participate in the circadian rhythm in the intrinsic heart rate. This was confirmed in the isolated, denervated sinus node dissected at four different time points. Block of I_f by 2 mM Cs^+ (selective blocker of I_f^{20}) decreased the heart rate as expected (Fig. 5A, top), but the effect of Cs^+ on heart rate was greater at ZT 12 than ZT 0 (Figure 5A, bottom). Furthermore, in the presence of Cs^+ , the circadian rhythm in the intrinsic heart rate was abolished (Figure 5A, top). Analogous results were obtained *in vivo*. In the conscious mouse, the heart rate at ZT 0 and ZT 12 was measured using an ECGenie (a platform with embedded ECG electrodes). An intraperitoneal injection of 6 mg/kg ivabradine (selective blocker of I_f^{21}) was given to block I_f . Once again, block of I_f decreased the heart rate as expected (Figure 5C) and the effect of ivabradine on heart rate was greater at ZT 12 than ZT 0 (Figure 5D). Furthermore, in the presence of ivabradine, the circadian rhythm in the heart rate *in vivo* was abolished (Figure 5C). It is concluded that HCN4 and I_f participate in the circadian rhythm in the intrinsic heart rate as well as the heart rate *in vivo*.

The so-called membrane and Ca^{2+} clocks are known to be responsible for pacemaking in the sinus node²². The membrane clock comprises ionic currents carried by a variety of ion channels. Although I_f carried by HCN channels is considered to be the most important, other ion channels also play a significant role. Various K^+ channels showed a circadian rhythm (Figure 2G) and their potential role is considered in the Discussion. The Ca^{2+} clock involves spontaneous Ca^{2+} releases (Ca^{2+} sparks) from the sarcoplasmic reticulum during diastole. This drives Ca^{2+} extrusion via the electrogenic Na^+ - Ca^{2+} exchanger (*Slc8a1*; NCX1). The resulting inward current generated by NCX1 helps drive pacemaking. Although a circadian rhythm was not detected in any of the principal components of the Ca^{2+} clock (apart from Ca^{2+} /calmodulin-dependent protein kinase II δ – *Camk2d*), the potential role of the Ca^{2+} clock in the circadian rhythm in the intrinsic heart rate was

investigated by recording Ca^{2+} sparks from isolated sinus node cells and measuring the effect of incapacitating the Ca^{2+} clock by 2 μM ryanodine on the intrinsic heart rate measured in the isolated sinus node (Figures S4-S6). Measurement of Ca^{2+} sparks showed that at ZT 12 as compared to ZT 0 there was an increase in spark duration and a non-significant increase in spark amplitude and consequently there was an increase in spark mass (calculated as amplitude $\times$ 1.206 $\times$ spark full width at half maximum amplitude³)²³ ($P=0.056$), although there was a decrease in spark frequency and diastolic Ca^{2+} (Figure S5). The increase in spark mass at ZT 12 could increase the impact of the Ca^{2+} clock on pacemaking (although the effect could be mitigated by the reduction of Ca^{2+} spark frequency). In the isolated sinus node, the effect on the intrinsic heart rate of incapacitating the Ca^{2+} clock by ryanodine was greater at ZT 12 than ZT 0 (Figure S6A). In the presence of ryanodine, the circadian rhythm in the intrinsic heart rate was abolished (Figure S6A, top) and it is concluded that the Ca^{2+} clock also participates in the circadian rhythm in the intrinsic heart rate.

Link between local circadian clock in sinus node and membrane and Ca^{2+} clocks

To investigate a possible link between the local circadian clock in the sinus node and the circadian rhythm in the intrinsic heart rate, experiments were conducted on a transgenic mouse in which the *Bmal1* gene had been knocked out in the heart only (driven by the α myosin heavy chain promoter)²⁴. Knockout of *Bmal1* is known to incapacitate the circadian clock²⁵. Figure 6A confirms that *Bmal1* mRNA was effectively knocked out in the sinus node at both ZT 0 and ZT 12 and Figure 6B shows that this disrupted the circadian rhythm in the expression of *Clock* mRNA – evidence that the circadian clock in the sinus node had been disrupted as expected. In the cardiac-specific *Bmal1* knockout mouse, from ZT 0 to ZT 12, there was no longer a significant variation in expression of *Hcn4* mRNA (Figure 6C) and HCN4 protein (Figure 3D,E). Consistent with this, the variation in I_f density from ZT 0 to ZT 12 was blunted (Figure 6D,E); this is best shown by Figure 6F, which compares the variation in I_f density from ZT 0 to ZT 12 in wild-type and cardiac-specific *Bmal1* knockout mice. Finally, Figure 5B (top) shows the intrinsic heart rate as measured in the isolated sinus node: whereas there was a circadian rhythm in the wild-type mice with the intrinsic heart rate peaking at $\sim$ ZT 12 (Figure 5A, top), in cardiac-specific *Bmal1* knockout mice this normal pattern was lost (Figure 5B, top). Furthermore, whereas there was a circadian rhythm in the reduction of the intrinsic heart rate on block of I_f by Cs^+ in wild-type mice (with the effect peaking at

~ZT 12; Figure 5A, bottom), in cardiac-specific *Bmal1* knockout mice, once again, this pattern was lost (Figure 5B, bottom). In contrast, cardiac-specific *Bmal1* knockout had little effect on the circadian rhythm in the Ca^{2+} clock-control of the intrinsic heart rate (Figure S6B). It is concluded that the local circadian clock in the sinus node is controlling HCN4 and I_f (but perhaps not the Ca^{2+} clock) and thereby the intrinsic heart rate.

Evidence that clock protein, BMAL1, controls *Hcn4* transcription

The CLOCK:BMAL1 heterodimer acts as a transcriptional activator or enhancer by binding to E-box binding sites in the promoter, intron or exon of a gene^{26,27}. Using RVISTA (<https://rvista.dcode.org>) we found eight canonical E-box binding sites in the *Hcn4* gene and 20 kb of its 5' flanking region (Figure 6H). *In vitro* ChIP was used to test whether BMAL1 specifically binds to these sites: 3T3-L1 cells were UV cross-linked following transfection with His-tagged *Bmal1*. His-tagged BMAL1 bound to its DNA targets was then immunoprecipitated using antibodies directed against the His-tag. DNA pulled down by ChIP was analysed by qPCR using primers mapping to the various E-box binding sites (Figure 6G). Figure 6G shows E-box binding sites D and G (within introns of the *Hcn4* gene; Figure 6H) were significantly more abundant than background signal (red dashed line).

Role for the autonomic nervous system in circadian rhythm in heart rate *in vivo*?

This study highlights the role of intrinsic factors in the circadian rhythm in intrinsic heart rate. Nevertheless, a role for the autonomic nervous system must be considered. In this study, *in vivo*, the heart rate was high during the awake period when the mice were physically active (Figure 1). If the level of physical activity is high, it would be expected to influence the heart rate via the autonomic nervous system (via an increase in sympathetic activity and possibly a decrease in vagal activity). However, in caged-housed mice with no access to a running wheel, activity levels will be low. A light pulse when mice are active is known to cause the mice to freeze and be inactive¹². Figure 7A shows that a 1 h light pulse from ZT13 to ZT14 caused the physical activity of the mice to fall to baseline values, whereas the heart rate remained relatively high. In contrast, a 1 h light pulse from ZT1 to ZT2 was again associated with a baseline level of physical activity, but the heart rate was relatively low. Therefore, in this experiment, the heart rate was primarily influenced by the time of day rather than the activity level. Figure S7 shows that there is no

discernible relationship between heart rate and physical level over 24 h (Figure 7A) and before, during and after the light pulses (Figure 7B). *In vivo*, the heart rate in the absence of physical activity can be obtained by recording the ECG from the anaesthetised mouse (although anaesthetic is known to depress the intrinsic pacemaker activity of the sinus node²⁸ as well as cardiac vagal and sympathetic tone²⁹). The heart rate in the anaesthetised mouse also varied in a daily manner – the heart rate was highest at ~ZT 12 and it varied by 51 ± 15 beats/min over 24 h (Figure 7B; Table S1). It is concluded that in this study the effect of activity on the heart rate of the mice was slight and not discernible. Next, the effect of vagotomy was investigated. There are right and left vagal nerves and sectioning of both is lethal. However, vagal nerves to the sinus node are primarily, but not exclusively, from the right vagus³⁰. The right vagus was sectioned in a group of rats (rat was studied, because we have experience of vagotomy in the rat³¹). Figure 7C shows that following the vagotomy the circadian rhythm in heart rate persisted.

We further tested the involvement of the autonomic nervous system by blocking cardiac muscarinic and β receptors by atropine and propranolol. Experiments were conducted on anaesthetised mice, and intraperitoneal injections of 1 mg/kg atropine and 1 mg/kg propranolol were given, similar doses to those used by others³²⁻³⁴ and by us in an earlier study of exercise training-induced bradycardia in the mouse³⁵. After autonomic blockade, the circadian rhythm in heart rate persisted, although it was reduced in amplitude (Figure 7D). In our earlier study, similar concentrations of atropine and propranolol also failed to abolish the exercise training-induced bradycardia³⁵; we concluded that the autonomic nervous system is not responsible for the exercise training-induced bradycardia³⁵ and based on Figure 7D we could conclude that it is also not solely responsible for the circadian rhythm in heart rate *in vivo*. However, following the publication of our study of exercise training-induced bradycardia, Aschar-Sobbi *et al.*³⁶ used intraperitoneal injections of higher concentrations of atropine and propranolol (2 mg/kg atropine and 10 mg/kg propranolol) and stated that the exercise training-induced bradycardia in the mouse is abolished on autonomic blockade. Therefore, we repeated the experiment in Figure 7D, but we used intraperitoneal injections of 2 mg/kg atropine and 10 mg/kg propranolol as did Aschar-Sobbi *et al.*³⁶. This time, after autonomic blockade, the circadian rhythm in heart rate was lost (Figure 7E). Although this suggests that the autonomic nervous system is involved in the circadian rhythm in heart rate *in*

vivo, the effect of the higher concentrations of atropine and propranolol must be taken into consideration. Propranolol blocks β -receptors with an IC_{50} of 12 nM (<http://www.selleckchem.com/products/propranolol-hcl.html>). However, work on heterologously expressed ion channels in cell lines has shown that propranolol blocks the cardiac Na^+ channel, $Na_v1.5$ (*Scn5a*) with an IC_{50} of 2.7 μM ³⁷ and HCN4 with an IC_{50} of 50.5 μM ³⁸. It was confirmed that propranolol also blocks I_f in native heart cells: in mouse sinus node cells, 68.4 μM propranolol blocked I_f by 54% (Figure 7F); typical current traces and current-voltage relationships are shown in Figure S8. Assuming that only 10% of propranolol is free³⁹, an injection of 10 mg/kg propranolol is estimated to give a propranolol concentration of 6 μM if it partitions in all body water (~60% of body mass in the mouse⁴⁰) and 66 μM if it only partitions into the blood (58.5 ml/kg in the mouse; <https://www.nc3rs.org.uk/mouse-decision-tree-blood-sampling>). Therefore, based on these estimates, it is possible that 10 mg/kg propranolol may block, to some extent, $Na_v1.5$ and HCN4, both of which vary in a circadian manner (Figure 2G). If propranolol did block HCN4 to a significant extent, it is not surprising that propranolol abolished the circadian rhythm in heart rate (Figure 7E), because ivabradine also did (Figure 5C). Propranolol at a dose of 1 mg/kg as used for Figure 7D is expected to cause less block of $Na_v1.5$ and little or no block of HCN4 and yet complete block of β -receptors (the expected plasma concentration is still many times greater than the IC_{50} for block of β -receptors). As an aside, calculation shows that atropine at a dose of 1 or 2 mg/kg is sufficient for complete block of M2 muscarinic receptors. Based on Figure 7D, the autonomic nervous system cannot be solely responsible for the circadian rhythm in heart rate *in vivo*.

DISCUSSION

For the first time, we show here that: (i) there is a circadian or daily rhythm in the *intrinsic* heart rate set by the pacemaker of the heart, the sinus node, (ii) there is a functioning circadian clock in the sinus node, (iii) there is a circadian rhythm in the expression of a variety of cardiac ion channels, including *Hcn4* and its corresponding ionic current, I_f , in the sinus node, (iv) *Hcn4* transcription is directly controlled by the clock transcription factor, BMAL1, (v) the circadian rhythm in *Hcn4* plays an important role in the circadian rhythm in the *intrinsic* heart rate.

280 **Local circadian clock in heart**

281 Previously, it has been demonstrated that there a functioning circadian clock in the heart^{e.g.41}. This
282 is the first report of a functioning circadian clock in the sinus node: qPCR showed the presence of
283 transcripts for many key circadian clock components and many showed the expected circadian
284 rhythm and expected phase relationship; for example, *Bmal1* was downregulated, but *Cry2*, *Per1*
285 and *Per2* were upregulated at ZT 12 compared to ZT 0 (Figure 2C). We also showed a circadian
286 rhythm in *Per1* using a bioluminescent reporter gene (Figure 2F).

287 **Circadian rhythm in cardiac ion channel expression is common**

288 Here we show a circadian rhythm in *Hcn4*, Na_v1.5 (*Scn5a*), K_v1.4 (*Kcna4*), K_v4.2 (*Kcnd2*), K_v1.5
289 (*Kcna5*), ERG (K_v11.1; *Kcnh2*), K_i6.1 (*Kcnj8*), K_i6.2 (*Kcnj11*), SUR1 (*Abcc8*), SUR2 (*Abcc9*),
290 TASK-1 (K2p3.1; *Kcnk3*) and Cl⁻ voltage-gated channel 2 (*Clcn2*) in the sinus node (Figure 2G).
291 Further experiments have shown that *Hcn1* also shows a circadian rhythm in the sinus node
292 (Yanwen Wang, unpublished data). Curiously, *Hcn2* has been reported to show a circadian rhythm
293 in liver⁴². A circadian rhythm in cardiac ion channel expression is not only seen in the sinus node –
294 it is likely to be occurring throughout the heart. In atrial muscle, a circadian rhythm has been
295 reported in K_v4.2 (*Kcnd2*)^{41,43,44}, KChIP2 (*Kcnip2*)^{41,44}, K_v1.5 (*Kcna5*)^{41,43}, TASK-1 (K2p3.1;
296 *Kcnk3*)⁴¹, Cx40 (*Gja5*)⁴⁵ and Cx43 (*Gja1*)⁴⁵. In ventricular muscle, a circadian rhythm has been
297 reported in Na_v1.5 (*Scn5a*)⁴⁶, Ca_v1.2 (*Cacna1c*)⁴⁷, K_v4.2 (*Kcnd2*)^{41,43,48}, KChIP2 (*Kcnip2*)^{41,44}, K_v1.5
298 (*Kcna5*)^{41,43}, ERG (K_v11.1; *Kcnh2*)⁴⁸, TASK-1 (K2p3.1; *Kcnk3*)⁴¹, Cx40 (*Gja5*)⁴⁵ and Cx43 (*Gja1*)⁴⁵.
299 Some circadian varying ion channels are common to the sinus node and the atrial or ventricular
300 muscle: Na_v1.5 (*Scn5a*), K_v4.2 (*Kcnd2*), K_v1.5 (*Kcna5*), ERG (K_v11.1; *Kcnh2*) and TASK-1 (K2p3.1;
301 *Kcnk3*). Of these, all were significantly more highly expressed at ZT 12 than ZT 0 in the sinus node
302 (Figure 2G); in atrial and ventricular muscle, there was a qualitatively similar circadian rhythm in
303 Na_v1.5 (*Scn5a*)⁴⁶, K_v4.2 (*Kcnd2*)^{41,43,48}, K_v1.5 (*Kcna5*; in ventricular muscle, but not atrial
304 muscle)^{41,43} and ERG (K_v11.1; *Kcnh2*)⁴⁸. However, the opposite circadian rhythm in TASK-1
305 (K2p3.1; *Kcnk3*) has been reported in atrial and ventricular muscle⁴¹.

306 This study has demonstrated that *Hcn4* is likely to be under the control of BMAL1
307 generated by the local clock (Figure 6). In the ventricle, it has been reported that: Nav1.5 (*Scn5a*)
308 and ERG (K_v11.1; *Kcnh2*) are under the control of CLOCK:BMAL1 generated by the local

clock^{46,48}; and K_v4.2 (*Kcnd2*), K_v1.5 (*Kcna5*), TASK-1 (K2p3.1; *Kcnk3*), Cx40 (*Gja5*) and Cx43 (*Gja1*) are under the control of the suprachiasmatic nucleus (not the local clock) via the autonomic nervous system⁴¹. Confusingly, KChIP2 (*Kcnip2*) has been reported to be under the control of a BMAL1-dependent (i.e. local clock-dependent) oscillator, krüppel-like factor 15 (*Klf15*)⁴⁴, and yet under the control of the suprachiasmatic nucleus (not the local clock) via the autonomic nervous system⁴¹.

Circadian rhythm in intrinsic heart rate

Cardiac-specific knockout of *Bmal1* abolished the normal circadian rhythm in the intrinsic heart rate (Figure 5A,B), proving that it is under the control of the local circadian clock (however, it is interesting that some type of rhythm, albeit abnormal, remained after cardiac-specific knockout of *Bmal1* – Fig. 5B). We measured a circadian rhythm in HCN4 protein and *I_f* (Figures 3B-E, 4 and S2) roughly in phase with the circadian rhythm in the intrinsic heart rate (in contrast *Hcn4* mRNA was out of phase – Figure 3A; Table S1). It is likely that the circadian rhythm in *I_f* plays an important role in the intrinsic heart rate: computer modelling suggested that the changes in *I_f* are sufficient to explain the changes in intrinsic heart rate (Figure 4G); block of HCN4 and *I_f* by Cs⁺ abolished the circadian rhythm in the intrinsic heart rate (Figure 5A); and cardiac-specific knockout of *Bmal1* abolished the normal circadian rhythm in *Hcn4* mRNA and protein, *I_f* and the effect of Cs⁺, as well the intrinsic heart rate (Figures 3E, 5B and 6C,E,F). However, it is likely that *I_f* is not the only mechanism involved: there was a circadian rhythm in CaMKIIδ (*Camk2d*) and Ca²⁺ sparks (Figures 2G and S5), and incapacitating the Ca²⁺ clock with ryanodine eliminated the circadian rhythm in the intrinsic heart rate (Figure S6A). In addition, it is possible that the circadian rhythm detected in various K⁺ channels, particularly ERG (K_v11.1; *Kcnh2*) (Figure 2G), also plays a role; in the sinus node, ERG (K_v11.1; *Kcnh2*), responsible for the rapid delayed rectifier K⁺ current, *I_{K,r}*, sets the maximum diastolic potential and blocking *I_{K,r}* abolishes pacemaking⁴⁹.

Circadian rhythm in heart rate *in vivo* – is it multifactorial?

This study has demonstrated a circadian rhythm in the *intrinsic* heart rate. Implications of this for the circadian rhythm in heart rate *in vivo* can be speculated on. Previously, the circadian rhythm in heart rate *in vivo* has been attributed to the autonomic nervous system and in particular to high vagal tone during the sleep period⁴. This primarily has been based on heart rate variability as a

surrogate measure of autonomic tone⁵; however, as discussed above, this cannot be used in any simple way as a measure of autonomic tone⁸. We report that autonomic blockade utilising high concentrations of atropine and propranolol abolished the circadian rhythm in heart rate *in vivo* (Figure 7E). Abolition of the circadian rhythm in heart rate *in vivo* on autonomic blockade has also been reported by Tong *et al.*⁴¹ However, Oosting *et al.*⁵⁰ have reported that autonomic blockade does not abolish the circadian rhythm in heart rate. Furthermore, caution must be used in interpreting the effect of autonomic blockade involving atropine and propranolol, because propranolol is non-specific and also blocks Na_v1.5 (*Scn5a*) and *I*_{Na} and HCN4 and *I*_f (Figures 7F and S8)^{37,38}. Because *I*_f at least is greater during the awake period, propranolol will be expected to have a greater depressing effect on heart rate during the awake period. This is an alternative explanation of why the circadian rhythm in heart rate *in vivo* was abolished by autonomic blockade in this study when utilising high concentrations of atropine and propranolol (Figure 7E). Nevertheless, if it is assumed that the autonomic nervous system is involved, the right vagus nerve (primarily, but not exclusively, responsible for the innervation of the sinus node³⁰) cannot be solely responsible, because sectioning the right vagus did not impact the circadian rhythm in heart rate *in vivo* in the rat (Figure 7C). This suggests that a circadian rhythm in sympathetic nerve activity or plasma catecholamine levels is more important (if it is assumed that the autonomic nervous system is involved). However, it has been reported that transgenic knockout of the M2 receptor or β 1, β 2 and β 3 adrenoceptors has no effect on the circadian rhythm in heart rate *in vivo*⁵¹. In addition, cardiac transplant patients with autonomic denervation have a preserved nocturnal bradycardia 7-36 months after transplantation^{52,53}. This work does not support a role for the autonomic nervous system in the circadian rhythm in heart rate *in vivo*.

This study has shown that there is a circadian rhythm in the intrinsic heart rate of the appropriate amplitude (72-107 beats/min) and phase (peaking at ~ZT 12; Figures 2A and 5A; Table S1) to be able to explain the circadian rhythm in heart rate *in vivo* (Figure 1). Furthermore, block of HCN4 and *I*_f by ivabradine had a bigger effect on heart rate during the awake period and abolished the circadian rhythm in heart rate *in vivo* in the mouse (Figure 5C). Data consistent with this have been obtained by Ptaszynski *et al.*⁵⁴ and Grigoryan *et al.*⁵⁵ who studied the effect of ivabradine on the heart rate of patients either with inappropriate sinus node tachycardia or

ischaemic heart disease and heart failure; whereas ivabradine caused a large decrease in heart rate during the day, it caused little decrease at night. In other words, the effect of ivabradine on heart rate showed a circadian rhythm and was greater when the subjects were awake. This is consistent with the work on the mice (Figure 5C,D)⁵⁴. However, it is known that cardiac-specific knockout of *Bmal1*⁴⁶ or cardiac-specific expression of a dominant negative *Clock* mutant⁵⁶ does not abolish the circadian rhythm in heart rate *in vivo* (although it does reduce it). Furthermore, we show here that cardiac-specific knockout of *Bmal1* abolishes the *normal* circadian rhythm in *Hcn4* (mRNA and protein), *I_f* and intrinsic heart rate (Figures 3E, 5 and 6C,E,F). Therefore, HCN4 and *I_f* cannot be the only mechanism controlling the circadian rhythm in heart rate *in vivo* – perhaps the reduction of the circadian rhythm in heart rate *in vivo*⁴⁶ on cardiac-specific knockout of *Bmal1* represents the contribution of the circadian rhythm in the intrinsic heart rate to the circadian rhythm in heart rate *in vivo*. What is responsible for the circadian rhythm in heart rate *in vivo* that remains after cardiac-specific *Bmal1* knockout? This could be due to the autonomic nervous system. Consistent with this, Figure 7D shows that the circadian rhythm in heart rate *in vivo* was reduced in amplitude after autonomic blockade (achieved with the lower doses of atropine and propranolol). However, the contribution may not be in the way originally conceived (acute control of heart rate via changes in ionic conductance): Tong *et al.*^{41,45} have shown that the circadian rhythm of K_v4.2 (*Kcnd2*), KChIP2 (*Kcnip2*), K_v1.5 (*Kcna5*), TASK-1 (K2p3.1; *Kcnk3*), Cx40 (*Gja5*) and Cx43 (*Gja1*) in the atria and ventricles is lost after autonomic blockade suggesting transcriptional regulation mediated by the autonomic nervous system. It is interesting that β -agonists affect both *Bmal1* and *Per2* in heart^{57,58}.

Summary

In summary, this study has shown that there is a circadian rhythm in the *intrinsic* heart rate and this is likely to contribute to the circadian rhythm in heart rate *in vivo*. Our findings provide new mechanistic insight into the fundamental question of why the heart rate of a mammal is lower when asleep and also explains the nocturnal occurrence of bradyarrhythmias^{1,3,59-63}.

METHODS

Extended methods are given in the Supplementary Information.

Animals

The telemetry study was approved by the Norwegian Council for Animal Research, in accordance with the Guide for the Care and Use of Laboratory Animals from the European Commission Directive 86/609/EEC. Ethical approval for vagotomy in rats was from University College London, in accordance with the UK Animals (Scientific Procedures) Act 1986. All remaining procedures were approved by the University of Manchester and were in accordance with the UK Animals (Scientific Procedures) Act 1986. Experiments were conducted on adult male mice: C57BL/6J mice obtained from Harlan Laboratories; *Per1::LUC* mice (in which luciferase expression is driven by the mouse *Per1* promoter and 5'-UTR elements⁶⁴) on a *Cry1^{+/+}Cry2^{+/+}* or *Cry1^{-/-}Cry2^{-/-}* background; and cardiac-specific *Bmal1* knockout mice.

Electrophysiology

ECGs or ECG-like electrograms were recorded for measurement of heart rate: from conscious unrestrained mice using either subcutaneously implanted radio telemetry transmitters or an ECGenie; from isofluorane-anaesthetised mice using a conventional three-lead system; from isolated Langendorff-perfused hearts; and from isolated right atrial preparations encompassing the sinus node. In the Langendorff-perfused hearts, a suitable pacing protocol was also used to determine the corrected sinus node recovery time. Strips of sinus node tissue were dissociated into single cells by a standard enzymatic and mechanical procedure and *I_f* was investigated using the patch clamp technique in the whole-cell mode. A previously developed biophysically-detailed mathematical model of the mouse sinus node cell action potential⁶⁵ was used to assess the effect of changes in the density of *I_f* on the spontaneous action potential.

mRNA and protein expression

RNA was isolated from frozen 1 mm sinus node punch biopsies and gene expression measured using quantitative PCR (qPCR) either using medium throughput custom-designed Taqman Low Density Array Cards or individual SYBR green assays. HCN4 protein was investigated using either sinus node homogenates and western blotting or tissue cryosections and immunohistochemistry. *Per1* expression was also measured using the *Per1::LUC* mouse: total bioluminescence was recorded for 96 h from freshly dissected intact sinus node preparations from *Per1::LUC* mice using a photomultiplier tube assembly.

In vitro chromatin immunoprecipitation (ChIP)

ChIP was performed on 3T3-L1 cells using the SimpleChIP chromatin immunoprecipitation kit according to the manufacturer's instructions. Prior to cross-linking, 3T3-L1 cultures at a density of 10^4 cells/cm² were transfected with plasmid containing His-tagged *Bmal1* for 48 h. Antibody directed against the His-tag was used for immunoprecipitation. DNA obtained from ChIP was analysed by qPCR using primers mapping to canonical E-box binding sites, i.e., consensus binding sites for the CLOCK::BMAL1 heterodimer. E-box binding sites on the *Hcn4* gene and in 20 kb of the 5' flanking region were obtained using the RVISTA function within ECR browser (<http://ecrbrowser.dcode.org/>). Quantification of *Hcn4* E-box binding sites was conducted by, first, normalisation to the housekeeping genes *Gapdh* and *L7* and, secondly, normalisation to data from 3T3-L1 cells subjected to transfection treatment without plasmid

ACKNOWLEDGEMENTS

We thank Ms Rayna Samuels for technical assistance.

AUTHOR CONTRIBUTIONS

Mark R. Boyett, Alicia D'Souza and George Hart conceived the project. Mark R. Boyett, Alicia D'Souza, Hugh D. Piggins, Halina Dobrzynski and Elizabeth Cartwright obtained funding for the project. Anne Berit Johnsen, Alicia D'Souza, Ulrik Wisloff, Eleanor Gill, Charlotte Cox, Yanwen Wang and Nick Ashton were involved in biotelemetry recordings. Yanwen Wang carried out electrophysiological experiments with assistance from Dario DiFrancesco and Annalisa Bucchi in patch clamp recordings. Alicia D'Souza carried out transcriptional profiling, bioluminescence recording and autonomic block experiments with assistance from Cali Anderson, Sven Wegner and Sunil Logantha. Alicia D'Souza, Yu Zhang and Yanwen Wang carried out western blotting experiments. Yanwen Wang and Nicholas Black performed immunofluorescent labelling studies. Haibo Ni and Henggui Zhang performed computer modelling studies using experimental data collected by Yanwen Wang. Yanwen Wang was responsible for the generation, breeding and maintenance of transgenic animals and all experiments on these animals (with assistance from Cheryl Petit, Hugh D. Piggins and Alicia D'Souza). Serve Olieslagers, Alicia D'Souza and Paula da Costa Martins were responsible for chromatin immunoprecipitation experiments. Svetlana Mastitskaya performed vagotomy experiments. Mark R. Boyett and Alicia D'Souza produced the first version of the manuscript, although subsequently all authors contributed.

454

455

SOURCES OF FUNDING

456

This work was supported by the British Heart Foundation (RG/11/18/29257; PG/15/16/31330) and

457

the CARIPLO Foundation (ACROSS 2014-0728).

458

459

COMPETING INTERESTS

460

None declared.

REFERENCES

- 1 Northcote, R. J., Canning, G. P. & Ballantyne, D. Electrocardiographic findings in male veteran endurance athletes. *British Heart Journal* **61**, 155-160 (1989).
- 2 Durgan, D. J. & Young, M. E. The cardiomyocyte circadian clock: emerging roles in health and disease. *Circulation Research* **106**, 647-658 (2010).
- 3 Otsuka, K. *et al.* Experimental study on the relationship between cardiac arrhythmias and sleep states by ambulatory ECG-EEG monitoring. *Clin Cardiol* **9**, 305-313 (1986).
- 4 Vandewalle, G. *et al.* Robust circadian rhythm in heart rate and its variability: influence of exogenous melatonin and photoperiod. *Journal of Sleep Research* **16**, 148-155 (2007).
- 5 Sztajzel, J. Heart rate variability: a noninvasive electrocardiographic method to measure the autonomic nervous system. *Swiss Medical Weekly* **134**, 514-522 (2004).
- 6 Massin, M. M., Maeyns, K., Withofs, N., Ravet, F. & Gerard, P. Circadian rhythm of heart rate and heart rate variability. *Archives of disease in childhood* **83**, 179-182 (2000).
- 7 Nakagawa, M. *et al.* Circadian rhythm of the signal averaged electrocardiogram and its relation to heart rate variability in healthy subjects. *Heart* **79**, 493-496 (1998).
- 8 Monfredi, O. *et al.* Biophysical characterisation of the under-appreciated and important relationship between heart rate variability and heart rate. *Hypertension* **64**, 1334-1343 (2014).
- 9 Degaute, J. P., van de Borne, P., Linkowski, P. & Van Cauter, E. Quantitative analysis of the 24-hour blood pressure and heart rate patterns in young men. *Hypertension* **18**, 199-210 (1991).
- 10 Dilaveris, P. E., Farbom, P., Batchvarov, V., Ghuran, A. & Malik, M. Circadian behavior of P-wave duration, P-wave area, and PR interval in healthy subjects. *Annals of Noninvasive Electrocardiology* **6**, 92-97 (2001).
- 11 Bonnemeier, H. *et al.* Circadian profile of QT interval and QT interval variability in healthy volunteers. *Pacing and Clinical Electrophysiology* **26**, 377-382 (2003).
- 12 LeGates, T. A. & Altimus, C. M. Measuring circadian and acute light responses in mice using wheel running activity. *Journal of Visualized Experiments* **48**, e2463 (2011).
- 13 Veasey, S. C. *et al.* An automated system for recording and analysis of sleep in mice.

490 *Sleep* **23**, 1025-1040 (2000).

491 14 Mohawk, J. A., Green, C. B. & Takahashi, J. S. Central and peripheral circadian clocks in
492 mammals. *Annual Review of Neuroscience* **35**, 445-462 (2012).

493 15 van der Horst, G. T. *et al.* Mammalian Cry1 and Cry2 are essential for maintenance of
494 circadian rhythms. *Nature* **398**, 627-630 (1999).

495 16 DiFrancesco, D. The role of the funny current in pacemaker activity. *Circulation Research*
496 **106**, 434-446 (2010).

497 17 Sinturel, F. *et al.* Diurnal Oscillations in Liver Mass and Cell Size Accompany Ribosome
498 Assembly Cycles. *Cell* **169**, 651-663 e614 (2017).

499 18 Petsakou, A., Sapsis, T. P. & Blau, J. Circadian Rhythms in Rho1 Activity Regulate
500 Neuronal Plasticity and Network Hierarchy. *Cell* **162**, 823-835 (2015).

501 19 Herrero, A., Duhart, J. M. & Ceriani, M. F. Neuronal and glial clocks underlying structural
502 remodeling of pacemaker neurons in drosophila. *Front Physiol* **8**, 918 (2017).

503 20 Nikmaram, M. R., Boyett, M. R., Kodama, I., Suzuki, R. & Honjo, H. Variation in the effects
504 of Cs⁺, UL-FS 49 and ZD7288 within the sinoatrial node. *American Journal of Physiology*
505 **272**, H2782-H2792 (1997).

506 21 DiFrancesco, D. Funny channels in the control of cardiac rhythm and mode of action of
507 selective blockers. *Pharmacological Research* **53**, 399-406 (2006).

508 22 Monfredi, O. J., Dobrzynski, H., Mondal, T., Boyett, M. R. & Morris, G. L. The anatomy and
509 physiology of the sinoatrial node - a contemporary review. *Pacing and Clinical*
510 *Electrophysiology* **33**, 1392-1406 (2010).

511 23 Hilliard, F. A. *et al.* Flecainide inhibits arrhythmogenic Ca²⁺ waves by open state block of
512 ryanodine receptor Ca²⁺ release channels and reduction of Ca²⁺ spark mass. *Journal of*
513 *Molecular and Cellular Cardiology* **48**, 293-301 (2010).

514 24 Dobrzynski, H. *et al.* Expression of Kir2.1 and Kir6.2 transgenes under the control of the α-
515 MHC promoter in the sinoatrial and atrioventricular nodes in transgenic mice. *Journal of*
516 *Molecular and Cellular Cardiology* **41**, 855-867 (2006).

517 25 Bunger, M. K. *et al.* Mop3 is an essential component of the master circadian pacemaker in
518 mammals. *Cell* **103**, 1009-1017 (2000).

- 519 26 Ripperger, J. A. & Schibler, U. Rhythmic CLOCK-BMAL1 binding to multiple E-box motifs
520 drives circadian Dbp transcription and chromatin transitions. *Nat Genet* **38**, 369-374 (2006).
- 521 27 Lee, Y. J., Han, D. H., Pak, Y. K. & Cho, S. H. Circadian regulation of low density
522 lipoprotein receptor promoter activity by CLOCK/BMAL1, Hes1 and Hes6. *Exp Mol Med* **44**,
523 642-652 (2012).
- 524 28 Bosnjak, Z. J. & Kampine, J. P. Effects of halothane, enflurane, and isoflurane on the SA
525 node. *Anesthesiology* **58**, 314-321 (1983).
- 526 29 Skovsted, P. & Saphthavichaikul, S. The effects of isoflurane on arterial pressure, pulse rate,
527 autonomic nervous activity, and barostatic reflexes. *Canadian Anaesthetists' Society*
528 *Journal* **24**, 304-314 (1977).
- 529 30 Ng, G. A., Brack, K. E. & Coote, J. H. Effects of direct sympathetic and vagus nerve
530 stimulation on the physiology of the whole heart - a novel model of isolated Langendorff
531 perfused rabbit heart with intact dual autonomic innervation. *Experimental Physiology* **86**,
532 319-329 (2001).
- 533 31 Basalay, M. V. *et al.* Glucagon-like peptide-1 (GLP-1) mediates cardioprotection by remote
534 ischaemic conditioning. *Cardiovascular Research* **112**, 669-676 (2016).
- 535 32 Leoni, A. L. *et al.* Chronic heart rate reduction remodels ion channel transcripts in the
536 mouse sinoatrial node but not in the ventricle. *Physiological Genomics* **24**, 4-12 (2005).
- 537 33 Gehrman, J. *et al.* Electrophysiological characterization of murine myocardial ischemia
538 and infarction. *Basic Research in Cardiology* **96**, 237-250 (2001).
- 539 34 Berul, C. I. *et al.* Ventricular arrhythmia vulnerability in cardiomyopathic mice with
540 homozygous mutant Myosin-binding protein C gene. *Circulation* **104**, 2734-2739 (2001).
- 541 35 D'Souza, A. *et al.* Exercise training reduces resting heart rate via downregulation of the
542 funny channel HCN4. *Nature Communications* **5**, 3775 (2014).
- 543 36 Aschar-Sobbi, R. *et al.* Increased atrial arrhythmia susceptibility induced by intense
544 endurance exercise in mice requires TNF α . *Nature Communications* **6**, 6018 (2015).
- 545 37 Wang, D. W. *et al.* Propranolol blocks cardiac and neuronal voltage-gated sodium
546 channels. *Frontiers in Pharmacology* **1**, 144 (2010).
- 547 38 Tamura, A. *et al.* Effects of antiarrhythmic drugs on the hyperpolarization-activated cyclic

548 nucleotide-gated channel current. *Journal of Pharmacological Sciences* **110**, 150-159
549 (2009).

550 39 Taylor, E. A. & Turner, P. The distribution of propranolol, pindolol and atenolol between
551 human erythrocytes and plasma. *British Journal of Clinical Pharmacology* **12**, 543-548
552 (1981).

553 40 Chapman, M. E., Hu, L., Plato, C. F. & Kohan, D. E. Bioimpedance spectroscopy for the
554 estimation of body fluid volumes in mice. *American Journal of Physiology - Renal
555 Physiology* **299**, F280-283 (2010).

556 41 Tong, M. *et al.* Circadian expressions of cardiac ion channel genes in mouse might be
557 associated with the central clock in the SCN but not the peripheral clock in the heart.
558 *Biological Rhythm Research* **44**, 519-530 (2013).

559 42 Koike, N. *et al.* Transcriptional architecture and chromatin landscape of the core circadian
560 clock in mammals. *Science* **338**, 349-354 (2012).

561 43 Yamashita, T. *et al.* Circadian variation of cardiac K⁺ channel gene expression. *Circulation*
562 **107**, 1917-1922 (2003).

563 44 Jeyaraj, D. *et al.* Circadian rhythms govern cardiac repolarization and arrhythmogenesis.
564 *Nature* **483**, 96-99 (2012).

565 45 Tong, M. Q. *et al.* Circadian expression of connexins in the mouse heart. *Biological Rhythm
566 Research* **47**, 631-639 (2016).

567 46 Schroder, E. A. *et al.* The cardiomyocyte molecular clock, regulation of Scn5a, and
568 arrhythmia susceptibility. *American Journal of Physiology-Cell Physiology* **304**, C954-965
569 (2013).

570 47 Chen, Y. *et al.* CLOCK-BMAL1 regulate the cardiac L-type calcium channel subunit
571 CACNA1C through PI3K-Akt signaling pathway. *Canadian Journal of Physiology and
572 Pharmacology* **94**, 1023-1032 (2016).

573 48 Schroder, E. A. *et al.* The cardiomyocyte molecular clock regulates the circadian
574 expression of Kcnh2 and contributes to ventricular repolarization. *Heart Rhythm* **12**, 1306-
575 1314 (2015).

576 49 Kodama, I. *et al.* Regional differences in the effects of E-4031 within the sinoatrial node.

- 577 *American Journal of Physiology* **276**, H793-H802 (1999).
- 578 50 Oosting, J., Struijker-Boudier, H. A. & Janssen, B. J. Autonomic control of ultradian and
579 circadian rhythms of blood pressure, heart rate, and baroreflex sensitivity in spontaneously
580 hypertensive rats. *Journal of hypertension* **15**, 401-410 (1997).
- 581 51 Swoap, S. J. *et al.* Vagal tone dominates autonomic control of mouse heart rate at
582 thermoneutrality. *American Journal of Physiology-Heart and Circulatory Physiology* **294**,
583 H1581-1588 (2008).
- 584 52 Kotsis, V. T. *et al.* Impact of cardiac transplantation in 24 hours circadian blood pressure
585 and heart rate profile. *Transplantation Proceedings* **37**, 2244-2246 (2005).
- 586 53 Idema, R. N. *et al.* Decreased circadian blood pressure variation up to three years after
587 heart transplantation. *American Journal of Cardiology* **73**, 1006-1009 (1994).
- 588 54 Ptaszynski, P. *et al.* The effect of ivabradine administration on the night drop of heart rate in
589 patients with inappropriate sinus tachycardia. *Journal of the American College of*
590 *Cardiology* **71**, 392-392 (2018).
- 591 55 Grigoryan, S., Hazarapetyan, L. G. & Kocharyan, S. P. The influence of ivabradine on
592 circadian pattern of heart rate and ischemic episodes in patients with ischemic heart
593 disease and heart failure. *European Heart Journal* **35**, 1012-1012 (2014).
- 594 56 Bray, M. S. *et al.* Disruption of the circadian clock within the cardiomyocyte influences
595 myocardial contractile function, metabolism, and gene expression. *American Journal of*
596 *Physiology-Heart and Circulatory Physiology* **294**, H1036-1047 (2008).
- 597 57 Durgan, D. J. *et al.* The intrinsic circadian clock within the cardiomyocyte. *American Journal*
598 *of Physiology-Heart and Circulatory Physiology* **289**, H1530-1541 (2005).
- 599 58 Beesley, S., Noguchi, T. & Welsh, D. K. Cardiomyocyte circadian oscillations are cell-
600 autonomous, amplified by β -adrenergic signaling, and synchronized in cardiac ventricle
601 tissue. *PLoS One* **11**, e0159618 (2016).
- 602 59 Gula, L. J., Krahn, A. D., Skanes, A. C., Yee, R. & Klein, G. J. Clinical relevance of
603 arrhythmias during sleep: guidance for clinicians. *Heart* **90**, 347-352 (2004).
- 604 60 Alboni, P., Holz, A. & Brignole, M. Vagally mediated atrioventricular block: pathophysiology
605 and diagnosis. *Heart* **99**, 904-908 (2013).

606 61 Manfredini, R. *et al.* Morning preference in onset of symptomatic third-degree
607 atrioventricular heart block. *Chronobiology international* **19**, 785-791 (2002).

608 62 Ector, H. *et al.* Bradycardia, ventricular pauses, syncope, and sports. *Lancet* **2**, 591-594
609 (1984).

610 63 Northcote, R. J., Rankin, A. C., Scullion, R. & Logan, W. Is severe bradycardia in veteran
611 athletes an indication for a permanent pacemaker? *British Medical Journal* **298**, 231-232
612 (1989).

613 64 Yamaguchi, S. *et al.* The 5' upstream region of mPer1 gene contains two promoters and is
614 responsible for circadian oscillation. *Current Biology* **10**, 873-876 (2000).

615 65 Kharche, S., Yu, J., Lei, M. & Zhang, H. A mathematical model of action potentials of
616 mouse sinoatrial node cells with molecular bases. *American Journal of Physiology* **301**
617 (2011).

618

FIGURE LEGENDS

Figure 1. Circadian rhythm in heart rate and other ECG parameters. Circadian rhythm of heart rate, PR interval, QRS duration, QT interval and physical activity (measured using telemetry) in conscious mice (n=9) over ~6 days. Light and shaded regions represent light and dark phases in this and all similar figures – there was a 12 h light/12 h dark cycle for the first three days and constant darkness subsequently. Timing of 1 h light pulses shown. In this and all similar figures, means±SEM are shown and data are fit with a standard sine wave shown in red.

Figure 2. Circadian rhythm in intrinsic heart rate, local circadian clock in sinus node, and circadian rhythm in expression of ion channels and other molecules underlying pacemaker activity of sinus node. **A**, Intrinsic heart rate measured from the Langendorff-perfused heart isolated at ZT 0 and ZT 12 (n=9 and 8 mice). *P<0.05; two-tailed unpaired t test with Welch's correction. **B**, Corrected sinus node recovery time (cSNRT) at three pacing cycle lengths measured from the Langendorff-perfused heart isolated at ZT 0 and ZT 12 (n=9 and 8 mice). *ZT 12 versus ZT 0, P<0.05; two-way ANOVA with Holm-Šídák post-hoc test. **C**, Relative expression of transcripts encoding key clock components in the sinus node at ZT 12 (as compared to ZT 0; n=7 and 9 mice). The vertical line corresponds to 1, i.e. no change. Values <1 correspond to a decrease at ZT 12 and >1 an increase. *ZT 12 versus ZT 0, P<0.05; limma test followed by Benjamini-Hochberg False Discovery Rate correction (set at 5%). **D** and **E**, Expression of *Bmal1* and *Clock* mRNA in the sinus node at four time points over 24 h (n=6 mice for ZT 0; n=6 for ZT 6; n=7 for ZT 12; n=8 for ZT 18). **F**, *Per1* activity (reported by luciferase bioluminescence) in the isolated sinus node from *Per1::LUC* mice on a *Cry1^{+/+}Cry2^{+/+}* or *Cry1^{-/-}Cry2^{-/-}* background. Similar data obtained from two other *Cry1^{+/+}Cry2^{+/+}* mice and two other *Cry1^{-/-}Cry2^{-/-}* mice. **G**, Relative expression of transcripts encoding ion channels, Na⁺-K⁺ pump subunits, intracellular Ca²⁺-handling molecules, gap junction channels and key transcription factors in the sinus node at ZT 12 (as compared to ZT 0; n=7 and 9 mice). *ZT 12 versus ZT 0, P<0.05; Limma test followed by Benjamini-Hochberg False Discovery Rate correction (set at 5%).

Figure 3. Diurnal rhythm in *Hcn4* mRNA and protein in sinus node. **A**, Expression of *Hcn4*

648 mRNA in the sinus node at six time points over 24 h (n=6 mice for ZT 0; n=6 for ZT 4; n=7 for ZT 8;
649 n=7 for ZT 12; n=7 for ZT 16; n=5 for ZT 20). **B**, Western blot for HCN4 and the housekeeper, β -
650 actin, in the sinus node of 5 mice culled at ZT 0 and 5 mice culled at ZT 12. **C**, Mean HCN4 protein
651 expression (relative to the expression of the housekeeper) in the sinus node determined by
652 western blot at ZT 0 and ZT 12. * $P < 0.05$; two-tailed unpaired t test with Welch's correction. **D**,
653 Immunolabelling of HCN4 protein (red signal) in sections through the sinus node dissected from
654 control wild-type (top) and cardiac-specific *Bmal1* knockout mice culled at ZT 0 and ZT 12. **E**,
655 Mean HCN4 protein expression determined by immunohistochemistry in the sinus node of control
656 wild-type and cardiac-specific *Bmal1* knockout mice at ZT 0 and ZT 12 (wild-type mice, n=56/42
657 sections from 3/3 mice; cardiac-specific *Bmal1* knockout mice, n=38/46 sections from 3/3 mice).
658 * $P < 0.05$; two-way ANOVA with Holm-Šídák post-hoc test.

659

660 **Figure 4. Circadian rhythm in I_f in sinus node.** **A**, Families of recordings of I_f from sinus node
661 cells made at ZT 2 and ZT 12. **B**, Current-voltage relationships for I_f recorded at ZT 2 (n=10 cells/3
662 mice) and ZT 12 (n=16 cells/3 mice). *ZT 12 versus ZT 0, $P < 0.05$; two-way ANOVA with Holm-
663 Šídák post-hoc test. **C**, Density of I_f at -125 mV over 24 h (n=10 cells for ZT 2; n=11 for ZT 3; n=10
664 for ZT 4; n=12 for ZT 5; n=8 for ZT 6; n=16 for ZT 7; n=14 for ZT 8; n=14 for ZT 9; n=27 for ZT 10;
665 n=16 for ZT 11; n=16 for ZT 12; n=18 for ZT 13; n=13 for ZT 14; n=15 for ZT 15; n=17 for ZT 16;
666 n=16 for ZT 17; n=11 for ZT 18; from 24 mice); I_f density is plotted against the time of recording. **D**,
667 Amplitude of I_f at -125 mV over 24 h (from same experiments as for C). **E**, C_m over 24 h (from
668 same experiments as for C). **F**, Computed mouse sinus node action potentials at ZT 2 and ZT 12
669 based on a model incorporating the measured differences in the density of I_f . **G**, Computed 'heart
670 rate' of a mouse sinus node cell over 24 h based on a model incorporating the measured
671 differences in density of I_f .

672

673 **Figure 5. Effect of block of HCN4 and I_f on intrinsic heart rate and heart rate *in vivo*.** **A**,
674 Intrinsic heart rate before and after the application of 2 mM Cs^+ (top) and change in intrinsic heart
675 rate after application of Cs^+ (bottom) measured in the isolated sinus node from wild-type mice
676 (n=10 mice for ZT 0; n=5 for ZT 6; n=9 for ZT 12; n=4 for ZT 18). **B**, Intrinsic heart rate before and

after the application of 2 mM Cs⁺ (top) and change in intrinsic heart rate after application of Cs⁺ (bottom) measured in the isolated sinus node from cardiac-specific *Bmal1* knockout mice (n=8 mice for ZT 0; n=5 for ZT 6; n=4 for ZT 12; n=3 for ZT 18). **C**, *In vivo* heart rate of wild-type mice measured at ZT 0 and ZT 12 before and after the administration of 6 mg/kg ivabradine (n=5 mice for ZT 0; n=5 for ZT 12). *P<0.05; two-way ANOVA with Dunnett's post-hoc test. **D**, Change in *in vivo* heart rate after administration of ivabradine (from same experiments as for C). *P<0.05; two-tailed paired t-test.

Figure 6. Cardiac-specific knockout of *Bmal1* stops or blunts circadian rhythm in *Hcn4* and *I_f*. **A, B and C**, Expression of *Bmal1*, *Clock* and *Hcn4* mRNA in the sinus node at ZT 0 and ZT 12 in the sinus node of wild-type mice (n=7 mice for ZT 0 and ZT 12) and cardiac-specific *Bmal1* knockout mice (n=7 mice for ZT 0 and ZT 12). *ZT 12 versus corresponding ZT 0, P<0.05; two-way ANOVA with Tukey's post-hoc test. **D**, Families of recordings of *I_f* made at ZT 2 and ZT 12 from sinus node cells isolated from cardiac-specific *Bmal1* knockout mice. **E**, Current-voltage relationships for *I_f* recorded at ZT 2 and ZT 12. Data shown in black were obtained from cardiac-specific *Bmal1* knockout mice (n=15 cells/3 mice at ZT 0; n=16 cells/3 mice at ZT 2). Data shown in grey were obtained from wild-type mice and have already been shown in Figure 4B. **F**, Density of *I_f* recorded at ZT 2 and ZT 12 from sinus node cells isolated from wild-type mice (n=16 cells/4 mice for ZT 2; n=16 cells/5 mice for ZT 12) and cardiac-specific *Bmal1* knockout mice (n=14 cells/3 mice for ZT 2; n=15 cells/3 mice for ZT 12). *I_f* expressed as a percentage of *I_f* at ZT 0. *P<0.05; two-way ANOVA with Holm-Šídák post-hoc test. **G**, Eight potential E-box binding sites pulled down on immunoprecipitation of His-tagged BMAL1 from cells transfected with His-tagged *Bmal1*. Data are normalised to immunoprecipitation from untransfected control cells; the red dashed line equals one and is the baseline level. *binding site of interest versus binding site A, P<0.05; n=2; one-way ANOVA with Dunnett's post-hoc test. **H**, Diagram of the *Hcn4* gene and 20 kb of the 5' flanking region showing the position of the eight potential E-box binding sites, A-H.

Figure 7. Circadian rhythm in heart rate *in vivo*. **A**, *In vivo* heart rate and physical activity measured by telemetry at the times shown during 24 h darkness (subjective day and night is

706 shown) with the exception of a 1 h light pulse delivered towards the start of the day (left) or night
 707 (right). The dotted red lines highlight the heart rate and physical activity at the end of the day-time
 708 light pulse and the red arrows highlight the heart rate and physical activity at the end of the night-
 709 time light pulse. From the same experiment as Figure 1. **B**, *In vivo* heart rate measured from
 710 anaesthetised mice at four time points over 24 h (n=12 mice for ZT 0; n=10 for ZT 6; n=10 for ZT
 711 12; n=10 for ZT 18). **C**, *In vivo* heart rate measured by telemetry at ZT 0 and ZT 12 in sham-
 712 operated and vagotomised rats at baseline (pre-surgery) and at 1, 3 and 7 days post-surgery
 713 *P<0.05. two-way ANOVA with Sidak's multiple comparisons test. **D**, *In vivo* heart rate measured
 714 from anaesthetised mice at ZT 0 (n=9) and ZT 12 (n=9) before (Control) and after autonomic block
 715 by intraperitoneal injection of 1 mg/kg atropine and 1 mg/kg propranolol. *ZT 12 versus
 716 corresponding ZT 0, P<0.05; two-way ANOVA with Tukey's post-hoc test. **E**, *In vivo* heart rate
 717 measured from anaesthetised mice at ZT 0 (n=5) and ZT 12 (n=5) before (Control) and after
 718 autonomic block by intraperitoneal injection of 2 mg/kg atropine and 10 mg/kg propranolol. *ZT 12
 719 versus corresponding ZT 0, P<0.05; two-way ANOVA with Tukey's post-hoc test. **F**, Density of I_f at
 720 -125 mV measured from 12 sinus node cells (from 3 mice) before and after application of 68.4 μ M
 721 propranolol. *P<0.05; paired t test.

In vivo

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7