

# Rare protein-altering variants in *ANGPTL7* lower intraocular pressure and protect against glaucoma

5 Yosuke Tanigawa<sup>1</sup>, Michael Wainberg<sup>1</sup>, Juha Karjalainen<sup>2,3,4</sup>, Tuomo Kiiskinen<sup>4,5</sup>,  
Susanna Lemmelä<sup>4,5</sup>, Joni A. Turunen<sup>6,7</sup>, Robert Graham<sup>8</sup>, Aki S. Havulinna<sup>4,5</sup>, Markus Perola<sup>5</sup>,  
Aarno Palotie<sup>2,4</sup>, FinnGen, Mark J. Daly<sup>2,3,4+</sup> and Manuel A. Rivas<sup>1+</sup>

1. Department of Biomedical Data Science, School of Medicine, Stanford University, Stanford, CA, USA.
2. Program in Medical and Population Genetics and Stanley Center for Psychiatric Research, Broad Institute of Harvard and MIT, Cambridge, MA, USA.
- 10 3. Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, MA, USA.
4. Institute for Molecular Medicine Finland (FIMM), University of Helsinki, Helsinki, Finland.
5. Finnish Institute for Health and Welfare, Helsinki, Finland
6. Department of Ophthalmology, University of Helsinki and Helsinki University Hospital, Helsinki, Finland
7. Folkhälsan Research Center, Biomedicum Helsinki, Helsinki, Finland
- 15 8. Maze Therapeutics, San Francisco, CA, USA.
- + Corresponding authors, contributed equally to the work

**Protein-altering variants that are protective against human disease provide *in vivo* validation of therapeutic targets. Here we use genotyping data in UK Biobank and FinnGen to conduct a search for protein-altering variants conferring lower intraocular pressure (IOP) and protection against glaucoma. Through protein-altering variant association analysis we find a missense variant in UK Biobank (rs28991009 (MAF=0.8%) genotyped in 81,527 individuals with measured IOP and an independent set of 4,269 glaucoma patients and 251,355 controls) that significantly lowers IOP ( $\beta = -0.73$  mmHg for heterozygotes,  $-2.96$  mmHg for homozygotes,  $P = 1 \times 10^{-13}$ ) and is associated with 34% reduced risk of glaucoma ( $P = 0.005$ ). In FinnGen, we identify an *ANGPTL7* missense variant at a greater than 50-fold increased frequency in Finland compared with other populations (rs147660927, p.Arg220Cys, MAF Finland = 4.1%), which was genotyped in 5,177 glaucoma patients and 130,461 controls and is associated with 30% lower glaucoma risk ( $P = 1 \times 10^{-9}$ ). We further find three rarer variants in UK Biobank, including a protein-truncating variant, which confer a strong composite lowering of IOP ( $P = 0.002$ ), suggesting the protective mechanism likely resides in the loss of an interaction or function. Our results support inhibition or down-regulation of *ANGPTL7* as a therapeutic strategy for glaucoma.**

## Introduction

35 Intraocular pressure is currently the sole modifiable risk factor and predictive measure for  
glaucoma<sup>1,2,3,4</sup> (**Supplementary Figure S1**). Genome-wide association studies (GWAS) have  
commonly used genetic associations to this endophenotype that exhibits high genetic  
correlation ( $r_g = 0.71$ ) to glaucoma, as an approach to prioritize genetic variants likely to  
40 contribute to disease risk<sup>5</sup>. A total of 53 independent loci have been unequivocally implicated in  
glaucoma<sup>5</sup>. For these discoveries, like most GWAS results, it has proven challenging to infer the  
functional consequences of common variant associations beyond cases where protein-altering  
variants have been directly implicated. Protein-altering variants, generally the strongest-acting  
genetic variants in medical genetics, include nonsynonymous substitutions and  
45 protein-truncating variants, and understanding their functional consequences provides insight  
into the therapeutic effects of inhibiting or down-regulating the gene in which they reside<sup>6</sup>. Thus,  
identifying protein-altering variants that confer protection from disease holds particular promise  
for identifying therapeutic targets.

Here we leverage two cohorts that provide complementarity for glaucoma gene discovery. First,  
UK Biobank has obtained intraocular pressure (IOP) measurements in approximately 128,000  
50 individuals in addition to case-control status for glaucoma from hospital in-patient and verbal  
questionnaire data in over 500,000 individuals<sup>7,8</sup>. Second, FinnGen has directly genotyped and  
aggregated disease outcomes in over 135,000 individuals from Finland, an isolated population  
with recent bottlenecks that offers an unprecedented advantage for studying rare variants in  
complex diseases<sup>9</sup>. With clinic-based recruitment focused on several areas including  
55 ophthalmology, and with 32.4% of the collection above age 70, FinnGen is particularly  
well-powered for aging-associated endpoints. We therefore conduct targeted association  
analysis with IOP in UK Biobank ( $N = 81,534$ ) to identify protein-altering variants that reduce  
IOP, and test whether those variants, or others in the same genes, also confer protection to  
glaucoma in FinnGen (5,177 cases and 130,461 controls) and UK Biobank (4,269 cases and  
60 251,355 controls not included in the IOP sample). Analysis of an allelic series of protein-altering  
variants in *ANGPTL7* in 9,446 glaucoma patients and over 350,000 controls identifies a  
significant lowering effect on IOP and protective association with glaucoma. By analyzing  
putative loss-of-function variants, we find concordant effect directions with the nonsynonymous  
substitutions, suggesting that the protective mechanism may reside in the loss of gene function.

## 65 Results

### Protein-altering variant association analysis

We conducted protein-altering variant association analysis with IOP, as measured via  
Goldmann-correlated tonometry, in 81,527 British individuals in UK Biobank dataset

(Methods)<sup>10</sup>. Across 41,632 rare (0.01 % < MAF < 1%) protein-altering variants outside of the MHC region with genotyping array data in UK Biobank, we used a generalized linear model implemented in PLINK<sup>11</sup> to scan for variants with IOP-lowering effects. We identified one protein-altering variant significantly associated with lower IOP below a Bonferroni-corrected  $P$  value <  $1.0 \times 10^{-6}$ , a nonsynonymous substitution (p.Gln175His) in *ANGPTL7* ( $P = 1.47 \times 10^{-13}$ ,  $\beta = -0.200$  SD 95% CI: [-0.253, -0.147], -0.73 mmHg for heterozygotes, -2.96 mmHg for homozygotes, **Table 1,2, Supplementary Figure S2**).

Based on this finding, we assessed whether any additional rare variant associations in *ANGPTL7* were present. We found three additional independent rare protein-altering variants in *ANGPTL7* (MAF < 0.25%) including a premature stop-gain allele (p.Arg177Ter). Collectively, these three variants showed a marginally significant association with lower IOP ( $P = 0.0023$ , **Table 1**), with the stop-gain allele p.Arg177Ter also showing a marginally significant effect on its own ( $P = 0.038$ ). Genotyping intensity plots and the concordance of genotype calls from array and whole-exome sequencing data were manually inspected to ensure high quality and consistent genotyping (**Supplementary Figure S3, Supplementary Table S1, Methods**) and alleles were confirmed to be independent (pairwise  $r^2 < 10^{-4}$  for each, **Supplementary Table S2, Methods**). Using inverse variance weighted meta-analysis method, the combined effect of those four variants are estimated to be -0.19 ([95% CI: -0.23 - -0.14] SD,  $P = 3.4 \times 10^{-16}$ ; -0.67 mmHg [95% CI: -0.85 - -0.49], **Supplementary Figure S4, Supplementary Tables S3-4, Methods**).

Given these findings, we next asked whether any of these putative IOP-lowering genetic variants showed effects consistent with reducing glaucoma risk in an independent set of British individuals that do not have IOP measures (4,269 cases and 251,355 controls). For p.Gln175His in *ANGPTL7*, we estimated that the variant lowers glaucoma risk by 34% ( $P = 0.00543$ ; OR = 0.66 [95% CI: 0.366 - 0.954]). The three additional protein-altering variants did not significantly confer protection against glaucoma (burden test  $P = 0.55$ ), although we expect this to be a result of limited power in the binary case-control setting.

Variant	rsID	HGVSp	MAF	Beta SD [95% CI]	$P$
1:11252357:A:G	rs200058074	p.Gln136Arg	.054 %	-.038 [-.26, .18]	$7.3 \times 10^{-1}$
1:11252369:G:A	rs28991002	p.Arg140His	.25 %	-.16 [-.25, -.061]	$1.3 \times 10^{-3}$
1:11253684:G:T	rs28991009	p.Gln175His	.81 %	-.20 [-.25, -.15]	$1.4 \times 10^{-13}$
1:11253688:C:T	rs143435072	p.Arg177Ter	.041 %	-.26 [-.51, -.014]	$3.8 \times 10^{-2}$

**Table 1. *ANGPTL7* IOP protein-altering variant association in UK Biobank.** Variant includes chromosome, position, reference, and alternate allele (hg19). rsID - the rs identifier of the genetic variant. HGVSp - the HGVS

protein sequence name. MAF - the minor allele frequency in UK Biobank British population. Beta - estimated regression coefficient with 95% confidence intervals. *P* - p-value of association.

Variant	Dosage	rsID	Carrier frequency in UK	Carrier frequency in Finland	N with IOP (UK Biobank)	IOP (mmHg) [95% CI]	OR for glaucoma
Gln175His	Gln/His	rs28991009	1.43%	0.24%	1355	-0.73 [-0.95, -0.52] ***	0.66 **
Gln175His	His/His		0.01%	NA	5	-2.95 [-6.43, 0.530] .	NS
Arg177Ter	Arg/Ter	rs143435072	0.07%	NA	62	-0.94 [-1.93, 0.0445] .	NS
Arg140His	Arg/His	rs28991002	0.51%	0.35%	427	-0.48 [-0.86, -0.10] *	NS
Gln136Arg	Gln/Arg	rs200058074	0.11%	NA	80	-0.21 [-1.08, 0.66]	NS
Arg220Cys	Arg/Cys	rs147660927	NA	7.82%	NA	NA	0.72 ***
Arg220Cys	Cys/Cys		NA	0.21%	NA	NA	0.47 .

100 Significant codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 '.' 1

**Table 2. *ANGPTL7* allelic series summary.** Variant: HGVS amino acid nomenclature. Dosage - genotype of individuals. rsID - rs identifier. Carrier frequency in UK - carrier frequency in UK Biobank for the respective genotype dosage. Carrier frequency in Finland - carrier frequency in FinnGen for the respective genotype dosage. N with IOP (UK Biobank) - number of individuals in UK Biobank with intraocular pressure measurements corresponding to the genotype dosage. IOP (mmHg) [95% CI] - unstandardized estimates of effect size on intraocular pressure measurements (NB: standardized estimated effect sizes may have lower p-values due to normalization procedure). OR for glaucoma - estimate odds ratio on glaucoma risk for the respective genotype dosage. NS non-significant ( $p > 0.1$ ).

## Independent Finnish-enriched missense allele in *ANGPTL7*

110 We then sought evidence from the FinnGen dataset that either the same or novel Finnish-enriched protein-altering variants would confirm the association of *ANGPTL7* with protection from glaucoma. This additional *ANGPTL7* association data, obtained in 5,177 glaucoma patients and 130,461 controls, provided strong support that protein-altering variants in *ANGPTL7* protect against glaucoma (case definitions described in [Supplementary Table S5](#)).

115 Specifically, we found that the p.Gln175His substitution has nominal evidence of association ( $P = 0.031$ , OR = 0.49) despite the variant only being present at a minor allele frequency of 0.1% in this Finnish cohort (8-fold depleted compared to UK Biobank). The remaining protein-altering

120 variants in *ANGPTL7* tested in UK Biobank were not found in the FinnGen dataset. Confirmation of an *ANGPTL7* effect on glaucoma risk was seen in data from an independent Finnish-specific protein-altering nonsynonymous substitution, p.Arg220Cys, which was strongly associated with protection from glaucoma ( $P = 1.0 \times 10^{-9}$ , OR = 0.72 [95% CI: 0.64 - 0.80], **Supplementary Figure S5**). Of note, this observation is advantaged by the property that p.Arg220Cys is found at a greater than 50-fold increased frequency in Finland compared with other populations<sup>12</sup>, reinforcing the value of isolated, bottlenecked populations in which the allele frequency spectrum is intensely concentrated on the minority of variants passing through the bottleneck.

130 While registry-based diagnoses in FinnGen do not yet contain detailed ophthalmologic records, a subset of 2695 glaucoma patients had been recorded as having primary open-angle glaucoma (POAG). In this sub-group a stronger effect was observed (OR=0.68) versus those glaucoma cases without a definitive record of POAG (OR=0.77 [95% CI: 0.67 - 0.88]), reminiscent of the stronger risk effects seen at the myocilin (*MYOC*) gene and other established genes in the POAG subgroup. Given the Finnish enrichment of the known strong glaucoma risk allele, p.Gln368Ter, in *MYOC* (MAF in Finland = 0.3%, MAF in Non-Finnish European = 0.16%, reference sequence: NM\_00026), we next asked whether carriers have risk reduced if they carry *ANGPTL7* p.Arg220Cys. In FinnGen, we estimate that 7.7% of individuals carriers for *MYOC* p.Gln368Ter variant are POAG cases in comparison to 2% for non-carriers. In the presence of *ANGPTL7* p.Arg220Cys, only 1.3% of individuals are POAG cases, and only 1 of 70 (1.4%) who carry both *MYOC* risk and *ANGPTL7* protective variants were POAG cases (**Supplementary Table S6**). This suggests *ANGPTL7* protection extends to the *MYOC* risk group but the small counts preclude any definitive statement regarding interaction - given the limited number of double-carriers, larger case-control series are needed to refine our understanding as to whether *ANGPTL7* p.Arg220Cys variant modifies the glaucoma risk conferred by p.Gln368Ter in *MYOC*.

145 Access to genotype data in over 300,000 individuals from the UK and 135,000 from the Finnish group enabled us to identify rare protein-altering homozygotes. In UK Biobank we found 28 individuals homozygous for the 175His allele, consistent with Hardy-Weinberg expectation ( $n=22.6$ ), where we estimated a 2.96 mm Hg drop in IOP compared to the mean IOP levels. Furthermore, the oldest reached the age of 80 and one of the 28 died (age 65). In FinnGen we found 268 individuals homozygous for the 220C allele, the oldest reached the age of 98, with no depletion of homozygotes compared with Hardy-Weinberg equilibrium expectation. There was no significant association of the homozygous genotype with a decreased lifespan. Together this indicates that having two copies with 175His or 220Cys in *ANGPTL7* is compatible with normal lifespan. To assess the potential impacts of those protein-truncating variants on reproductive fitness we assessed the association of p.Gln175His with the number of live births and number of children fathered and found no significant association ( $P > 0.05/4$ , **Supplementary Table S7**). Hence, we did not find any severe medical consequences that would be of obvious concern in developing a therapeutic to mimic the effect of these alleles.

The combined significance of *ANGPTL7* protein-altering variants across more than 435,000 samples tested is  $P = 3.4 \times 10^{-16}$  for IOP and  $P = 1.9 \times 10^{-11}$  for glaucoma using inverse-variance weighted method (**Supplementary Figure S6**).

## 160 Transcript and protein expression

*ANGPTL7*, a five-exon protein-coding gene, encodes the Angiotensin-related protein 7, which is expressed in several human tissues including the trabecular meshwork, cornea and retina<sup>13–15</sup>. Recently, *ANGPTL7* overexpression in primary human trabecular meshwork cells was found to alter the expression of relevant trabecular meshwork proteins of the extracellular matrix, including fibronectin, collagens type I, IV, and V, myocilin, versican, and MMP1, and *ANGPTL7* protein was increased as the disease progressed in POAG beagle dogs<sup>13</sup>. We examined proteomics expression data in normal tissues and cell lines from ProteomicsDB and MOPED<sup>16,17</sup> and found vitreous humor tissue-specific expression of *ANGPTL7* ( $\log_{10}$  ppm = 1, **Supplementary Figure S7**). These data suggest that the eye is the relevant tissue type in understanding the functional consequences of the variants discovered in this study. Further work is dissecting the role of *ANGPTL7* in all possible cell types in the eye is warranted.

## Discussion

This study establishes strong genetic evidence for the involvement of *ANGPTL7* in glaucoma risk in which a powerful allelic series, including all low-frequency nonsynonymous substitutions and a single premature stop-gain substitution, is conclusively associated with reduced disease risk and endophenotype-lowering effects. Increasing evidence is associating *ANGPTL7* proteins to cardiometabolic phenotypes<sup>18–22</sup>. Although it has been recently proposed that *ANGPTL7* levels are increased in obesity and reduced after physical exercise, we do not observe any evidence of genetic association in either UK Biobank or FinnGen to support this hypothesis<sup>23</sup>. In the context of the other established variants in glaucoma, including the protein-truncating variants in *MYOC*, p.Gln175His and the 57-fold Finnish enriched p.Arg220Cys variant in *ANGPTL7* exert a comparable protective effect. Because of the strong protective effect associated with the *ANGPTL7* coding variants, studies of *ANGPTL7* inhibition and the specific action of these variant proteins should be useful in understanding the mechanism by which protection to glaucoma disease occurs and whether this reveals a promising therapeutic opportunity similar to that which has been realized from the examples of *PCSK9*, *APOC3* and cardiovascular disease<sup>24–26</sup>. The phenotype and longevity profile of *ANGPTL7* homozygotes suggest that this is likely to be a safe strategy for therapeutic intervention.

## Acknowledgments

190 This research has been conducted using the UK Biobank Resource under Application Number 24983, “Generating effective therapeutic hypotheses from genomic and hospital linkage data” (<http://www.ukbiobank.ac.uk/wp-content/uploads/2017/06/24983-Dr-Manuel-Rivas.pdf>). Based on the information provided in Protocol 44532 the Stanford IRB has determined that the research does not involve human subjects as defined in 45 CFR 46.102(f) or 21 CFR 50.3(g).

195 All participants of UK Biobank provided written informed consent (more information is available at <https://www.ukbiobank.ac.uk/2018/02/gdpr/>). For the Finnish Institute of Health and Welfare (THL) driven FinnGen preparatory project (here called FinnGen), all patients and control subjects had provided informed consent for biobank research, based on the Finnish Biobank Act. Alternatively, older cohorts were based on study specific consents and later transferred to the THL Biobank after approval by Valvira, the National Supervisory Authority for Welfare and Health. Recruitment protocols followed the biobank protocols approved by Valvira. The Ethical Review Board of the Hospital District of Helsinki and Uusimaa approved the FinnGen study protocol Nr HUS/990/2017. The FinnGen preparatory project is approved by THL, approval numbers THL/2031/6.02.00/2017, amendments THL/341/6.02.00/2018, THL/2222/6.02.00/2018

200 and THL/283/6.02.00/2019. All DNA samples and data in this study were pseudonymized. We thank all the participants in the UK Biobank and Finnish Biobanks used in FinnGen studies. This work was supported by National Human Genome Research Institute (NHGRI) of the National Institutes of Health (NIH) under awards R01HG010140 (M.A.R.). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. We thank Sirpa Soini, Fatima Rodriguez and David Amar for invaluable feedback on the manuscript. Y.T. is supported by Funai Overseas Scholarship from Funai Foundation for Information Technology and the Stanford University School of Medicine. FinnGen is supported by Abbvie, Astra Zeneca, Biogen, Celgene, Genentech, GSK, Merck, Pfizer, and Sanofi. M.A.R. is supported by Stanford University and a National Institute of Health center for

205 Multi- and Trans-ethnic Mapping of Mendelian and Complex Diseases grant (5U01 HG009080).

210

215

## Author information

### Author contributions

M.A.R. and M.J.D. conceived and designed the study. M.P. designed the FinnGen preparatory project in collaboration with A.P and M.J.D. M.A.R., Y.T., J.K., and M.J.D. carried out the

220 statistical and computational analyses. M.A.R., Y.T., J.K., T.K., J.A.T., and M.J.D. carried out quality control of the data. The manuscript was written by M.A.R., Y.T., and M.J.D.; and revised by all the co-authors. All co-authors have approved of the final version of the manuscript.

## Competing financial interests

Robert Graham is an employee of MazeTx.

225

## Data availability:

Data is displayed in the Global Biobank Engine (<https://biobankengine.stanford.edu>). Analysis scripts and notebooks are available on GitHub at <https://github.com/rivas-lab/ANGPTL7>.

## References

1. Cantor, L. B. Brimonidine in the treatment of glaucoma and ocular hypertension.  
230 *Therapeutics and Clinical Risk Management* **2**, 337–346 (2006).
2. Kass, M. A. *et al.* The Ocular Hypertension Treatment Study: a randomized trial determines that topical ocular hypotensive medication delays or prevents the onset of primary open-angle glaucoma. *Arch. Ophthalmol.* **120**, 701–13; discussion 829–30 (2002).
3. Kazemian, P., Lavieri, M. S., Van Oyen, M. P., Andrews, C. & Stein, J. D. Personalized  
235 Prediction of Glaucoma Progression Under Different Target Intraocular Pressure Levels Using Filtered Forecasting Methods. *Ophthalmology* **125**, 569–577 (2018).
4. Group, T. J. A. of M. D. in G. (jamdig) C. & The Japanese Archive of Multicentral Database in Glaucoma (JAMDIG) construction group. A novel method to predict visual field progression more accurately, using intraocular pressure measurements in glaucoma  
240 patients. *Scientific Reports* **6**, (2016).
5. MacGregor, S. *et al.* Genome-wide association study of intraocular pressure uncovers new pathways to glaucoma. *Nat. Genet.* **50**, 1067–1071 (2018).
6. Leshchiner, E. S. *et al.* Small-molecule inhibitors directly target CARD9 and mimic its protective variant in inflammatory bowel disease. *Proc. Natl. Acad. Sci. U. S. A.* **114**,



- 245 11392–11397 (2017).
7. Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data. *Nature* **562**, 203–209 (2018).
  8. DeBoever, C. *et al.* Medical relevance of protein-truncating variants across 337,205 individuals in the UK Biobank study. *Nat. Commun.* **9**, 1612 (2018).
  - 250 9. Lim, E. T. *et al.* Distribution and medical impact of loss-of-function variants in the Finnish founder population. *PLoS Genet.* **10**, e1004494 (2014).
  10. Chan, M. P. Y. *et al.* Associations with Intraocular Pressure in a Large Cohort: Results from the UK Biobank. *Ophthalmology* **123**, 771–782 (2016).
  11. Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer  
255 datasets. *Gigascience* **4**, 7 (2015).
  12. Lek, M. *et al.* Analysis of protein-coding genetic variation in 60,706 humans. *Nature* **536**, 285–291 (2016).
  13. Comes, N., Buie, L. K. & Borrás, T. Evidence for a role of angiopoietin-like 7 (ANGPTL7) in extracellular matrix formation of the human trabecular meshwork: implications for  
260 glaucoma. *Genes Cells* **16**, 243–259 (2011).
  14. Santulli, G. Angiopoietin-like proteins: a comprehensive look. *Front. Endocrinol.* **5**, 4 (2014).
  15. Kuchtey, J. *et al.* Angiopoietin-like 7 secretion is induced by glaucoma stimuli and its concentration is elevated in glaucomatous aqueous humor. *Invest. Ophthalmol. Vis. Sci.* **49**,  
265 3438–3448 (2008).
  16. Kolker, E. *et al.* MOPED: Model Organism Protein Expression Database. *Nucleic Acids Res.* **40**, D1093–9 (2012).
  17. Schmidt, T. *et al.* ProteomicsDB. *Nucleic Acids Res.* **46**, D1271–D1281 (2018).

18. Musunuru, K. & Kathiresan, S. Cardiovascular endocrinology: Is ANGPTL3 the next  
270 PCSK9? *Nat. Rev. Endocrinol.* **13**, 503–504 (2017).
19. Musunuru, K. *et al.* Exome sequencing, ANGPTL3 mutations, and familial combined  
hypolipidemia. *N. Engl. J. Med.* **363**, 2220–2227 (2010).
20. Gusarova, V. *et al.* Genetic inactivation of ANGPTL4 improves glucose homeostasis and is  
associated with reduced risk of diabetes. *Nat. Commun.* **9**, 2252 (2018).
- 275 21. Clapham, K. R. *et al.* A null mutation in ANGPTL8 does not associate with either plasma  
glucose or type 2 diabetes in humans. *BMC Endocr. Disord.* **16**, 7 (2016).
22. Peloso, G. M. *et al.* Association of low-frequency and rare coding-sequence variants with  
blood lipids and coronary heart disease in 56,000 whites and blacks. *Am. J. Hum. Genet.*  
**94**, 223–232 (2014).
- 280 23. Abu-Farha, M. *et al.* Plasma and adipose tissue level of angiopoietin-like 7 (ANGPTL7) are  
increased in obesity and reduced after physical exercise. *PLoS One* **12**, e0173024 (2017).
24. Jørgensen, A. B., Frikke-Schmidt, R., Nordestgaard, B. G. & Tybjærg-Hansen, A.  
Loss-of-function mutations in APOC3 and risk of ischemic vascular disease. *N. Engl. J.*  
*Med.* **371**, 32–41 (2014).
- 285 25. TG and HDL Working Group of the Exome Sequencing Project, National Heart, Lung, and  
Blood Institute *et al.* Loss-of-function mutations in APOC3, triglycerides, and coronary  
disease. *N. Engl. J. Med.* **371**, 22–31 (2014).
26. Cohen, J. C., Boerwinkle, E., Mosley, T. H., Jr & Hobbs, H. H. Sequence variations in  
PCSK9, low LDL, and protection against coronary heart disease. *N. Engl. J. Med.* **354**,  
290 1264–1272 (2006).
27. Tanigawa, Y. *et al.* Components of genetic associations across 2,138 phenotypes in the UK  
Biobank highlight novel adipocyte biology. *bioRxiv* 442715 (2019). doi:10.1101/442715

28. McCarthy, M. & O'Brien, E. McCarthy Group Front page. *McCarthy Group* (2018). Available at: <http://mccarthy.well.ox.ac.uk/>. (Accessed: 20th May 2019)
- 295 29. Van Hout, C. V. *et al.* Whole exome sequencing and characterization of coding variation in 49,960 individuals in the UK Biobank. *bioRxiv* 572347 (2019). doi:10.1101/572347

## Methods

### Genome-wide association analysis in UK Biobank

For British individuals ( $n = 337,151$ ) in UK Biobank as described elsewhere<sup>8</sup>, we applied genome-wide association analysis for directly genotyped variants were applied to Goldmann-correlated intraocular pressure described more detailed earlier<sup>10</sup> (right, UK Biobank Field ID 5255, Global Biobank Engine phenotype ID: INI5255) and glaucoma (Global Biobank Engine phenotype ID: HC276), using generalized linear model association analysis implemented in PLINK v2.00aLM (2 April 2019) with age, sex, types of genotyping array, and the first 4 genotype principal components as described elsewhere<sup>11,27</sup>. The glaucoma phenotype was previously defined as a part of “high confidence” disease outcome phenotypes by combining disease diagnoses (UK Biobank Field ID 41202) from the UK National Health Service Hospital Episode Statistics (ICD10 codes: H40.[0-6,8,9], H42.8, and Q15.0) with self-reported non-cancer diagnosis questionnaire (UK Biobank Field ID 20002)<sup>8</sup>.

310

### Genotyping quality control in UK Biobank

#### Manual inspection of intensity plots

For the identified rare ( $0.01\% < \text{MAF} < 1\%$ ) protein-altering variants in *ANGPTL7* (reference sequence: NM\_021146), we generated and inspected intensity plots with McCarthy Group's ScatterShot using “UKB - All Participants” module<sup>28</sup>.

315

#### Variant-calling consistency analysis

For individuals with whole-exome sequencing data ( $n = 49,960$ ), we extracted the genotype calls of coding variants in *ANGPTL7* using PLINK v2.00aLM (2 April 2019) and compared the consistency between array-genotyped dataset and whole-exome sequencing dataset<sup>11,29</sup>.

## Independence analysis of alleles

320

### Pairwise $r^2$ computation within British individuals in UK Biobank

We computed pairwise  $r^2$  for the identified rare protein-altering variants in *ANGPTL7* within British individuals in UK Biobank using PLINK v1.90b6.7 64-bit (2 Dec 2018) with --ld <variant\_ID\_1> <variant\_ID\_1> hwe-midp subcommand<sup>11</sup>.

### Number of individuals with the combination of genotypes in UK Biobank

325

Using the extracted genotype calls from for the identified rare protein-altering variants in *ANGPTL7* (see Variant-calling consistency analysis section), we counted the number of British individuals by the combination of genotypes. We computed the expected number of individuals under Hardy-Weinberg equilibrium model and the independence assumption:

330

- The expected frequencies of REF/REF, REF/ALT, and ALT/ALT carriers are  $(1 - AF)^2$ ,  $2 \times AF(1-AF)$ , and  $AF^2$ , respectively.
- The expected genotyping rate is independently estimated by the observed genotyping rate for each variant.
- The expected frequency of the combination of genotypes is computed under the independent assumption among alleles

335

### Combined effect size estimates of rare protein-altering variants

Using the inverse-variance method for fixed-effects model implemented in R meta package, we performed a meta-analysis of the estimated effect sizes (BETAs) of protein-altering variants on intraocular pressure and glaucoma. For glaucoma, the results are reported as odds ratio. 95% confidence intervals are reported for IOP and glaucoma.

340

### Association analysis with reproductive fitness

Using the number of live births (UK Biobank Field ID: 2734, Global Biobank Engine phenotype ID: INI2734) and the number of children fathered (UK Biobank Field ID: 2405, Global Biobank Engine phenotype ID: INI2405), we performed association analysis for the four identified protein-altering variants using R script with age, types of genotyping array, and the first 4 genotype principal components as covariates.


345

### Homozygous carrier analysis

For UK Biobank British individuals, we extracted the genotype calls with PLINK v2.00aLM (2 April 2019) and identified homozygous carrier of 175H allele<sup>11</sup>. We examined the year of birth (UK Biobank Field ID 34) and age at death (UK Biobank Field ID 40007)<sup>7</sup>.


350

## Supplementary Tables and Figures




**Supplementary Figure S1.** Phenotype distribution of Intraocular pressure stratified by glaucoma disease status displayed as a Tukey's box plot. The middle bold horizontal line represents the median, the lower and upper hinges show the first and third quartiles, the lower and upper whiskers represent  $1.5 \times$  interquartile range from the hinges. The data points beyond whiskers are plotted individually.

355




**Supplementary Figure S2.** The protein-altering variant GWAS QQ plot for intraocular pressure. The variants outside of MHC region with 0.01 % < MAF < 1% are included in the analysis.

**A**




**B**




360

**C**



**D**



**Supplementary Figure S3.** The intensity plots for *ANGPTL7* protein-altering variants with  $0.01\% < \text{MAF} < 1\%$ . **(A)** rs200058074 (p.Gln136Arg). **(B)** rs28991002 (p.Arg140His). **(C)** rs28991009 (p.Gln175His). **(D)** rs143435072 (p.Arg177Ter)




365 **Supplementary Table S1.** Consistency of the genotype call for four protein-altering variants in  
*ANGPTL7* between genotyping array and exome sequencing data. Variant including  
 chromosome, position, reference, and alternate allele (hg19), the rs identifier of the genetic  
 variant (rsID), consequence of the variation (Consequence), genotype call from array (Array)  
 and exome data (Exome), and the number of individuals (N). Inconsistent variant calls are  
 370 highlighted in bold font.

Variant	rsID	Consequence	Array	Exome	N
1:11252357:A:G	rs200058074	p.Gln136Arg	A/G	A/G	42
1:11252357:A:G	rs200058074	p.Gln136Arg	A/A	A/A	49837
1:11252357:A:G	rs200058074	p.Gln136Arg	NA	A/G	1
1:11252357:A:G	rs200058074	p.Gln136Arg	NA	A/A	29
1:11252369:G:A	rs28991002	p.Arg140His	A/A	A/A	1
1:11252369:G:A	rs28991002	p.Arg140His	G/A	G/A	218
1:11252369:G:A	rs28991002	p.Arg140His	<b>G/A</b>	<b>G/G</b>	<b>1</b>
1:11252369:G:A	rs28991002	p.Arg140His	G/G	G/G	49,663
1:11252369:G:A	rs28991002	p.Arg140His	NA	G/G	26
1:11253684:G:T	rs28991009	p.Gln175His	T/T	T/T	4
1:11253684:G:T	rs28991009	p.Gln175His	G/T	G/T	720
1:11253684:G:T	rs28991009	p.Gln175His	G/G	G/G	49,093
1:11253684:G:T	rs28991009	p.Gln175His	NA	G/T	1
1:11253684:G:T	rs28991009	p.Gln175His	NA	G/G	91
1:11253688:C:T	rs143435072	p.Arg177Ter	C/T	C/T	41
1:11253688:C:T	rs143435072	p.Arg177Ter	<b>C/T</b>	<b>C/C</b>	<b>1</b>
1:11253688:C:T	rs143435072	p.Arg177Ter	C/C	C/C	49,857
1:11253688:C:T	rs143435072	p.Arg177Ter	NA	C/C	10


375 **Supplementary Table S2.** Number of individuals stratified by genotype of rare (0.01% < MAF < 1%) protein-altering variants in *ANGPTL7*. The combination of genotypes is shown in the first four columns (rs200058074, rs28991002, rs28991009, and rs143435072) as well as the number of British individuals with the genotype combination in UK Biobank (n\_observed). The expected number of individuals are computed under Hardy-Weinberg equilibrium model and the independence assumption (n\_expected, Method).

rs200058074	rs28991002	rs28991009	rs143435072	n_observed	n_expected
A/A	G/G	G/G	C/C	328,492.00	329,401.21
A/A	G/G	G/T	C/C	5,395.00	5,390.72
A/A	G/A	G/G	C/C	1,693.00	1,672.69
A/A	G/G	NA	C/C	471.00	470.71
A/G	G/G	G/G	C/C	356.00	353.03
A/A	G/G	G/G	C/T	270.00	267.80
A/A	NA	G/G	C/C	194.00	192.47
NA	G/G	G/G	C/C	192.00	188.77
A/A	G/G	G/G	NA	37.00	38.13
A/A	G/G	T/T	C/C	28.00	22.06
A/A	G/A	G/T	C/C	8.00	27.37
A/G	G/G	G/T	C/C	4.00	5.78
A/A	NA	G/T	C/C	2.00	3.15
A/A	G/A	G/G	NA	2.00	0.19
A/A	G/G	NA	C/T	2.00	0.38
NA	G/A	G/G	C/C	1.00	0.96
A/A	A/A	G/G	C/C	1.00	2.12
A/A	G/A	G/G	C/T	1.00	1.36
A/A	G/G	G/T	C/T	1.00	4.38
A/G	G/G	NA	C/C	1.00	0.50

**A**



**B**



**Supplementary Figure S4:** The combined effect size estimates of rare protein-altering variants. The estimates of effect size on unstandardized intraocular pressure measurements (IOP [mmHg]) and normalized intraocular pressure measurements (SD[IOP]) are shown for the four protein-altering variants in *ANGPTL7*. The combined effect size estimate is summarized using inverse-variance method under the fixed effect model (Methods).

380

**Supplementary Table S3.** The combined effect size estimates of rare protein-altering variants.

385 The estimates of effect size on normalized intraocular pressure measurements (SD [IOP]) and their 95% confidence interval (95% CI) are shown for the four protein-altering variants in *ANGPTL7*. The combined effect size estimate is summarized using inverse-variance method under the fixed effect model (Methods). %W (fixed) column indicates the relative weights used in the fixed effect model. The bottom right cell indicates the P-value of combined effect size model.

Variant	SD (IOP)	95% CI	%W(fixed)
Gln175His	-0.2003	[-0.2534; -0.1471]	70.5
Arg177Ter	-0.2626	[-0.5107; -0.0144]	3.2
Arg140His	-0.1555	[-0.2502; -0.0607]	22.1
Gln136Arg	-0.0380	[-0.2564; 0.1804]	4.2
<b>Fixed effect model</b>	-0.1856	[-0.2302; -0.1410]	P-value: $3.4 \times 10^{-16}$

**Supplementary Table S4.** The combined effect size estimates of rare protein-altering variants.

395 The estimates of effect size on unstandardized intraocular pressure measurements (IOP [mmHg]) and their 95% confidence interval (95% CI) are shown for the four protein-altering variants in *ANGPTL7*. The combined effect size estimate is summarized using inverse-variance method under the fixed effect model (Methods). %W (fixed) column indicates the relative weights used in the fixed effect model. The bottom right cell indicates the P-value of combined effect size model.

Variant	IOP (mmHg)	95% CI	%W(fixed)
Gln175His	-0.7430	[-0.9547; -0.5314]	70.5
Arg177Ter	-0.9444	[-1.9334; 0.0445]	3.2
Arg140His	-0.4816	[-0.8593; -0.1039]	22.1
Gln136Arg	-0.2078	[-1.0785; 0.6630]	4.2
<b>Fixed effect model</b>	-0.6694	[-0.8471; -0.4917]	P-value: $1.6 \times 10^{-13}$

400 **Supplementary Table S5:** Glaucoma definitions in FinnGen. ICD-codes are used in the Finnish hospital discharge and cause-of-death registries. ATC-codes are used in the Social Insurance Institution prescription drug purchase registry. All endpoint definitions in the FinnGen phenome-wide association analysis are available online (<https://www.finnngen.fi/fi/node/68>).

Phenotype name	ICD-10	ICD-9	ICD-8	ICD-8 exclude	ATC (drug purchases)
Glaucoma (broad definition, combination of all the subphenotypes below)	Any of the below	Any of the below	Any of the below		S01EA
Glaucoma suspect	H40.0	365.0	375.97		
Primary open-angle glaucoma	H40.1	365.1	375.1[0-3]	375.11	
Primary angle-closure glaucoma	H40.2	365.2	375.00 375.11		
Glaucoma secondary to eye trauma	H40.3	365.6E			
Glaucoma secondary to eye inflammation	H40.4	365.6F 365.6B	364.01		
Glaucoma secondary to other eye disorders	H40.5	365.5 365.6A			
Glaucoma secondary to drugs	H40.6	365.3A			
Other and unspecified glaucoma	H40.[8-9]	365.6X 365.6C 365.[8-9]	375.2 375.9	375.97	
Glaucoma in diseases classified elsewhere	H42	365.4A			
Use of antiglaucoma preparations and miotics					S01EA



**Supplementary Figure S5:** Phenome-wide association analysis of p.R220C in FinnGen.  $-\log_{10}(P\text{-value})$  is displayed in y-axis. Disease endpoints grouped by disease categories are displayed on the x-axis. Highlighted associations with  $P < 0.001$  are shown.

405 **Supplementary Table S6.** FinnGen summary of *MYOC* p.Gln168Ter and *ANGPTL7* p.Arg220Cys carriers in primary open glaucoma cases.

Numbers of POA Glaucoma cases/controls by genotype:

ANGPTL7

wt . R220C+

410 MYOC- 2,499/119,192 = 2.0% . 146/10,570 = 1.3%

MYOC+ 49/630 = 7.7% . 1/69 = 1.4%


415 **Supplementary Table S7.** *ANGPTL7* protein-altering variant association with reproductive fitness, (A) the number of live births and (B) the number of children fathered. Variant includes chromosome, position, reference, and alternate allele (hg19). rsID - the rs identifier of the genetic variant. HGVS - the HGVS protein sequence name. MAF - the minor allele frequency in UK Biobank British population. Beta - estimated regression coefficient with 95% confidence intervals. P - p-value of association.

### A. Number of live births

Variant	rsID	HGVS	MAF	Dosage	Beta [95% CI]	P
1:11252357:A:G	rs200058074	p.Gln136Arg	.054 %	Gln/Arg	0.089 [-0.064, 0.24]	0.26
1:11252369:G:A	rs28991002	p.Arg140His	.25 %	Arg/His	0.29 [-1.9, 2.5]	0.80
1:11253684:G:T	rs28991009	p.Gln175His	.81 %	Gln/His	-0.0076 [-0.050, 0.035]	0.72
				His/His	0.23 [-0.48, 0.93]	0.52
1:11253688:C:T	rs143435072	p.Arg177Ter	.041 %	Arg/Ter	0.2 [0.021, 0.38]	0.029

### B. Number of children fathered

Variant	rsID	HGVS	MAF	Dosage	Beta [95% CI]	P
1:11252357:A:G	rs200058074	p.Gln136Arg	.054 %	Gln/Arg	0.0064 [-0.19, 0.21]	0.95
1:11252369:G:A	rs28991002	p.Arg140His	.25 %	Arg/His	0.0030 [-0.081, 0.087]	0.94
1:11253684:G:T	rs28991009	p.Gln175His	.81 %	Gln/His	-0.0052 [-0.054, 0.044]	0.83
				His/His	0.41 [-0.17, 0.98]	0.16
1:11253688:C:T	rs143435072	p.Arg177Ter	.041 %	Arg/Ter	-0.052 [-0.27, 0.17]	0.64



420 **Supplementary Figure S6:** The combined effect size estimates of rare protein-altering variants. The estimates of effect size (odds ratio, OR) on glaucoma is shown in log scale for the five protein-altering variants in *ANGPTL7*. The combined effect size estimate is summarized using inverse-variance method under the fixed effect model and 95% confidence intervals are shown (Methods).





425 **Supplementary Figure S7:** Protein expression in normal tissues and cell lines from ProteomicsDB and MOPED for ANGPTL7

## Supplementary Note

### FinnGen

FinnGen consists of the following people:

430

#### Steering Committee

Aarno Palotie	University of Helsinki / FIMM
Mark Daly	University of Helsinki / FIMM

#### Pharma

Howard Jacob	Abbvie
435 Athena Matakidou	Astra Zeneca
Heiko Runz	Biogen
Sally John	Biogen
Robert Plenge	Celgene
Julie Hunkapiller	Genentech
440 Meg Ehm	GSK
Dawn Waterworth	GSK
Caroline Fox	Merck
Anders Malarstig	Pfizer
Kathy Klinger	Sanofi
445 Kathy Call	Sanofi

#### UH & Biobanks

Tomi Mäkelä	University of Helsinki / FIMM
Jaakko Kaprio	University of Helsinki / FIMM
Petri Virolainen	Auria BB / Univ. of Turku /VSSHP
450 Kari Pulkki	Auria BB / Univ. of Turku /VSSHP
Terhi Kilpi	THL Biobank (BB) / THL
Markus Perola	THL Biobank (BB) / THL
Jukka Partanen	Finnish Red Cross Blood Service/FHRB
Anne Pitkäranta	HUS/Univ Hosp Districts
455 Riitta Kaarteenaho	Borealis BB/Univ. of Oulu/PPSHP
Seppo Vainio	Borealis BB/Univ. of Oulu/PPSHP
Kimmo Savinainen	Tampere BB/Univ Tampere/PSHP
Veli-Matti Kosma	Eastern Finland BB/UEF/PSSHP
Urho Kujala	Central Finland BB /UJy/KSSHP

460

## Scientific Committee

### Pharma

	Jeff Waring	Abbvie
	Bridget Riley-Gillis	AbbVie
	Athena Matakidou	Astra Zeneca
465	Heiko Runz	Biogen
	Jimmy Liu	Biogen
	Shameek Biswas	Celgene
	Julie Hunkapiller	Genentech
	Dawn Waterworth	GSK
470	Meg Ehm	GSK
	Josh Hoffman	GSK
	Dorothee Diogo	Merck
	Caroline Fox	Merck
	Anders Malarstig	Pfizer
475	Catherine Marshall	Pfizer
	Xinli Hu	Pfizer
	Kathy Call	Sanofi
	Kathy Klinger	Sanofi

### UH & Biobanks

480	Samuli Ripatti	University of Helsinki / FIMM
	Johanna Schleutker	Auria BB / Univ. of Turku /VSSHP
	Markus Perola	THL Biobank (BB) / THL
	Tiina Wahlfors	Finnish Red Cross Blood Service/FHRB
	Olli Carpen	HUS/Univ Hosp Districts
485	Johanna Myllyharju	Borealis BB/Univ. of Oulu/PPSHP
	Johannes Kettunen	Borealis BB/Univ. of Oulu/PPSHP
	Reijo Laaksonen	Tampere BB/Univ Tampere/PSHP
	Arto Mannermaa	Eastern Finland BB/UEF/PSSHP
	Juha Paloneva	Central Finland BB /UJy/KSSHP
490	Urho Kujala	Central Finland BB /UJy/KSSHP

## Clinical Groups

### Neurology Group

	Hilkka Soininen	LEAD: Kuopio
	Valtteri Julkunen	Kuopio

495	Anne Remes	Oulu
	Reetta Kälviäinen	Kuopio
	Mikko Hiltunen	Kuopio
	Jukka Peltola	Tampere
	Pentti Tienari	Helsinki
500	Juha Rinne	Turku
	Adam Ziemann	AbbVie
	Jeffrey Waring	AbbVie
	Sahar Esmaeeli	AbbVie
	Nizar Smaoui	AbbVie
505	Anne Lehtonen	AbbVie
	Susan Eaton	Biogen
	Heiko Runz	Biogen
	Sanni Lahdenperä	Biogen
	Janet van Adelsberg	Celgene
510	Shameek Biswas	Celgene
	John Michon	Genentech
	Geoff Kerchner	Genentech
	Julie Hunkapiller	Genentech
	Natalie Bowers	Genentech
515	Edmond Teng	Genentech
	John Eicher	Merck
	Vinay Mehta	Merck
	Padhraig Gormley	Merck
	Kari Linden	Pfizer
520	Christopher Whelan	Pfizer
	Fanli Xu	GSK
	David Pulford	GSK

#### Gastroenterology Group

	Martti Färkkilä LEAD:	Helsinki
525	Sampsa Pikkarainen	HUS
	Airi Jussila	Tampere
	Timo Blomster	Oulu
	Mikko Kiviniemi	Kuopio
	Markku Voutilainen	Turku
530	Bob Georgantas	AbbVie
	Graham Heap	AbbVie
	Jeffrey Waring	AbbVie
	Nizar Smaoui	AbbVie
	Fedik Rahimov	AbbVie
535	Anne Lehtonen	AbbVie

	Keith Usiskin	Celgene
	Tim Lu	Genentech
	Natalie Bowers	Genentech
	Danny Oh	Genentech
540	John Michon	Genentech
	Vinay Mehta	Merck
	Dermot Reilly	Merck
	Kirsi Kalpala	Pfizer
	Melissa Miller	Pfizer
545	Xinli Hu	Pfizer
	Linda McCarthy	GSK
Rheumatology Group		
	Kari Eklund	LEAD:Helsinki
	Antti Palomäki	Turku
550	Pia Isomaki	Tampere
	Laura Pirilä	Turku
	Oili Kaipainen-Seppänen	Kuopio
	Johanna Huhtakangas	Oulu
	Bob Georgantas	AbbVie
555	Jeffrey Waring	AbbVie
	Fedik Rahimov	AbbVie
	Apinya Lertratanakul	AbbVie
	Nizar Smaoui	AbbVie
	Anne Lehtonen	AbbVie
560	David Close	AstraZeneca
	Marla Hochfeld	Celgene
	Natalie Bowers	Genentech
	John Michon	Genentech
	Dorothee Diogo	Merck
565	Vinay Mehta	Merck
	Kirsi Kalpala	Pfizer
	Nan Bing	Pfizer
	Xinli Hu	Pfizer
	Jorge Esparza Gordillo	GSK
570	Nina Mars	University of Helsinki / FIMM

#### Pulmonology Group

	Tarja Laitinen	LEAD:Tampere
	Margit Pelkonen	Kuopio
	Paula Kauppi	Helsinki
575	Hannu Kankaanranta	Tampere

	Terttu Harju	Oulu
	Nizar Smaoui	AbbVie
	David Close	AstraZeneca
	Steven Greenberg	Celgene
580	Hubert Chen	Genentech
	Natalie Bowers	Genentech
	John Michon	Genentech
	Vinay Mehta	Merck
	Jo Betts	GSK
585	Soumitra Ghosh	GSK

#### Cardiometabolic Diseases Group

	Veikko Salomaa	Lead: THL
	Teemu Niiranen	THL
	Markus Juonala	Turku
590	Kaj Metsärinne	Turku
	Mika Kähönen	Tampere
	Juhani Juntila	Oulu
	Markku Laakso	Kuopio
	Jussi Pihlajamäki	Kuopio
595	Juha Sinisalo	Helsinki
	Marja-Riitta Taskinen	Helsinki
	Tiinamaija Tuomi	Helsinki
	Jari Laukkanen	Keski-Suomen Keskussairaala
	Ben Challis	AstraZeneca
600	Keith Usiskin	Celgene
	Andrew Peterson	Genentech
	Julie Hunkapiller	Genentech
	Natalie Bowers	Genentech
	John Michon	Genentech
605	Dorothee Diogo	Merck
	Dermot Reilly	Merck
	Audrey Chu	Merck
	Vinay Mehta	Merck
	Jaakko Parkkinen	Pfizer
610	Melissa Miller	Pfizer
	Anthony Muslin	Sanofi
	Dawn Waterworth	GSK

#### Oncology Group

	Heikki Joensuu	Lead: Helsinki
615	Tuomo Meretoja	Helsinki

	Olli Carpen	Helsinki
	Lauri Aaltonen	Helsinki
	Annika Auranen	Tampere
	Peeter Karihtala	Oulu
620	Saila Kauppila	Oulu
	Päivi Auvinen	Kuopio
	Klaus Elenius	Turku
	Relja Popovic	AbbVie
	Jeffrey Waring	AbbVie
625	Bridget Riley-Gillis	AbbVie
	Anne Lehtonen	AbbVie
	Athena Matakidou	AstraZeneca
	Jennifer Schutzman	Genentech
	Julie Hunkapiller	Genentech
630	Natalie Bowers	Genentech
	John Michon	Genentech
	Vinay Mehta	Merck
	Andrey Loboda	Merck
	Aparna Chhibber	Merck
635	Heli Lehtonen	Pfizer
	Stefan McDonough	Pfizer
	Marika Crohns	Sanofi
	Diptee Kulkarni	GSK
	Ophthalmology Group	
640	Kai Kaarniranta	Lead: Kuopio
	Joni Turunen	HUS/ Secretary
	Terhi Ollila	HUS
	Sanna Seitsonen	HUS
	Hannu Uusitalo	Tampere
645	Vesa Aaltonen	Turku
	Hannele Uusitalo-Järvinen	PSHP
	Marja Luodonpää	Oulu
	Nina Hautala	Oulu
	Heiko Runz	Biogen
650	Erich Strauss	Genentech
	Natalie Bowers	Genentech
	Hao Chen	Genentech
	John Michon	Genentech
	Anna Podgornaia	Merck
655	Vinay Mehta	Merck
	Dorothee Diogo	Merck

Joshua Hoffman GSK

### Dermatology Group

660 Kaisa Tasanen Oulu  
Laura Huilaja Oulu  
Katariina Hannula-Jouppi HUS  
Teea Salmi Tampere  
Sirku Peltonen Turku  
Leena Koulu Turku  
665 Ilkka Harvima Kuopio  
Kirsi Kalpala Pfizer  
Ying Wu Pfizer  
David Choy Genentech  
John Michon Genentech  
670 Nizar Smaoui AbbVie  
Fedik Rahimov AbbVie  
Anne Lehtonen AbbVie  
Dawn Waterworth GSK

### FinnGen Teams

675 Administration Team

Anu Jalanko University of Helsinki / FIMM  
Risto Kajanne University of Helsinki / FIMM  
Ulrike Lyhs University of Helsinki / FIMM

### Communication

680 Mari Kaunisto University of Helsinki / FIMM

### Analysis Team

Justin Wade Davis Abbvie  
Bridget Riley-Gillis Abbvie  
Danjuma Quarless Abbvie  
685 Slavé Petrovski Astra Zeneca  
Jimmy Liu Biogen  
Chia-Yen Chen Biogen  
Paola Bronson Biogen  
Robert Yang Celgene  
690 Joseph Maranville Celgene  
Shameek Biswas Celgene  
Diana Chang Genentech



	Julie Hunkapiller	Genentech
	Tushar Bhangale	Genentech
695	Natalie Bowers	Genentech
	Dorothee Diogo	Merck
	Emily Holzinger	Merck
	Padhraig Gormley	Merck
	Xulong Wang	Merck
700	Xing Chen	Pfizer
	Åsa Hedman	Pfizer
	Joshua Hoffman	GSK
	Clarence Wang	Sanofi
	Ethan Xu	Sanofi
705	Franck Auge	Sanofi
	Clement Chatelain	Sanofi
	Mitja Kurki	University of Helsinki / FIMM/ Broad Institute
	Samuli Ripatti	University of Helsinki / FIMM
	Mark Daly	University of Helsinki / FIMM
710	Juha Karjalainen	University of Helsinki / FIMM/ Broad Institute
	Aki Havulinna	University of Helsinki / FIMM
	Anu Jalanko	University of Helsinki / FIMM
	Kimmo Palin	University of Helsinki
	Priit Palta	University of Helsinki / FIMM
715	Pietro della Briotta Parolo	University of Helsinki / FIMM
	Wei Zhou	Broad Institute
	Susanna Lemmelä	University of Helsinki / FIMM
	Manuel Rivas	University of Stanford
	Jarmo Harju	University of Helsinki / FIMM
720	Aarno Palotie	University of Helsinki / FIMM
	Arto Lehisto	University of Helsinki / FIMM
	Andrea Ganna	University of Helsinki / FIMM
	Vincent Llorens	University of Helsinki / FIMM
	Antti Karlsson	Auria BB / Univ. of Turku /VSSHP
725	Kati Kristiansson	THL BB / THL
	Mikko Arvas	Finnish Red Cross Blood Service BB /FHRB
	Kati Hyvärinen	Finnish Red Cross Blood Service BB /FHRB
	Jarmo Ritari	Finnish Red Cross Blood Service BB /FHRB
	Tiina Wahlfors	Finnish Red Cross Blood Service BB /FHRB
730	Miika Koskinen	Helsinki BB/HUS/Univ Hosp Districts
	Olli Carpen	Helsinki BB/HUS/Univ Hosp Districts
	Johannes Kettunen	Borealis BB/Univ. of Oulu/PPSHP
	Katri Pylkäs	Borealis BB/Univ. of Oulu/PPSHP
	Marita Kalaoja	Borealis BB/Univ. of Oulu/PPSHP
735	Minna Karjalainen	Borealis BB/Univ. of Oulu/PPSHP

Tuomo Mantere Borealis BB/Univ. of Oulu/PPSHP  
Eeva Kangasniemi Tampere BB/Univ Tampere/PSHP  
Sami Heikkinen Eastern Finland BB/UEF/PSSHP  
Arto Mannermaa Eastern Finland BB/UEF/PSSHP  
740 Eija Laakkonen Central Finland BB /UJy/KSSHP  
Juha Kononen Central Finland BB /UJy/KSSHP

#### Sample Collection Coordination

Anu Loukola Helsinki BB/HUS/Univ Hosp Districts

#### Sample Logistics

745 Päivi Laiho THL BB / THL  
Tuuli Sistonen THL BB / THL  
Essi Kaiharju THL BB / THL  
Markku Laukkanen THL BB / THL  
Elina Järvensivu THL BB / THL  
750 Sini Lähteenmäki THL BB / THL  
Lotta Männikkö THL BB / THL  
Regis Wong THL BB / THL

#### Registry Data Operations

Kati Kristiansson THL BB / THL  
755 Hannele Mattsson THL BB / THL  
Susanna Lemmelä University of Helsinki / FIMM  
Tero Hiekkalinna THL BB / THL  
Manuel González Jiménez THL BB / THL

#### Genotyping

760 Kati Donner University of Helsinki / FIMM

#### Sequencing Informatics

Priit Palta University of Helsinki / FIMM  
Kalle Pärn University of Helsinki / FIMM  
Javier Nunez-Fontarnau University of Helsinki / FIMM  
765

#### Data Management and IT Infrastructure

Jarmo Harju University of Helsinki / FIMM  
Elina Kilpeläinen University of Helsinki / FIMM  
Timo P. Sipilä University of Helsinki / FIMM  
Georg Brein University of Helsinki / FIMM  
770 Alexander Dada University of Helsinki / FIMM

Ghazal Awaisa            University of Helsinki / FIMM  
Anastasia Shcherban   University of Helsinki / FIMM  
Tuomas Sipilä            University of Helsinki / FIMM

#### Clinical Endpoint Development

775    Hannele Laivuori        University of Helsinki / FIMM  
       Aki Havulinna         University of Helsinki / FIMM  
       Susanna Lemmelä      University of Helsinki / FIMM  
       Tuomo Kiiskinen      University of Helsinki / FIMM

#### Trajectory Team

780    Tarja Laitinen          Tampere University Hospital  
       Harri Siirtola          University of Tampere  
       Javier Gracia Tabuenca   University of Tampere

#### Biobank Directors

785    Lila Kallio              Auria Biobank  
       Sirpa Soini             THL Biobank  
       Jukka Partanen         Blood Service Biobank  
       Kimmo Pitkänen        Helsinki Biobank  
       Seppo Vainio            Northern Finland Biobank Borealis  
       Kimmo Savinainen     Tampere Biobank  
790    Veli-Matti Kosma        Biobank of Eastern Finland  
       Teijo Kuopio            Central Finland Biobank