The flagellar substrate specificity switch protein FlhB assembles onto the extra-membrane export gate to regulate type three secretion

Lucas Kuhlen^{1,2}, Steven Johnson¹, Andreas Zeitler³, Sandra Bäurle^{3,6}, Justin C. Deme^{1,4}, Rebecca Debo^{3,7}, Joseph Fisher¹, Samuel Wagner^{3,5} & Susan M. Lea^{1,4*}

¹Sir William Dunn School of Pathology, University of Oxford, Oxford OX13RE, UK. ²Department of Chemistry, University of Oxford, Oxford, UK. ³University of Tübingen, Interfaculty Institute of Microbiology and Infection Medicine (IMIT), Elfriede-Aulhorn-Str. 6, 72076 Tübingen, Germany. ⁴Central Oxford Structural Microscopy and Imaging Centre, University of Oxford, Oxford OX1 3RE, UK. ⁵German Center for Infection Research (DZIF), partner-site Tübingen, Elfriede-Aulhorn-Str. 6, 72076 Tübingen, Germany. ⁶Current address University Hospital Tübingen, Department of Thoracic & Cardiovascular Surgery, 72076 Tübingen, Germany ⁷Current address University of Tübingen Department of Geosciences, Centre for Applied Geosciences (ZAG), 72076 Tübingen, Germany

*Correspondence to: Susan M. Lea, susan.lea@path.ox.ac.uk.

Abstract

Export of proteins through type three secretion systems (T3SS) is critical for motility and virulence of many major bacterial pathogens. Proteins are transported through an export gate complex consisting of three proteins (FlpQR in flagellar systems, SctRST in virulence systems) that were initially annotated as membrane proteins, but which we have recently shown assemble into an extra-membranous helical assembly. A fourth putative membrane protein (FlhB/SctU) is essential to the export process, and also functions to “switch” secretion substrate specificity once the growing hook/needle structures reach their determined length. Here we present the structure of an export gate containing the switch protein from a *Vibrio* polar flagellar system at 3.2 Å resolution by cryo-electron microscopy. The structure reveals that the FlhB/SctU further extends the helical export gate assembly with its four putative transmembrane helices adopting an out-of-membrane location, wrapped around the other export gate components at the base of the structure. The unusual topology of the switch protein helices creates a loop that wraps around the bottom of the closed export gate complex. Structure-informed mutagenesis suggests that this loop is critical in gating secretion and we propose that a series of conformational changes in the type 3 secretion system trigger opening of the export gate through the interactions between FlhB/SctU and FlpQR/SctRST.

Main

Type three secretion is a mechanism of bacterial protein secretion across both inner and outer bacterial membranes. It is found in the virulence-associated injectisome (vT3SS), a molecular syringe, and the bacterial flagellum (fT3SS), a motility organelle (1). Both families contribute in significant ways to bacterial pathogenesis. vT3SS facilitate secretion not only across the bacterial envelope but also insert translocon proteins at the tip of the needle into the eukaryotic host plasma membrane, allowing direct injection of virulence factors in the host cytoplasm. The fT3SS is responsible for construction of the flagellar filament in both Gram-negative and Gram-positive bacteria, and hence imparts pathogenicity (2) for example via the ability to swim towards favourable environments or sense environmental conditions (3).

T3SS are multi-megadalton protein complexes that are capable of bridging from the bacterial cytoplasm to the extracellular space. At the core of the secretion system is the highly conserved export apparatus (EA) (4, 5), which is made up of five predicted transmembrane (TM) proteins (SctR, SctS, SctT, SctU and SctV in the vT3SS; FlhP, FlhQ, FlhR, FlhB and FlhA in the fT3SS). FlhA/SctV has been shown to form a nonameric ring (6), consisting of a large cytoplasmic domain situated below a hydrophobic domain predicted to contain 72 helices. This structure was proposed to surround an “export gate” through which substrates would enter the secretion pathway. This export gate is constructed from the other 4 EA proteins and was predicted to lie in the inner membrane. However, our recently determined structures of the *S. enterica* serovar Typhimurium FlhP₅Q₄R₁ and the *Shigella flexneri* SctR₅S₄T₁ complexes (7, 8) demonstrated that the export gate is actually embedded within the proteinaceous core of the T3SS basal body, placing it above the predicted location of the inner membrane. Furthermore, the helical structure of the export gate makes it likely that it is responsible for nucleating the helical filaments that assemble above it (9). Interestingly, the export gate complex has also recently been proposed to facilitate inward transport across the inner membrane associated with nanotubes (10, 11). The final component of the EA, FlhB/SctU, has long been known to be essential for all T3SS-mediated protein secretion. In addition, FlhB/SctU has a major role in switching the specificity of secretion substrates, mediating the transition from the early components necessary to build the flagellar hook in fT3SS and injectisome needle in vT3SS, to the later subunits required for flagellar filament or injectisome translocon assembly. The FlhB/SctU family of proteins all

contain an N-terminal hydrophobic sequence that is predicted to form 4 TM helices (FlhB_{TM}) and a smaller cytoplasmic C-terminal domain (FlhB_C). Crystal structures of the FlhB/SctU cytoplasmic domain from a range of species and systems (12-14) demonstrated a compact fold with an unusual autocatalytic cleavage site in a conserved NPTH sequence. Cleavage between the Asn and Pro residues, splitting FlhB_C into FlhB_{CN} and FlhB_{CC}, is required for the switching event to occur and a variety of mechanisms have been proposed to explain the need for this unusual mechanism (15). Little was known about the predicted TM portion of FlhB/SctU. Co-evolution analysis and molecular modelling led to suggestions that it forms a 4-helix bundle in the membrane (16), while crosslinks (17) and partial co-purification of FlhB with FliPQR were consistent with FlhB/SctU interacting with the export gate via a conserved site on the FliP₅Q₄R₁ complex (7). However, given the inaccuracy of the TM predictions for the other export gate components revealed by the PQR structure, we sought to determine the molecular basis of the interaction of FlhB with FliPQR. Here we present the structure of the TM region of FlhB bound to the FliPQR complex, in addition to two novel structures of the FliPQR homologues from *Vibrio mimicus* and *Pseudomonas savastanoi*. The structure reveals a unique topology that presents a loop that wraps around the base of the closed export gate. Mutagenesis studies confirm the crucial role played by the FlhB loop in the export process and suggest potential mechanisms of regulation of opening of the assembly.

Results

Conservation of the FliPQR structure

Our previously determined structures of *S. Typhimurium* FliPQR (7) and the vT3SS homologue SctRST from *S. flexneri* (8) demonstrated that the stoichiometry of the core structure (FliP₅Q₄R₁) is conserved between virulence and flagellar systems. However, classification of the SctRST data revealed variable occupancy of the SctS component (up to a maximum of 4 copies), consistent with our earlier native mass spectrometry measurements (7). Furthermore, our native mass spectrometry had also demonstrated that a small proportion of the *P. savastanoi* FliPQR complex contained 5 copies of FliQ, with the predicted 5th FliQ binding site beginning to encroach on the predicted FlhB interaction site. In order to further analyse the structural conservation and stoichiometry of the export apparatus core FliPQR we chose the homologous complexes from the fT3SS of two other bacterial species for structural studies: the *V. mimicus* polar flagellum FliPQR complex, which has a longer FliP sequence including an N-terminal domain conserved in the *Vibrionales* order (Supplementary Fig. 1), and the *P. savastanoi* FliPQR complex, that is a mixture of FliP₅Q₅R₁ and FliP₅Q₄R₁ complexes by native mass spectrometry (7). We determined the structures of both complexes using cryo-EM

and single particle analysis to 4.1 Å and 3.5 Å respectively (Fig. 1a, Table 1, Supplementary Fig. 2 and Supplementary Fig. 3). Both structures are highly similar to *S. Typhimurium* FliPQR (7) (RMSD=1.6 Å over all chains) and *S. flexneri* SctRST (*V. mimicus* FliPQR and SctRST RMSD=1.9 Å and *P. savastanoi* FliPQR and SctRST RMSD=2.3 Å) (7, 8).

Fig. 1 Conservation of the structure of the FliPQR export gate in the closed state

a, Cryo-EM volumes calculated in Relion using data from *S. Typhimurium* FliPQR (left, EMD-4173), *S. flexneri* SctRST (centre left, SctR₅S₄T₁ class (8) (EMD-4734)), *P. savastanoi* FliPQR (centre right) and *V. mimicus* FliPQR (right). FliQ₂ and FliQ₄ are coloured orange and FliQ₁, FliQ₃ and FliQ₅ are coloured red. **b**, Immunodetection of SctU^{FLAG} on Western blots of SDS PAGE-separated crude membrane samples of the indicated *S. Typhimurium* SctS pBpa mutants (denoted with X). Each sample is shown with and without UV-irradiation to induce photocrosslinking of pBpa to neighbouring interaction partners. **c**, As in (b) but testing interactions to SctU with pBpa in SctT and SctR. **d**, Mapping of the confirmed contact points, including those previously identified (17) on the structure of FliPQR (*S. Typhimurium*) and a model with a fifth FliQ subunit which is based on the structure of *P. savastanoi* FliPQR.

Consistent with our previous native mass spectrometry data, the structure of *P. savastanoi* revealed an additional FliQ subunit in the complex. In the *S. Typhimurium* and *V. mimicus* FliPQR structures there are four FliP-FliQ units, each the structural equivalent of a FliR subunit (7), but the fifth FliP is missing a FliQ. In the *P. savastanoi* structure, FliQ₅ binds the remaining FliP subunit in the same way as in the other FliP-FliQ units. This FliQ₅ subunit is located close to the site

on FliPQR we previously identified as important for interaction with FlhB/SctU (7, 17). Mapping of more a extensive *in vivo* photocrosslinking analysis based on covariance (16) between SctU and SctR, SctS, and SctT supports a binding site for SctU that involves large parts of helix 2 of SctS and helix 4 of SctT (Fig. 1b-c and Supplementary Fig. 4). Mapping of the residues on the structure of FliPQR and a model of FliPQR containing a fifth FliQ subunit reveal a large binding site in the complex containing four FliQ subunits and a more compact binding site when a fifth FliQ subunit is modelled (Fig. 1d). In this way FliQ₅/SctS₅ might be compatible with FlhB/SctU binding, depending on the unknown structure of the FlhB/SctU transmembrane domain (FlhB/SctU_{TM}), but addition of a sixth Fli/SctQ using the same helical parameters, would block this site.

Table 1 Cryo-EM statistics

	EMD-10095, PDB 6S3R	EMD-10096, PDB 6S3S	EMD-10093, PDB 6S3L
Data collection and processing			
Magnification	165,000	165,000	165,000
Voltage (kV)	300	300	300
Electron exposure (e ⁻ /Å ²)	48	48	48
Defocus range (μm)	0.5–4	0.5–4	0.5–4
Pixel size (Å)	0.822	0.822	0.822
Symmetry imposed	C1	C1	C1
Initial particle images (no.)	503,177	1,050,955	1,386,230
Final particle images (no.)	97,987	243,489	162,408
Map resolution (Å)	3.5	4.1	3.2
FSC threshold	0.143	0.143	0.143
Refinement			
Initial model used	EMDB-4173	EMDB-4173	EMDB-4173
Model resolution (Å)	3.5	4.1	3.2
FSC threshold	0.143	0.143	0.143
Map sharpening B factor (Å ²)	-101	-214	-97
Model composition			
Non-hydrogen atoms	12,855	12,321	13,849
Protein residues	1,669	1,569	1,768
Ligands	0	0	0
B factors (Å²)			
Protein	36.94	101.85	43.55
Ligand	N/A	N/A	N/A
r.m.s. deviations			
Bond lengths (Å)	0.0058	0.0066	0.01
Bond angles (°)	0.88	0.87	0.93
Validation			
MolProbity score	2.54	2.25	2.45
Clashscore	18	18.28	15.56
Poor rotamers (%)	2.22	0.07	2.72
Ramachandran plot			
Favored (%)	90.82	91.86	93.64
Allowed (%)	8.39	7.76	5.84
Disallowed (%)	0.79	0.39	0.52

Architecture of the FlpQR-FlhB export gate complex

We have observed four FliQ subunits in the *S. Typhimurium* and *V. mimicus* FlpQR and the *S. flexneri* SctRST structures but as we have previously observed FliQ to be sensitive to dissociation by detergent in the purification process (7, 8), it was possible that the 5th FliQ is a genuine component of the complex but was lost in the purification of less stable homologues. As the stoichiometry of FliQ has potentially large implications for the placement of FlhB in the system (Fig. 1d), we endeavoured to produce the more physiologically relevant FlpQR-FlhB complex.

After extensive screening of detergents, constructs with different placement of affinity tags and sequences from a variety of species for co-expression and co-purification of FlhB with FlpQR, we were able to prepare a monodisperse sample of the complex from *V. mimicus* (Fig. 2a). We analysed this sample by cryo-EM and determined the 3.2 Å structure of the complex (Fig. 2b, Table 1 and Supplementary Fig. 5), revealing a single copy of FlhB added to the previously observed Flp₅Q₄R₁ complex. The FlpQR subcomplex in this structure is very similar to the structure of the FlpQR complex in the absence of FlhB (RMSD=0.6 Å), while FlhB is observed to contain four long helices in the putative TM domain (FlhB_{TM}), forming two distinct hairpins that are wrapped around the outside of the FlpQR complex. This extensive interaction surface between FlhB and FlpQR reveals FlhB to be an integral part of the core of the export apparatus instead of an accessory factor. The opened out structure of the 4 predicted TM helices of the FlhB_{TM} domain once again highlights the potential dangers in predicting complex structures in the absence of some of the subunits. The soluble, globular, cytoplasmic domain (FlhB_C) is not visible, likely due to flexibility in the linker between the two domains. We tested whether this disorder of FlhB_C relative to FlpQR-FlhB_{TM} is due to the presence of the detergent micelle in our sample by imaging the sample in the amphipol A8-35, perhaps a better mimic of the proteinaceous environment relevant to the assembled T3SS (7), but we did not observe any additional density resulting from ordering of FlhB_C (data not shown).

Intriguingly, the two helical hairpins of FlhB are joined by a loop (FlhB_L) that literally loops around the (closed) entrance of the FlpQR gate (Fig. 2c). Consistent with our previous prediction (7) and crosslinking analysis (Fig. 1), FlhB contacts the site across Flp₅ and FliR, but it additionally contacts cross-linkable residues in the Flp₄ subunit (Fig. 2d). As previously predicted (7), hydrophobic cavities between Flp and FliQ, in addition to lateral cavities between the FlhB

hairpins and the FlhB/FliQ interface, are observed to contain densities consistent with lipid or detergent molecules although these could not be modelled unambiguously in the current volume (Supplementary Fig. 6).

Fig. 2 Architecture of the FlhPQR-FlhB export gate complex

a, SDS-PAGE gel of the *V. mimicus* FlhPQR-FlhB complex. **b**, Cryo-EM density map of the FlhPQR-FlhB complex with the density corresponding to FlhB coloured in green. **c**, Zoom of the FlhB loop at the bottom of the complex. **d**, Model of FlhPQR (surface) and FlhB (cartoon, green) with residues in FliP/SctR and FliR/SctT cross-linking to FlhB/SctU highlighted in blue as in Fig. 1d.

Structure of the hydrophobic domain of FlhB

The density corresponding to FlhB was of sufficient quality to build an atomic model of the structure using only sequence information (Fig. 3a and Supplementary Fig. 5). The topology of FlhB_{TM} is unusual; the helices 1 and 4 neighbour each other in the middle of the structure, while helix 2 and 3 flank the central pair on either side (Fig. 3b,c). In order to further validate the topology of FlhB we compared our model to contacts derived from evolutionary co-variation (Fig. 3d and Supplementary Fig. 7). This shows strong contacts between helices 1 and 2, 3 and 4 and 1 and 4 but an absence of contacts between helices 2 and 3, which is inconsistent with a helical bundle, but consistent with our more extended and topologically unusual structure.

Despite observing up to five FliQ subunits in FliPQR structures, there are only four FliQ molecules in this structure. In fact, the hairpin composed of FlhB helices 1 and 2 is bound to the site occupied by FliQ₅ in our *P. savastanoi* FliPQR structure, packing on FliP₅, whilst helices 3 and 4 pack on FliR. Thus the presence of FliQ₅ would block binding of FlhB (Fig. 3e), suggesting that the fifth FliQ binds to the complex in a non-native fashion due to the absence of FlhB in the overexpression system. This superposition of FlhB and FliQ₅ also reveals that the hairpin of helices 1 and 2 of FlhB adopts a very similar structure to FliQ despite the fact they are topologically distinct, with helix 2 of FlhB being structurally equivalent to helix 1 of FliQ and vice versa, i.e. the directionality of the hairpin is reversed along the long axis. Modelling FliQ₅ and FliQ₆ using the same helical parameters by which FliQ₁ to FliQ₄ are related reveals that FlhB continues the spiral of FliQ subunits and even helices 3 and 4 follow the same parameters (Fig. 3e) despite not interacting with a FliP subunit. Given the very different topologies of the two proteins, the level to which FlhB helices 1 and 2 and FliQ superpose is surprising. An evolutionary relationship between FlhB and FliQ is unlikely due to the topology differences, suggesting that the similarity of the structures is a result of convergent evolution and the need to form this helical assembly.

Fig. 3 Structural analysis of the FlhB hydrophobic domain

a, Quality of the cryo-EM volume corresponding to FlhB. Zoom box shows the fit to density of the model. **b,c**, Rainbow colouring of the FlhB model with numbers indicating the N and C-termini of the four helices (*V. mimicus* numbering). **d**, Evolutionary co-variation within FlhB calculated using RaptorX (18). Only contacts with a probability greater than 0.5 are plotted. Red boxes highlight the interaction between helix 1 and 2 (i), helix 1 and 4 (ii), helix 3 and 4 (iii), the N-terminus and FlhB_{CN} (iv), within the N-terminus (v) and between FlhB_L and the N-terminus (vi). **e**, Overlay of FlhB (green) and a modelled FliQ₅ and FliQ₆ following the same helical parameters as FliQ₁ to FliQ₄ in *V. mimicus*. **f**, Zoomed view of the interaction between the FlhB loop and FliQ, highlighting the intercalation of conserved hydrophobic residues in FlhB between the FliQ subunits.

An extended loop between helices is essential for secretion

The unusual topology of FlhB_{TM} means that a long loop, FlhB_L, between helix 2 and 3 (residues 110-139) connects the two hairpins of the structure. Most unexpectedly, this loop, which contains the most highly conserved residues within FlhB (Supplementary Fig. 8), is seen to wrap around the base of the PQR complex, contacting each of the FliQ subunits in turn and inserting conserved hydrophobic residues into the cavities between the FliQs (Fig. 3f). The loop structure also reveals how a single FliQ residue can co-evolve with multiple FlhB_L residues. Although FlhB_L in isolation doesn't

further constrict the base of the already closed PQR complex, the aperture does become significantly smaller when taking into account the poorly resolved termini of FlhB_{TM} (Supplementary Fig. 9). Therefore FlhB_L could contribute to export gate closure via trapping of the FlhB N-terminus and the linker connecting FlhB_{TM} to FlhB_C, in the direct line of the export pathway or by pinning the FliQ subunits closed. A mutation in the FlhB N-terminus had been reported (19) to act as a *ΔfliHI* bypass mutant (the ATPase and its regulator), presumed to be involved in controlling the opening of the export channel. In the FliPQR-FlhB complex the equivalent residue (FlhB_{P28} in *S. Typhimurium*, FlhB_{A28} in *V. mimicus*) locates very close to the pore entrance (Supplementary Fig. 9). In the *S. Typhimurium* SctRSTU complex the corresponding residue strongly photocrosslinks to SctS (Supplementary Fig. 4), supporting the notion that SctU mediates gating of the export apparatus core complex.

We decided to further probe the function of FlhB_L using mutagenesis in the motile *E. coli* strain W (Fig. 4a and Supplementary Fig. 10). Given that opening of the FliPQR-FlhB aperture would require a conformational change in FlhB_L, we hypothesised two mechanisms for FlhB_L such conformational changes. FlhB_L could either move away from the entrance to the gate through a hinging motion like a lid, or it could extend into a structure with less secondary structure in order to stay in contact with the binding sites on the opening FliQ subunits, reminiscent of a sphincter.

Mutations in either the conserved hydrophobic residues of FlhB_L that insert between the FliQ subunits (Fig. 3f) or the highly conserved loop of FliQ severely reduced motility (Fig. 4a) without affecting binding of FlhB to FliPQR (Supplementary Fig. 11). Although substitution with the bulky, hydrophobic amino acid tryptophan and removal of bulky sidechains only reduced motility, introduction of charged residues completely abolished motility, suggesting that secretion can proceed at lower efficiency when the FliQ-FlhB_L interaction is only reduced rather than completely disrupted as in the aspartate mutations.

We performed an extensive *in vivo* photocrosslinking analysis to validate the interactions and functional relevance of the corresponding SctU_L in the vT3SS-1 of *S. Typhimurium*. While no crosslinks to SctS could be identified with the artificial crosslinking amino acid *p*-benzoyl-phenylalanine (*p*Bpa) introduced into SctU_L itself (Supplementary Fig. 12), numerous crosslinks were identified with *p*Bpa in the lower part of SctS that faces SctU_L (Fig. 4b). Using 2-dimensional blue native/SDS PAGE, we could show that the crosslink observed with SctS_{Q41X} occurred not only in the SctRSTU assembly intermediate but also in the assembled needle complex (Fig. 4c), adding further support to the idea that the

structure of the isolated complex represents the structure of the complex in the full assembly. The observed crosslinks were independent of functional secretion of the vT3SS, indicating that assembly of needle adapter, the inner rod, and needle filament does not lead to a conformational change of this part of the SctRSTU complex (Fig. 4b). Strikingly, introduction of *pBpa* at several positions of SctS led to a strong defect in secretion but not SctS-SctU_L interaction, highlighting the relevance of this site for secretion function of T3SS (Fig. 4d), while *pBpa* substitutions within SctU_L were more functionally neutral. In total, we found a large number of residues at the FlhQ/SctS-FlhB_L/SctU_L interface that are required for type three secretion (Fig. 4e).

Strong cross-linking between SctS and SctU even in the assembled needle complex and loss of function in more disruptive mutations in which the interaction between FlhQ and FlhB_L is altered through the introduction of charged residues suggest that this interaction is important for activity and FlhB_L is not simply one of the closure points of the complex in assembly intermediates. If this interaction is maintained in the open state of the export gate, a more extended conformation of FlhB_L would be required. Consistent with this idea, deletions of six or more residues in FlhB_L led to loss of motility (Fig. 4a).

We further investigated the function of FlhB_L through more targeted mutations. FlhB_L is a largely extended polypeptide with little secondary structure, but a short stretch at its C-terminus is helical. Interestingly, a mutation of a glycine in this α -helix in the FlhB homologue YscU disrupted secretion in the *Yersinia* vT3SS (20). The equivalent mutation in *E. coli* FlhB and mutation of the conserved positively charged residue R135 (*E. coli* W numbering) had little effect (Fig. 4a) assessed at the level of motility although we cannot rule out more subtle effects on the efficiency with which secretion occurs.

Fig. 4 Functional analysis of FlhB_L

a, Motility in soft agar of *E. coli* Δ FliOPQR complemented with plasmids expressing FliOPQR with the indicated mutations in FliQ (left) and *E. coli* Δ FlhB complemented with plasmids expressing FlhB with the indicated mutations (*E. coli* numbering) (right). **b**, Immunodetection of SctU^{FLAG} on Western blots of SDS-Page separated crude membrane samples of the indicated *S. Typhimurium* SctS *pBpa* mutants (denoted with X). Each sample is shown with and without UV-irradiation to induce photocrosslinking of *pBpa* to neighbouring interaction partners. Crosslinks between SctS_{*pBpa*} and SctU^{FLAG} are indicated. Crosslinking analysis was performed in the wild type and in an ATP hydrolysis-deficient SctN_{K165E} mutant that is incapable of type III secretion. **c**, Immunodetection of SctU^{FLAG} on Western blots of 2D blue native/SDS-PAGE separated crude membrane samples of the indicated *S. Typhimurium* SctS *pBpa* mutant. The sample is shown with and without UV-irradiation to induce photocrosslinking of *pBpa* to neighbouring interaction partners. Crosslinks between SctS_{*pBpa*} and SctU^{FLAG} in the SctRSTU assembly intermediate and in the assembled needle complex (NC) are indicated. **d**, Immunodetection of the early T3SS substrate SctP on Western blots of SDS-PAGE separated culture supernatants of the indicated *S. Typhimurium* SctS *pBpa* mutants. **e**, Structure of FliPQR-FlhB highlighting mutation sites that impaired motility or secretion in red and mutation sites that had no or only a small effect in yellow.

Discussion

In this study we show that FlhB_{TM} is part of the export gate complex together with FliPQR. Two pairs of helices of FlhB bind to FliPQR through a structure mimicking the shape of FliQ, despite topological reversal, an example of molecular convergent evolution. The unusual topology of FlhB places helices 2 and 3 apart from each other allowing them to mount a loop, FlhB_L, onto the cytoplasm-facing surface of the export gate. Although the way in which FlhB_L wraps around the closed pore suggests a role in maintaining the closed state, our structures of FliPQR/SctRST in the absence of

FlhB/SctU are also closed (7, 8), as is the complex in the context of the assembled T3SS (21), suggesting instead that FlhB may be involved in opening of the gate rather than locking it closed. We further propose that in the process of opening FlhB_L forms a more extended structure, implying that it undergoes cycles of extension and contraction as the export gate opens and closes.

The location and the topology of FlhB_{TM} place the N-terminus, FlhB_{CN} and FlhB_L in close proximity just underneath the aperture of the gate. Although the resolution of the map is poor in the region of the cytoplasmic face of the complex, it is possible to trace the approximate position of the FlhB N-terminus and FlhB_{CN} (Supplementary Fig. 9). The close association of the N-terminus and FlhB_{CN} is consistent with the strong contacts derived from evolutionary co-variation between the N-terminus and the N-terminal part of the cytoplasmic domain (FlhB_{CN}) (Fig. 3c) and a genetic interaction in *S. Typhimurium* between E11 and E230, in the N-terminus of FlhB and FlhB_{CN} respectively (22). Furthermore, the direction in which FlhB_{CN} leaves the export gate implies that, in the context of the assembled T3SS nanomachine, FlhB_C could be located anywhere between the FlpQR-FlhB gate and the nonameric ring of the FlhA cytoplasmic domain below (Fig. 5a). It is conceivable that the conformational changes required for opening of the export gate are propagated via pulling forces imparted on helix 4 of FlhB_{TM}, which is linked to the other helices of FlhB_{TM} via a conserved network of buried charged residues (Fig. 5b) including D208 that has been demonstrated to play a role in motility (23). Such forces could initiate in the cytoplasmic domain of FlhA, which binds to substrate-chaperone complexes and has been demonstrated to exist in multiple conformations (24, 25) marking it as the only component of the export apparatus observed in multiple conformational states to date. As FlhB_C is thought to interact with substrates just before they pass through the export gate (26), it is possible that transition of FlhA_C between the different conformations pulls on FlhB_C, either directly or via substrate, thereby pulling on the FlhB_{TM} network. Alternatively, changes in the FlhA_{TM} domain that are triggered via FlhA_C or by the proton-motive force (27) could directly influence the conformation of the FlhB_{TM} domain.

The mechanism of suppressor mutations in the N-terminal residues of FlhB, such as the P28T mutation that rescues motility of a strain deleted for the ATPase complex ($\Delta fliHI$) (19, 28), has long been mysterious. Our structure, demonstrating the clustering of the N-terminus, FlhB_{CN} and FlhB_L at the cytoplasmic entrance of the gate, suggests that they may rescue function by altering the dynamics of the closure point in order to facilitate opening of the export gate. This notion is further supported by the interaction we observe of SctU_{F24pBpa} (equivalent to P28T) with SctS. This direct

1 functional link between the ATPase and the gating mechanism, in conjunction with a host of other mutational data in
2 FlhA and FlhB, suggests that cycles of ATP hydrolysis may induce conformational changes in the export gate, presumably
3 via the FliJ stalk of the ATPase complex interacting with the FlhA ring that is positioned between the ATPase and the
4 export gate.

5 Finally, FlhB/SctU is known to play a key role in substrate switching, an event which requires autocatalytic cleavage of
6 the NPTH sequence in FlhB_C/SctU_C (14, 15, 26, 29). Although we do not observe the residues involved in switching in our
a

export gate structure, the fact that we are able to cross-link SctU in
fully assembled basal bodies, using the same residues as in the
purified complex, suggests that the gating mechanism discussed here
is likely applicable regardless of substrate. Clearly future studies will
need to focus on observing gating and switching events.

In summary, our structure of FlhB as part of the export gate complex
extends our understanding of the regulation of the T3SS export
apparatus and suggests possible mechanisms of export gate opening.

Fig. 5 Position of FlhB_C in the complete T3SS

a, Placement of FliPQR-FlhB in a tomographic reconstruction of the *Salmonella* SPI-1 vT3SS (EMD-8544) (30). FliPQR is shown in grey, FlhB_{TM} in green, two possible positions of FlhB_C (PDB: 3b0z) (13) in yellow and light blue and the density corresponding to the FlhA homologue is highlighted in blue. b, Network of salt bridges formed by conserved charged residues in FlhB (*V. mimicus* numbering).

Methods

Materials

Chemicals were from Sigma-Aldrich unless otherwise specified. The detergents n-dodecylmaltoside (DDM) and lauryl maltose neopentyl glycol (LMNG) and the amphipol A8-35 were from Anatrace. *p*-benzoyl-phenylalanine was from Bachem. Primers are listed in Supplementary Table 2 and were synthesized by Invitrogen or Eurofins.

Bacterial strains and plasmids

Bacterial strains and plasmids used in this study are listed in Supplementary Table 3. Plasmids were generated by Gibson assembly of PCR fragments using the NEBuilder HiFi Master Mix (NEB) or in vivo assembly (31). Fragments were created by PCR with the relevant primers using Q5 polymerase (NEB) and genomic DNA templates obtained from DSMZ (*Vibrio mimicus* strain DSM 19130 and *Pseudomonas savastanoi*, pv. phaseolicola 1448A strain DSM 21482). Gibson assembly and PCR were carried out following the manufacturer's recommendations. *Escherichia coli* W for motility assays was obtained from DSMZ (DSM 1116). Bacterial cultures were supplemented as required with ampicillin (100 µg/mL) or kanamycin (30 µg/mL or 60 µg/mL for large scale expression in TB medium). *S. Typhimurium* strains were cultured with low aeration at 37°C in Luria Bertani (LB) broth supplemented with 0.3 M NaCl to induce expression of genes of SPI-1. As required, bacterial cultures were supplemented with tetracycline (12.5 µg/ml), streptomycin (50 µg/ml), chloramphenicol (10 µg/ml), ampicillin (100 µg/ml) or kanamycin (25 µg/ml). Low-copy plasmid-based expression of SctRSTU^{FLAG} was induced by the addition of 500 µM rhamnose to the culture medium.

Generation of chromosomal deletion mutants

Electrocompetent *E. coli* W expressing λ Red recombinase from plasmid pKD46 were transformed with DNA fragments containing a chloramphenicol resistance cassette surrounded by sequences homologous to the gene of interest as described in Supplementary Table 3. Colonies were selected on LB agar containing chloramphenicol (20 µg/mL) and

transformed again with pCP20 and grown on LB agar containing ampicillin (100 µg/mL) at 30 °C. Finally, clones were grown in LB media at 37 °C. Deletion mutations were confirmed by PCR. All *Salmonella* strains were derived from *Salmonella enterica* serovar Typhimurium strain SL1344 (Hoiseth and Stocker, 1981) and created by allelic exchange as previously described (Kaniga *et al.*, 1994).

Purification of export gate complexes

FliOPQR or FliOPQR-FliH were expressed in *Escherichia coli* BL21 (DE3) as a single operon from a pT12 vector (Supplementary Table 3) as described previously (7). Briefly, cells were grown at 37 °C in TB media containing rhamnose monohydrate (0.1%), harvested by spinning at 4,000 g, resuspended in TBS (100 mM Tris, 150 mM NaCl, 1 mM EDTA, pH 8) and lysed in an EmulsiFlex C5 homogenizer (Avestin). Membranes were prepared from the cleared lysate by ultracentrifugation at 40,000 rpm in a 45 Ti rotor (Beckmann) for 3 hours. Membranes were solubilized in 1% (w/v) LMNG and proteins were purified using a StrepTrap column (GE Healthcare). The resin was washed in TBS containing 0.01% (w/v) LMNG and proteins were eluted in the wash buffer supplemented with 10 mM desthiobiotin. Intact complexes were separated from aggregate by size-exclusion chromatography in TBS containing 0.01% (w/v) LMNG (S200 10/300 increase or Superose 6 increase, GE Healthcare).

For preparation of FliPQR-FliH solubilised by the amphipol A8-35, the protein was purified in DDM using 1% (w/v) for extraction from the membrane and 0.02% (w/v) subsequently. Eluate from the StrepTrap column was mixed with amphipol at a ratio of amphipol to protein of 10:1. After incubating for one hour, the sample was dialysed into TBS using a 10,000 MWCO Slide-A-Lyzer device (ThermoFisher Scientific) overnight followed by size-exclusion chromatography on a Superose 6 increase column using TBS as the running buffer.

Sample preparation for cryo-EM

3 µl of purified complex at 1 to 3.6 mg/ml were applied to glow-discharged holey carbon-coated grids (Quantifoil 300 mesh, Au R1.2/1.3). Grids were blotted for 3 s at 100% humidity at 22 °C and frozen in liquid ethane using a Vitrobot Mark IV (FEI). For samples solubilised in detergent, blotting was preceded by a wait time of 5 to 10 seconds. *V. mimicus* FliPQR was supplemented with 0 mM, 0.05 mM, 0.5 mM or 3 mM fluorinated Fos-Choline prior to grid preparation.

EM data acquisition and model building

All data contributing to the final models were collected on a Titan Krios (FEI) operating at 300 kV. All movies were recorded on a K2 Summit detector (Gatan) in counting mode at a sampling of 0.822 Å/pixel, 2.4 e⁻ Å⁻²/frame, 8 s exposure, total dose 48 e⁻ Å⁻², 20 fractions written. Motion correction and dose weighting were performed using MotionCor implemented in Relion 3.0 (32) (*V. mimicus* FlIPQR-FlhB and *P. savastanoi* FlIPQR) or using Simple-unblur (33) (*V. mimicus* FlIPQR). CTFs were calculated using CTFFIND4 (34). Particles were picked in Simple and processed in Relion 2.0 (35) and 3.0 (32) as described in Supplementary Fig. 2, Supplementary Fig. 3 and Supplementary Fig. 5. Atomic models of FlIPQR and FlhB were built using Coot (36) and refined in Phenix (37).

Motility assays

E. coli W strains WL1 or WL2 (Supplementary Table 3) were transformed with plasmids encoding FlIPQR or FlhB containing the mutations to be tested. 3 µl of a saturated overnight culture were injected into soft agar plates (0.28% agar, 2YT media, containing ampicillin (100 µg/mL) or kanamycin (30 µg/mL) and 0.1% arabinose or 0.5% rhamnose monohydrate as appropriate) and incubated at room temperature.

S. Typhimurium vT3SS secretion assay

Proteins secreted into the culture medium via the vT3SS-1 were analyzed as described previously (Monjarás Feria *et al.*, 2015). *S. Typhimurium* strains were cultured with low aeration in LB broth supplemented with 0.3 M NaCl at 37°C for 5 h. Bacterial suspensions were centrifuged at 10,000 x *g* for 2 min and 4°C to separate whole cells and supernatants. Whole cells were resuspended in SDS PAGE loading buffer. Supernatants were passed through 0.2 µm pore size filters and supplemented with 0.1 % Na-desoxycholic acid. Proteins in the supernatant were precipitated with 10 % trichloroacetic acid (TCA) for 30 min at 4°C and pelleted via centrifugation at 20,000 x *g* for 20 min and 4°C. Pellets containing precipitated proteins were washed with acetone and resuspended in SDS PAGE loading buffer. Whole cell samples and secreted proteins were analyzed by SDS PAGE, Western blotting, and immunodetection.

In vivo photocrosslinking

In vivo photocrosslinking was carried out as described previously (Farrell *et al.*, 2005; Dietsche *et al.*, 2016) with minor modifications. In order to enhance expression of vT3SS-1, *S. Typhimurium* strains expressed Hila, the master transcriptional regulator of SPI-1 T3SS, from a high copy plasmid under the control of an arabinose-inducible P_{araBAD} promoter (Guzman *et al.*, 1995). Bacterial cultures were grown in LB broth supplemented with, 0.3 M NaCl, 1 mM pBpa and 0.05 % arabinose at 37°C for 5 h. 5 ODU of bacterial cells were harvested and washed once with 5 ml chilled PBS to remove residual media. Bacteria were pelleted by centrifugation at 4,000 x *g* for 3 min and 4°C and afterwards resuspended in 1 ml of chilled PBS. Bacterial suspensions were transferred into 6-well cell culture dishes and irradiated for 30 min with UV light ($\lambda=365$ nm) on a UV transilluminator table (UVP). Subsequently, bacterial cells were pelleted by centrifugation at 10,000 x *g* for 2 min and 4°C. Samples were stored at -20°C until use.

Crude membrane preparation

Crude membranes were purified following the published protocol (Dietsche *et al.*, 2016). 5 ODU of bacterial cells were resuspended in 750 μ l lysis buffer (50 mM triethanolamine pH 7.5, 250 mM sucrose, 1 mM EDTA, 1 mM MgCl₂, 10 μ g/ml DNase, 2 μ g/ml lysozyme, 1:100 protease inhibitor cocktail), and incubated on ice for 30 minutes. Cell slurries were lysed via continuous bead milling. Intact cells, beads and debris were removed by centrifugation for 10 min at 10,000 x *g* and 4°C. Supernatants were centrifuged for 50 min at 52,000 rpm and 4°C in a Beckman TLA 55 rotor to pellet bacterial membranes. Pellets containing crude membranes were store at -20°C until use. Samples were analyzed by SDS PAGE, Western blotting, and immunodetection.

Western blotting and immunodetection

Samples were loaded onto SERVAGel™ TG PRiME 8-16% precast gels and transferred on PVDF membranes (Bio-RAD). Proteins were detected with primary antibodies anti-St₁SctP (InvJ) (Monjarás Feria *et al.*, 2015) (1:2000) or M2 anti-FLAG (1:10,000) (Sigma-Aldrich). Secondary antibodies (Thermo-Fisher) were goat anti-mouse IgG Dylight 800 conjugate (1:5000). Scanning of the PVDF membranes and image analysis was performed with a Li-Cor Odyssey system and Image Studio 3.1 (Li-Cor).

Acknowledgements

We thank E. Johnson and A. Costin of the Central Oxford Structural Microscopy and Imaging Centre for assistance with data collection. H. Elmlund (Monash) is thanked for assistance with access to SIMPLE code ahead of release. The Central Oxford Structural Microscopy and Imaging Centre is supported by the Wellcome Trust (201536), the EPA Cephalosporin Trust, the Wolfson Foundation, and a Royal Society/Wolfson Foundation Laboratory Refurbishment Grant (WL160052). Work performed in the lab of S. M. Lea was supported by a Wellcome Trust Investigator Award (100298) and an MRC Program Grant (M011984). L.K. is a Wellcome Trust PhD student (109136).

Author Contributions

L.K. performed experiments, did strain and plasmid construction, complex purification, native mass spectrometry, cryo-EM grid optimization, cryo-EM data analysis, and model building and analysis. J.D. performed experiments, cryo-EM grid optimization, and data collection. J.F. performed experiments, complex purification. Work in the laboratory of SW related to this project was funded by the German Center for Infection Research (DZIF), project TTU06.801/808. A.Z., S.B., and R.D. generated pBpa mutants, performed crosslinking experiments and secretion assays, and analyzed data. S.W. designed injectisome functional experiments, analyzed data S.J. and S.M.L. supervised experimental work and wrote the first draft of the paper with L.K. All authors contributed to and commented on the final manuscript.

References

1. M. Erhardt, K. Namba, K. T. Hughes, Bacterial nanomachines: the flagellum and type III injectisome. *Cold Spring Harbor perspectives in biology* **2**, a000299 (Nov, 2010).
2. Q. Duan, M. Zhou, L. Zhu, G. Zhu, Flagella and bacterial pathogenicity. *Journal of basic microbiology* **53**, 1 (Jan, 2013).
3. Z. Liu *et al.*, Mucosal penetration primes *Vibrio cholerae* for host colonization by repressing quorum sensing. *Proceedings of the National Academy of Sciences of the United States of America* **105**, 9769 (Jul 15, 2008).
4. S. Wagner *et al.*, Bacterial type III secretion systems: a complex device for the delivery of bacterial effector proteins into eukaryotic host cells. *Fems Microbiol Lett* **365**, (Oct, 2018).
5. S. Wagner *et al.*, Organization and coordinated assembly of the type III secretion export apparatus. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 17745 (Oct 12, 2010).
6. P. Abrusci *et al.*, Architecture of the major component of the type III secretion system export apparatus. *Nature structural & molecular biology* **20**, 99 (Jan, 2013).
7. L. Kuhlen *et al.*, Structure of the core of the type III secretion system export apparatus. *Nature structural & molecular biology* **25**, 583 (Jul, 2018).
8. S. Johnson, L. Kuhlen, J. C. Deme, P. Abrusci, S. M. Lea, The structure of an injectisome export gate demonstrates conservation of architecture in the core export gate between flagellar and virulence type three secretion systems. *bioRxiv*, (2019).

9. C. E. Torres-Vargas *et al.*, The inner rod of virulence-associated type III secretion systems constitutes a needle adapter of one helical turn that is deeply integrated into the system's export apparatus. *Molecular microbiology*, (Jun 10, 2019).
10. R. R. Pal *et al.*, Pathogenic E. coli Extracts Nutrients from Infected Host Cells Utilizing Injectisome Components. *Cell* **177**, 683 (Apr 18, 2019).
11. S. Bhattacharya *et al.*, A Ubiquitous Platform for Bacterial Nanotube Biogenesis. *Cell reports* **27**, 334 (Apr 9, 2019).
12. J. E. Deane *et al.*, Crystal structure of Spa40, the specificity switch for the Shigella flexneri type III secretion system. *Molecular microbiology* **69**, 267 (Jul, 2008).
13. V. A. Meshcheryakov, A. Kitao, H. Matsunami, F. A. Samatey, Inhibition of a type III secretion system by the deletion of a short loop in one of its membrane proteins. *Acta crystallographica. Section D, Biological crystallography* **69**, 812 (May, 2013).
14. R. Zarivach *et al.*, Structural analysis of the essential self-cleaving type III secretion proteins EscU and SpaS. *Nature* **453**, 124 (May 1, 2008).
15. J. V. Monjaras Feria, M. D. Lefebvre, Y. D. Stierhof, J. E. Galan, S. Wagner, Role of autocleavage in the function of a type III secretion specificity switch protein in Salmonella enterica serovar Typhimurium. *mBio* **6**, e01459 (Oct 13, 2015).
16. W. R. Taylor, T. R. Matthews-Palmer, M. Beeby, Molecular Models for the Core Components of the Flagellar Type-III Secretion Complex. *PloS one* **11**, e0164047 (2016).
17. T. Dietsche *et al.*, Structural and Functional Characterization of the Bacterial Type III Secretion Export Apparatus. *PLoS pathogens* **12**, e1006071 (Dec, 2016).
18. S. Wang, S. Sun, Z. Li, R. Zhang, J. Xu, Accurate De Novo Prediction of Protein Contact Map by Ultra-Deep Learning Model. *PLoS computational biology* **13**, e1005324 (Jan, 2017).
19. T. Minamino, K. Namba, Distinct roles of the Flil ATPase and proton motive force in bacterial flagellar protein export. *Nature* **451**, 485 (Jan 24, 2008).
20. K. E. Riordan, O. Schneewind, YscU cleavage and the assembly of Yersinia type III secretion machine complexes. *Molecular microbiology* **68**, 1485 (Jun, 2008).
21. J. Hu *et al.*, Cryo-EM analysis of the T3S injectisome reveals the structure of the needle and open secretin. *Nat Commun* **9**, 3840 (Sep 21, 2018).
22. Y. Inoue, M. Kinoshita, K. Namba, T. Minamino, Mutational analysis of the C-terminal cytoplasmic domain of FlhB, a transmembrane component of the flagellar type III protein export apparatus in Salmonella. *Genes to cells : devoted to molecular & cellular mechanisms*, (Apr 8, 2019).
23. M. Erhardt *et al.*, Mechanism of type-III protein secretion: Regulation of FlhA conformation by a functionally critical charged-residue cluster. *Molecular microbiology* **104**, 234 (Apr, 2017).
24. Q. Xing *et al.*, Structures of chaperone-substrate complexes docked onto the export gate in a type III secretion system. *Nat Commun* **9**, 1773 (May 2, 2018).
25. Y. Inoue *et al.*, Structural Insights into the Substrate Specificity Switch Mechanism of the Type III Protein Export Apparatus. *Structure* **27**, 965 (Jun 4, 2019).
26. L. D. Evans, S. Poulter, E. M. Terentjev, C. Hughes, G. M. Fraser, A chain mechanism for flagellum growth. *Nature* **504**, 287 (Dec 12, 2013).
27. T. Minamino, Y. V. Morimoto, N. Hara, P. D. Aldridge, K. Namba, The Bacterial Flagellar Type III Export Gate Complex Is a Dual Fuel Engine That Can Use Both H⁺ and Na⁺ for Flagellar Protein Export. *PLoS pathogens* **12**, e1005495 (Mar, 2016).
28. T. Minamino, B. Gonzalez-Pedrajo, M. Kihara, K. Namba, R. M. Macnab, The ATPase Flil can interact with the type III flagellar protein export apparatus in the absence of its regulator, FliH. *Journal of bacteriology* **185**, 3983 (Jul, 2003).
29. G. M. Fraser *et al.*, Substrate specificity of type III flagellar protein export in Salmonella is controlled by subdomain interactions in FlhB. *Molecular microbiology* **48**, 1043 (May, 2003).
30. B. Hu, M. Lara-Tejero, Q. Kong, J. E. Galan, J. Liu, In Situ Molecular Architecture of the Salmonella Type III Secretion Machine. *Cell* **168**, 1065 (Mar 9, 2017).
31. J. Garcia-Nafria, J. F. Watson, I. H. Greger, IVA cloning: A single-tube universal cloning system exploiting bacterial In Vivo Assembly. *Scientific reports* **6**, 27459 (Jun 6, 2016).
32. J. Zivanov, T. Nakane, S. H. W. Scheres, A Bayesian approach to beam-induced motion correction in cryo-EM single-particle analysis. *IUCr* **6**, 5 (Jan 1, 2019).

- 1 33. C. F. Reboul, M. Eager, D. Elmlund, H. Elmlund, Single-particle cryo-EM-Improved ab initio 3D reconstruction
2 with SIMPLE/PRIME. *Protein science : a publication of the Protein Society* **27**, 51 (Jan, 2018).
- 3 34. A. Rohou, N. Grigorieff, CTFFIND4: Fast and accurate defocus estimation from electron micrographs. *Journal of*
4 *structural biology* **192**, 216 (Nov, 2015).
- 5 35. D. Kimanius, B. O. Forsberg, S. H. Scheres, E. Lindahl, Accelerated cryo-EM structure determination with
6 parallelisation using GPUs in RELION-2. *eLife* **5**, (Nov 15, 2016).
- 7 36. P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, Features and development of Coot. *Acta crystallographica.*
8 *Section D, Biological crystallography* **66**, 486 (Apr, 2010).
- 9 37. P. V. Afonine *et al.*, Real-space refinement in PHENIX for cryo-EM and crystallography. *Acta crystallographica.*
10 *Section D, Structural biology* **74**, 531 (Jun 1, 2018).