SICM6A: Identifying m⁶A Site across Species by Transposed GRU Network

Wenzhong Liu^{1*}

ABSTRACT

N6-methyladenosine (m⁶A) is the most prevalent cross-species RNA methylation modification and plays a pivotal role in various biological processes. The biochemical methods to find m⁶A sites are expensive and time-consuming, and the false positive rate of identified sites is high relatively. Meanwhile, the current computations are complex, and the prediction performance is relatively low both on little data sets and large data sets. This paper, at this point, presents a deep learning model with a transposed operation in the middle of GRU layers, SICM6A, for identifying m⁶A sites across-species. It adopts the mixed precision training manner to improve the speed and performance, and predicts m⁶A sites only by directly reading the 3-mer encoding of the m⁶A short sequence. The cross-validation and independent test verification show SICM6A is more accurate than the state-of-the-art methods. This, therefore, makes SICM6A provide new idea for predicting other modification sites of RNA sequences. The prediction software SICM6A is on github (<https://github.com/lwzyb/SICM6A>).

Key words: N6-methyladenosine; *k*-mer; deep learning; GRU Network

INTRODUCTION

N6-methyladenosine (m⁶A) is a methylation modification of the sixth nitrogen of the post-transcriptional adenosine base¹. As the studies have shown, m⁶A modification involves a variety of biological activities, such as mRNA splicing and stabilization, protein translation and localization, stem cell differentiation and reprogramming, biological clock, cerebellar development, and tumorigenesis²⁻⁴. The m⁶A site is evolutionarily conserved and has the consensus topology: DRACH motif for *Mus musculus* and *Homo sapiens*⁵ (where, D = A, G or U; R = A or G; and H = A, C or U), RRACH motif⁶ for *Arabidopsis thaliana* and RGAC motif⁷ for *Saccharomyces cerevisiae*. In the large range genome, methylated RNA immunoprecipitation sequencing techniques such as MeRIP-seq or m⁶A-seq can identify m⁶A sites in many species^{8,9}. Thus, in the whole genome analysis, high-throughput sequencing and wet experiments are the most expensive biochemical methods, hence the relatively high positive rate of false site identification¹⁰. Therefore, the use of computational tools to establish accurate models for predicting m⁶A sites is essential.

Current studies indicate that it is feasible to predict m⁶A sites from RNA sequences. The *Saccharomyces cerevisiae* methods of m⁶A site modification such as iRNA-Methyl¹¹, iDNA6mA-PseKNC¹², TargetM6A¹³, RNA-MethylPred¹⁴, M6APred¹⁵ appeared earlier. M6APred-EL¹⁶ is the recent way to predict M6A sites of *Saccharomyces cerevisiae*. RFATHM6A¹⁷ is a tool for predicting m⁶A sites in *Arabidopsis thaliana*. The tools encode RNA sequences using physical and chemical properties, motifs, and space structural features. The traditional machine learning techniques such as support vector and random forest are some of the methods used for classification or prediction. The feature extraction process of

¹Department of Software Engineering, School of Computer Science, Sichuan University of Science & Engineering. 180 Xueyuan Street, Huixing Road, Zigong City, Sichuan Province, China. 643000. *Corresponding author: Wenzhong Liu Telephone: +086 18708318095 Email: newwenzhong@hotmail.com

these methods is very complex, and the training time is too long normally. Therefore, these methods are not suitable for large data sets, for example, *Mammalia* data sets of SRAMP. SRAMP¹⁸ also has improved traditional machine learning methods to predict m⁶A sites of *Mammalia* by combining the scores of several random forest classifiers. If the RNA secondary structure is analyzed before SRAMP prediction, and the calculation amount is very large, so the feature extraction time is very long. WHISTLE uses support vector machines to predict N6-methyladenosine sites in human epithelial cells, and provides query service for m⁶A sites¹⁹. BERMP²⁰ introduced deep learning BGRU net to the prediction of m⁶A sites on SRAMP data sets for the first time, the calculation time can be reduced and the deep learning model shows a particular application prospect. Its prediction accuracy of m⁶A sites is higher in *Arabidopsis* and *Mammalia* datasets, but lower for *Saccharomyces cerevisiae*. Therefore, only the method combined the scores between random forest and GRU can help more precisely predict the m⁶A sites of *Saccharomyces cerevisiae*. DeepM6Aseq²¹ applies one-dimensional convolution layers to m⁶A site prediction of *Mammalia*, the *Mammalia* data set for SRAMP has had much removal of redundancy. Compared with traditional machine learning, the existing deep learning classifiers are superior for species with large data sets of *Mammalia*, but inferior to those with small data sets of *Saccharomyces cerevisiae*. It means that the compatibility of existing deep learning models in different species remains a problem, and the deep learning models have not been fully applied to the prediction of m⁶A sites across species. Therefore, this paper has constructed a deep learning model named SICM6A (Figure 1) for predicting m⁶A sites across species.

METHODS

The datasets of SICM6A are from the BERMP data sets and DeepM6Aseq. The BERMP data sets are sorted from SRAMP data sets and divided into three types: *Saccharomyces cerevisiae*, *Arabidopsis thaliana*, and *Mammalia*, wherein the *Mammalia* data set is included the full transcription and mature mRNA. The BERMP data set for each species includes a training set and an independent test set. The DeepM6Aseq datasets include a mammalian training set that is extracted from SRAMP and an independent test set that is retrieved from the real peak data of HepG2 and human brain cell line. This paper adopts BERMP datasets to test the cross-species prediction ability of SICM6A model and DeepM6Aseq datasets to verify the identification accuracy of experimental data.

Sequence fragments are extracted from every mRNA sequence of the data sets, where DRACH, RGAC or GAC motif is in the middle. The window length n of a sequence fragment is 51 bits for *Saccharomyces cerevisiae*, 101 bits for *Arabidopsis thaliana* and HepG2- the human brain, 251 bits for *Mammalia* mature RNA and 501 bits for *Mammalia* full transcription. Next, SICM6A uses 3-mer encoding (Figure 1.A) to encode the sequence fragments. Each 3 consecutive nucleotides are extracted from left to right in turn, and are encoded as integer vector. The gap symbol “-” and the four types of nucleotides, A, G, C and U are commonly composed of 5 symbols: 0, 1, 2, 3, 4. 3-mer nucleotides are converted to three numbers that combined as a integer. For example, as U->4, A->1, G->2, then 3-mer string UAG->412. After that, 3-mer nucleotides are encoded one by one from left to right on the sequence fragment, so each 3-mer nucleotides are therefore encoded as an integer and the sequence fragment changes to a short vector.

The SICM6A net what it named the transposed GRU network (Figure 1.B) analyses the short integer vector. SICM6A uses the word-vector embedding layer to convert the short vector into a two-dimensional matrix ($n \times 125$). The features of the two-dimensional matrix are processed in turn by three GRU layers: BGRU, UGRU and UGRU. The num_embeddings and embedding_dim of the word-vector embedding layer are 1024, 125 separately. The first GRU layer’s parameters like as: the number of layers is 2, input_size is 125, hidden_size is 125, dropout is 0.5 and bidirectional is equal to “True”. The second GRU layer has 2 layers, input_size is n , hidden_size is 128, dropout is 0.5. The third GRU layer has 2 layers, input_size is

16, hidden_size is 128, dropout is 0.5. The dimensional transpose operation is in the middle of the front two GRU layer and the dimensional reshape operation is in the middle of the last two UGRU layers. The BGRU layer outputs $n \times 250$ dimensional matrix, next the matrix is transposed and the UGRU generates 1×128 dimensional matrix. After the matrix is reshaped to 8×16 dimension, the third GRU layer outputs 1×128 dimensional matrix to the fully connected layer FC1 (Output=16) and FC2 (Output=2). The three layers of LeakyReLU (negative_slope=0.2), BatchNorm1d (momentum=0.5) and Dropout(p=0.25) are in the middle of FC1 and FC2. Finally, the log_Softmax layer computes the score of the sequence fragment. The sequence fragment is positive while the score is greater than a threshold, otherwise is negative. The thresholds are generally determined based on the scores of the independent test sets, corresponding to three grades of high (95%), medium (90%), low (85%) specificity (**Sp**).

The SICM6A model adopts Adam optimizer, with the learning rate of 0.0001 and the CrossEntropyLoss. The batch numbers of full transcription, mature mRNA, *Arabidopsis thaliana*, *Saccharomyces cerevisiae* and HepG2_human brain cell line are 400, 1000, 200, 200, 85 respectively. The model was trained by the Pytorch framework with the mixed precision training plugin “apex” using the GPU card GTX980TI 6G. The opt_level of apex is “O2” for SICM6A. To optimize Pytorch, apex uses FP16 to speed up tensor calculation, FP32 to calculate loss value, and completes loss scaling by multiplying or dividing the scaling factor. Therefore, a small amount of GPU RAM can improve the speed and performance of the model.

RESULTS

This paper performed five cross-validations on each *Mammalia* training data set and ten cross-validations on *Saccharomyces cerevisiae*, *Arabidopsis* training data sets respectively. The AUC (Area Under Curve) values of cross-validation are 0.9080, 0.8314, 0.9376, 0.8035 for full transcription, mature mRNA, *Arabidopsis thaliana* and *Saccharomyces cerevisiae* (Figure 2.a). In addition, the four models were also trained to predict the four corresponding independent test data sets. The AUC values are 0.9122, 0.8323, 0.9433, 0.8170 on the four types of independent test data sets (Figure 2.b). The results of cross-validation and independent test have similar AUC values. The AUC of SICM6A on independent test data sets is higher than state-of-the-art methods like BERMP, DeepM6A, SRAMP and M6APred-EL etc. too (Table 1). The figures indicate that the SICM6A model has better robustness and can be feasible for predicting m⁶A sites across species.

By comparing with advanced methods, SICM6A exhibits excellent predictive performance too. The **Acc** (Accuracy), **Sn** (Sensitivity), **Sp** (Specificity), **MCC** (Matthews Correlation Coefficient) of SICM6A are: 72.71, 55.56, 89.86, 48.3, 81.7 on *Saccharomyces cerevisiae* data set; 89.11, 88.28, 89.95, 78.2 on *Arabidopsis thaliana* data set; 86.33, 48.86, 90.01, 32.8 on mature mRNA data set; 88.30, 71.22, 90.00, 48.5 on full transcription (Table 1). The metric data **Acc**, **Sn**, **Sp**, **MCC** are better than existing methods on *Saccharomyces cerevisiae* and *Arabidopsis thaliana* data sets. The **Acc**, **Sn**, **Sp**, **MCC** of SICM6A are greater than SRAMP and DeepM6A on mature mRNA and full transcription data sets. The **Sn**, **MCC** of SICM6A are better than BERMP on full transcription data set and **Acc** is greater on mature mRNA data set, other metric values are close to BERMP. The metrics show that the SICM6A model has good stability not only for a small amount of data, *Saccharomyces cerevisiae*, and *Arabidopsis thaliana*, but also for *Mammalia* with a large amount of data. Meanwhile, the **Acc**, **F1-score**, **MCC**, **AUC** of SICM6A are 78.0, 0.828, 52.6, 83.21 (Table 2) on the HepG2 and human brain cell line data sets, and greater than DeepM6ASeq and SRAMP. It represents SICM6A could identify m⁶A sites from the experimental data.

CONCLUSIONS

All in all, a universal classifier can help identify m⁶A sites of multiple species, so it is important to establish a predictive model with such a function. However, the current state of research is that m⁶A site prediction methods of *Saccharomyces cerevisiae* are often unable to accurately predict mammalian m⁶A sites, and vice versa¹⁸. In view of this situation, this paper made innovations in coding, model and training methods, which were different from traditional methods, and constructed the prediction method of m⁶A sites based on deep learning, SICM6A. In terms of coding, this paper adopted a very efficient 3-mer coding. Compared with complex traditional coding, the coding method of SICM6A was not only very simple and intuitive, but also did not spend a lot of time on feature coding. On the model side, SICM6A used the word vector layer to learn the 3-mer encoding value to find its best feature vector. In the processing of the feature vector, the SICM6A introduced the transpose operation of feature into the middle of the first two GRU layers, namely BGRU layer and UGRU layer. This novel approach is similar to extracting features by horizontal and vertical filtering, the first GRU layer could capture features from the mammalian large data set, and the second GRU layer could capture features from the *Saccharomyces cerevisiae* small data set. In training, in order to solve the bottleneck problem of training large data sets on medium and low level GPU graphics cards - time consuming, this paper used the mixed precision training method, which greatly accelerated the training speed and improved the prediction accuracy. Therefore, through the above improved methods, this paper subtly solved the cross-species prediction problem of deep learning, so that the SICM6A model can quickly capture the m⁶A site features from large and small data sets.

Lastly, to facilitate the identification of m⁶A sites by biologists, it developed the pytorch based software SICM6A (Figure 3), which integrates the deep learning model SICM6A proposed in this paper. The source codes are on github (<https://github.com/lwzyb/SICM6A>).

REFERENCE

1. Liu, J. & Jia, G. Methylation Modifications in Eukaryotic Messenger RNA. *Journal of Genetics & Genomics* 41, 21-33 (2014).
2. Karikó, K., Buckstein, M., Ni, H. & Weissman, D. Suppression of RNA Recognition by Toll-like Receptors: The Impact of Nucleoside Modification and the Evolutionary Origin of RNA. *Immunity* 23, 165-175 (2005).
3. Jia, G. et al. N6-methyladenosine in nuclear RNA is a major substrate of the obesity-associated FTO. *Nature Chemical Biology* 7, 885-887 (2011).
4. Nilsen, T.W. Molecular biology. Internal mRNA methylation finally finds functions. *Science* 343, 1207-1208 (2014).
5. Dan, D. et al. Topology of the human and mouse m6A RNA methylomes revealed by m6A-seq. *Nature* 485, 201 (2012).
6. Levis, R. & Penman, S. 5'-Terminal structures of poly(A)+ cytoplasmic messenger RNA and of poly(A)+ and poly(A)- heterogeneous nuclear RNA of cells of the dipteran *Drosophila melanogaster*. *Journal of Molecular Biology* 120, 487-515 (1978).
7. Schraga, S. et al. High-resolution mapping reveals a conserved, widespread, dynamic mRNA methylation program in yeast meiosis. *Cell* 155, 1409-1421 (2013).
8. Meyer, K. et al. Comprehensive Analysis of mRNA Methylation Reveals Enrichment in 3' UTRs and near Stop Codons. *Cell* 149, 1635-1646 (2012).
9. Dan, D., Sharon, M.M., Mali, S.D., Ninette, A. & Gideon, R. Transcriptome-wide mapping of N(6)-methyladenosine by m(6)A-seq based on immunocapturing and massively parallel sequencing. *Nature Protocols* 8, 176-189 (2013).
10. Zhang, J., Feng, P., Lin, H. & Chen, W. Identifying RNA N6-Methyladenosine Sites in *Escherichia coli* Genome. *Frontiers in Microbiology* 9, 955- (2018).

11. Chen, W., Feng, P., Ding, H., Lin, H. & Chou, K.C. iRNA-Methyl: Identifying N 6 -methyladenosine sites using pseudo nucleotide composition. *Analytical Biochemistry* 490, 26-33 (2015).
12. Feng, P. et al. iDNA6mA-PseKNC: Identifying DNA N 6 -methyladenosine sites by incorporating nucleotide physicochemical properties into PseKNC. *Genomics* 111, S0888754318300090 (2018).
13. Li, G.Q., Liu, Z., Shen, H.B. & Yu, D.J. TargetM6A: Identifying N6-methyladenosine Sites from RNA Sequences via Position-Specific Nucleotide Propensities and a Support Vector Machine. *IEEE Trans Nanobioscience* PP, 1-1 (2016).
14. Jia, C.Z., Zhang, J.J. & Gu, W.Z. RNA-MethylPred: A High Accuracy Predictor to Identify N6-methyladenosine in RNA. *Analytical Biochemistry* 510, 72-75 (2016).
15. Chen, W., Hong, T., Liang, Z., Lin, H. & Zhang, L. Identification and analysis of the N6-methyladenosine in the *Saccharomyces cerevisiae* transcriptome. *Scientific Reports* 5, 13859 (2015).
16. Wei, L., Chen, H. & Su, R. M6APred-EL: A Sequence-Based Predictor for Identifying N6-methyladenosine Sites Using Ensemble Learning. *Mol Ther Nucleic Acids* 12, 635-644 (2018).
17. Wang, X. & Yan, R. RFathM6A: a new tool for predicting m 6 A sites in *Arabidopsis thaliana*. *Plant Molecular Biology* 96, 327-337 (2018).
18. Zhou, Y., Zeng, P., Li, Y.-H., Zhang, Z. & Cui, Q. SRAMP: prediction of mammalian N6-methyladenosine (m6A) sites based on sequence-derived features. *Nucleic Acids Res* 44, e91-e91 (2016).
19. Chen, K. et al. WHISTLE: a high-accuracy map of the human N6-methyladenosine (m6A) epitranscriptome predicted using a machine learning approach. *Nucleic Acids Res* 47, e41-e41 (2019).
20. Huang, Y., He, N., Chen, Y., Chen, Z. & Li, L. BERMP: a cross-species classifier for predicting m(6)A sites by integrating a deep learning algorithm and a random forest approach. *Int J Biol Sci* 14, 1669-1677 (2018).
21. Zhang, Y. & Hamada, M. DeepM6ASeq: prediction and characterization of m6A-containing sequences using deep learning. *BMC Bioinformatics* 19, 524-524 (2018).
22. Liu, Z. et al. pRNAm-PC: Predicting N(6)-methyladenosine sites in RNA sequences via physical-chemical properties. *Analytical Biochemistry* 497, 60-67 (2016).
23. Zhang, M. et al. Improving m(6)A sites prediction with heuristic selection of nucleotide physical-chemical properties. *Analytical Biochemistry* 508, S0003269716301154 (2016).

Legend

Figure 1. SICM6A model. **A.** 3-mer encoding. The string -, A, G, C and U are represented of 5 symbols: 0, 1, 2, 3, 4 respectively, then AUG->142. Each 3-mer nucleotides, are encoded from left to right on the sequence fragment, and the sequence fragment changes to a short vector. **B.** SICM6A net. The word-vector embedding layer converts the short vector into a two-dimensional vector ($n \times 125$). The two-dimensional vector are input into a transposed and reshape GRU structure---three GRU layers in turn. The dimensional transpose operation in the middle of the front two GRU layer and the dimensional reshape operation in the middle of the last two UGRU layers. Then 1×128 dimensional matrix is output to the fully connected layer FC1 and FC2. Next, the score of the sequence fragment is computed by the log_Softmax layer. At last, a threshold of score is applied to segment, the sequence fragment is classified as positive while the score is greater than the threshold, otherwise is negative.

Figure 2. Performance of SICM6A. **A.** Cross-validation Performance of SICM6A on the training data sets. Five cross validation is used to full transcription and mature mRNA data sets, and ten cross validation to *Saccharomyces cerevisiae*, *Arabidopsis thaliana* data sets. **B.** Prediction performance of SICM6A on the independent testing data sets.

SICM6A for Predicting M6A sites

Input your genomic/mRNA sequences with FASTA format.

Or upload a file(<=1M):

Mode

- ☐ Full transcript
- ☐ Mature mRNA
- ☐ *Arabidopsis thaliana*
- ☒ *Saccharomyces cerevisiae*
- ☐ hepg2_brain

Threshold

- ☒ High(95%)
- ☐ Midum(90%)
- ☐ Low(85%)

Figure 3. Interface of SICM6A. The user can paste the sequence into the text box and load the sequence text file in fasta format. Users can choose four species of full transcription, mature mRNA, *Arabidopsis thaliana* and *Saccharomyces cerevisiae*, and set three thresholds of high, medium and low. After the user submits, the software generates the scores of short sequence fragments.

Table 1. Prediction results of different classifiers on independent datasets.

Species	Classifiers	<i>Acc</i>	<i>Sn</i>	<i>Sp</i>	<i>MCC</i>	<i>AUC</i>
<i>Mammalia</i> Full transcript	SICM6A ¹	88.30	71.22	90.00	48.5	91.2
	BERMP ²	88.34	70.72	90.10	48.4	90.7
	DeepM6ASeq ³	85.81	44.14	89.98	29.0	82.9
	SRAMP ⁴	-	64.50	88.10	40.5	87.1
<i>Mammalia</i> Mature mRNA	SICM6A ¹	86.33	48.86	90.01	32.8	83.2
	BERMP ²	86.32	49.42	90.00	33.1	82.9
	DeepM6ASeq ³	84.77	32.39	89.99	19.8	73.3
	SRAMP ⁴	-	44.00	90.00	29.4	79.4
<i>Arabidopsis thaliana</i> mRNA	SICM6A ¹	89.11	88.28	89.95	78.2	94.3
	BERMP ²	87.20	84.21	90.19	74.5	93.4
	DeepM6ASeq ³	87.80	85.65	89.95	75.7	93.0
	RFathM6A ⁵	-	82.20	90.00	72.5	93.6
<i>Saccharomyces cerevisiae</i> mRNA	SICM6A ¹	72.71	55.56	89.86	48.3	81.7
	BERMP ²	68.84	47.34	90.34	41.7	80.0
	DeepM6ASeq ³	70.29	50.72	89.85	44.1	79.7
	M6APred-EL ⁶	71.26	52.66	89.86	45.8	80.8
	pRNA _m -PC ⁷	69.74	69.72	69.75	40.0	76.2
	M6A-HPCS ⁸	67.33	71.77	69.75	35.0	71.3
	RAM-NPPS ⁹	70.77	72.46	69.08	42.0	78.0

Note: As it is hard to get the codes of BERMP, SRAMP, RFathM6A, M6APred-EL, pRNA_m-PC, M6A-HPCS and RAM-NPPS, the comparison data is excerpted from their papers. 1. The data sets come from BERMP²⁰. 2. The data is excerpted from BERMP²⁰. 3. DeepM6ASeq is re-implemented using the data sets same as SICM6A. 4. The data is excerpted from SRAMP¹⁸. 5. The data is excerpted from RFathM6A¹⁷. 6. M6APred-EL is re-implemented using the data sets same as SICM6A. 7. The data is excerpted from pRNA_m-PC²² and BERMP²⁰. 8. The data is excerpted from M6A-HPCS²³ and BERMP²⁰. 9. The data excerpted from BERMP²⁰.

Table 2. Performance comparison on the HepG2 and human brain cell line data set ¹

Classifiers	<i>Acc</i>	<i>F1-score</i>	<i>MCC</i>	<i>AUC</i>
SICM6A	78.0	0.828	52.6	83.21
DeepM6ASeq ²	76.3	0.808	49.9	82.80
SRAMP-Mature ²	73.2	0.787	42.8	78.50
SRAMP-Full ²	76.2	0.824	48.3	81.80

Note: 1. The training and independent test data sets comes from DeepM6ASeq²¹. 2. The metric data is excerpted from DeepM6ASeq²¹.