Aryana Zardkoohi

Centro Nacional de Referencia en Entomología

Instituto Costarricense de Investigación y

Enseñanza en Nutrición y Salud

Tres Ríos, Cartago, Costa Rica

Phone: +506 8924 5528

Email: azardkoohi@inciensa.sa.cr

Co-occurrence of *kdr* mutations V1016I and F1534C in pyrethroid-resistant *Aedes aegypti* (Diptera: Culicidae) populations from Costa Rica

Aryana Zardkoohi¹, David Castañeda², Carmen Castillo², Juan C Lol², Francisco Lopez², Rodrigo Marín Rodríguez³, and Norma Padilla²

¹ Centro Nacional de Referencia en Entomología, Instituto Costarricense de Investigación y Enseñanza en Nutrición y Salud, Apartado Postal 4-2250, Tres Ríos, Cartago, Costa Rica.

² Grupo de Biología y Control de Vectores, Centro de Estudios en Salud, Universidad del Valle de Guatemala (CES-UVG), 18 Avenida 11-95, Zona 15, Vista Hermosa III, Ciudad de Guatemala, Guatemala.

³ Programa Nacional de Control de Vectores, Ministerio de Salud, San José, Costa Rica.

Abstract

Aedes aegypti (Linnaeus, 1762) is considered the most important mosquito vector species for several arboviruses (e.g., dengue, chikungunya, Zika) in Costa Rica. The main strategy for the control and prevention of *Aedes*-borne diseases relies on insecticide-based vector control. However, the emergence of insecticide resistance in the mosquito populations present a big threat for the prevention actions. The characterization of the mechanisms driving the insecticide resistance in *Ae. aegypti* are vital for decision making in vector control programs. Therefore, we analyzed the voltage-gated sodium channel gene for the presence of the V1016I and F1534C *kdr* mutations in pyrethroid-resistant *Ae. aegypti* populations from Puntarenas and Limon provinces, Costa Rica. The CDC bottle bioassays showed that both Costa Rican *Ae. aegypti* populations were resistant to permethrin and deltamethrin. In the case of *kdr* genotyping, results revealed the co-occurrence of V1016I and F1534C mutations in permethrin and deltamethrin-resistant populations, as well as the fixation of the 1534C allele. Therefore, our findings make an urgent call to expand the knowledge about the insecticide resistance status and mechanisms in the Costa Rican populations of *Ae. aegypti* which must be a priority to develop an effective resistance management plan.

Keywords

Aedes aegypti, insecticide resistance, voltage-gated sodium channel gene, *kdr* mutation, Costa Rica

Introduction

The region of the Americas has experienced the emergence and re-emergence of *Aedes*-borne diseases (ABD; e.g., dengue, chikungunya, Zika) (Weaver and Reisen 2010, Espinal et al. 2019), with the mosquito *Aedes aegypti* (Diptera: Culicidae, Linnaeus) widely distributed across the region (Leta et al. 2018). Costa Rica reported one of the highest numbers of confirmed cases of Zika virus (ZIKV) during the Zika emergency from 2015-2018 in the Central America region (PAHO 2018). Despite control efforts, ABD persists in Limon and Puntarenas provinces, where the highest number cases of ZIKV of Costa Rica were reported (Sanchez et al. 2019).

One of the main strategies for the control and prevention of ABD in Costa Rica relies on insecticide-based vector control, targeting *Aedes* populations. For the adult populations, several pyrethroids have been used sequentially (deltamethrin, cyfluthrin, cypermethrin) (Marín Rodríguez et al. 2009, Bisset et al. 2013), and currently permethrin and lambda-cyhalothrin are being used (Ministerio de Salud de Costa Rica 2019). However, focal studies have reported a decrease in the susceptibility against pyrethroids in *Ae. aegypti* from Puntarenas and Limon (Bisset et al. 2013, Calderón-Arguedas and Troyo 2014, 2016). Moreover, biochemical and synergists assays suggest that carboxylesterases and cytochrome P450s could be involved in pyrethroid resistance. Besides from biochemical activity, one of the main mechanisms of pyrethroid resistance in insects is the target-site insensitivity (Moyes et al. 2017), which has been poorly described in Costa Rica (Chaves et al. 2015). Non-synonymous mutations on the knockdown resistance (*kdr*) region of the voltage-gated sodium channel (VGSC) gene are responsible for this pyrethroid insensitivity (Du et al. 2016). A study published in 2003 reported seven novel mutations in *Ae. aegypti* strains from Asia, Africa, the Caribbean, South America and Oceania (Brenques et al. 2003). Although many non-synonymous mutations have been

discovered in the *Ae. aegypti* VGSC gene, only five have been associated with pyrethroid resistance: V410L, S989P, I1011M, V1016G, and F1534C (Du et al. 2013, Hirata et al. 2014, Haddi et al. 2017).

Molecular screening of the Latin American populations of *Ae. aegypti* evidenced the emergence of the *kdr* mutation V1016I in the region (Saavedra-Rodriguez et al. 2007) and its co-occurrence with the I1011M mutation, which was recently found in Panama (Murcia et al. 2019). The first report of the F1534C mutation was detected in 2011 in the *Ae. aegypti* population from Thailand (Yanola et al. 2011). Since then, the F1534C mutation has been documented worldwide (Linss et al. 2014, Kushwah et al. 2015, Hamid et al. 2018, Maestre-Serrano et al. 2019, Sombié et al. 2019), except from Central America. To the best of our knowledge, *kdr* mutations have not been studied in Costa Rican areas where pyrethroid resistance has been reported for *Ae. aegypti*. In order to fill the gap about the mechanisms associated with insecticide resistance, we analyzed the VGSC gene for the presence of *kdr* mutations in pyrethroid-resistant *Ae. aegypti* populations from Limon and Puntarenas.

Material and Methods

Mosquito sampling and rearing

Aedes mosquito eggs were collected as part of the National Entomological Surveillance of the Ministry of Health from Costa Rica in two provinces (Figure 1): Puntarenas (in the Pacific coast) and Limon (in the Caribbean coast). For each site, 50 ovitraps were placed and visited weekly to collect the mosquito eggs during May 2017. Eggs collected from the ovitraps for each site were pooled, hatched and the larvae reared into adults under insectary conditions ($28 \pm 2^\circ\text{C}$; $70 \pm 10\%$ RH, and 12:12 photoperiod). A F1 generation was obtained to get enough

individuals to perform the bioassays. All emerged adults were provided access to cotton pads soaked in 7% sugar solution until bioassays.

Adult bioassays

Insecticide susceptibility against permethrin (type I pyrethroid) and deltamethrin (type II pyrethroid) in the Costa Rican *Ae. aegypti* populations were assessed with the CDC bottle bioassay (Brogdon and McAllister 1998). From each population, a total of 100 F1 females (2-5-days old, non-blood-fed) were exposed to diagnostic dose of deltamethrin (10 µg/bottle) or permethrin (15 µg/bottle) for 30 min as diagnostic time. The New Orleans strain was used as insecticide-susceptible reference strain to support the appropriate performance of the bioassays. Based on the incapacitation rate at 30 min, each population was classified according to the standard WHO criteria as follows (WHO 2016): a population with an incapacitation rate >98% was classified as susceptible (S), an incapacitation rate between 90-98% suggested possible resistance (PR) requiring further tests, and an incapacitation rate <90% indicated confirmed resistance (R). Resistant and incapacitated mosquitoes were preserved individually at -20°C for molecular assays.

***Kdr* genotyping**

Genomic DNA (gDNA) was extracted from permethrin and deltamethrin-resistant mosquitoes using DNAzol (Thermo Fisher Scientific) according to manufacturer's instructions with modifications. Briefly, 100 µL of DNAzol was used to homogenize a whole mosquito and DNA pellets were resuspended in 50 µL of DEPC water. Subsequently, *kdr* genotyping was performed to detect the presence of V1016I and F1534C mutations in the pyrethroid-resistant mosquitoes. The genotype of each mutation was determined by a melt curve analysis of a real-time PCR using allele-specific primers for the V1016I mutation (Val1016f, Ile1016f and

Ile1016r) reported by Saavedra-Rodriguez et al. (2007), and for the F1534C mutation (F1534-f, C1534-f and CP-r) reported by Yanola et al. (2011).

PCR reaction for the V1016I mutation contained 10 µL of 2X PowerUp SYBR Green Master Mix (Thermo Fisher Scientific), 0.5 µL of each primer (50 pmol/µL), 1 µL of gDNA (5 to 15 ng) in a final volume of 20 µL. The PCR conditions were as followed: 95°C for 2 min, 40 cycles of 95°C for 10 sec, 60°C for 10 sec, and 72°C for 30 sec, with a final stage of 10 sec at 95°C. The melt curve conditions were 65°C to 95°C with an increase of 0.2°C each 10 sec. Based on the melting temperatures, samples were considered as mutant homozygous if they presented a unique peak of 79°C, wildtype homozygous if they presented a unique peak of 85°C, and heterozygous if they presented both of the previous peaks. In the case of F1534C mutation, each reaction contained 10 µL of 2X PowerUp SYBR Green Master Mix (Thermo Fisher Scientific), 0.066 µL of C1534-f primer (50 pmol/µL), 0.2 µL of primers F1534-f (50 pmol/µL) and CP-r (50 pmol/µL), 1 µL of gDNA (5 to 15 ng) in a final volume of 20 µL. The PCR conditions were as followed: 95°C for 3 min, 40 cycles of 95°C for 10 sec, 57°C for 10 sec, and 72°C for 30 sec, with a final stage of 10 sec at 95°C. The melt curve conditions were 65°C to 95°C with an increase of 0.5°C every 5 sec. Based on the melting temperatures, samples were considered as mutant homozygous if they presented a unique peak of 85°C, wildtype homozygous if they presented a unique peak of 80°C, and heterozygous if they presented both of the previous peaks. Previously genotyped *Ae. aegypti* by sequencing were used as positive controls for the PCR assays. The genotypic and allelic frequencies were calculated for each insecticide and population.

Sequencing of the VGSC gene

To validate the genotypes obtained from the *kdr* genotyping, two different PCR assays were used to amplify a partial fragment of domain IIS6 and domain IIIS6 of the VGSC gene in a subsample of 13 mosquitoes which represents the different genotypes identified by the real-time PCR. The first assay amplified the domain IIS6 using the primers IIS5-6 F and IIS5-6 R (spanning the codons 989, 1011 and 1016) originally reported by Al Nazawi et al. (2017). Each PCR reaction contained 1X Colorless GoTaq Flexi Buffer, 1.5 mM MgCl₂, 0.2 mM dNTPs, 2.5 µM of each primer (IIS5-6F and IIS5-6R), 1.25 U of GoTaq Hot Start Polymerase (Promega), and 1.25 ng of gDNA in a total volume of 50 µL. The cycling parameters included an initial step of denaturation at 95°C for 2 min, followed by 40 cycles of 94°C for 30 sec, 50°C for 30 sec, and 72°C for 1 min, with a final extension step of 10 min at 72°C. For the second assay, a region of the domain IIIS6 was amplified using the primers AaNa31F and AaNa31R (spanning the codon 1534) reported by Harris et al. (2010). Each PCR reaction contained 1X Colorless GoTaq Flexi Buffer, 1.5 mM MgCl₂, 0.2 mM dNTPs, 0.5 µM of each primer (AaNa31F and AaNa31R), 2.5 U of GoTaq Hot Start Polymerase (Promega), and 5 ng of gDNA in a total volume of 50 µL. The cycling parameters included an initial denaturation at 95°C for 2 min, followed by 35 cycles of 94°C for 30 sec, 59°C for 30 sec, and 72°C for 1 min, with a final extension step of 10 min at 72°C. Both PCR assays were performed using the SimpliAmp Thermal Cycler (Applied Biosystems).

All PCR products were visualized on a 2% agarose gel stained with ethidium bromide under UV light. Then, 50 µL of PCR product of each sample was purified with SpinPrep PCR Clean-Up Kit (EMD Millipore Corp) according to manufacturer's instructions. The PCR products purified were sequenced by MacroGen Inc. (Maryland, USA) with the primers used in the PCR

assays. Sequences were assembled and aligned using MEGA 7 (Kumar et al. 2016), and all codons were analyzed for non-synonymous substitutions.

Results and Discussion

The results of the CDC bottle bioassays are presented in Table 1. Incapacitation rates varied between 35 and 87% for deltamethrin and permethrin, confirming resistance to these pyrethroids in both Costa Rican *Ae. aegypti* populations. Puntarenas exhibited lower incapacitation rate (between 35 and 62%) for both insecticides compared with Limon (between 79 and 87%). Incapacitation rates for permethrin (between 35 and 79%) were lower than deltamethrin (between 62 and 87%) in the two *Ae. aegypti* populations from Costa Rica. A previous study in 2013, found suspected resistance to deltamethrin (incapacitation rate of 90.5%) in a Puntarenas *Ae. aegypti* population (Bisset et al. 2013). Therefore, our results suggest that deltamethrin resistance persists as well as an increase in the deltamethrin resistance from Puntarenas *Ae. aegypti* population.

As part of the insecticide resistance management, early detection of insecticide resistance mechanisms is critical for decision making in vector control programs (Corbel et al. 2019). To our knowledge, this is the first study to found and report a co-occurrence of both F1534C and V1016I mutations in resistant populations from Costa Rica (Table 2). The three different possible genotypes (V/V, V/I and I/I) for the *kdr* mutation V1016I were found in Puntarenas and Limon. However, for the 1534 codon, the homozygous mutant genotype C/C was the only one detected in the 72 mosquitoes analyzed. This result suggests that the mutation could be fixed in the field populations. The *kdr* allele frequency for 1016I varied between 0.167 and 0.980, and the *Ae. aegypti* population from Puntarenas showed a higher allele frequency than

Limon. In addition, DNA sequencing confirmed the presence of all genotypes observed by the allele-specific real-time PCR in the two Costa Rican *Ae. aegypti* populations.

The *kdr* mutation V1016I was originally reported as a widespread mutation across many Latin American *Ae. aegypti* populations, including a population from Guanacaste province, Costa Rica (Saavedra-Rodriguez et al. 2007). Although, functional conformation experiments do not confirm the role of the V1016I mutation in pyrethroid resistance, many studies have correlated the presence of this mutation to phenotypic resistance (Deming et al. 2016, Brito et al. 2018, Maestre-Serrano et al. 2019, Ryan et al. 2019). In contrast, *kdr* mutation F1534C has been strongly associated with DDT and pyrethroid resistance and confirmed by functional experiments in *Xenopus oocytes* (Du et al. 2013). Higher levels of pyrethroid resistance in mosquitoes have been observed in populations with the co-occurrence of the mutations V1016I and F1534C in other countries (Aponte et al. 2013, Kawada et al. 2014, Chapadense et al. 2015, Dusfour et al. 2015, Sombié et al. 2019). This co-occurrence has been observed approximately in the 97% of mosquitoes analyzed from Puntarenas, whereas a 32% of the mosquitoes from the province of Limon presented both mutations V1016I and F1534C (Table 2). These results could explain the different levels of pyrethroid resistance between the two *Ae. aegypti* populations analyzed (Table 1). In agreement with Linss et al. (2014), the V1016I mutation was not present without the F1534C mutation. Based on recent evidence, fixation of F1534C mutation in the populations analyzed represents a major threat for the insecticide-based vector control in Costa Rica (Vera-Maloof et al. 2015). Since this fixated mutation could favor the emergence of other *kdr* mutations, like the V1016I. Multiple *kdr* mutations could cause an additive effect increasing the levels of insecticide resistance and decreasing significantly the efficiency of vector control programs (Li et al. 2015, Plernsub et al. 2016, Saavedra-Rodriguez et al. 2018).

Overall, these results evidenced the presence of *kdr* mutations in permethrin and deltamethrin-resistant *Ae. aegypti* individuals from Costa Rica as well as the first report of the F1534C mutation and its co-occurrence with V1016I. Therefore, our findings make an urgent call to expand the knowledge about the insecticide resistance status and mechanisms in the Costa Rican populations of *Ae. aegypti* which must be a priority to develop an effective resistance management plan.

Acknowledgments

We thank Dr. Lissette Navas, INCIENSA's President, for facilitating the development of this research. This work was supported by Cooperative Agreement Number U51GH000970 funded by the Centers for Disease Control and Prevention (CDC). Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention from USA or the Department of Health and Human Services from the Costa Rica's Ministry of Health. The authors declare that they have not conflicts of interest.

References

- Aponte, H. A., R. P. Penilla, F. Dzúl-Manzanilla, A. Che-Mendoza, A. D. López, F. Solís, P. Manrique-Saide, H. Ranson, A. Lenhart, P. J. McCall, and A. D. Rodríguez. 2013. The pyrethroid resistance status and mechanisms in *Aedes aegypti* from the Guerrero state, Mexico. *Pestic. Biochem. Physiol.* 107: 226–234.
- Bisset, J. A., R. Marín, M. M. Rodríguez, D. W. Severson, Y. Ricardo, L. French, M. Díaz, and O. Pérez. 2013. Insecticide resistance in two *Aedes aegypti* (Diptera: Culicidae) strains from Costa Rica. *J. Med. Entomol.* 50: 352–361.
- Brengues, C., N. J. Hawkes, F. Chandre, L. McCarroll, S. Duchon, P. Guillet, S. Manguin,

248 **J. C. Morgan, and J. Hemingway. 2003.** Pyrethroid and DDT cross-resistance in *Aedes*
249 *aegypti* is correlated with novel mutations in the voltage-gated sodium channel gene. Med.
250 Vet. Entomol. 17: 87–94.

251 **Brito, L. P., L. Carrara, R. M. De Freitas, J. B. P. Lima, and A. J. Martins. 2018.** Levels
252 of resistance to pyrethroid among distinct *kdr* alleles in *Aedes aegypti* laboratory lines and
253 frequency of *kdr* alleles in 27 natural populations from Rio de Janeiro, Brazil. Biomed
254 Res. Int. 2018: 2410819.

255 **Brogdon, W. G., and J. C. McAllister. 1998.** Simplification of adult mosquito bioassays
256 through use of time-mortality determinations in glass bottles. J. Am. Mosq. Control Assoc.
257 14: 159–164.

258 **Calderón-Arguedas, Ó., and A. Troyo. 2014.** Perfil de resistencia a insecticidas en una cepa
259 de *Aedes aegypti* (Linnaeus) de la región Caribe de Costa Rica. Rev. Cubana Med. Trop.
260 66: 351–359.

261 **Calderón-Arguedas, Ó., and A. Troyo. 2016.** Evaluación de la resistencia a insecticidas en
262 cepas de *Aedes aegypti* (Diptera: Culicidae) de la Región Caribe de Costa Rica. Rev.
263 Cubana Med. Trop. 68: 95–104.

264 **Chapadense, F. G. G., E. K. K. Fernandes, J. B. P. Lima, A. J. Martins, L. C. Silva, W. T.**
265 **da Rocha, A. H. dos Santos, and P. Cravo. 2015.** Phenotypic and genotypic profile of
266 pyrethroid resistance in populations of the mosquito *Aedes aegypti* from Goiânia, Central
267 West Brazil. Rev. Soc. Bras. Med. Trop. 48: 607–609.

268 **Chaves, L. F., E. Kawashima, K. Futam, N. Minakawa, and R. Marín Rodríguez. 2015.**
269 Lack of *kdr* mutations in a population of Asian tiger mosquitoes from Costa Rica. Bull.
270 Insectology. 68: 61–63.

271 **Corbel, V., C. Durot, N. L. Achée, F. Chandre, M. B. Coulibaly, J.-P. David, G. J. Devine,**
272 **I. Dusfour, D. M. Fonseca, J. Griego, W. Juntarajumnong, A. Lenhart, S. Kasai, A.**

J. Martins, C. Moyes, L. C. Ng, J. Pinto, J. F. Pompon, P. Muller, K. Raghavendra, D. Roiz, H. Vatandoost, J. Vontas, and D. Weetman. 2019. Second WIN International Conference on “Integrated approaches and innovative tools for combating insecticide resistance in vectors of arboviruses”, October 2018, Singapore. *Parasit. Vectors.* 12: 331.

Deming, R., P. Manrique-Saide, A. Medina Barreiro, E. U. K. Cardenã, A. Che-Mendoza, B. Jones, K. Liebman, L. Vizcaino, G. Vazquez-Prokopec, and A. Lenhart. 2016. Spatial variation of insecticide resistance in the dengue vector *Aedes aegypti* presents unique vector control challenges. *Parasit. Vectors.* 9: 67.

Du, Y., Y. Nomura, G. Satar, Z. Hu, R. Nauen, S. Y. He, B. S. Zhorov, and K. Dong. 2013. Molecular evidence for dual pyrethroid-receptor sites on a mosquito sodium channel. *Proc. Natl. Acad. Sci.* 110: 11785–11790.

Du, Y., Y. Nomura, B. S. Zhorov, and K. Dong. 2016. Sodium channel mutations and pyrethroid resistance in *Aedes aegypti*. *Insects.* 7: E60.

Dusfour, I., P. Zorrilla, A. Guidez, J. Issaly, R. Girod, L. Guillaumot, C. Robello, and C. Strode. 2015. Deltamethrin resistance mechanisms in *Aedes aegypti* populations from three French overseas territories worldwide. *PLoS Negl. Trop. Dis.* 9: e0004226.

Espinal, M. A., J. K. Andrus, B. Jauregui, S. H. Waterman, D. M. Morens, J. I. Santos, O. Horstick, L. A. Francis, and D. Olson. 2019. Emerging and reemerging *Aedes*-transmitted arbovirus infections in the region of the Americas: Implications for health policy. *Am. J. Public Health.* 109: 387–392.

Haddi, K., H. V. V. Tomé, Y. Du, W. R. Valbon, Y. Nomura, G. F. Martins, K. Dong, and E. E. Oliveira. 2017. Detection of a new pyrethroid resistance mutation (V410L) in the sodium channel of *Aedes aegypti*: A potential challenge for mosquito control. *Sci. Rep.* 7: 46549.

Hamid, P. H., V. I. Ninditya, J. Prastowo, A. Haryanto, A. Taubert, and C. Hermosilla.

2018. Current status of *Aedes aegypti* insecticide resistance development from Banjarmasin, Kalimantan, Indonesia. Biomed Res. Int. 2018: 1735358.

Harris, A. F., S. Rajatileka, and H. Ranson. 2010. Pyrethroid resistance in *Aedes aegypti* from Grand Cayman. Am. J. Trop. Med. Hyg. 83: 277–284.

Hirata, K., O. Komagata, K. Itokawa, A. Yamamoto, T. Tomita, and S. Kasai. 2014. A single crossing-over event in voltage-sensitive Na⁺ channel genes may cause critical failure of Dengue mosquito control by insecticides. PLoS Negl. Trop. Dis. 8: e3085.

Kawada, H., S. Z. M. Oo, S. Thaung, E. Kawashima, Y. N. M. Maung, H. M. Thu, K. Z. Thant, and N. Minakawa. 2014. Co-occurrence of point mutations in the voltage-gated sodium channel of pyrethroid-resistant *Aedes aegypti* populations in Myanmar. PLoS Negl. Trop. Dis. 8: e3032.

Kumar, S., G. Stecher, and K. Tamura. 2016. MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. Mol. Biol. Evol. 33: 1870–1874.

Kushwah, R. B. S., C. L. Dykes, N. Kapoor, T. Adak, and O. P. Singh. 2015. Pyrethroid-resistance and presence of two knockdown resistance (*kdr*) mutations, F1534C and a novel mutation T1520I, in Indian *Aedes aegypti*. PLoS Negl. Trop. Dis. 9: e3332.

Leta, S., T. J. Beyene, E. M. De Clercq, K. Amenu, M. U. G. Kraemer, and C. W. Revie. 2018. Global risk mapping for major diseases transmitted by *Aedes aegypti* and *Aedes albopictus*. Int. J. Infect. Dis. 67: 25–35.

Li, C. X., P. E. Kaufman, R. De Xue, M. H. Zhao, G. Wang, T. Yan, X. X. Guo, Y. M. Zhang, Y. De Dong, D. Xing, H. D. Zhang, and T. Y. Zhao. 2015. Relationship between insecticide resistance and *kdr* mutations in the dengue vector *Aedes aegypti* in Southern China. Parasit. Vectors. 8: 325.

Linss, J. G. B., L. P. Brito, G. A. Garcia, A. S. Araki, R. V. Bruno, J. B. P. Lima, D. Valle, and A. J. Martins. 2014. Distribution and dissemination of the Val1016Ile and

Phe1534Cys *Kdr* mutations in *Aedes aegypti* Brazilian natural populations. *Parasit. Vectors*. 7: 25.

Maestre-Serrano, R., P. Pareja-Loaiza, D. Gomez Camargo, G. Ponce-García, and A. E. Flores. 2019. Co-occurrence of V1016I and F1534C mutations in the voltage-gated sodium channel and resistance to pyrethroids in *Aedes aegypti* (L.) from the Colombian Caribbean region. *Pest Manag. Sci.* 75: 1681–1688.

Marín Rodríguez, R., M. del C. Marquetti Fernández, and M. Díaz Ríos. 2009. Índices larvales de *Aedes aegypti* antes y después de intervenciones de control en Limón, Costa Rica. *Rev. Cubana Med. Trop.* 61.

Ministerio de Salud de Costa Rica. 2019. Plan nacional de monitoreo y manejo de la resistencia a los insecticidas en vectores de arbovirosis y malaria en Costa Rica.

Moyes, C. L., J. Vontas, A. J. Martins, L. C. Ng, S. Y. Koou, I. Dusfour, K. Raghavendra, J. Pinto, V. Corbel, J. P. David, and D. Weetman. 2017. Contemporary status of insecticide resistance in the major *Aedes* vectors of arboviruses infecting humans. *PLoS Negl. Trop. Dis.* 11: e0005625.

Murcia, O., B. Henríquez, A. Castro, S. Koo, J. Young, R. Márquez, D. Pérez, L. Cáceres, and A. Valderrama. 2019. Presence of the point mutations Val1016Gly in the voltage-gated sodium channel detected in a single mosquito from Panama. *Parasit. Vectors*. 12: 62.

Al Nazawi, A. M., J. Aqili, M. Alzahrani, P. J. McCall, and D. Weetman. 2017. Combined target site (*kdr*) mutations play a primary role in highly pyrethroid resistant phenotypes of *Aedes aegypti* from Saudi Arabia. *Parasit. Vectors*. 10: 161.

(PAHO) Pan American Health Organization. 2018. Zika cases and congenital syndrome associated with Zika virus reported by countries and territories in the Americas, 2015 - 2018. Cumulative cases. *Pan Am. Heal. Organ.*

348 **Plernsub, S., J. Saingamsook, J. Yanola, N. Lumjuan, P. Tippawangkosol, K. Sukontason,**
349 **C. Walton, and P. Somboon. 2016.** Additive effect of knockdown resistance mutations,
350 S989P, V1016G and F1534C, in a heterozygous genotype conferring pyrethroid resistance
351 in *Aedes aegypti* in Thailand. *Parasit. Vectors.* 9: 417.

352 **Ryan, S. J., S. J. Mundis, A. Aguirre, C. A. Lippi, E. Beltrán, F. Heras, V. Sanchez, M. J.**
353 **Borbor-Cordova, R. Sippy, A. M. Stewart-Ibarra, and M. Neira. 2019.** Seasonal and
354 geographic variation in insecticide resistance in *Aedes aegypti* in southern Ecuador. *PLoS*
355 *Negl. Trop. Dis.* 13: 1–18.

356 **Saavedra-Rodriguez, K., F. V. Maloof, C. L. Campbell, J. Garcia-Rejon, A. Lenhart, P.**
357 **Penilla, A. Rodriguez, A. A. Sandoval, A. E. Flores, G. Ponce, S. Lozano, and W. C.**
358 **Black. 2018.** Parallel evolution of vgsc mutations at domains IS6, IIS6 and IIS6 in
359 pyrethroid resistant *Aedes aegypti* from Mexico. *Sci. Rep.* 8: 6747.

360 **Saavedra-Rodriguez, K., L. Urdaneta-Marquez, S. Rajatileka, M. Moulton, A. E. Flores,**
361 **I. Fernandez-Salas, J. Bisset, M. Rodriguez, P. J. McCall, M. J. Donnelly, H. Ranson,**
362 **J. Hemingway, and W. C. Black IV. 2007.** A mutation in the voltage-gated sodium
363 channel gene associated with pyrethroid resistance in Latin American *Aedes aegypti*.
364 *Insect Mol. Biol.* 16: 785–798.

365 **Sanchez, F., L. A. Barboza, and P. Vásquez. 2019.** Parameter estimates of the 2016-2017
366 Zika outbreak in Costa Rica: An Approximate Bayesian Computation (ABC) approach.
367 *Math. Biosci. Eng.* 16: 2738–2755.

368 **Sombié, A., E. Saiki, F. Yaméogo, T. Sakurai, T. Shirozu, S. Fukumoto, A. Sanon, D.**
369 **Weetman, P. J. McCall, H. Kanuka, and A. Badolo. 2019.** High frequencies of F1534C
370 and V1016I *kdr* mutations and association with pyrethroid resistance in *Aedes aegypti*
371 from Somgandé (Ouagadougou), Burkina Faso. *Trop. Med. Health.* 47: 2.

372 **Vera-Maloof, F. Z., K. Saavedra-Rodriguez, A. E. Elizondo-Quiroga, S. Lozano-Fuentes,**

and W. C. Black IV. 2015. Coevolution of the Ile1,016 and Cys1,534 mutations in the voltage gated sodium channel gene of *Aedes aegypti* in Mexico. PLoS Negl. Trop. Dis. 9: e0004263.

Weaver, S. C., and W. K. Reisen. 2010. Present and future arboviral threats. Antiviral Res. 85: 328–345.

(WHO) World Health Organization. 2016. Test procedures for insecticide resistance monitoring in malaria vector mosquitoes: Second edition. World Heal. Organ. Tech. Rep. Ser.

Yanola, J., P. Somboon, C. Walton, W. Nachaiwieng, P. Somwang, and L. Prapanthadara. 2011. High-throughput assays for detection of the F1534C mutation in the voltage-gated sodium channel gene in permethrin-resistant *Aedes aegypti* and the distribution of this mutation throughout Thailand. Trop. Med. Int. Heal. 16: 501–509.

Figure legend

Fig 1. *Aedes* mosquito collection sites. A map of Costa Rica shows the location of Puntarenas (in the Pacific coast) and Limon (in the Caribbean coast) provinces.

Figure 1

Table 1. Pyrethroid susceptibility in Costa Rican *Aedes aegypti* populations assessed by CDC bottle bioassay.

Population	n	% incapacitation [95% CI]	Status
Permethrin			
Puntarenas	100	35 [16.23-53.77]	R
Limon	100	79 [77.04-80.96]	R
Deltamethrin			
Puntarenas	100	62 [54.49-69.51]	R
Limon	100	87 [85.04-88.96]	R

Table 2. Genotypic and allelic frequencies of the V1016I and F1534C *kdr* mutations in pyrethroid-resistant mosquitoes from Costa Rican *Aedes aegypti*.

Population	n	V1016I genotypes			F1534C genotypes			<i>kdr</i> allele frequency	
		V/V	V/I	I/I	F/F	F/C	C/C	1016I	1534C
Permethrin-resistant									
Puntarenas	13	1	5	7	0	0	13	0.731	1
Limon	21	16	3	2	0	0	21	0.167	1
Deltamethrin-resistant									
Puntarenas	25	0	1	24	0	0	25	0.980	1
Limon	13	7	3	3	0	0	13	0.347	1