Structural basis of DNA targeting by a transposon-encoded CRISPR-Cas system

Tyler S. Halpin-Healy¹, Sanne E. Klompe¹, Samuel H. Sternberg^{*1} and Israel S.
Fernández^{*1}

¹Department of Biochemistry and Molecular Biophysics, Columbia University,
New York, NY, 10032, USA.

^{*}To whom correspondence should be addressed at shs2107@cumc.columbia.edu (SHS) and
isf2106@cumc.columbia.edu (ISF)

Abstract

Bacteria have evolved adaptive immune systems encoded by Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) and the CRISPR-associated (Cas) genes to maintain genomic integrity in the face of relentless assault from pathogens and mobile genetic elements [1–3]. Type I CRISPR-Cas systems canonically target foreign DNA for degradation via the joint action of the ribonucleoprotein complex Cascade and the helicase-nuclease Cas3 [4, 5] but nuclease-deficient Type I systems lacking Cas3 have been repurposed for RNA-guided transposition by bacterial Tn7-like transposons [6, 7]. How CRISPR- and transposon-associated machineries collaborate during DNA targeting and insertion has remained elusive. Here we determined structures of a novel TniQ-Cascade complex encoded by the *Vibrio cholerae* Tn6677 transposon using single particle electron cryo-microscopy (cryo-EM), revealing the mechanistic basis of this functional coupling. The quality of the cryo-EM maps allowed for de novo modeling and refinement of the transposition protein TniQ, which binds to the Cascade complex as a dimer in a head-to-tail configuration, at the interface formed by Cas6 and Cas7 near the 3' end of the crRNA. The natural Cas8-Cas5 fusion protein binds the 5' crRNA handle and contacts the TniQ dimer via a flexible insertion domain. A target DNA-bound structure reveals critical interactions necessary for protospacer adjacent motif (PAM) recognition and R-loop formation. The present work lays the foundation for a structural understanding of how DNA targeting by TniQ-Cascade

leads to downstream recruitment of additional transposon-associated proteins, and
will guide protein engineering efforts to leverage this system for programmable
DNA insertions in genome engineering applications.

31

32 We previously demonstrated that a transposon derived from *Vibrio cholerae* Tn6677 undergoes
 33 programmable transposition in *E. coli* directed by a CRISPR RNA (crRNA), and that this activ-
 34 ity requires four transposon- and three CRISPR-associated genes in addition to a CRISPR array
 35 (**Fig. 1a** [7]). Whereas TnsA, TnsB, and TnsC exhibit functions that are consistent with their
 36 homologs from a related and well-studied cut-and-paste DNA transposon, *E. coli* Tn7 (reviewed in
 37 citePeters:2014aa), we showed that TniQ, a homolog of *E. coli* TnsD, forms a co-complex with the
 38 Cascade ribonucleoprotein complex encoded by the Type I-F variant CRISPR-Cas system. This
 39 finding suggested an alternative role for TniQ, as compared to the role of *Eco*TnsD in identifying
 40 target sites during Tn7 transposition. Rather, we proposed that RNA-guided DNA targeting by
 41 Cascade could deliver TniQ to DNA in a manner compatible with downstream transpososome for-
 42 mation, and that TniQ might interact with Cascade near the 3' end of the crRNA, consistent with
 43 RNA-guided DNA insertion occurring approx. 49-bp downstream from the PAM-distal edge of the
 44 target site. To determine this unambiguously, we purified the *V. cholerae* TniQ-Cascade complex
 45 loaded with a native crRNA and determined its structure by cryo-EM. The overall complex adopts
 46 a helical architecture with protuberances at both ends (**Fig. 1** and **Extended Data Fig. 1**
 47 and **2**). The global architecture is similar to previously determined structures of Cascade from I-E
 48 and I-F systems (Extended Data Fig. 3) [8–11] with the exception of a large mass of additional
 49 density attributable to TniQ (see below). Maximum likelihood classification methods implemented
 50 in Relion3 [12] allowed us to identify significant dynamics in the entire complex, which appears to

51 "breathe", widening and narrowing the distance between the two protuberances (**Extended Data**
52 **Fig. 1d** and **Supplementary Movie 1**). The large subunit encoded by a natural Cas8-Cas5
53 fusion protein (hereafter referred to simply as Cas8) forms one protuberance and recognizes the
54 5' end of the crRNA via base- and backbone-specific contacts (**Extended Data Fig. 4, 5a-**
55 **c, 6a**), akin to the canonical roles played by Cas8 and Cas5 (**Extended Data Fig. 3**). Cas8
56 exhibits two primary subdomains formed mainly by α -helices, along with a third domain of ap-
57 proximately 100 residues (residues 277 to 385) that is predicted to form three α -helices but could
58 not be built in our maps due to its intrinsic flexibility (**Fig. 1c**). However, low-pass filtered maps
59 revealed that this flexible domain connects with the TniQ protuberance at the opposite end of the
60 crescent-shaped complex (**Extended Data Fig. 2e**). Additionally, there seemed to be a loose
61 coupling between the Cas8 flexible domain and overall "breathing" of the complex, as stronger
62 density for that domain could be observed in the closed state (**Extended Data Fig. 1d** and
63 **Supplementary Movie 1**). Six Cas7 subunits protect much of the crRNA by forming a helical
64 filament along its length (**Fig. 1b** and **d**), similar to other Type I Cascade complexes (**Extended**
65 **Data Fig. 3** [8–11]. A "finger" motif in Cas7 clamps the crRNA in regular intervals, causing
66 every sixth nucleotide (nt) of the 32-nt spacer to flip out while leaving the flanking nucleotides
67 available for DNA recognition (**Extended Data Fig. 4f**). These bases are pre-ordered in short
68 helical segments, with a conserved phenylalanine stacking below the first base of every segment.
69 Cas7.1, the monomer furthest away from Cas8, interacts with Cas6 (also known as Csy4), which
70 is the ribonuclease responsible for processing of the precursor RNA transcript derived from the

CRISPR locus. The Cas6-Cas7.1 interaction is mediated by a β -sheet formed by the contribution of a β -strands from Cas6 and the two β -strands that form the "finger" of Cas7.1 (**Extended Data Fig. 5f**). Cas6 also forms extensive interactions with the conserved stem-loop in the repeat-derived 3' crRNA handle (**Fig. 1 and Extended Data Fig. 5d and e**), with an arginine-rich α -helix (residues 110 to 128) docked in the major groove, positioning multiple basic residues within interaction distance of the negatively charged RNA backbone. The interaction established between Cas6 and Cas7.1 forms a continuous surface where TniQ is docked, forming the other protuberance of the crescent. The intrinsic flexibility of the complex rendered lower local resolutions in this area of the maps, which we overcame using local alignments masking the area comprising TniQ, Cas6, Cas7.1 and the crRNA handle (**Extended Data Fig. 7**). The enhanced maps allowed for de novo modeling and refinement of TniQ, for which no previous structure or homology model has been reported (**Fig. 2**). Notably, TniQ binds to Cascade as a dimer with head-to-tail configuration (**Fig. 2**), a surprising result given the expectation that *EcoTnsD* functions as a monomer during Tn7 transposition [13]. TniQ is composed of two domains: an N-terminal domain of approximately 100 residues formed by three short α -helices and a second, larger domain of approximately 300 residues with signature sequence for the TniQ family. A DALI search [14] using the refined TniQ model as a probe yielded significant structural similarity of the N-terminal domain to proteins containing Helix-Turn-Helix (HTH) domains (**Extended Data Fig. 8**). This domain is often involved in nucleic acid recognition, however there are reported examples where it has been re-purposed for protein-protein interactions [15]. The remaining C-terminal TniQ-domain is formed by 10 α -helices

of variable length and is predicted to contain two tandem zinc finger motifs, though this region was poorly defined in the maps (**Fig. 2**). Overall, the double domain composition of TniQ results in an elongated structure, bent at the junction of the HTH and the TniQ-domain (**Fig. 2**). The HTH domain of one monomer engages the TniQ-domain of the other monomer via interactions between α -helix 3 (H3) and α -helix 11 (H11), respectively, in a tight protein-protein interaction (**Fig. 2c**). This reciprocal interaction is complemented by multiple interactions established between the TniQ-domains from both monomers (up to 45 non-covalent interactions as reported by PISA [16]). Tethering of the TniQ dimer to Cascade is accomplished by specific interactions established with both Cas6 and Cas7.1 (**Fig. 3**). One monomer of TniQ interacts with Cas6 via its C-terminal TniQ-domain, while the other TniQ monomer contacts Cas7.1 through its N-terminal HTH domain (**Fig. 2b, 3**). The loop connecting alpha-helices H6 and H7 of the TniQ-domain of the first TniQ monomer is inserted in a hydrophobic cavity formed at the interface of two α -helices of Cas6 (**Fig. 3b, d**). The TniQ histidine residue 265 is involved in rearranging the hydrophobic loop connecting H6 and H7 (**Fig. 3d**), which is inserted in the hydrophobic pocket of Cas6 formed by residues L20, Y74, M78, Y83 and F84. The HTH domain of the other TniQ monomer interacts with Cas7.1 through a network of interactions established mainly by α -helix H2 and the linker connecting H2 and H3 (**Fig. 3c, e**). Thus, both the HTH domain and the TniQ-domain exert dual roles to drive TniQ dimerization and dock onto Cascade. In order to explore the structural determinants of DNA recognition by the TniQ-Cascade complex, we determined the structure of the complex bound to a double-stranded DNA (dsDNA) substrate containing the 32-bp target sequence, 5'-CC-3' PAM,

and 20-bp of flanking dsDNA on both ends (**Fig. 4** and **Extended Data Fig. 9**). Density for 28 nucleotides of the target strand (TS) and 8 nucleotides for the non-target strand (NTS) could be confidently assigned in the reconstructed maps (**Fig.4c**). As with previous I-F Cascade structures, Cas8 recognizes the double-stranded PAM within the minor groove (**Extended Data Fig. 10** [10]), and an arginine residue (R246) establishes a stacking interaction with a guanine nucleotide on the TS, which acts like a wedge to separate the double-stranded PAM from the neighboring unwound DNA where base-pairing with the crRNA begins (**Fig. 4b**). Twenty-two nucleotides of the TS within the 32-bp target showed clear density, but surprisingly, the terminal nine nucleotides were not ordered. The TS base-pairs with the spacer region of the crRNA in short, discontinuous, helical segments, as observed previously for I-E and I-F DNA-bound Cascade complexes [10,11] with every 6th base flipped out of the heteroduplex by the insertion of a Cas7 finger (**Extended Data Fig. 6b**). The observed 22-bp heteroduplex is stabilized by the four Cas7 monomers proximal to the PAM (Cas7.6-7.3), but even after local masked refinements, no density could be observed for any TS nucleotides that would base-pair with the 3' end of the crRNA spacer bound by Cas7.2 and Cas7.1. These two Cas7 monomers are proximal to Cas6 and in the region previously described to exhibit dynamics due to the interaction of the Cas8 flexible domain with the inner face of the TniQ-dimer. In addition, the disordered nucleotides also correspond to positions 25-28 of the target site where RNA-DNA mismatches are detrimental for RNA-guided DNA integration [7]. Thus, we propose the possibility that the partial R-loop structure we observed may represent an intermediate conformation refractory to integration, and that further structural rearrangements may be critical

for further stabilization of an open conformation, possibly driven by recruitment of the TnsC ATPase. Here we present the first cryo-EM structures of a CRISPR-Cas effector complex bound to the transposition protein TniQ, with and without target DNA. These structures reveal the unexpected presence of TniQ as a dimer that forms bipartite interactions with Cas6 and Cas7.1 within the Cascade complex, forming a likely recruitment platform for downstream-acting transposition proteins (Fig. 4d). Our structures furthermore reveal a possible fidelity checkpoint, whereby formation of a complete R-loop requires conformational rearrangements that may depend on extensive RNA-DNA complementarity and/or downstream factor recruitment; this proofreading step could account for the highly specific RNA-guided DNA integration we previously reported for the *V. cholerae* transposon [7]. In light of recent work demonstrating exaptation of Type V-K CRISPR-Cas systems by similar Tn7-like transposons that also encode TniQ [17, 18], it will be interesting to determine whether tethering of TniQ to evolutionarily distinct CRISPR RNA effector complexes - Cascade or Cas12k - is a general theme of RNA-guided transposition.

Methods

TniQ-Cascade purification.

Protein components of TniQ-Cascade were expressed from a pET-derivative vector containing the native *V. cholerae* tniQ-cas8-cas7-cas6 operon with an N-terminal His10-MBP-TEVsite fusion on TniQ. The crRNA was expressed separately from a pACYC-derivative vector containing a minimal

repeat-spacer-repeat CRISPR array encoding a spacer from the endogenous *V. cholerae* CRISPR array. The TniQ-Cascade complex was overexpressed and purified as described previously [7], and was stored in Cascade Storage Buffer (20 mM Tris-Cl, pH 7.5, 200 mM NaCl, 1 mM DTT, 5% glycerol).

Sample preparation for electron microscopy.

For negative staining, 3 μ L of purified TniQ-Cascade ranging from 100 nM to 2 μ M was incubated with plasma treated (H₂/O₂ gas mix, Gatan Solarus) CF400 carbon-coated grids (EMS) for 1 minute. Excess solution was blotted and 3 μ L of 0.75% uranyl formate was added for an additional minute. Excess stain was blotted away and grids were air-dried overnight. Grid screening for both negative staining and cryo conditions was performed on a Tecnai-F20 microscope (FEI) operated at 200 KeV and equipped with a Gatan K2-Summit direct detector. Microscope operation and data collection were carried out using the Leginon/Appion software. Initial negative staining grid screening allowed determination of a suitable concentration range for cryo conditions. Several grid geometries were tested in the 1-4 μ M concentration range for cryo conditions using a Vitrobot Mark-II operated at 4 C, 100% humidity, blot force 3, drain time 0, waiting time 15 seconds, and blotting times ranging from 3-5 seconds. The best ice distribution and particle density was obtained with 0.6/1 UltrAuFoil grids (Quantifoil).

Electron microscopy.

A preliminary dataset of 300 images in cryo was collected with the Tecnai-F20 microscope using a pixel size of 1.22 Å/pixel with illumination conditions adjusted to 8 e-/pixel/second with a frame window of 200 ms. Preprocessing and image processing were integrally done in Relion3 [12] with ctf estimation integrated via a wrapper to Gctf [19]. An initial model computed using the SGD algorithm [20] implemented in Relion3 was used as initial reference for a refine 3D job that generated a sub-nanometric reconstruction with approximately 10,000 selected particles. Clear secondary structure features in the 2D averages and the 3D reconstruction could be identified. For the DNA-bound TniQ-Cascade complex containing DNA, we pre-incubated two complementary 74-nt oligonucleotides (NTS: 5' TTCATCAAGCCATTGGACCGCCTTACAGGACGCTTTGGCTTCATTGCTTTTCAC 3', TS: 5' TTTTGGCCGTCAAGGCGAAGCTGAAAAGCAATGAAGCCAAAGCGTCCTGTAAGGCGGT 3') for 5 minutes at 95° C in hybridization buffer (20 mM Tris-Cl, pH 7.5, 100 mM KCl, 5 mM MgCl₂) to form dsDNA, which was subsequently aliquoted and flash frozen. Complex formation was performed by incubating a 3x molar excess of dsDNA with TniQ-Cascade at 37° C for 5 minutes prior to vitrification, which followed the conditions optimized for the apo complex (defined as TniQ-Cascade with crRNA but no DNA ligand). High resolution data for the apo complex were collected in a Tecnai-Polara-F30 microscope operated at 300 KeV equipped with a K3 direct detector (Gatan). A 30 µm C2 aperture was used with a pixel size of 0.95 Å/pixel and illumination conditions in microprobe mode adjusted to a fluence of 16e-/pixel/second. Four-second images with a frame width of 100 ms (1.77 e-/2/frame) were collected in counting mode. For the DNA-bound

complex, high resolution data were collected in a Titan Krios microscope (FEI) equipped with an energy filter (20 eV slit width) and a K2 direct detector (Gatan) operated at 300 KeV. A 50 μm C2 aperture was used with a pixel size of 1.06 $\text{\AA}/\text{pixel}$ and illumination conditions adjusted in nanoprobe mode to a fluence of 8e-/pixel/second. Eight-second images with a frame width of 200 ms (1.42 e-/2/frame) were collected in counting mode.

Image processing.

Motion correction was performed for every micrograph applying the algorithm described for Motioncor2 [21] implemented in Relion3 with 5 by 5 patches for the K2 data and 7 by 5 patches for the K3 data. Parameters of the contrast transfer function for each motion-corrected micrograph were obtained using Gctf integrated in Relion3. Initial particle picking of a subset of 200 images randomly chosen was performed with the Laplacian tool of the Auto-picking module of Relion3, using an estimated size for the complex of 200 $\text{\AA}$. 15,000 particles were extracted in a 300 pixels box size and binned 3 times for an initial 2D classification job. Selected 2D averages from this job were used as templates for Auto-picking of the full dataset. The full dataset of binned particles was subjected to a 2D classification job to identify particles able to generate averages with clear secondary structure features. The selected subgroup of binned particles after the 2D classification selection was refined against a 3D volume obtained by SGD with the F20 data. This "consensus" volume was inspected to localize areas of heterogeneity which were clearly identified at both ends of the crescent shape characteristic of this complex. Both ends were then individually masked using

soft masks of around 20 pixels that were subsequently used in classification jobs without alignments in Relion3. The T parameter used for this classification job was 6 and the total number of classes was 10. This strategy allowed us to identify two main population of particles which correspond to an "open" and "closed" state of the complex. Particles from both subgroups were separately re-extracted to obtain unbinned datasets for further refinement. New features implemented in Relion3, namely Bayesian polishing and ctf parameters refinement, allowed the extension of the resolution to 3.4, 3.5 and 2.9 Å for the two apo and the DNA-bound complexes, respectively. Post processing was performed with a soft-mask of 5 pixels being the B-factor estimated automatically in Relion3 following standard practice. A final set of local refinements was performed with the masks used for classification. The locally aligned maps exhibit very good quality for the ends of the C-shape. These maps were used for de novo modeling and initial model refinement.

Model building and refinement.

For the Cas7 and Cas6 monomers, the E. coli homologs (PDB accession code 4TVX) were initially docked with Chimera [22] and transformed to poly-alanine models. Substantial rearrangement of the finger region of Cas7 monomers, as well as other secondary structure elements of Cas6, were performed manually in COOT [23] before amino acid substitution of the poly-alanine model. Well-defined bulky side chains of aromatic residues allowed a confident assignment of the register. The crRNA was also well defined in the maps and was traced de novo with COOT. For Cas8 and TniQ in particular, no structural similarity was found in the published structures able to

224 explain our densities. Locally refined maps using soft masks at both ends of the crescent-shaped
 225 complex rendered well-defined maps below 3.5 Å resolution. These maps were used for manual de
 226 novo tracing of a poly-alanine model in COOT that was subsequently mutated to the *V. cholerae*
 227 sequences. Bulky side chains for aromatic residues showed excellent density and were used as
 228 landmarks to adjust the register of the sequence. For refinement, an initial step of real space
 229 refinement against the cryo-EM maps was performed with the phenix real space refinement tool of
 230 the Phenix package [24], with secondary structure restraints activated. A second step of reciprocal
 231 space refinement was performed in Refmac5 [25], with secondary restraints calculated with Prosmart
 232 [26] and LibG [27]. Weight of the geometry term versus the experimental term was adjusted to
 233 avoid overfitting of the model into cryo-EM map, as previously reported³⁰. Model validation was
 234 performed in Molprobit [28].

References

- [1] Ron L Dy, Corinna Richter, George P C Salmond, and Peter C Fineran. Remarkable mechanisms in microbes to resist phage infections. *Annu Rev Virol*, 1(1):307–31, Nov 2014.
- [2] Frank Hille, Hagen Richter, Shi Pey Wong, Majda Bratovič, Sarah Ressel, and Emmanuelle Charpentier. The biology of crispr-cas: Backward and forward. *Cell*, 172(6):1239–1259, 03 2018.
- [3] Shany Doron, Sarah Melamed, Gal Ofir, Azita Leavitt, Anna Lopatina, Mai Keren, Gil Amitai, and Rotem Sorek. Systematic discovery of antiphage defense systems in the microbial pangenome. *Science*, 359(6379), 03 2018.
- [4] Tomas Sinkunas, Giedrius Gasiunas, Saktham P Waghmare, Mark J Dickman, Rodolphe Barrangou, Philippe Horvath, and Virginijus Siksnys. In vitro reconstitution of cascade-mediated crispr immunity in streptococcus thermophilus. *EMBO J*, 32(3):385–94, Feb 2013.
- [5] Sy Redding, Samuel H Sternberg, Myles Marshall, Bryan Gibb, Prashant Bhat, Chantal K Guegler, Blake Wiedenheft, Jennifer A Doudna, and Eric C Greene. Surveillance and processing of foreign dna by the escherichia coli crispr-cas system. *Cell*, 163(4):854–65, Nov 2015.
- [6] Joseph E Peters, Kira S Makarova, Sergey Shmakov, and Eugene V Koonin. Recruitment of crispr-cas systems by tn7-like transposons. *Proc Natl Acad Sci U S A*, 114(35):E7358–E7366, 08 2017.

- [7] Sanne E Klompe, Phuc L H Vo, Tyler S Halpin-Healy, and Samuel H Sternberg. Transposon-encoded crispr-cas systems direct rna-guided dna integration. *Nature*, 571(7764):219–225, Jul 2019.
- [8] Ryan N Jackson, Sarah M Golden, Paul B G van Erp, Joshua Carter, Edze R Westra, Stan J J Brouns, John van der Oost, Thomas C Terwilliger, Randy J Read, and Blake Wiedenheft. Structural biology. crystal structure of the crispr rna-guided surveillance complex from escherichia coli. *Science*, 345(6203):1473–9, Sep 2014.
- [9] Saikat Chowdhury, Joshua Carter, MaryClare F Rollins, Sarah M Golden, Ryan N Jackson, Connor Hoffmann, Lyn’Al Nosaka, Joseph Bondy-Denomy, Karen L Maxwell, Alan R Davidson, Elizabeth R Fischer, Gabriel C Lander, and Blake Wiedenheft. Structure reveals mechanisms of viral suppressors that intercept a crispr rna-guided surveillance complex. *Cell*, 169(1):47–57.e11, 03 2017.
- [10] Tai Wei Guo, Alberto Bartesaghi, Hui Yang, Veronica Falconieri, Prashant Rao, Alan Merk, Edward T Eng, Ashleigh M Raczkowski, Tara Fox, Lesley A Earl, Dinshaw J Patel, and Sriram Subramaniam. Cryo-em structures reveal mechanism and inhibition of dna targeting by a crispr-cas surveillance complex. *Cell*, 171(2):414–426.e12, Oct 2017.
- [11] Sabin Mulepati, Annie Héroux, and Scott Bailey. Structural biology. crystal structure of a crispr rna-guided surveillance complex bound to a ssdna target. *Science*, 345(6203):1479–84, Sep 2014.

- [12] Jasenko Zivanov, Takanori Nakane, Björn O Forsberg, Dari Kimanius, Wim Jh Hagen, Erik Lindahl, and Sjors Hw Scheres. New tools for automated high-resolution cryo-em structure determination in relion-3. *Elife*, 7, 11 2018.
- [13] Jason W Holder and Nancy L Craig. Architecture of the tn7 posttransposition complex: an elaborate nucleoprotein structure. *J Mol Biol*, 401(2):167–81, Aug 2010.
- [14] Liisa Holm and Laura M Laakso. Dali server update. *Nucleic Acids Res*, 44(W1):W351–5, 07 2016.
- [15] L Aravind, Vivek Anantharaman, Santhanam Balaji, M Mohan Babu, and Lakshminarayan M Iyer. The many faces of the helix-turn-helix domain: transcription regulation and beyond. *FEMS Microbiol Rev*, 29(2):231–62, Apr 2005.
- [16] Eugene Krissinel. Stock-based detection of protein oligomeric states in jspisa. *Nucleic Acids Res*, 43(W1):W314–9, Jul 2015.
- [17] Guilhem Faure, Sergey A Shmakov, Winston X Yan, David R Cheng, David A Scott, Joseph E Peters, Kira S Makarova, and Eugene V Koonin. Crispr-cas in mobile genetic elements: counter-defence and beyond. *Nat Rev Microbiol*, 17(8):513–525, Aug 2019.
- [18] Jonathan Strecker, Alim Ladha, Zachary Gardner, Jonathan L Schmid-Burgk, Kira S Makarova, Eugene V Koonin, and Feng Zhang. Rna-guided dna insertion with crispr-associated transposases. *Science*, 365(6448):48–53, 07 2019.

- [19] Kai Zhang. Gctf: Real-time ctf determination and correction. *J Struct Biol*, 193(1):1–12, Jan 2016.
- [20] Ali Punjani, John L Rubinstein, David J Fleet, and Marcus A Brubaker. cryosparc: algorithms for rapid unsupervised cryo-em structure determination. *Nat Methods*, 14(3):290–296, 03 2017.
- [21] Shawn Q Zheng, Eugene Palovcak, Jean-Paul Armache, Kliment A Verba, Yifan Cheng, and David A Agard. Motioncor2: anisotropic correction of beam-induced motion for improved cryo-electron microscopy. *Nat Methods*, 14(4):331–332, 04 2017.
- [22] Eric F Pettersen, Thomas D Goddard, Conrad C Huang, Gregory S Couch, Daniel M Greenblatt, Elaine C Meng, and Thomas E Ferrin. Ucsf chimera—a visualization system for exploratory research and analysis. *J Comput Chem*, 25(13):1605–12, Oct 2004.
- [23] P Emsley, B Lohkamp, W G Scott, and K Cowtan. Features and development of coot. *Acta Crystallogr D Biol Crystallogr*, 66(Pt 4):486–501, Apr 2010.
- [24] Pavel V Afonine, Billy K Poon, Randy J Read, Oleg V Sobolev, Thomas C Terwilliger, Alexandre Urzhumtsev, and Paul D Adams. Real-space refinement in phenix for cryo-em and crystallography. *Acta Crystallogr D Struct Biol*, 74(Pt 6):531–544, 06 2018.
- [25] G N Murshudov, A A Vagin, and E J Dodson. Refinement of macromolecular structures by the maximum-likelihood method. *Acta Crystallogr D Biol Crystallogr*, 53(Pt 3):240–55, May 1997.

- 308 [26] Robert A Nicholls, Marcus Fischer, Stuart McNicholas, and Garib N Murshudov.
309 Conformation-independent structural comparison of macromolecules with prosmart. *Acta Crys-*
310 *tallogr D Biol Crystallogr*, 70(Pt 9):2487–99, Sep 2014.
- 311 [27] Alan Brown, Fei Long, Robert A Nicholls, Jaan Toots, Paul Emsley, and Garib Murshudov.
312 Tools for macromolecular model building and refinement into electron cryo-microscopy recon-
313 structions. *Acta Crystallogr D Biol Crystallogr*, 71(Pt 1):136–53, Jan 2015.
- 314 [28] Christopher J Williams, Jeffrey J Headd, Nigel W Moriarty, Michael G Prisant, Lizbeth L
315 Videau, Lindsay N Deis, Vishal Verma, Daniel A Keedy, Bradley J Hintze, Vincent B Chen,
316 Swati Jain, Steven M Lewis, W Bryan Arendall, 3rd, Jack Snoeyink, Paul D Adams, Simon C
317 Lovell, Jane S Richardson, and David C Richardson. Molprobity: More and better reference
318 data for improved all-atom structure validation. *Protein Sci*, 27(1):293–315, 01 2018.
- 319 [29] Israel S Fernández, Xiao-Chen Bai, Garib Murshudov, Sjors H W Scheres, and V Ramakrish-
320 nan. Initiation of translation by cricket paralysis virus ires requires its translocation in the
321 ribosome. *Cell*, 157(4):823–31, May 2014.
- 322 [30] Alp Kucukelbir, Fred J Sigworth, and Hemant D Tagare. Quantifying the local resolution of
323 cryo-em density maps. *Nat Methods*, 11(1):63–5, Jan 2014.

324 **Data availability.**

325 Maps and models have been deposited in the EMDB (accession codes 20349, 20350 and 20351) and
326 the PDB (accession codes 6PIF, 6PIG and 6PIJ).

327 **Acknowledgements.**

328 We acknowledge Bob Grassucci and Zhening Zhang for technical assistance in cryo-EM data acqui-
 329 sition. Part of this work was performed at the Simons Electron Microscopy Center and National
 330 Resource for Automated Molecular Microscopy located at the New York Structural Biology Center,
 331 supported by grants from the Simons Foundation (SF349247), NYSTAR, and the NIH National
 332 Institute of General Medical Sciences (GM103310).

Figure 1

Fig.1. Overall architecture of the *V. cholerae* TniQ-Cascade complex.

a, Genetic architecture of the Tn6677 transposon (top), and plasmid constructs used to express and purify the TniQ-Cascade co-complex. Selected cryo-EM reference-free 2D classes in multiple orientations are shown on the right. **b**, Orthogonal views of the cryo-EM map for the TniQ-Cascade complex, showing Cas8 (pink), six Cas7 monomers (green), Cas6 (salmon), crRNA (grey), and TniQ monomers (blue, yellow). The complex adopts a helical architecture with protuberances at both ends. **c**, A flexible domain in Cas8 comprising residues 277-385 (grey) could only be visualized in

low-pass filtered maps. The unsharpened map is shown as semi-transparent, grey map overlaid on the post-processed map segmented and colored according to a. **d**, Refined model for the TniQ-Cascade complex derived from the cryo-EM maps shown in **b**.

Figure 2

Fig.2. TniQ binds Cascade in a dimeric, head-to-tail configuration.

a, Left, overall view of the TniQ-Cascade cryo-EM unsharpened map (grey) overlaid on the post-processed map segmented and colored as in **Fig.1**. Right, cryo-EM map (top) and refined model (bottom) of the TniQ dimer. The two monomers interact with each other in a head-to-tail configuration and are anchored to Cascade via Cas6 and Cas7.1. **b**, Secondary structure diagram of the TniQ dimer: eleven α -helices are organized into an N-terminal Helix-Turn-Helix (HTH) domain and a C-terminal TniQ-domain. Dimer interactions between H3 and H11 are indicated, as are interaction sites with Cas6 and Cas7.1. **c**, Cryo-EM density for the H3-H11 interaction shows clear side-chain features (top), allowing accurate modeling of the interaction (bottom). **d**, Schematic of the dimer interaction, showing the important dimerization interface between the HTH and TniQ-domain.

Figure 3

Fig.3. Cas6 and Cas7.1 form a binding platform for TniQ.

a, Top, zoomed area showing the interaction site of Cascade and the TniQ dimer. Cas6 and Cas7.1 are displayed as molecular Van der Waals surfaces, the crRNA is shown as grey spheres, and the TniQ monomers as ribbons. **b**, The loop connecting TniQ.1 α -helices H6 and H7 (blue) binds within a hydrophobic cavity of Cas6. **c**, Cas7.1 interacts via with the HTH domain of the TniQ.2 monomer (yellow), mainly through H2 and the loop connecting H2 and H3. **d-e**, Experimental cryo-EM densities observed for the TniQ-Cas6 (textbfd) and TniQ-Cas7.1 (textbfe) interaction.

Figure 4

Fig.4. DNA-bound structure of the TniQ-Cascade complex.

a, Schematic of crRNA and the portion of the dsDNA substrate that was experimentally observed within the electron density map for DNA-bound TniQ-Cascade. Target Strand (TS), non-target strand (NTS), as well as the PAM and seed regions are indicated. **b**, Selected cryo-EM reference-free 2D classes for DNA-bound TniQ-Cascade; density corresponding to dsDNA could be directly observed protruding from the Cas8 component in the 2D averages (white arrows). **c**, Cryo-EM map

for DNA-bound TniQ-Cascade. The crRNA is in dark grey and the DNA is in red. On the right and bottom, detailed views for the PAM and seed recognition regions of the map, with refined models represented as sticks within the electron density. Cas8 is shown in pink, Cas7 in green, crRNA in grey, and DNA in red. **d**, The *V. cholerae* transposon encodes a TniQ-Cascade co-complex that utilizes the sequence content of the crRNA to bind complementary DNA target sites (left). We propose that the incomplete R-loop observed in our structure (middle) represents an intermediate state that may precede a downstream "locking" step involving proofreading of the RNA-DNA complementarity. TniQ is positioned at the PAM-distal end of the DNA-bound Cascade complex, where it likely interacts with TnsC during downstream steps of RNA-guided DNA insertion.

EXTENDED DATA FIGURES

Extended Data Figure 1

381 **Extended Data Fig. 1. Cryo-EM sample optimization and image pro-** 382 **cessing workflow.**

383 **a**, Representative negatively stained micrograph for 500 nM TniQ-Cascade. **b**, Left, representative
384 cryo-EM image for 2 μ M TniQ-Cascade. A small dataset of 200 images was collected in a Tecnai-F20
385 microscope equipped with a Gatan K2 camera. Right, reference-free 2D class averages for this initial
386 cryo-EM dataset. **c**, Left, representative image from a large dataset collected in a Tecnai Polara
387 microscope equipped with a Gatan K3 detector. Middle, detailed 2D class averages were obtained
388 that were used for initial model generation using the SGD algorithm implemented in Relion3 (right).
389 **d**, Image processing workflow used to identify the two main classes of the TniQ-cascade complex in
390 open and closed conformations. Local refinements with soft masks were used to improve the quality
391 of the map within the terminal protuberances of the complex. These maps were instrumental for
392 *de novo* modeling and initial model refinement.

Extended Data Figure 2

393

Extended Data Fig. 2. Fourier Shell Correlation (FSC) curves, local resolution, and unsharpened filter maps for the TniQ-Cascade complex in closed conformation.

a, Gold-standard FSC curve using half maps; the global resolution estimation is 3.4 Å by the FSC 0.143 criterion. **b**, Cross-validation model-vs-map FSC. Blue curve, FSC between the shacked model refined against half map 1; red curve, FSC against half map 2, not included in the refinement; black curve, FSC between final model against the final map. The overlap observed between the blue and red curves guarantees a non-overfitted model [29]. **c**, Unsharpened map colored according to local resolutions, as reported by RESMAP [30]. **d**, Final model colored according to B-factors calculated by REFMAC. **e**, A flexible Cas8 domain encompassing residues 277-385 contacts the TniQ dimer at the other side of the crescent shape. Applying a Gaussian filter of increasing width to the unsharpened map allows for a better visualization of this flexible region.

Extended Data Figure 3

407 **Extended Data Fig. 3. Superposition of TniQ-Cascade with structurally**
 408 **similar Cascade complexes.**

409 The *V. cholerae* I-F variant TniQ-Cascade complex (left) was superposed with *Pseudomonas aerug-*
 410 *inosa* I-F Cascade11 (also known as Csy complex; middle, PDB ID: 6B45) and *Escherichia coli* I-E
 411 Cascade9 (right, PDB ID: 4TVX). Shown are superpositions of the entire complex (top), the Cas8
 412 and Cas5 subunits with the 5' crRNA handle (middle top), the Cas7 subunit with a fragment of
 413 crRNA (middle bottom), and the Cas6 subunit with the 3' crRNA handle (bottom).

Extended Data Figure 4

414

415 Extended Data Fig. 4. Representative cryo-EM densities for all the
416 components of the TniQ-Cascade complex in closed conformation.

417 **a**, Final refined model of TniQ-Cascade, with Cas8 in purple, Cas7 monomers in green, Cas6 in red,
418 the TniQ monomers in blue and yellow, and the crRNA in grey. **b-h**, Final refined model inserted

419 in the final cryo-EM density for select regions of all the molecular components of the TniQ-Cascade
420 complex. Residues are numbered.

Extended Data Figure 5

Extended Data Fig. 5. Cas8 and Cas6 interaction with the crRNA.

a, Refined model for the TniQ-Cascade shown as ribbons inserted in the semitransparent Van der Waals surface, colored as in Fig 1. **b** and **c**, Zoomed view of Cas8, which interacts with the 5' end of crRNA. **d** and **e**, Zoomed view of Cas6 interacting with crRNA. **f**, Detailed view of the crRNA structure.

425 end of the crRNA. The inset shows electron density for the highlighted region, where the base of
 426 nucleotide C1 is stabilized by stacking interactions with arginine residues R584 and R424. **d**, Cas6
 427 interacts with the 3' end of the crRNA "handle" (nucleotides 45-60). **e**, An arginine-rich α -helix
 428 is deeply inserted within the major groove of the terminal stem-loop. This interaction is mediated
 429 by electrostatic interactions between basic residues of Cas6 and the negatively charged phosphate
 430 backbone of the crRNA. **f**, Cas6 (red) also interacts with Cas7.1 (green), establishing a β -sheet
 431 formed by β -strands contributed from both proteins.

crRNA

Cas8

Cas7.6

Cas7.5

Cas7.4

Cas7.3

Cas7.2

Cas7.1

TniQ2/TniQ1

Cas6

Cas7.6 finger

Cas7.5 finger

Cas7.4 finger

Cas7.3 finger

Cas7.2 finger

Cas7.1 finger

39

433 **Extended Data Fig. 6. Schematic representation of crRNA and target**

434 **DNA recognition by TniQ-Cascade.**

435 **a**, TniQ-Cascade residues that interact with the crRNA are indicated. Approximate location for all
436 protein components of the complex are also shown, as well as the position of each Cas7 "finger".?

437 **b**, TniQ-Cascade residues that interact with crRNA and target DNA, shown as in **a**.

Extended Data Figure 7

438

439 **Extended Data Fig. 7. Fourier Shell Correlation (FSC) curves, local**
 440 **resolution, and local refined maps for the TniQ-Cascade complex in open**
 441 **conformation.**

442 **a**, Gold-standard FSC curve using half maps; the global resolution estimation is 3.5 Å by the FSC
 443 0.143 criterion. **b**, Cross-validation model-vs-map FSC. Blue curve, FSC between shacked model

444 refined against half map 1; red curve, FSC against half map 2, not included in the refinement;
 445 black curve, FSC between final model against the final map. The overlapping between the blue
 446 and red curves guarantees a non-overfitted model. **c**, Unsharpened map colored according to local
 447 resolutions, as reported by RESMAP. Right, slice through the map shown on the left. **d**, Local
 448 refinements with soft masks improved the maps in flexible regions. Shown the region of the map
 449 corresponding to the TniQ dimer. Unsharpened maps colored according to the local resolution
 450 estimations are shown before (left) and after (right) masked refinements. **e**, Final model for the
 451 TniQ dimer region, colored according to the local B-factors calculated by REFMAC.

Extended Data Figure 8

Extended Data Fig. 8. TniQ harbors a HTH domain involved in protein-protein interactions within the TniQ dimer.

A DALI search¹⁵ using the refined TniQ model as probe found significant similarity between the N-terminal domain of TniQ with PDB entries 4r24 (**a**) and 3ucs (**b**) (Z score 4.1/4.1, r.m.s.d. 3.8/5.1). Both proteins contain Helix-Turn-Helix (HTH) domains and HTH domains are often involved in nucleic acid recognition and mediate protein-protein interactions [15]. **c**, The TniQ dimer is stabilized in a head-to-tail configuration by reciprocal interactions mediated by the HTH domain and the TniQ-domains from both monomers.

Extended Data Figure 9

461

462 **Extended Data Fig. 9. Fourier Shell Correlation (FSC) curves, local res-**
463 **olution, and unsharpened filter maps for the DNA-bound TniQ-Cascade**
464 **complex complex.**

465 **a**, Gold-standard FSC curve using half maps; the global resolution estimation is 2.9 Å by the FSC
466 0.143 criterion. **b**, Cross-validation model-vs-map FSC. Blue curve, FSC between the shacked model
467 refined against half map 1; red curve, FSC against half map 2, not included in the refinement; black
468 curve, FSC between final model against the final map. The overlap observed between the blue and
469 red curves guarantees a non-overfitted model. **c**, Left, unsharpened map colored according to local
470 resolutions, as reported by RESMAP. dsDNA is visible at the top right projecting outside of the
471 complex. Right, final model colored according to B-factors calculated by REFMAC.

Extended Data Figure 10

**Extended Data Fig. 10. Superposition of DNA-bound TniQ-Cascade
with structurally similar Cascade complexes.**

The DNA-bound structure of *V. cholerae* I-F variant TniQ-Cascade complex (left) was superposed with DNA-bound structures of *Pseudomonas aeruginosa* I-F Cascade11 (also known as Csy complex; middle, PDB ID: 6B44) and *Escherichia coli* I-E Cascade9 (right, PDB ID: 5H9F). Shown are superpositions of the entire complex (top), the Cas8 and Cas5 subunits with the 5' crRNA handle and double-stranded PAM DNA (middle top), the Cas7 subunit with a fragment of crRNA (middle bottom), and the Cas6 subunit with the 3' crRNA handle (bottom).