Analysis of stochastic timing of intracellular events with gene switching*

Khem Raj Ghusinga¹ and Abhyudai Singh²

Abstract—An important step in execution of several cellular processes is accumulation of a regulatory protein up to a specific threshold level. Since production of a protein is inherently stochastic, the time at which its level crosses a threshold exhibits cell-to-cell variation. A problem of interest is to characterize how the statistics of event timing is affected by various steps of protein expression. Our previous work studied this problem by considering a gene expression model where gene was always active. Here we extend our analysis to a scenario where gene stochastically switches between active and inactive states. We formulate event timing as the first-passage time for a protein's level to cross a threshold and investigate how the rates of gene activation/inactivation affect the distribution and moments of the first-passage time. Our results show that both the time-scale of gene switching with respect to the protein degradation rate as well as the ratio of the gene inactivation to gene activation rates are important parameters in shaping the event-timing distribution.

I. INTRODUCTION

In the ever-changing cellular environment, a cell has to actively sense and respond to both internal and external cues [1], [2]. The environment is represented via transcription factors. A change in the environment leads to localization of appropriate transcription factors which subsequently activate expression of target genes [3]–[5]. The temporal aspect of the target gene activity plays important role in cellular function as cells often couple important decisions or events to accumulation of specific regulatory proteins up to critical thresholds. Many examples of such events appear in context of development [6]–[11], cell-cycle control [12]–[18], cell differentiation [19]–[23], sporulation [24], [25], apoptosis [26]–[28], lysis of infected bacterial cells [29]–[31], etc.

As the protein levels are subject to molecular fluctuations due to inherent noise in gene expression [32]–[42], the timing of an event that triggers at a critical threshold is expected to exhibit cell-to-cell variation. Indeed, recent single-cell experiments have shown considerable cell-to-cell variability in timing of cellular events [29], [43].

Our recent work has characterized the stochasticity in the timing of events originating from the inherent gene expression noise by formulating event timing as a first-passage time problem [44], [45]. The models of gene expression considered therein however only accounted for transcription

(synthesis of mRNA from gene) and translation (synthesis of protein from mRNA) while assuming that the gene was always active. However, single-cell measurements have revealed that genes are not always active. Instead, they switch between active and inactive states [46]–[54].

Here we explore the effects of gene switching on the first-passage time statistics. We model the gene dynamics as a telegraph process that switches between ON (active) and OFF (inactive) states at exponentially distributed times. We specifically examine the effect of the average time period of the pulsing, and the average duty cycle (average fraction of time for which the signal stays in ON state) on the FPT statistics. In addition to gene switching, our model considers production of mRNAs when gene is in the ON state. Regardless of the state of the gene, each mRNA can synthesize proteins before degrading. For analytical tractability, we assume that mRNA half-life is small as compared to that of the protein of our interest. Therefore, we can ignore its dynamics and assume that each transcription event produces protein molecules in bursts that follow geometric distribution [55]–[57].

Our results show that two important parameters that affect the shape of the distribution as well as the moments of event timing are: the time-scale of switching with respect to the protein degradation rate, and the ratio of gene activation rate to gene inactivation rate. The effect of time-scale is more prominent when the gene activation rate is smaller than or comparable to the gene inactivation rate.

The paper is organized as follows. Section II provides detailed description of our model and then discusses the calculations for first-passage time for the model considered herein. Section III explores the effects of gene switching parameters on the first-passage time statistics, and finally Section IV concludes the paper.

II. STOCHASTIC DESCRIPTION OF EVENT TIMING

In this section, we describe a model of gene expression and use it to compute statistics of event timing.

A. Model Formulation

Let $g(t) \in \{0, 1\}$ denote the state of the gene such that $g(t) = 0$ implies that the gene is inactive (OFF) and $g(t) = 1$ represents that the gene is active (ON). The stochastic switching of the gene between these two states is modeled as a telegraph process

$$g(t) = 0 \xrightleftharpoons[k_b]{k_f} g(t) = 1. \quad (1)$$

*This work was supported by NIH through grant 1R01GM124446-01 to AS

¹Khem Raj Ghusinga is with the Department of Biology, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA khem@email.unc.edu

²Abhyudai Singh is with the Department of Electrical and Computer Engineering, University of Delaware, Newark, DE 19716, USA absingh@udel.edu

Once the gene is in the active state, it produces mRNAs with a rate k . In other words, the mRNA synthesis rate is given by $kg(t)$. Each mRNA molecule produces proteins with a rate k_p and it degrades with rate γ_m . The protein degradation rate is denoted by γ_p (Fig. 1). Both mRNA transcripts and the protein molecules degradation events are considered to be single step processes.

For analytical tractability, we assume that mRNA half-life is small as compared to that of the protein of our interest, i.e., $\gamma_m \gg \gamma_p$. Therefore, we can ignore its dynamics and assume that each transcription event produces protein molecules in bursts that follow geometric distribution with mean burst size $b = k_p/\gamma_m$ [55]–[57]. More precisely, let $x(t)$ denote the protein level in a cell at time t then the probabilities of gene switching, protein production, and degradation events taking place in an infinitesimal time interval $(t, t + dt)$ as

$$\mathbb{P}(g(t) = 0|g(t) = 1) = k_b dt, \quad (2a)$$

$$\mathbb{P}(g(t) = 1|g(t) = 0) = k_f dt, \quad (2b)$$

$$\mathbb{P}(x(t) = i + B|x(t) = i) = kg(t)dt, \quad (2c)$$

$$\mathbb{P}(x(t) = i - 1|x(t) = i) = i\gamma dt, \quad (2d)$$

$$\mathbb{P}(B = i) = \frac{b^i}{(b+1)^{i+1}}, b \geq 0, i \in \{0, 1, 2, \dots\}. \quad (2e)$$

Here, $\mathbb{P}$ is notation for probability and B denotes the burst size which follows a geometric distribution with mean b .

Having defined the model, we can now characterize the statistics of the time at which protein level crosses a specific threshold.

B. Event Timing Distribution

We are interested in computing distribution of the first time at which $x(t)$ crosses a threshold X (Fig. 1). The time to this event can be conveniently formulated as first-passage time (FPT) defined as

$$T := \inf\{t \geq 0 : x(t) \geq X | x(0) = 0\}. \quad (3)$$

The pdf of T can be computed by constructing an auxiliary process $x_{aux}(t)$ on the state-space $\{0, 1, \dots, X\}$. The initial conditions as well as the probabilities of occurrences for this process are exactly same as $x(t)$ in (2), except that here X is an absorbing state, i.e., once the process reaches the state X , it remains there forever. In terms of the first-passage time, both processes $x(t)$, and $x_{aux}(t)$ are identical.

The probability that the state X is reached by $x_{aux}(t)$ in the time-interval $(t, t + dt)$ is given by

$$\begin{aligned} \mathbb{P}(T \in (t, t + dt)) = \\ \sum_{i=0}^{X-1} k dt \times \mathbb{P}(B \geq X - i) \times \mathbb{P}(g(t) = 1, x_{aux}(t) = i). \end{aligned} \quad (4)$$

Denoting the pdf of T by $f_T(t)$, using the fact that $\mathbb{P}(B \geq X - i) = (b/b + 1)^{X-i}$, and denoting $P_{1,i}(t) := \mathbb{P}(g(t) = 1, x_{aux}(t) = i)$, we get

$$f_T(t) = \sum_{i=0}^{X-1} k \left(\frac{b}{b+1} \right)^{X-i} P_{1,i}(t). \quad (5)$$

The terms $P_i(t)$ can be determined from the Chemical Master Equation (CME) which can be written as a system of differential equations

$$f_T(t) = \sum_{i=0}^{X-1} k \left(\frac{b}{b+1} \right)^{X-i} P_{1,i}(t). \quad (6)$$

The terms $P_{1,i}(t)$ can be computed from the master equation for the auxiliary process that can be written as

$$\begin{aligned} \frac{dP_{0,i}(t)}{dt} = -k_f P_{0,i}(t) + k_b P_{1,i}(t) \\ + \gamma [(i+1)P_{0,i+1}(t) - iP_{0,i}(t)], \quad 0 \leq i \leq X-2, \end{aligned} \quad (7a)$$

$$\begin{aligned} \frac{dP_{0,X-1}(t)}{dt} = -k_f P_{0,X-1}(t) + k_b P_{1,X-1}(t) \\ - (X-1)\gamma P_{0,X-1}(t), \end{aligned} \quad (7b)$$

$$\begin{aligned} \frac{dP_{1,i}(t)}{dt} = k_f P_{0,i}(t) - k_b P_{1,i}(t) \\ + \gamma [(i+1)P_{1,i+1}(t) - iP_{1,i}(t)] \\ + \sum_{n=0}^{i-1} k \frac{b^{i-n}}{(b+1)^{i-n+1}} P_{1,n}(t) \\ - k \frac{b}{b+1} P_{1,i}(t), \quad 0 \leq i \leq X-2, \end{aligned} \quad (7c)$$

$$\begin{aligned} \frac{dP_{1,X-1}(t)}{dt} = k_f P_{0,X-1}(t) - k_b P_{1,X-1}(t) \\ - (X-1)\gamma P_{1,X-1}(t) \\ + \sum_{n=0}^{X-2} k \frac{b^{X-n}}{(b+1)^{X-n+1}} P_{1,n}(t) \\ - k \frac{b}{b+1} P_{1,X-1}(t), \end{aligned} \quad (7d)$$

where we have used the following notation

$$P_{0,i}(t) = \mathbb{P}(g(t) = 0, x_{aux}(t) = i), \quad (8)$$

$$P_{1,i}(t) = \mathbb{P}(g(t) = 1, x_{aux}(t) = i) \quad (9)$$

[58]–[61]. Stacking the probabilities in a single vector

$$\mathbf{P}(t) = \begin{bmatrix} P_{0,0}(t) \\ \vdots \\ P_{0,X-1}(t) \\ P_{1,0}(t) \\ \vdots \\ P_{1,X-1}(t) \end{bmatrix}. \quad (10)$$

leads to a convenient form of the CME as the following linear time invariant dynamical system

$$\frac{d\mathbf{P}(t)}{dt} = \mathbf{A}\mathbf{P}(t). \quad (11)$$

The matrix $\mathbf{A}$ in the above equation can be written as

$$\mathbf{A} = \begin{bmatrix} \mathbf{A}_{00} & \mathbf{A}_{01} \\ \mathbf{A}_{10} & \mathbf{A}_{11} \end{bmatrix}, \quad (12)$$

Fig. 1. *Event timing as a first-passage time problem.* Left: A sketch of the gene expression model that includes production and degradation of mRNA/protein molecules. The gene dynamics is represented by $g(t)$. Right: An event of interest is triggered when the protein level reaches a critical event threshold for the first time. Each protein trajectory is representative of protein level over time inside individual cells. As a consequence of stochastic expression of the protein, the threshold is attained at different times in different cells. The corresponding time at which the event happens denotes the first-passage time/event time.

where (i, j) -th elements of the $X \times X$ block matrices are given by

$$(\mathbf{A}_{00})_{ij} = \begin{cases} (i-1)\gamma, & j = i+1 \\ -k_f - (i-1)\gamma, & j = i \\ 0, & \text{otherwise.} \end{cases} \quad (13a)$$

$$(\mathbf{A}_{01})_{ij} = \begin{cases} k_b, & j = i \\ 0, & \text{otherwise.} \end{cases} \quad (13b)$$

$$(\mathbf{A}_{10})_{ij} = \begin{cases} k_f, & j = i \\ 0, & \text{otherwise.} \end{cases} \quad (13c)$$

$$(\mathbf{A}_{11})_{ij} = \begin{cases} 0, & j > i+1 \\ (i-1)\gamma, & j = i+1 \\ -k_b - k_{\frac{b}{b+1}} - (i-1)\gamma, & j = i \\ k_{\frac{b^{i-j}}{(b+1)^{i-j+1}}}, & j < i. \end{cases} \quad (13d)$$

where $i, j \in \{1, \dots, X\}$.

Using the solution of the linear time invariant system of equations in (10), the pdf of T can be written as

$$f_T(t) = k\mathbf{U}^\top \mathbf{P}(t), \quad (14)$$

where $\mathbf{U}$ represents the following $2X$ -dimensional vector

$$\mathbf{U} = \begin{bmatrix} 0 & \dots & 0 & \left(\frac{b}{b+1}\right)^X & \dots & \frac{b}{b+1} \end{bmatrix}^\top. \quad (15)$$

The solution $\mathbf{P}(t)$ can be analytically computed as

$$\mathbf{P}(t) = \exp(\mathbf{A}t)\mathbf{P}(0). \quad (16)$$

Here the initial condition is taken as $\mathbf{P}(0) = [0 \dots 0 \ 1 \ 0 \dots 0]^\top$ which corresponds to $g(0) = 1$,

$x_{aux}(0) = x(0) = 0$ with probability one. As a result, the pdf of T takes the form

$$f_T(t) = k\mathbf{U}^\top \exp(\mathbf{A}t)\mathbf{P}(0). \quad (17)$$

We can now use (17) to compute the moments of T . In particular, we study the first two moments in this work.

C. Moments of Event Timing

We can use (17) to compute a m^{th} order moment as

$$\langle T^m \rangle = \int_0^\infty t^m f_T(t) dt = k\mathbf{U}^\top \left(\int_0^\infty \exp(\mathbf{A}t) dt \right) \mathbf{P}(0). \quad (18)$$

Though we do not provide a formal proof here, the matrix $\mathbf{A}$ has the property that its diagonal elements are negative and dominant (i.e., sum of absolute values of all other elements in a column is less than absolute value of the diagonal element). Thus, we can use the result from [62, Thm. 1, pp 48–49] to argue that $\mathbf{A}$ is full-rank and eigenvalues with negative real parts. Therefore, the above integral simplifies to [45]

$$\langle T^m \rangle = (-1)^{m+1} m! k\mathbf{U}^\top (\mathbf{A}^{-1})^{m+1} \mathbf{P}(0). \quad (19)$$

To compute $\mathbf{A}^{-1}$, we now use the standard formulas for inverse of a block diagonal matrix as [63]

$$\mathbf{A}^{-1} = \begin{bmatrix} \mathbf{A}'_{00} & \mathbf{A}'_{01} \\ \mathbf{A}'_{10} & \mathbf{A}'_{11} \end{bmatrix} \quad (20)$$

where

$$\mathbf{A}'_{00} = (\mathbf{A}_{00} - \mathbf{A}_{01}\mathbf{A}_{11}^{-1}\mathbf{A}_{10})^{-1} \quad (21)$$

$$\mathbf{A}'_{01} = (\mathbf{A}_{00} - \mathbf{A}_{01}\mathbf{A}_{11}^{-1}\mathbf{A}_{10})^{-1} \mathbf{A}_{01}\mathbf{A}_{11}^{-1} \quad (22)$$

$$\mathbf{A}'_{10} = -(\mathbf{A}_{11} - \mathbf{A}_{10}\mathbf{A}_{00}^{-1}\mathbf{A}_{01})^{-1} \mathbf{A}_{10}\mathbf{A}_{00}^{-1} \quad (23)$$

$$\mathbf{A}'_{11} = -(\mathbf{A}_{11} - \mathbf{A}_{10}\mathbf{A}_{00}^{-1}\mathbf{A}_{01})^{-1} \quad (24)$$

Typically, there are multiple ways to write the inverse, depending upon which of the block matrices are non-singular. Due to sheer complexity of the expressions, we do not provide explicit form of $\mathbf{A}^{-1}$. At any rate, we can numerically invert these matrices and obtain $\mathbf{A}^{-1}$ to compute the moments.

So far, we have found an expression for the distribution and moments of T in terms of the model parameters. In the next section, we explore the effect of model parameters on these.

III. EFFECT OF MODEL PARAMETERS

We begin by exploring the effect of gene switching parameters on the distribution of T given by (17). To keep the analysis relative to the protein time-scale, we set $\gamma_p = 1$. In Fig. 2, we plot the distribution for various values of k_f and k_b . Since the time-scale of gene turning off is set by $1/k_b$, we look at its values that are orders of magnitude different. Furthermore, the average time for which the gene remains active is given by $1/(1 + k_b/k_f)$. Therefore, we look at various values of k_f that span a couple of orders of magnitude of k_b . We note that when k_b is small as compared to γ , i.e., the switching occurs at a slow rate, then the ratio of k_f to k_b doesn't affect the distribution much (blue curves). On the other hand, if k_b is much higher than γ , then the ratio greatly impacts the distribution shape (magenta curves). In particular, when k_b is higher than k_f , the distribution has a heavy-tail (top, right). Likewise, when k_f is much larger than k_b , then the relative value of k_b with respect to γ doesn't affect the distribution much (bottom row). The value of k_b starts to impact the distribution when k_f is either similar or much smaller than k_b (middle and top rows).

In our previous work [64], we noted that the relative positions of the study-state protein level with respect to the event threshold can affect the noise in event timing in non-trivial ways. In this model, the steady-state protein level is given by

$$x_{ss} = \frac{1}{1 + \frac{k_b}{k_f}} \frac{k \times b}{\gamma}. \quad (25)$$

Indeed when the distribution has a long-tail (as in top row, right), $x_{ss} = 50/11$ which is much less than the event threshold $X = 20$. Therefore, the relative positions of x_{ss} and X seem to matter in this case as well.

We can also use the expression of moments in (19) to look at the effect of moments. To illustrate this, we look at the effect of duty cycle $r = \frac{k_{on}}{k_{on} + k_{off}}$, i.e., fraction of time the gene is in ON state on the moments of event timing. Interestingly, while the mean time expectedly decreases with increase in the duty cycle, the noise (quantified by coefficient of variation squared) shows a non-monotonic behavior (Fig. 3). Note that while the decrease in noise can be explained by the relative positions of the event threshold and x_{ss} , the earlier increase in the noise cannot be explained this way since x_{ss} increases with duty-cycle.

IV. CONCLUSIONS

In this paper, we studied the effect of gene switching on the distribution and statistics of timing of an event that triggers upon attainment of a specific protein level. We showed that both the ratio of switching rates as well as the time-scale of switching as compared to the protein degradation rate affect the statistics. Future work will explore effect of feedback strategies on event timing, wherein the feedback affects the rate of activation or inactivation.

REFERENCES

- [1] E. Kussell and S. Leibler, "Phenotypic diversity, population growth, and information in fluctuating environments," *Science*, vol. 309, pp. 2075–2078, 2005.
- [2] T. J. Perkins and P. S. Swain, "Strategies for cellular decision-making," *Molecular systems biology*, vol. 5, p. 326, 2009.
- [3] M. S. Cyert, "Regulation of nuclear localization during signaling," *Journal of Biological Chemistry*, vol. 276, pp. 20805–20808, 2001.
- [4] F. Estruch, "Stress-controlled transcription factors, stress-induced genes and stress tolerance in budding yeast," *FEMS microbiology reviews*, vol. 24, pp. 469–486, 2000.
- [5] J. M. Kyriakis, "The integration of signaling by multiprotein complexes containing raf kinases," *Biochimica et Biophysica Acta (BBA)-Molecular Cell Research*, vol. 1773, pp. 1238–1247, 2007.
- [6] H. Salz, "Male or female? The answer depends on when you ask," *PLOS Biology*, vol. 5, p. e335, 2007.
- [7] Y. Goldschmidt, E. Yurkovsky, A. Reif, R. Rosner, A. Akiva, and I. Nachman, "Control of relative timing and stoichiometry by a master regulator," *PLOS ONE*, vol. 10, pp. 1–14, 2015.
- [8] H. H. McAdams and A. P. Arkin, "Stochastic mechanisms in gene expression," *Proceedings of the National Academy of Sciences*, vol. 94, pp. 814–819, 1997.
- [9] J. M. Pedraza and J. Paulsson, "Random timing in signaling cascades," *Molecular Systems Biology*, vol. 3, p. 81, 2007.
- [10] I. Nachman, A. Regev, and S. Ramanathan, "Dissecting timing variability in yeast meiosis," *Cell*, vol. 131, pp. 544–556, 2007.
- [11] S. Gupta, J. Varennes, H. C. Korswagen, and A. Mugler, "Temporal precision of regulated gene expression," *PLoS Computational Biology*, vol. 14, p. e1006201, 2018.
- [12] K. C. Chen, L. Calzone, A. Csikasz-Nagy, F. R. Cross, B. Novak, and J. J. Tyson, "Integrative analysis of cell cycle control in budding yeast," *Molecular Biology of the Cell*, vol. 15, pp. 3841–3862, 2004.
- [13] J. M. Bean, E. D. Siggia, and F. R. Cross, "Coherence and timing of cell cycle start examined at single-cell resolution," *Molecular Cell*, vol. 21, pp. 3–14, 2006.
- [14] D. L. Satinover, D. L. Brautigan, and P. T. Stukenberg, "Aurora-A kinase and inhibitor-2 regulate the cyclin threshold for mitotic entry in xenopus early embryonic cell cycles," *Cell Cycle*, vol. 5, pp. 2268–2274, 2006.
- [15] X. Liu, X. Wang, X. Yang, S. Liu, L. Jiang, Y. Qu, L. Hu, Q. Ouyang, and C. Tang, "Reliable cell cycle commitment in budding yeast is ensured by signal integration," *eLife*, vol. 4, p. e03977, 2015.
- [16] K. R. Ghusinga, C. A. Vargas-Garcia, and A. Singh, "A mechanistic stochastic framework for regulating bacterial cell division," *Scientific Reports*, p. 30229, 2016.
- [17] B. Duriez, S. Chilaka, J.-F. Bercher, E. Hercule, and M.-N. Prioleau, "Replication dynamics of individual loci in single living cells reveal changes in the degree of replication stochasticity through S phase," *Nucleic Acids Research*, vol. 47, pp. 5155–5169, 2019.
- [18] D. Antunes and A. Singh, "Quantifying gene expression variability arising from randomness in cell division times," *Journal of Mathematical Biology*, vol. 71, pp. 437–463, 2015.
- [19] D. T. Champlin and J. W. Truman, "Ecdysteroid control of cell proliferation during optic lobe neurogenesis in the moth manduca sexta," *Development*, vol. 125, pp. 269–277, 1998.
- [20] T. Koyama, M. Iwami, and S. Sakurai, "Ecdysteroid control of cell cycle and cellular commitment in insect wing imaginal discs," *Molecular and Cellular Endocrinology*, vol. 213, pp. 155–166, 2004.
- [21] J. H. Levine, M. E. Fontes, J. Dworkin, and M. B. Elowitz, "Pulsed feedback defers cellular differentiation," *PLoS Biol*, vol. 10, p. e1001252, 2012.

Fig. 2. Effect of gene switching parameters on the probability density function of T in (17). We have chosen $\gamma = 1$, so as to normalize all parameters with respect to the protein degradation rate. The values of k_f and k_b for each plot are mentioned therein. Rest of the parameter values are chosen as $k = 50$, and $X = 20$, and $b = 1$.

Fig. 3. Effect of gene switching parameters on the moments of T in (19). We have chosen $\gamma = 1$, so as to normalize all parameters with respect to the protein degradation rate. Rest of the parameter values are chosen as $k = 50$, and $X = 20$, and $b = 1$.

- [22] J. H. Levine and M. B. Elowitz, "Polyphasic feedback enables tunable cellular timers," *Current Biology*, vol. 24, pp. R994–R995, 2014.
- [23] K. H. Aull, E. J. Tanner, M. Thomson, and L. S. Weinberger, "Transient thresholding: A mechanism enabling noncooperative transcriptional circuitry to form a switch," *Biophysical Journal*, vol. 112, no. 11, pp. 2428 – 2438, 2017.
- [24] K. Carniol, P. Eichenberger, and R. Losick, "A threshold mechanism governing activation of the developmental regulatory protein σ^F in *bacillus subtilis*," *Journal of Biological Chemistry*, vol. 279, pp. 14860–14870, 2004.
- [25] P. J. Piggot and D. W. Hilbert, "Sporulation of *bacillus subtilis*," *Current Opinion in Microbiology*, vol. 7, pp. 579–586, 2004.
- [26] S. L. Spencer, S. Gaudet, J. G. Albeck, J. M. Burke, and P. K. Sorger, "Non-genetic origins of cell-to-cell variability in TRAIL-induced apoptosis," *Nature*, vol. 459, pp. 428–432, 2009.
- [27] J. Roux, M. Hafner, S. Bandara, J. J. Sims, H. Hudson, D. Chai, and P. K. Sorger, "Fractional killing arises from cell-to-cell variability in overcoming a caspase activity threshold," *Molecular Systems Biology*, vol. 11, p. 803, 2015.
- [28] M. Krackova, G. Akiri, A. George, R. Sachidanandam, and S. Aaronson, "A threshold mechanism mediates p53 cell fate decision between growth arrest and apoptosis," *Cell Death & Differentiation*, vol. 20, pp. 576–588, 2013.
- [29] J. J. Dennehy and I.-N. Wang, "Factors influencing lysis time stochasticity in bacteriophage λ ," *BMC Microbiology*, vol. 11, p. 174, 2011.
- [30] R. White, S. Chiba, T. Pang, J. S. Dewey, C. G. Savva, A. Holzenburg, K. Pogliano, and R. Young, "Holin triggering in real time," *Proceedings of the National Academy of Sciences*, vol. 108, pp. 798–803, 2011.
- [31] A. Singh and J. J. Dennehy, "Stochastic holin expression can account for lysis time variation in the bacteriophage λ ," *Journal of the Royal Society Interface*, vol. 11, p. 20140140, 2014.
- [32] L. Cai and N. F. X. S. Xie, "Stochastic protein expression in individual cells at the single molecule level," *Nature*, vol. 440, pp. 358–362, Sept. 2006.
- [33] D. K. Wells, W. L. Kath, and A. E. Motter, "Control of stochastic and induced switching in biophysical networks," *Physical Review X*, vol. 5, p. 031036, 2015.
- [34] J. M. Raser and E. K. O'Shea, "Noise in gene expression: Origins, consequences, and control," *Science*, vol. 309, pp. 2010 – 2013, 2005.
- [35] A. Raj and A. van Oudenaarden, "Nature, nurture, or chance: stochastic gene expression and its consequences," *Cell*, vol. 135, pp. 216–226, 2008.
- [36] M. Kærn, T. C. Elston, W. J. Blake, and J. J. Collins, "Stochasticity in gene expression: from theories to phenotypes," *Nature Reviews Genetics*, vol. 6, pp. 451–464, 2005.
- [37] A. M. Corrigan, E. Tunnacliffe, D. Cannon, and J. R. Chubb, "A continuum model of transcriptional bursting," *eLife*, vol. 5, p. e13051, 2016.
- [38] R. D. Dar, B. S. Razoosky, A. Singh, T. V. Trimeloni, J. M. McCollum, C. D. Cox, M. L. Simpson, and L. S. Weinberger, "Transcriptional burst frequency and burst size are equally modulated across the human genome," *Proceedings of the National Academy of Sciences*, vol. 109, pp. 17454–17459, 2012.
- [39] M. Hinczewski and D. Thirumalai, "Cellular signaling networks function as generalized wiener-kolmogorov filters to suppress noise," *Physical Review X*, vol. 4, p. 041017, 2014.
- [40] A. Singh, B. Razoosky, C. D. Cox, M. L. Simpson, and L. S.

- Weinberger, "Transcriptional bursting from the HIV-1 promoter is a significant source of stochastic noise in HIV-1 gene expression," *Biophysical Journal*, vol. 98, pp. L32–L34, 2010.
- [41] A. Raj, C. Peskin, D. Tranchina, D. Vargas, and S. Tyagi, "Stochastic mRNA synthesis in mammalian cells," *PLOS Biology*, vol. 4, p. e309, 2006.
- [42] R. D. Dar, S. M. Shaffer, A. Singh, B. S. Razooky, M. L. Simpson, A. Raj, and L. S. Weinberger, "Transcriptional bursting explains the noise-versus-mean relationship in mRNA and protein levels," *PLOS ONE*, vol. 11, p. e0158298, 2016.
- [43] E. Yurkovsky and I. Nachman, "Event timing at the single-cell level," *Briefings in Functional Genomics*, vol. 12, pp. 90–98, 2013.
- [44] M. C. Lagomarsino, M. Caselle, M. Osella, *et al.*, "Stochastic timing in gene expression for simple regulatory strategies," *Nucleic Acids Research*, p. gkw1235, 2016.
- [45] K. R. Ghusinga, J. J. Dennehy, and A. Singh, "First-passage time approach to controlling noise in the timing of intracellular events," *Proceedings of the National Academy of Sciences*, vol. 114, pp. 693–698, 2017.
- [46] B. Daigle, M. Soltani, L. Petzold, and A. Singh, "Inferring single-cell gene expression mechanisms using stochastic simulation," *Bioinformatics*, vol. 31, pp. 1428–1435, 2015.
- [47] T. Fukaya, B. Lim, and M. Levine, "Enhancer control of transcriptional bursting," *Cell*, vol. 166, pp. 358–368, 2015.
- [48] D. M. Suter, N. Molina, D. Gatfield, K. Schneider, U. Schibler, and F. Naef, "Mammalian genes are transcribed with widely different bursting kinetics," *Science*, vol. 332, pp. 472–474, 2011.
- [49] A. M. Corrigan and J. R. Chubbemail, "Regulation of transcriptional bursting by a naturally oscillating signal," *Current Biology*, vol. 24, pp. 205–211, 2014.
- [50] S. Chong, C. Chen, H. Ge, and X. S. Xie, "Mechanism of transcriptional bursting in bacteria," *Cell*, vol. 158, pp. 314–326, 2014.
- [51] A. Singh, B. S. Razooky, R. D. Dar, and L. S. Weinberger, "Dynamics of protein noise can distinguish between alternate sources of gene-expression variability," *Molecular Systems Biology*, vol. 8, p. 607, 2012.
- [52] I. Golding, J. Paulsson, S. Zawilski, and E. Cox, "Real-time kinetics of gene activity in individual bacteria," *Cell*, vol. 123, pp. 1025–1036, 2005.
- [53] N. Kumar, A. Singh, and R. V. Kulkarni, "Transcriptional bursting in gene expression: analytical results for general stochastic models," *PLOS Computational Biology*, vol. 11, p. e1004292, 2015.
- [54] C. R. Bartman, S. C. Hsu, C. C.-S. Hsiung, A. Raj, and G. A. Blobel, "Enhancer regulation of transcriptional bursting parameters revealed by forced chromatin looping," *Molecular Cell*, vol. 62, pp. 237–247, 2016.
- [55] J. Paulsson, "Model of stochastic gene expression," *Physics of Life Reviews*, vol. 2, pp. 157–175, 2005.
- [56] V. Elgart, T. Jia, A. T. Fenley, and R. Kulkarni, "Connecting protein and mrna burst distributions for stochastic models of gene expression," *Physical Biology*, vol. 8, p. 046001, 2011.
- [57] V. Shahrezaei and P. S. Swain, "Analytical distributions for stochastic gene expression," *Proceedings of the National Academy of Sciences*, vol. 105, pp. 17256–17261, 2008.
- [58] B. Munsky and M. Khammash, "The finite state projection algorithm for the solution of the chemical master equation," *Journal of Chemical Physics*, vol. 124, p. 044104, 2006.
- [59] D. T. Gillespie, "A general method for numerically simulating the stochastic time evolution of coupled chemical reactions," *Journal of Computational Physics*, vol. 22, pp. 403–434, 1976.
- [60] D. A. McQuarrie, "Stochastic approach to chemical kinetics," *Journal of Applied Probability*, vol. 4, pp. 413–478, 1967.
- [61] F. Wilkinson, *Chemical kinetics and reaction mechanisms*. Van Nostrand Reinhold Co, 1980.
- [62] X. Liao, L. Wang, and P. Yu, *Stability of Dynamical Systems*, vol. 5. Elsevier, 2007.
- [63] T.-T. Lu and S.-H. Shiou, "Inverses of 2×2 block matrices," *Computers & Mathematics with Applications*, vol. 43, pp. 119–129, 2002.
- [64] K. R. Ghusinga and A. Singh, "Theoretical predictions on the first-passage time for a gene expression model," in *2015 54th IEEE Conference on Decision and Control (CDC)*, pp. 3864–3869, IEEE, 2015.