Epigenomic and functional characterization of a core DNA methyltransferase in the human pathogen *Clostridium difficile*

Pedro H. Oliveira¹, John W. Ribis², Elizabeth M. Garrett³, Dominika Trzilova³, Alex Kim¹, Ognjen Sekulovic², Edward A. Mead¹, Theodore Pak¹, Shijia Zhu¹, Gintaras Deikus¹, Marie Touchon^{4,5}, Colleen Beckford¹, Nathalie E. Zeitouni¹, Deena Altman^{1,6}, Elizabeth Webster¹, Irina Oussenko¹, Supinda Bunyavanich¹, Aneel K. Aggarwal⁷, Ali Bashir¹, Gopi Patel⁶, Camille Hamula⁶, Shirish Huprikar⁶, Eric E. Schadt¹, Robert Sebra¹, Harm van Bakel¹, Andrew Kasarskis¹, Rita Tamayo³, Aimee Shen^{2*}, Gang Fang^{1*}

¹ Department of Genetics and Genomic Sciences, Institute for Genomics and Multiscale Biology, Mount Sinai School of Medicine, New York, New York, United States of America

² Department of Molecular Biology and Microbiology, Tufts University School of Medicine, Boston, Massachusetts, United States of America

³ Department of Microbiology and Immunology, University of North Carolina at Chapel Hill School of Medicine, Chapel Hill, North Carolina, USA

⁴ Microbial Evolutionary Genomics, Institut Pasteur, 25–28 rue du Docteur Roux, Paris, 75015, France

⁵ CNRS, UMR3525, 25–28 rue du Docteur Roux, Paris, 75015, France

⁶ Department of Medicine, Infectious Disease, Mount Sinai School of Medicine, New York, New York, United States of America

⁷ Department of Pharmacological Sciences and Department of Oncological Sciences, Mount Sinai School of Medicine, New York, New York, United States of America

Email: Aimee Shen (aimee.shen@tufts.edu); Gang Fang (gang.fang@mssm.edu)

Keywords: DNA methylation; SMRT sequencing; sporulation; restriction-modification systems.

1 Abstract

2 *Clostridioides difficile* is a leading cause of health care-associated infections. Although
3 significant progress has been made in the understanding of its genome, the epigenome of *C.*
4 *difficile* and its functional impact has not been systematically explored. Here, we performed
5 the first comprehensive DNA methylome analysis of *C. difficile* using 36 human isolates and
6 observed great epigenomic diversity. We discovered an orphan DNA methyltransferase with
7 a well-defined specificity whose corresponding gene is highly conserved across our dataset
8 and in all ~300 global *C. difficile* genomes examined. Inactivation of the methyltransferase
9 gene negatively impacted sporulation, a key step in *C. difficile* disease transmission,
10 consistently supported by multi-omics data, genetic experiments, and a mouse colonization
11 model. Further experimental and transcriptomic analysis also suggested that epigenetic
12 regulation is associated with cell length, biofilm formation, and host colonization. These
13 findings open up a new epigenetic dimension to characterize medically relevant biological
14 processes in this critical pathogen. This work also provides a set of methods for comparative
15 epigenomics and integrative analysis, which we expect to be broadly applicable to bacterial
16 epigenomics studies.

1 Introduction

2 *Clostridioides* (formerly *Clostridium*) *difficile* is a spore-forming Gram-positive obligate
 3 anaerobe and the leading cause of nosocomial antibiotic-associated disease in the
 4 developed world¹. Clinical symptoms of *C. difficile* infection (CDI) in humans range in
 5 severity from mild self-limiting diarrhea to severe, life-threatening inflammatory conditions,
 6 such as pseudomembranous colitis or toxic megacolon. Since the vegetative form of *C.*
 7 *difficile* cannot survive in the presence of oxygen, the bacterium is transmitted via the
 8 fecal/oral route as a metabolically dormant spore². In the intestinal environment, these
 9 spores subsequently germinate into actively growing, toxin-producing vegetative cells that
 10 are responsible for disease pathology³. CDI progresses in an environment of host microbiota
 11 dysbiosis, which disrupts the colonization resistance typically provided by a diverse intestinal
 12 microbiota⁴. In the last two decades, there has been a dramatic rise in outbreaks with
 13 increased mortality and morbidity due in part to the emergence of epidemic-associated
 14 strains with enhanced growth^{5,6}, toxin production⁷, and antibiotic resistance⁸. *C. difficile* was
 15 responsible for half a million infections in the United States in 2011, with 29,000 individuals
 16 dying within 30 days of the initial diagnosis⁹. Those most at risk are older adults, particularly
 17 those who take antibiotics that perturb the normally protective intestinal microbiota.

18
 19 Despite the significant progress achieved in the understanding of *C. difficile* physiology,
 20 genetics, and genomic evolution^{10,11}, the roles played by epigenetic factors, namely DNA
 21 methylation, have not been systematically studied^{12,13}. In the bacterial kingdom, there are
 22 three major forms of DNA methylation: N6-methyladenine (6mA, the most prevalent form
 23 representing ~80%), N4-methylcytosine (4mC), and 5-methylcytosine (5mC). Although
 24 bacterial DNA methylation is most commonly associated with restriction-modification (R-M)
 25 systems that defend hosts against invading foreign DNA^{14,15}, increasing evidence suggests
 26 that DNA methylation also regulates a number of biological processes such as DNA

1 replication and repair, cell cycle, chromosome segregation and gene expression, among
 2 others¹⁶⁻²¹. Efficient high-resolution mapping of bacterial DNA methylation events has only
 3 recently become possible with the advent of Single Molecule Real-Time sequencing (SMRT-
 4 seq)^{22,23}, which can detect all three types of DNA methylation, albeit at different signal-to-
 5 noise ratios: high for 6mA, medium for 4mC, and low for 5mC. This technique enabled the
 6 characterization of the first bacterial methylomes^{24,25}, and since then, more than 2,100 (as of
 7 06/2019) have been mapped, heralding a new era of “bacterial epigenomics”²⁶.

8
 9 Herein, we mapped and characterized DNA methylomes of 36 human *C. difficile* isolates
 10 using SMRT-seq and comparative epigenomics. We observed great epigenomic diversity
 11 across *C. difficile* isolates, as well as the presence of a highly conserved methyltransferase
 12 (MTase). Inactivation of this MTase resulted in a functional impact on sporulation, a key step
 13 in *C. difficile* transmission, consistently supported by multi-omics data and genetic
 14 experiments. Further experimental and integrative transcriptomic analysis suggested that
 15 epigenetic regulation by DNA methylation also modulates *C. difficile* cell length, host
 16 colonization and biofilm formation. Finally, the epigenomic landscape of *C. difficile* also
 17 allowed us to perform a data-driven joint analysis of multiple defense systems and their
 18 contribution to gene flux. These discoveries are expected to stimulate future investigations
 19 along a new epigenetic dimension to characterize and potentially repress medically relevant
 20 biological processes in this critical pathogen.

Results

Methylome analysis reveals great epigenomic diversity in *C. difficile*

From an ongoing Pathogen Surveillance Project at Mount Sinai Medical Center, 36 *C. difficile* isolates were collected from fecal samples of infected patients (Supplementary Table 1). A total of 15 different MLST sequence types (STs) belonging to clades 1 (human and animal, HA1) and 2 (so-called hypervirulent or epidemic)²⁷ are represented in our dataset (Fig. 1a). Using SMRT-seq with long library size selection, *de novo* genome assembly was achieved at high quality (Supplementary Table 1). Methylation motifs were found using the SMRTportal protocol (Materials and Methods). We found a total of 17 unique high-quality methylation motifs in the 36 genomes (average of 2.6 motifs per genome) (Fig. 1a, Supplementary Table 2a). The large majority of target motifs were of 6mA type, one motif (TAACTG) belonged to the 4mC type, and no confident 5mC motifs were detected (Supplementary Text). Like most bacterial methylomes, >95% of the 6mA and 4mC motif sites were methylated (Fig. 1b, Supplementary Table 2a).

Genomes pertaining to the same ST tend to have more similar sets of methylation motifs relative to those from different STs. Those belonging to ST-2, ST-8, ST-21, and ST-110 showed the highest motif diversities. One 6mA motif, CAAAAA, was present across all genomes, which led us to hypothesize that 6mA methylation events at this motif, and its corresponding MTase, play an important and conserved function in *C. difficile*.

21

A DNA methyltransferase and its target motif are ubiquitous in *C. difficile*

Motivated by the consistent presence of the methylation motif CAAAAA across all the *C. difficile* isolates, we proceeded to examine the encoded MTases. From the 36 genome assemblies, we identified a total of 139 MTase genes (average of 3.9 per genome) (Fig. 1a, Supplementary Table 2b) representing all the four major types²⁸. 38 MTases (27%) belong

1 to Type I R-M systems, and 100 MTases (72%) belong to Type II (Fig. 1c). All but one of the
 2 Type II MTases are solitary, *i.e.*, devoid of a cognate restriction endonuclease (REase) (Fig.
 3 1d). The least abundant MTases (1%) belong to Type III R-M systems (Fig. 1c). A Type IV
 4 (no cognate MTase) McrBC REase gene was also present in all genomes (Supplementary
 5 Table 2b). 28% of MTase genes were located in mobile genetic elements (MGEs) (19% in
 6 prophages and 9% in integrative conjugative/mobile elements), while the large majority was
 7 encoded in other chromosomal regions (Fig. 1e, Supplementary Tables 2b-d). No MTase
 8 genes were found in plasmids. We further found multiple additional defense systems (*e.g.*,
 9 abortive infection systems, CRISPR-Cas, toxin-antitoxin), and performed an integrative
 10 analysis with R-M systems in relation to host defense and gene flux (Supplementary Figs.
 11 1a-e, Supplementary Fig. 2, Supplementary Tables 3a-f, Supplementary Text), such as that
 12 involving phages (Supplementary Fig. 3, Supplementary Table 3g, Supplementary Text).

13
 14 Consistent with the presence of a highly conserved CAAAAA motif, we identified a Type II
 15 6mA solitary DNA MTase (577 aa) present across isolates (Fig. 1f, Supplementary Table
 16 2b). This MTase is encoded by *CD2758* in *C. difficile* 630, a reference strain that was
 17 isolated from a *C. difficile* outbreak in Switzerland^{10,29}. Here we have named CD2758 as
 18 CamA (**C**. **d***ifficile* **a**denine **m**ethyltransferase **A**). The ubiquity of this MTase was not
 19 restricted to the 36 isolates, as we were able to retrieve orthologs in a list of ~300 global *C.*
 20 *difficile* isolates from GenBank (Supplementary Table 4). REBASE also showed functional
 21 orthologs of *camA* only in very few other *Clostridiales* and *Fusobacteriales* (Supplementary
 22 Figs. 4a, b), suggesting that this MTase is fairly unique to *C. difficile*. The genomic context of
 23 *camA* is largely conserved across strains (Supplementary Fig. 4c), located ~25 kb upstream
 24 of the S-layer biogenesis locus (Supplementary Fig. 4d). Several of the genes flanking *camA*
 25 (including itself) are part of the *C. difficile* core-genome (see below), suggesting that they
 26 may play biological roles fundamental to *C. difficile*. Hence, CamA is a solitary Type II 6mA
 27 MTase that specifically recognizes the CAAAAA motif and is ubiquitous and fairly unique to
 28 *C. difficile*.

1

2 **Inactivation of *camA* reduces sporulation levels *in vitro***

3 To discover possible functional roles of CamA, we searched among published transposon
 4 sequencing studies of *C. difficile*. From a recent study analyzing *C. difficile* gene essentiality
 5 and sporulation, we found a homolog of *camA* (*R20291_2646*) among ~800 genes whose
 6 mutation impacted sporulation in the 027 isolate R20291³⁰. Given the critical role of
 7 sporulation in the persistence and dissemination of *C. difficile* in humans and hospital
 8 settings³¹, we decided to test if *camA* inactivation could reduce spore purification efficiencies
 9 in the 630 Δ *erm* strain background as seen for *R20291_2646* transposon mutants. We used
 10 allele-coupled exchange³² to construct an in-frame deletion in 630 Δ *erm camA* and to
 11 complement Δ *camA* with either wild-type *camA* or a variant encoding a catalytic site
 12 mutation (N165A) of the MTase in single copy from the *pyrE* locus (Supplementary Fig. 5a;
 13 Supplementary Tables 5a, b; Supplementary Materials and Methods; Supplementary Text).
 14 We observed that spore purification efficiencies decreased by ~50% in the mutant relative to
 15 wild type (Fig. 2a, $P < 10^{-2}$, ANOVA, Tukey's test). Complementation of Δ *camA* with the wild
 16 type, but not the catalytic mutant construct, restored spore purification efficiencies to values
 17 similar to those observed in wild-type cells (Fig. 2a, Supplementary Table 5c). No
 18 differences in growth were observed between wild-type and mutant strains (Supplementary
 19 Fig. 5b). Hence, this complementation experiment supports that the loss of methylation
 20 events by CamA, rather than the loss of non-catalytic roles of this protein, leads to the
 21 decrease in spore yield.

22

23 The diminished spore purification efficiencies observed in the Δ *camA* mutants could be due
 24 to reduced number of cells that induce sporulation or defects in spore assembly³³. Visual
 25 inspection of samples before and after spore purification on a density gradient revealed
 26 qualitatively lower levels of mature, phase-bright spores (Supplementary Fig. 5c). Since
 27 purified wild-type and Δ *camA* spores had similar levels of chloroform resistance
 28 (Supplementary Fig. 5d), and wild-type and Δ *camA* spores germinated with similar efficiency

when plated on media containing germinant (Supplementary Fig. 5e), the reduced spore purification efficiencies of the MTase mutants likely reflect a defect in sporulation initiation rather than the sporulation process itself.

Consistent with this hypothesis, fewer $\Delta camA$ cells were qualitatively observed to be sporulating in phase-contrast microscopy analyses relative to wild type (Fig. 2b). To determine whether loss of CamA impairs the frequency rather than the progression of sporulation, we visualized and quantified the number of sporulating cells at different stages of spore assembly using the FM4-64 membrane stain and Hoechst nucleoid stain³⁴ (Fig. 2c).

While similar numbers of wild-type and $\Delta camA$ cells were observed at asymmetric division (the first morphological stage of sporulation), at the 9 h time point, 50% fewer $\Delta camA$ cells had initiated engulfment. Furthermore, 11 h after sporulation induction, ~2-fold more $\Delta camA$ cells were at the asymmetric division stage relative to wild type, whereas 50% fewer $\Delta camA$ cells had completed engulfment. Since similar numbers of sporulating cells were observed between wild type and $\Delta camA$ at the 11 h timepoint, $\Delta camA$'s sporulation defect appears to arise from fewer cells progressing beyond asymmetric division rather than a defect in sporulation induction.

To confirm that loss of CamA leads to a decrease in the number of cells producing functional spores, we compared the ability of $\Delta camA$ to form heat-resistant spores capable of germinating and outgrowing using a heat resistance assay³⁵. The $\Delta camA$ mutant and the catalytic mutant complementation strain produced ~50% fewer heat-resistant spores than wild-type and the wild-type complementation strain (Supplementary Fig. 5e, $P < 10^{-3}$, ANOVA, Tukey's test). Taken together, these findings suggest that CAAAAA methylation enhances sporulation *in vitro*. This functional difference prompted us to perform a comprehensive methylome and transcriptome analysis of the wild-type and MTase mutant strains.

1 **Comparative analysis of CAAAAA sites across *C. difficile* genomes**

2 The *C. difficile* genome has an average of 7,721 (± 197 , sd) CAAAAA motif sites
 3 (Supplementary Table 6a). Adjusted by the k-mer frequency of the AT-rich *C. difficile*
 4 genome (70.9%) using Markov models³⁶ (Materials and Methods), CAAAAA motif sites are
 5 significantly under-represented in the chromosome, particularly in intragenic regions
 6 (Supplementary Fig. 6a, Supplementary Table 6b). To evaluate if specific chromosomal
 7 regions are enriched or depleted for this motif, we used a multi-scale signal representation
 8 (MSR) approach³⁷ (Materials and Methods). Briefly, MSR uses wavelet transformation to
 9 examine the chromosome at a succession of increasing length scales by testing for
 10 enrichment or depletion of a given genomic signal. While scale values <10 are typically
 11 associated with regions <100 bp, genomic regions enriched for CAAAAA sites at scale
 12 values >20 corresponding to segments larger than 1 kb (*i.e.* gene and operon scale) were
 13 observed and included genes related to sporulation and colonization (Fig. 3a, regions A-E;
 14 Supplementary Tables 6c, d): stage 0 (*spo0A*), stage III (*spoIIIAA-AH*), and stage IV
 15 (*spoIVB*, *sigK*) of sporulation, membrane transport (PTS and ABC-type transport systems),
 16 transcriptional regulation (*e.g.*, *iscR*, *fur*), and multiple cell wall proteins (CWPs).

17
 18 To further characterize CAAAAA motif sites, we built upon our large collection of *C. difficile*
 19 methylomes by categorizing these motif sites on the basis of their positional conservation
 20 across genomes. We performed whole genome alignment of 37 *C. difficile* genomes (36
 21 isolates + *C. difficile* 630 as reference) (Materials and Methods, Supplementary Dataset 1),
 22 and classified each motif position in the alignment as either: (1) conserved orthologous
 23 (devoid of SNPs or indels); (2) variable orthologous (in which at least one genome contains
 24 a SNP or indel); and (3) non-orthologous (Fig. 3b). We found a total of 5,828 conserved
 25 orthologous motif positions, 1,050 variable orthologous positions (885 with SNPs and 165
 26 with indels), and an average of 843 non-orthologous positions per genome (Supplementary
 27 Table 6e). The latter were, as expected, largely mapped to MGEs. Among orthologous

positions, the variable ones represent a particularly interesting subset to study, since they contribute to variations of CAAAAA sites across genomes with subsequent methylation abrogation (Supplementary Table 6f). We used DAVID gene enrichment analysis and found cytoplasm- (e.g.: *pheA*, *fdhD*, *ogt1*, *spoIVA*) and motility-related genes (e.g.: *fliZ*, *fliN*, *fliM*, *flgL*) to over-represent orthologous variable CAAAAA positions (FDR < 5%, Fig. 3c). Concomitantly, the regions with the highest density of orthologous variable positions were found at the S-layer locus region (region D, Fig. 3a) and between positions 0.31-0.33 Mb, which is rich in flagellar genes. The very large number and dispersion of conserved orthologous positions precluded a similar functional analysis. To test whether homologous recombination (HR) contributes to the cross-genome variation of CAAAAA motif sites located in the core-genome, we also performed a systematic analysis of such events (Supplementary Figs. 6b-d, Supplementary Table 6g, Supplementary Text) and found that HR tracts indeed over-represent (O/E=1.40, $P < 10^{-3}$; Chi-square test) orthologous variable CAAAAA motif positions, while the core-genome without HR tracts underrepresents them (O/E=0.89, $P < 10^{-3}$; Chi-square test) (Supplementary Figs. 6e, f).

Collectively, genome-wide distribution analyses and across-genome comparative analyses suggest that CAAAAA sites are enriched in regions harboring genes related to sporulation and colonization, orthologous variable CAAAAA positions are enriched in regions harboring genes related to cytoplasm- and motility-related genes, and the former's variability is at least partially fueled by HR.

Non-methylated CAAAAA motif sites are enriched in regulatory elements

DNA methylation is highly motif driven in bacteria; i.e. in most cases, >95% of the occurrence of a methylation motif is methylated. However, a small fraction of methylation motif sites can be non-methylated. The on/off switch of DNA methylation in a bacterial cell can contribute to epigenetic regulation as a result of competitive binding between DNA MTases and other DNA binding proteins (e.g. transcription factors, TFs) as previously

described for *E. coli*^{21,38-40}. Previous bacterial methylome studies analyzing one or few genomes usually have insufficient statistical power to perform a systematic interrogation of non-methylated motifs sites. Building on our rich collection of 36 *C. difficile* methylomes, we performed a systematic detection and analysis of non-methylated CAAAAA sites. To classify a CAAAAA site as non-methylated, we adopted stringent filtering criteria at interpulse duration ratio (ipdR) and sequencing coverage, and found an average of 21.5 non-methylated CAAAAA motif sites per genome (Supplementary Fig. 7a; Supplementary Table 7a). Non-methylated motif sites were found dispersed throughout the full length of the *C. difficile* genome, yet were overrepresented in orthologous variable and non-orthologous CAAAAA positions (O/E=respectively 1.51 and 1.49) and underrepresented in orthologous conserved CAAAAA positions (O/E 0.84) ($P < 10^{-4}$; Chi-square test). This is consistent with the idea that variable positions are more likely to be non-methylated to provide breadth of expression variation. Most of the non-methylated positions (85.4% of 245) failed to conserve such status in more than three genomes at orthologous positions (Fig. 4a, Supplementary Table 7a), while a minor percentage of positions (5.5%) remained non-methylated in at least one third of the isolates, suggesting that competitive protein binding is expected to be more active in certain genomic regions (Fig. 4a), e.g. upstream of the pathogenicity locus (PaLoc) (position 786,216), in prophage genes (1,593,616), within the *atp* operon (3,430,190), and in the xylose operon (3,561,672) (Supplementary Fig. 7b).

The non-methylated CAAAAA positions detected across the 36 *C. difficile* genomes allowed a systematic search for evidence of overlap with TF binding sites (TFBSs) and transcription start sites (TSSs). To test this, we queried our genomes for putative binding sites of 21 TFs pertaining to 14 distinct families (Supplementary Table 7b; Materials and Methods) and found overlaps between prominent peaks of non-methylated CAAAAA positions (Fig. 4a) and the TFBSs of CodY and XylR (Fig. 4b, Supplementary Fig. 7b, Supplementary Table 7c). Performing the analysis at the genome level, both CodY and XylR binding sites showed significant enrichment ($P < 10^{-3}$, Mann-Whitney-Wilcoxon test) for non-methylated CAAAAA

(Fig. 4c; Supplementary Fig. 7c). In a similar enrichment analysis using 2,015 TSSs reconstructed from RNA-seq data coverage⁴¹ (Materials and Methods; Supplementary Table 7d), we found a genome-level enrichment: non-methylated CAAAAA sites preferentially overlapped with TSSs (Figs. 4d, e; Supplementary Figs. 7d, e; $P < 10^{-3}$, Mann-Whitney-Wilcoxon test,).

In addition to the on/off epigenetic switch driven by competitive binding between the MTase and other DNA binding proteins at CAAAAA sites, we hypothesized that epigenetic heterogeneity within a clonal population could also stem from DNA replication errors at CAAAAA sites, especially because homopolymer tracts are expected to be more error-prone⁴². To test this hypothesis, we investigated if CAAAAA had a larger than expected frequency of mutated reads (compared to baseline sequencing errors). Specifically, we tested if CAAAAA sites in *C. difficile* are more error prone during DNA replication, compared to the control motifs TAAAAA or GAAAAA (here called KAAAAA). To account for the confounding effect of mutation rate at different sequence contexts, we further computed the % mutated reads (SNPs + indels) in four distinct species with a broad dispersion of mutation rates (*Mycobacterium tuberculosis* < *E. coli* $\approx$ *C. difficile* < *Helicobacter pylori*). Interestingly, we indeed found a significantly higher % of mutated reads mapping to CAAAAA sites compared to KAAAAA sites in *C. difficile* (Fig. 4f, Supplementary Fig. 7f), while this difference was not observed in any of the control genomes. The observation of increased mutagenesis at 6mA target sites is in line with recent data obtained in *Neisseria meningitidis*⁴³ and is worth further investigation in future work.

Collectively, these results highlight two types of variations that affect CAAAAA methylation status in *C. difficile*: (i) on/off epigenetic switch of CAAAAA sites preferentially overlapping with putative TFBSs and TSSs, and (ii) higher mutation rates at the CAAAAA motif sites that contribute to cell-to-cell heterogeneity.

1 Loss of CAAAAA methylation impacts transcription of multiple gene 2 categories including sporulation

3 To study the functional significance of methylation at CAAAAA sites, we used RNA-seq to
4 compare the transcriptomes of wild-type *C. difficile* 630 Δ *erm* with that of Δ *camA* both in
5 liquid medium (exponential and stationary growth stage) and following sporulation induction
6 (9 and 10.5 h) (Supplementary Fig. 8, Supplementary Table 8a, Materials and Methods). Of
7 the 3,896 genes annotated in *C. difficile* 630, 36 – 361 (0.9 – 9.3%, depending on the time
8 point) were differentially expressed (DE) at a 5% FDR and $|\log_2FC| > 1$ (2-fold change in
9 gene expression) (Fig. 5a, Supplementary Tables 8b-d). DE genes in Δ *camA* relative to wild
10 type appeared to have a significant enrichment in CAAAAA motif sites compared to non-DE
11 genes ($P < 10^{-2}$, Mann-Whitney-Wilcoxon test) in broth culture; a qualitatively similar trend
12 was also observed during sporulation (Supplementary Fig. 9a). Consistent with our finding
13 that loss of CamA reduces spore formation, the transcriptome analyses revealed that 118
14 and 120 genes previously identified as being induced during sporulation^{44,45} were expressed
15 at $\geq 50\%$ lower levels in Δ *camA* cells relative to wild type at the 9 and 10.5 h timepoints,
16 respectively (Supplementary Table 8b).

17
18 The transcriptional program that mediates sporulation in *C. difficile* is controlled by a master
19 transcriptional activator, Spo0A, and four sporulation-specific sigma factors, σ^F , σ^E , σ^G and
20 σ^K . These factors activate distinct regulons that ultimately lead to the assembly of functional
21 spores^{37,38} (Fig. 5b). Activated Spo0A induces the expression of genes encoding σ^F , σ^E , and
22 σ^G as well as factors required for asymmetric division and the post-translational activation of
23 the early-stage sporulation sigma factors, σ^F and σ^E (Fig. 5b). σ^F is the first sporulation-
24 specific sigma factor to be fully activated, and it only becomes active in the forespore after
25 asymmetric division is completed⁴⁶. Activated σ^F subsequently induces the transcription of
26 genes whose products mediate σ^G activation in the forespore and partially mediates σ^E
27 activation in the mother cell⁴⁷. Activated σ^E induces the transcription of *sigK*⁴⁸ and factors

1 required for the excision of a prophage-like element from the *sigK* gene⁴⁹. Thus, *C. difficile*
 2 sporulation is controlled by a transcriptional hierarchy that is coupled to morphological
 3 events such that downstream sigma factors (σ^G and σ^K) depend on the activation of
 4 upstream sigma factors (σ^F and σ^E). Notably, genes in the regulons of all four sporulation-
 5 specific sigma factors were under-expressed in $\Delta camA$ relative to wild type, whereas a
 6 relatively small subset of Spo0A regulon genes exhibited this pattern of regulation
 7 (Supplementary Fig. 9b, Supplementary Table 8e). These observations suggested that loss
 8 of CamA reduces spore formation by targeting events early on during sporulation.

9
 10 To gain further insight into the regulatory stage of sporulation that CamA-mediated DNA
 11 methylation specifically impacts, we used qRT-PCR to analyze the expression of genes
 12 encoding Spo0A, the sporulation-specific sigma factors^{44,50}, and their individual regulons⁵¹⁻⁵³.
 13 These analyses were conducted on a separate set of RNA samples harvested at 9 and 11 h
 14 following sporulation induction. Consistent with our RNA-Seq analyses, Spo0A regulon
 15 genes, *spo0A*, *sigF*, and *sigE*^{44,51}, were expressed at similar levels between wild type and
 16 $\Delta camA$ at both 9 and 11 h, implying that the $\Delta camA$ mutant activates Spo0A at levels similar
 17 to wild type. In contrast, σ^F and σ^E regulon genes, *spoIIQ* and *spoIVA*^{44,54}, respectively, were
 18 under-expressed in $\Delta camA$ relative to wild type (Fig. 5b). Reduced SpoIIQ and SpoIVA
 19 levels were observed in $\Delta camA$ by western blot, confirming the transcriptional analyses
 20 (Supplementary Fig. 9c). Based on the hierarchical organization of the sporulation regulatory
 21 cascade, these results implicate σ^F activation as the earliest sporulation stage affected by
 22 CamA. This conclusion is supported by our morphological analyses, since fewer $\Delta camA$
 23 cells initiate and complete engulfment relative to wild type, and engulfment requires both σ^F -
 24 and σ^E -dependent gene products⁵⁵. Notably, similar numbers of $\Delta camA$ and wild-type cells
 25 are sporulating based on these morphological analyses (Fig. 2c), consistent with the similar
 26 levels of *spo0A* observed in WT and $\Delta camA$ (Fig. 5b). While a small subset of Spo0A
 27 regulon genes are under-expressed in $\Delta camA$ cells, at least some of these genes are dually

1 regulated by Spo0A and σ^F . For example, transcription of *spoIIIR*⁵², which encodes a
2 signaling protein required for σ^E activation, is activated by both Spo0A and σ^F ^{44,47}.

3

4 *In vivo* impacts of the *camA* mutation

5 To test whether the sporulation defect of $\Delta camA$ impacts *C. difficile* infection or transmission,
6 we analyzed the effect of the $\Delta camA$ mutation in an established mouse model of infection in
7 which a cocktail of antibiotics is used to sensitize the animals to *C. difficile* infection⁵⁶. When
8 inoculated with 630 Δerm strains, antibiotic-treated mice typically do not develop fulminant
9 disease and instead serve as a model of intestinal colonization and persistence by *C.*
10 *difficile*^{57,58}. Groups of mice (6 males, 6 females) were inoculated by oral gavage with
11 10^5 spores of the three genotypes: wild type, $\Delta camA$, and $\Delta camA$ -C. No mortality was
12 observed at the given doses of *C. difficile* spores as expected. Fecal samples were collected
13 every 24 h for seven days. All three *C. difficile* strains reached comparable levels in feces at
14 days 1 and 2 post-inoculation, indicating that they germinate and establish colonization
15 equally (Fig. 6a, Supplementary Materials and Methods). As expected, CFU levels
16 decreased steadily from day 2 post-inoculation to day 7. However, the $\Delta camA$ mutant
17 showed CFU levels 10-100 times lower than those observed in the wild-type and
18 complemented strains throughout this time frame (ANOVA, $P < 10^{-4}$). The bacteria declined
19 below the limit of detection in the feces 6 days post-inoculation for the MTase mutant, while
20 they remained detectable at days 6 and 7 for the wild-type and complemented strains.

21

22 To test whether loss of CamA leads to defects in virulence, we compared $\Delta camA$ and wild
23 type in a hamster model of infection. Clindamycin-treated golden Syrian hamsters are highly
24 susceptible to the effects of the *C. difficile* toxins and thus represent a model of acute
25 disease. Groups of 6 hamsters (3 male, 3 female) were inoculated by oral gavage with
26 10^3 spores of the wild-type, $\Delta camA$, and $\Delta camA$ -C strains. These *C. difficile* strains elicited
27 diarrheal symptoms and weight loss in this model, and we observed no difference in animal

1 survival times post inoculation (Fig. 6b, Supplementary Materials and Methods). This result
2 is consistent with the observation that the wild-type, $\Delta camA$, and $\Delta camA-C$ strains exhibit no
3 differences in toxin gene expression (Supplementary Table 8a) and produce comparable
4 levels of TcdA *in vitro* (Supplementary Fig. 9d). Together, these data indicate that CAAAAA
5 methylation by CamA does not influence toxin-mediated aspects of *C. difficile* pathogenesis
6 but instead impacts *C. difficile*'s ability to persist within the host intestinal tract.

8 Additional functional impacts of the *camA* mutation

9 Considering the high conservation of *camA* across *C. difficile* genomes, we asked if some
10 additional phenotypes could be impacted by the gene's inactivation. While analyzing images
11 of sporulating *C. difficile*, we noticed that $\Delta camA$ mutant cells appeared to be shorter on
12 average than wild-type cells. To test this possibility, we measured the lengths of wild-type
13 and $\Delta camA$ cells during log-phase and stationary-phase growth in broth culture and during
14 sporulation (9 and 11 h time points). These analyses revealed that $\Delta camA$ cells were ~15%
15 shorter than wild-type cells ($P < 0.001$, one-way ANOVA, Fig. 6c, d) even though no
16 difference in cell growth was observed (Supplementary Fig. 5b). Interestingly, genes
17 encoding putative cell wall remodeling enzymes, *spoVD*, *cwp16*, and *cwp17*, were over-
18 expressed in the $\Delta camA$ mutant relative to wild type during growth in broth culture
19 (Supplementary Fig. 9e).

21 In addition to the above experimental characterization, we attempted to build on the RNA-
22 seq data further beyond the sporulation phenotype, with a special focus on biological
23 processes critical to *C. difficile* infection. Specifically, we performed an overlap analysis
24 between the list of DE genes from our RNA-seq data (wild type vs. $\Delta camA$ mutant; four
25 different time points) and those from published studies focusing on the colonization and
26 infection by this pathogen (Materials and Methods). Out of the total 20 pairwise overlap
27 analyses, we observed 9 significant overlaps ($P < 0.01$ with Bonferroni correction) between
28 our dataset and those of others (Supplementary Table 8f). First, DE genes in the $\Delta camA$

mutant (sporulation phases) had a significant overlap to DE genes in conditions favoring the production of biofilm on a solid substrate⁵⁹ (Fig. 6e, Supplementary Table 8f) (O/E=1.3-1.6, $P < 10^{-9}$, Chi-Square test). Motivated by this significant overlap, we performed biofilm assays using crystal violet staining of adherent biofilm biomass, and made consistent observation that the $\Delta camA$ mutant produced more biofilm than wild type (Fig. 6f). These results suggest that methylation inhibits the expression of genes that promote biofilm formation, e.g., encoding factors involved in the adherence to a surface and/or stabilizing interbacterial interactions. The differences in biofilm production between $\Delta camA$ and $\Delta camA/N165A$ could be explained if the latter retained some DNA binding ability capable of altering transcription of some genes even in the absence of methylation. Second, by comparing DE genes in $\Delta camA$ relative to wild type with genes DE during infection in different murine gut microbiome compositions⁶⁰, we found a statistically significant overlap, particularly genes expressed at stationary stage (O/E=1.4, $P < 10^{-6}$, Chi-Square test) (Supplementary Fig. 10a, Supplementary Table 8f). Lastly, using DE genes obtained from murine gut isolates at increasing time points after infection⁶¹, we found significant overlaps with DE genes in the $\Delta camA$ mutant (O/E=1.5, $P < 10^{-4}$, Chi-Square test) (Supplementary Fig. 10b, Supplementary Table 8f). Collectively, these integrative overlap analyses provide additional evidence that DNA methylation events by CamA may directly and/or indirectly affect the expression of multiple genes involved in the *in vivo* colonization and biofilm formation of *C. difficile* and inspire future work to elucidate the mechanisms underlying the functional roles of CAAAAA methylation in *C. difficile* pathogenicity.

Discussion

C. difficile is responsible for one of the most common hospital-acquired infections and classified by the US Centers for Disease Control and Prevention as an urgent healthcare risk with significant morbidity and mortality⁹. Because CDI is spread by bacterial spores found within feces, extensive research has been devoted to better understand the genome

1 of this critical pathogen and its sporulation machinery. To address these common goals, we
 2 performed the first comprehensive characterization of the DNA methylation landscape
 3 across a diverse collection of clinical isolates. During our epigenome analysis, we identified
 4 an 6mA MTase (*camA*) conserved across all isolates (and in another ~300 published *C.*
 5 *difficile* genomes) sharing a common methylation motif (CAAAAA). Inactivation of the gene
 6 encoding this MTase resulted in a sporulation defect *in vitro* (Fig. 2). This represents, to the
 7 best of our knowledge, the first time that DNA methylation has been found to impact
 8 sporulation in any bacterium, opening a new dimension of study in *C. difficile*. Infection
 9 studies using the mouse model indicate a role for CamA in the persistence of *C. difficile* in
 10 the intestinal tract. Since enumeration of *C. difficile* recovered in feces of the infected
 11 animals reflects the number of *C. difficile* spores in the gut, the reduced burden of $\Delta camA$ in
 12 the mouse may be due to the mutant's defect in sporulation, as the ability to form spores
 13 was previously shown to be important for persistence². That $\Delta camA$ exhibited virulence
 14 comparable to the wild type in the hamster model suggests that DNA methylation does not
 15 impact toxin-mediated disease. However, due to the pleiotropic nature of the MTase it
 16 remains possible that multiple factors contribute to the more pronounced effect observed in
 17 the mouse model.

18
 19 The highly conserved *camA* and its flanking genes across *C. difficile* genomes suggest that
 20 additional phenotypes may be regulated by CamA beyond sporulation. Consistently,
 21 CAAAAA sites were overrepresented in a set of regions enriched in genes with functions
 22 linked to sporulation, motility, and membrane transport. Further supporting a broader
 23 regulatory network of CamA, is that its loss reduces cell length and results in statistically
 24 significant overlap between transcriptional signatures identified in our study (wild type vs
 25 $\Delta camA$ mutant) and those of others observed during the *in vivo* colonization and biofilm
 26 formation (Fig. 5b, Supplementary Figs. 7b, c).

27

1 The fact that *camA* is a solitary MTase gene without a cognate restriction gene further
 2 supports a view that widespread methylation in bacteria has a functional importance beyond
 3 that attributed to R-M systems. Previously, the most extensively characterized 6mA MTase
 4 was Dam targeting GATC in *E. coli*. Dam plays multiple important functions and is essential
 5 in some pathogens²¹. However, since it is conserved in the large diversity of γ-
 6 proteobacteria, it was not considered a promising drug target. In contrast, the uniqueness of
 7 *camA* in all *C. difficile* genomes and in just a few *Clostridiales* makes it a promising drug
 8 target that may inhibit *C. difficile* in a much more specific manner, which is particularly
 9 relevant since gut dysbiosis potentiates *C. difficile* infection^{62,63}. In addition, since this MTase
 10 seemingly does not impact the general fitness of *C. difficile*³⁰, a drug specifically targeting it
 11 may be developed with a lower chance for resistance.

12
 13 Considering the large number of genes differentially expressed in the Δ*camA* mutant, the
 14 functional impact of CAAAAA methylation is likely mediated by multiple genes that are either
 15 directly regulated by DNA methylation or indirectly regulated by a transcriptional cascade.
 16 Mechanistically, DNA methylation can either activate or repress a gene depending on other
 17 DNA binding proteins that compete with DNA MTases^{16,17,21,64}, so the competition between
 18 transcription factors and MTases may form an epigenetic switch to turn on/off a gene.

19
 20 With more than 2,200 bacterial methylomes published to date, it is becoming increasingly
 21 evident that epigenetic regulation of gene expression is highly prevalent across bacterial
 22 species. Despite the exciting prospects for studying epigenetic regulation, our ability to
 23 comprehensively analyze bacterial epigenomes is limited by a bottleneck in integratively
 24 characterizing methylation events, methylation motifs, transcriptomic data, and functional
 25 genomics data. In this regard, this work represents the first comprehensive comparative
 26 analysis of a large collection of a single bacterial species and thus provides a detailed
 27 roadmap that can be used by the scientific community to leverage the current status quo of
 28 epigenetic analyses.

1 **Materials and Methods**

2 *Data and code availability*

3 Genome assemblies are available via NCBI under BioProject ID PRJNA448390. RNA-Seq
4 data has been submitted to the NCBI Sequence Read Archive (SRA) under project
5 PRJNA445308. Scripts and a tutorial supporting all key analyses of this work will be made
6 publicly available at <http://github.com/fanglab/> upon the acceptance of the manuscript.

8 *Clostridium difficile* isolates and culture

9 36 clonal *C. difficile* isolates from CDI fecal samples were obtained using protocols
10 developed in an ongoing Pathogen Surveillance Program at Mount Sinai Hospital
11 (Supplementary Table 1). Additionally, 9 fully sequenced and assembled *C. difficile*
12 genomes were retrieved from Genbank Refseq (<ftp://ftp.ncbi.nih.gov/genomes>, last
13 accessed in November 2016) (Supplementary Table 1). Raw sequencing data from global
14 and UK collections comprising 291 *C. difficile* 027/BI/NAPI genomes were used¹¹
15 (Supplementary Table 4). *C. difficile* positive stool samples were frozen at -80 °C prior to
16 analysis. All stool samples underwent culture for *C. difficile* using ethanol shock culture
17 method, adapted from Griffiths et al. Briefly, approximately 80 mg of solid stool (50 µl liquid
18 stool samples) was added to 0.5 ml of 70% ethanol wash and the sample was vortex mixed
19 and incubated at room temperature for 20 min. A loopful was then cultured onto *C. difficile*
20 selective agar (CDSA, Becton Dickinson, Franklin Lakes, NJ) and the plates were incubated
21 anaerobically at 37 °C for up to 72 h. A single colony was subcultured onto a Trypticase™
22 soy agar with 5% defibrinated sheep blood plate (TSA II™, Becton Dickinson, Franklin
23 Lakes, NJ) and incubated anaerobically at 37 °C for 48 h, after which colonies giving the
24 characteristic *C. difficile* odor and fluorescence under UV illumination were obtained and
25 confirmed by MALDI on a Brucker biotyper. For long-term storage, individual colonies were
26 emulsified in tryptic soy broth containing 15% glycerol and stored at -80 °C.

1 **Single-molecule real-time (SMRT) sequencing**

2 Primer was annealed to size-selected (>8 kb) SMRTbells with the full-length libraries (80 °C
3 for 2 min and 30 s followed by decreasing the temperature by 0.1 °C increments to 25 °C).
4 The polymerase-template complex was then bound to the P6 enzyme using a ratio of 10:1
5 polymerase to SMRTbell at 0.5 nM for 4 h at 30 °C and then held at 4 °C until ready for
6 magbead loading, prior to sequencing. The magnetic bead-loading step was conducted at 4
7 °C for 60 min per manufacturer's guidelines. The magbead-loaded, polymerase-bound,
8 SMRTbell libraries were placed onto the RSII machine at a sequencing concentration of
9 125-175 pM and configured for a 240 min continuous sequencing run.

10

11 ***De novo* genome assembly and motif discovery**

12 The RS_HGAP3 protocol was used for *de novo* genome assembly, followed by custom
13 scripts for genome finishing and annotation. RS_Modification_and_Motif_Analysis.1 was
14 used for *de novo* methylation motif discovery. A custom script was used to examine each
15 motif to ensure its reliable methylation states. In brief, variations of a putative motif are
16 examined by comparing the ipdR distribution of each variation with non-methylated motifs.

17

18 **Identification of R-M systems**

19 Identification of R-M systems was performed as previously described¹⁴. Briefly, curated
20 reference protein sequences of Types I, II, IIC and III R-M systems and Type IV REases
21 were downloaded from the data set 'gold standards' of REBASE⁶⁵ (last accessed in
22 November 2016). All-against-all searches were performed for REase and MTase standard
23 protein sequences retrieved from REBASE using BLASTP v2.5.0+ (default settings, e value
24 < 10⁻³). The resulting e values were log-transformed and used for clustering into protein
25 families by Markov Clustering (MCL) v14-137⁶⁶. Each protein family was aligned with MAFFT
26 v7.305b⁶⁷ using the E-INS-i option, 1,000 cycles of iterative refinement, and offset 0.
27 Alignments were visualized in SEAVIEW v4.6.1⁶⁸ and manually trimmed to remove poorly

1 aligned regions at the extremities. Hidden Markov model (HMM) profiles were then built from
2 each multiple sequence alignment using the hmmbuild program from the HMMER v3.0
3 suite⁶⁹ (default parameters) (available at <https://github.com/pedrocas81>). Types I, II, and III
4 R-M systems were identified by searching genes encoding the MTase and REase
5 components at less than five genes apart.

6

7 **CAAAAA motif abundance and exceptionality**

8 We evaluated the exceptionality of the CAAAAA motif using R'MES³⁶ v3.1.0
9 (<http://migale.jouy.inra.fr/?q=rmes>). This tool computes scores of exceptionality for k-mers of
10 length l , by comparing observed and expected counts under Markov models that take
11 sequence composition under consideration. R'MES outputs scores of exceptionality, which
12 are, by definition, obtained from P -values through the standard one-to-one probit
13 transformation. Analysis of motif abundance was performed with a previous developed
14 framework³⁷ involving a multi-scale representation (MSR) of genomic signals. We created a
15 binary genomic signal for motif content, which was 1 at motif positions, and 0 otherwise. 50
16 length scales were used. Pruning parameter values were set to default and the P -value
17 threshold to 10^{-6} .

18

19 **Whole-genome multiple alignment and classification of CAAAAA positions**

20 Whole-genome multiple alignment of 37 genomes (36 *C. difficile* isolates and *C. difficile* 630)
21 was produced by the progressiveMauve program⁷⁰ v2.4.0 with default parameters. Since
22 progressiveMauve does not rely on annotations to guide the alignment, we first used the
23 Mauve Contig Mover⁷¹ to reorder and reorient draft genome contigs according to the
24 reference genome of *C. difficile* 630. A core alignment was built after filtering and
25 concatenating locally collinear blocks (LCBs) of size ≥ 50 bp using the stripSubsetLCBs
26 script (<http://darlinglab.org/mauve/snapshots/2015/2015-01-09/linux-x64/>). The lower value
27 chosen for LCB size accounts for the specific aim of maximizing the number of orthologous

1 motifs detected. The XMFA output format of Mauve was converted to VCF format using
 2 dedicated scripts, and VCFtools⁷² was used to parse positional variants (SNPs and indels).
 3 Orthologous occurrences of the CAAAAA motif were defined if an exact match to the motif
 4 was present in each of the 37 genomes (conserved orthologous positions), or if at least one
 5 motif (and a maximum of $n-1$, with n being the number of genomes) contained positional
 6 polymorphisms (maximum of two SNPs or indels per motif) (variable orthologous positions).
 7 Non-orthologous occurrences of CAAAAA were obtained from the whole genome alignment
 8 before the extraction of LCBs. The former correspond to those situations where the
 9 CAAAAA motif was absent in at least one genome. Typically, these correspond to regions
 10 containing MGEs or unaligned repetitive regions.

11

12 Identification of transcription factor binding sites, and transcription start sites

13 Identification of transcription factor binding sites (TFBS) was performed by retrieving *C.*
 14 *difficile* 630 regulatory sites in FASTA format from the RegPrecise database
 15 (<http://regprecise.lbl.gov>⁷³, last accessed July 2017). These were converted to PWMs using
 16 in-house developed scripts. This led to a total of 21 PWMs pertaining to 14 distinct
 17 transcription factor families (Supplementary Table 7b). Matches between these matrices and
 18 *C. difficile* genomes was performed with MAST⁷⁴ (default settings). MAST output was filtered
 19 on the basis of P -value. Hits with P -value $<10^{-9}$ were considered positive, while hits
 20 $>10^{-5}$ were considered negative. Hits with intermediate P -values were only considered
 21 positive if the P -value of the hit divided by the P -value of the worst positive hit was lower
 22 than 100. For the CcpA, LexA, NrdR, and CodY (which have shorter binding sites), we
 23 considered positive hits those with P -values $<10^{-8}$. Transcription start sites (TSSs) were
 24 predicted with Parseq⁴¹ under the 'fast' speed option from multiple RNA-seq datasets (see
 25 below). Transcription and breakpoint probabilities were computed using a background
 26 expression level threshold of 0.1 and a score penalty of 0.05. We kept only high-confidence
 27 5' breakpoint hits, located at a maximum distance of 200 bp from the nearest start codon. A

1 ± 5 bp window around the TSS was considered if only one single predicted value was
2 obtained; otherwise we considered an interval delimited by the minimum and maximum
3 values predicted by Parseq.

4

5 **RNA sequencing, read alignment, and differential expression analysis**

6 Purified RNA was extracted from three biological replicates of sporulating (9, 10.5 h) and
7 exponential and stationary grown cultures of *C. difficile* 630 Δ *erm* and *C. difficile*
8 630 Δ *erm* Δ *camA*, DNase-treated, ribosomal RNA-depleted, and converted to cDNA as
9 previously described⁴⁴ (more details in Supplementary Methods). RNA sequencing was
10 performed on a HiSeq 2500, yielding an average of 29.4 (± 4.5 , sd) million 100-bp single-end
11 reads per sample (exponential and stationary growth timepoints) and 26.9 (± 4.3 , sd) million
12 150-bp paired-end reads per sample (sporulation time points). Read quality was checked
13 using FastQC v0.11.5 (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc>). We used
14 Trimmomatic⁷⁵ v0.39 to remove adapters and low-quality reads (parameters: PE, -phred33,
15 ILLUMINACLIP:<adapters.fa>:2:30:10:8:True, SLIDINGWINDOW:4:15, LEADING:20
16 TRAILING:20, MINLEN:50). Subsequently, rRNA sequences were filtered from the data set
17 using SortMeRNA⁷⁶ v2.1, based on the SILVA 16s and 23s rRNA databases⁷⁷, and Rfam 5s
18 rRNA database⁷⁸. The resulting non-rRNA reads were mapped to the *C. difficile* 630
19 reference genome using BWA-MEM v0.7.17-r1198⁷⁹. The resulting bam files were sorted
20 and indexed with Samtools⁸⁰ v1.9, and read assignment was performed with featureCounts⁸¹
21 v1.6.4 (excluding multi-mapping and multi-overlapping reads). A gene was included for
22 differential expression analysis if it had more than one count in all samples. Normalization
23 and differential expression testing were performed using the Bioconductor package DESeq2
24 v1.18.1⁸². Genes with a false discovery rate (FDR) < 0.05 and $|\log_2FC| > 1$ were called as
25 differentially expressed. Functional classification of genes was performed using the DAVID
26 online database (<https://david.ncifcrf.gov>)⁸³ with Fisher's exact test enrichment statistics, a
27 Benjamini-Hochberg corrected *P*-value cutoff of 0.05, and FDR < 0.05. The reproducibility of

1 DAVID's functional classification was tested with Blast2GO v5.2⁸⁴ and Panther v14⁸⁵. Briefly,
2 for Blast2GO, we ran BLASTX searches of the *C. difficile* 630 genome against the entire
3 GenBank bacterial protein database (as of 09/2018). The output, in XML format, was loaded
4 into Blast2GO, and mapping, annotation and enrichment analysis was performed as
5 indicated (<http://docs.blast2go.com/user-manual/quick-start/>). For Panther, we downloaded
6 the most recent HMM library
7 (ftp.pantherdb.org/hmm_scoring/13.1/PANTHER13.1_hmmscoring.tgz), and annotated our
8 *C. difficile* 630 protein set with pantherScore2.1.pl. Both input and background gene lists
9 were formatted to the Panther Generic Mapping File type, as described in the website
10 (<http://www.pantherdb.org>). To assess the significance of the intersection between multiple
11 datasets of differentially expressed genes (typically observed during *C. difficile* colonization
12 and infection), we collected gene-expression data from *in vivo* and *in vitro* studies⁵⁹⁻⁶¹, in
13 which key factors for gut colonization (e.g., time post-infection, antibiotic exposure, and
14 spatial structure (planktonic, biofilm growth)) were tested. Differentially expressed genes
15 were called under the same conditions as described above. Statistical analyses and
16 graphical representation of multi-set intersections was performed with the R package
17 *SuperExactTest*⁸⁶.

18

References

1. Smits WK, Lyras D, Lacy DB, Wilcox MH, Kuijper EJ. *Clostridium difficile* infection. *Nat. Rev. Dis. Primers* **2**, 16020 (2016).
2. Deakin LJ, *et al.* The *Clostridium difficile* *spo0A* gene is a persistence and transmission factor. *Infect. Immun.* **80**, 2704-2711 (2012).
3. Paredes-Sabja D, Shen A, Sorg JA. *Clostridium difficile* spore biology: sporulation, germination, and spore structural proteins. *Trends Microbiol.* **22**, 406-416 (2014).
4. Seekatz AM, Young VB. *Clostridium difficile* and the microbiota. *J. Clin. Invest.* **124**, 4182-4189 (2014).
5. Zidaric V, Rupnik M. Sporulation properties and antimicrobial susceptibility in endemic and rare *Clostridium difficile* PCR ribotypes. *Anaerobe* **39**, 183-188 (2016).
6. Collins J, *et al.* Dietary trehalose enhances virulence of epidemic *Clostridium difficile*. *Nature* **553**, 291-294 (2018).
7. Lanis JM, Barua S, Ballard JD. Variations in TcdB activity and the hypervirulence of emerging strains of *Clostridium difficile*. *PLoS Pathog.* **6**, e1001061 (2010).
8. Valiente E, Cairns MD, Wren BW. The *Clostridium difficile* PCR ribotype 027 lineage: a pathogen on the move. *Clin. Microbiol. Infect.* **20**, 396-404 (2014).
9. Lessa FC, *et al.* Burden of *Clostridium difficile* infection in the United States. *N. Engl. J. Med.* **372**, 825-834 (2015).
10. Sebaihia M, *et al.* The multidrug-resistant human pathogen *Clostridium difficile* has a highly mobile, mosaic genome. *Nat. Genet.* **38**, 779-786 (2006).
11. He M, *et al.* Emergence and global spread of epidemic healthcare-associated *Clostridium difficile*. *Nat. Genet.* **45**, 109-113 (2013).
12. Herbert M, O'Keeffe TA, Purdy D, Elmore M, Minton NP. Gene transfer into *Clostridium difficile* CD630 and characterisation of its methylase genes. *FEMS Microbiol. Lett.* **229**, 103-110 (2003).
13. van Eijk E, *et al.* Complete genome sequence of the *Clostridium difficile* laboratory strain 630Deltaerm reveals differences from strain 630, including translocation of the mobile element CTn5. *BMC Genomics* **16**, 31 (2015).
14. Oliveira PH, Touchon M, Rocha EP. The interplay of restriction-modification systems with mobile genetic elements and their prokaryotic hosts. *Nucleic Acids Res.* **42**, 10618-10631 (2014).
15. Oliveira PH, Touchon M, Rocha EP. Regulation of genetic flux between bacteria by restriction-modification systems. *Proc. Natl. Acad. Sci. USA* **113**, 5658-5663 (2016).
16. Casadesus J, Low D. Epigenetic gene regulation in the bacterial world. *Microbiol. Mol. Biol. Rev.* **70**, 830-856 (2006).
17. Low DA, Weyand NJ, Mahan MJ. Roles of DNA adenine methylation in regulating bacterial gene expression and virulence. *Infect. Immun.* **69**, 7197-7204 (2001).

- 1 18. Cohen NR, *et al.* A role for the bacterial GATC methylome in antibiotic stress survival. *Nat.*
2 *Genet.* **48**, 581-586 (2016).
3
- 4 19. Manso AS, *et al.* A random six-phase switch regulates pneumococcal virulence via global
5 epigenetic changes. *Nat. Commun.* **5**, 5055 (2014).
6
- 7 20. Attack JM, *et al.* A biphasic epigenetic switch controls immuno-evasion, virulence and niche
8 adaptation in non-typeable *Haemophilus influenzae*. *Nat. Commun.* **6**, 7828 (2015).
9
- 10 21. Wion D, Casades J. N6-methyl-adenine: an epigenetic signal for DNA-protein interactions.
11 *Nat. Rev. Microbiol.* **4**, 183-192 (2006).
12
- 13 22. Flusberg BA, *et al.* Direct detection of DNA methylation during single-molecule, real-time
14 sequencing. *Nat. Methods* **7**, 461-465 (2010).
15
- 16 23. Beaulaurier J, Schadt EE, Fang G. Deciphering bacterial epigenomes using modern
17 sequencing technologies. *Nat. Rev. Genet.* **20**, 157-172 (2019).
18
- 19 24. Fang G, *et al.* Genome-wide mapping of methylated adenine residues in pathogenic
20 *Escherichia coli* using single-molecule real-time sequencing. *Nat. Biotechnol.* **30**, 1232-1239
21 (2012).
22
- 23 25. Murray IA, *et al.* The methylomes of six bacteria. *Nucleic Acids Res.* **40**, 11450-11462 (2012).
24
- 25 26. Davis BM, Chao MC, Waldor MK. Entering the era of bacterial epigenomics with single
26 molecule real time DNA sequencing. *Curr. Opin. Microbiol.* **16**, 192-198 (2013).
27
- 28 27. Smits WK. Hype or hypervirulence: a reflection on problematic *C. difficile* strains. *Virulence* **4**,
29 592-596 (2013).
30
- 31 28. Roberts RJ, *et al.* A nomenclature for restriction enzymes, DNA methyltransferases, homing
32 endonucleases and their genes. *Nucleic Acids Res.* **31**, 1805-1812 (2003).
33
- 34 29. Wust J, Sullivan NM, Hardegger U, Wilkins TD. Investigation of an outbreak of antibiotic-
35 associated colitis by various typing methods. *J. Clin. Microbiol.* **16**, 1096-1101 (1982).
36
- 37 30. Dembek M, *et al.* High-throughput analysis of gene essentiality and sporulation in *Clostridium*
38 *difficile*. *MBio* **6**, e02383 (2015).
39
- 40 31. Barra-Carrasco J, Paredes-Sabja D. *Clostridium difficile* spores: a major threat to the hospital
41 environment. *Future Microbiol.* **9**, 475-486 (2014).
42
- 43 32. Ng YK, *et al.* Expanding the repertoire of gene tools for precise manipulation of the
44 *Clostridium difficile* genome: allelic exchange using *pyrE* alleles. *PLoS ONE* **8**, e56051
45 (2013).
46
- 47 33. Donnelly ML, Fimlaid KA, Shen A. Characterization of *Clostridium difficile* spores lacking
48 either SpoVAC or dipicolinic acid synthetase. *J. Bacteriol.* **198**, 1694-1707 (2016).
49
- 50 34. Pogliano J, *et al.* A vital stain for studying membrane dynamics in bacteria: a novel
51 mechanism controlling septation during *Bacillus subtilis* sporulation. *Mol. Microbiol.* **31**, 1149-
52 1159 (1999).
53
- 54 35. Shen A, Fimlaid KA, Pishdadian K. Inducing and quantifying *Clostridium difficile* spore
55 formation. *Methods Mol. Biol.* **1476**, 129-142 (2016).
56
- 57 36. Schbath S, Hoebeke M. R'MES: a tool to find motifs with a significantly unexpected frequency
58 in biological sequences. In: *Advances in genomic sequence analysis and pattern discovery*
59 (ed^{ed}(eds L. Elnitskiy, OP, and L. Welch). World Scientific (2011).
60

- 1 37. Knijnenburg TA, *et al.* Multiscale representation of genomic signals. *Nat. Methods* **11**, 689-
2 694 (2014).
3
- 4 38. Lim HN, van Oudenaarden A. A multistep epigenetic switch enables the stable inheritance of
5 DNA methylation states. *Nat. Genet.* **39**, 269-275 (2007).
6
- 7 39. Ardisson S, *et al.* Cell cycle constraints and environmental control of local DNA
8 hypomethylation in alpha-Proteobacteria. *PLoS Genet.* **12**, e1006499 (2016).
9
- 10 40. Cota I, Bunk B, Sproer C, Overmann J, König C, Casades J. OxyR-dependent formation of
11 DNA methylation patterns in OpvABOFF and OpvABON cell lineages of *Salmonella enterica*.
12 *Nucleic Acids Res.* **44**, 3595-3609 (2016).
13
- 14 41. Mirauta B, Nicolas P, Richard H. Parseq: reconstruction of microbial transcription landscape
15 from RNA-Seq read counts using state-space models. *Bioinformatics* **30**, 1409-1416 (2014).
16
- 17 42. Moxon R, Bayliss C, Hood D. Bacterial contingency loci: the role of simple sequence DNA
18 repeats in bacterial adaptation. *Annu. Rev. Genet.* **40**, 307-333 (2006).
19
- 20 43. Sater MR, *et al.* DNA methylation assessed by SMRT Sequencing Is linked to mutations in
21 *Neisseria meningitidis* isolates. *PLoS ONE* **10**, e0144612 (2015).
22
- 23 44. Fimlaid KA, *et al.* Global analysis of the sporulation pathway of *Clostridium difficile*. *PLoS*
24 *Genet.* **9**, e1003660 (2013).
25
- 26 45. Pishdadian K, Fimlaid KA, Shen A. SpoIID-mediated regulation of sigmaK function during
27 *Clostridium difficile* sporulation. *Mol. Microbiol.* **95**, 189-208 (2015).
28
- 29 46. Pereira FC, *et al.* The spore differentiation pathway in the enteric pathogen *Clostridium*
30 *difficile*. *PLoS Genet.* **9**, e1003782 (2013).
31
- 32 47. Saujet L, Pereira FC, Henriques AO, Martin-Verstraete I. The regulatory network controlling
33 spore formation in *Clostridium difficile*. *FEMS Microbiol Lett* **358**, 1-10 (2014).
34
- 35 48. Pishdadian K, Fimlaid KA, Shen A. SpoIID-mediated regulation of sigmaK function during
36 *Clostridium difficile* sporulation. *Mol Microbiol* **95**, 189-208 (2015).
37
- 38 49. Serrano M, *et al.* A recombination directionality factor controls the cell type-specific activation
39 of sigmaK and the fidelity of spore development in *Clostridium difficile*. *PLoS Genet.* **12**,
40 e1006312 (2016).
41
- 42 50. Saujet L, *et al.* Genome-wide analysis of cell type-specific gene transcription during spore
43 formation in *Clostridium difficile*. *PLoS Genet.* **9**, e1003756 (2013).
44
- 45 51. Rosenbusch KE, Bakker D, Kuijper EJ, Smits WK. C. difficile 630Deltaerm Spo0A regulates
46 sporulation, but does not contribute to toxin production, by direct high-affinity binding to target
47 DNA. *PLoS ONE* **7**, e48608 (2012).
48
- 49 52. Saujet L, *et al.* Genome-wide analysis of cell type-specific gene transcription during spore
50 formation in *Clostridium difficile*. *PLoS Genet* **9**, e1003756 (2013).
51
- 52 53. Fimlaid KA, *et al.* Global analysis of the sporulation pathway of *Clostridium difficile*. *PLoS*
53 *Genet* **9**, e1003660 (2013).
54
- 55 54. Fimlaid KA, Jensen O, Donnelly ML, Siegrist MS, Shen A. Regulation of *Clostridium difficile*
56 Spore Formation by the SpoIIQ and SpoIIIA Proteins. *PLoS Genet* **11**, e1005562 (2015).
57
- 58 55. Ribis JW, Fimlaid KA, Shen A. Differential requirements for conserved peptidoglycan
59 remodeling enzymes during *Clostridioides difficile* spore formation. *Mol Microbiol* **110**, 370-
60 389 (2018).

- 1
- 2 56. Chen X, *et al.* A mouse model of *Clostridium difficile*-associated disease. *Gastroenterology*
- 3 **135**, 1984-1992 (2008).
- 4
- 5 57. Theriot CM, Koumpouras CC, Carlson PE, Bergin, II, Aronoff DM, Young VB. Cefoperazone-
- 6 treated mice as an experimental platform to assess differential virulence of *Clostridium*
- 7 *difficile* strains. *Gut Microbes* **2**, 326-334 (2011).
- 8
- 9 58. McKee RW, Aleksanyan N, Garrett EM, Tamayo R. Type IV pili promote *Clostridium difficile*
- 10 adherence and persistence in a mouse model of infection. *Infect. Immun.* **86**, (2018).
- 11
- 12 59. Maldarelli GA, *et al.* Type IV pili promote early biofilm formation by *Clostridium difficile*.
- 13 *Pathog. Dis.* **74**, (2016).
- 14
- 15 60. Jenior ML, Leslie JL, Young VB, Schloss PD. *Clostridium difficile* colonizes alternative
- 16 nutrient niches during infection across distinct murine gut microbiomes. *mSystems* **2**, (2017).
- 17
- 18 61. Fletcher JR, Erwin S, Lanzas C, Theriot CM. Shifts in the gut metabolome and *Clostridium*
- 19 *difficile* transcriptome throughout colonization and infection in a mouse model. *mSphere* **3**,
- 20 (2018).
- 21
- 22 62. Lewis BB, Pamer EG. Microbiota-Based Therapies for *Clostridium difficile* and Antibiotic-
- 23 Resistant Enteric Infections. *Annual review of microbiology* **71**, 157-178 (2017).
- 24
- 25 63. Abt MC, McKenney PT, Pamer EG. *Clostridium difficile* colitis: pathogenesis and host
- 26 defence. *Nat Rev Microbiol* **14**, 609-620 (2016).
- 27
- 28 64. Sanchez-Romero MA, Cota I, Casadesus J. DNA methylation in bacteria: from the methyl
- 29 group to the methylome. *Curr. Opin. Microbiol.* **25**, 9-16 (2015).
- 30
- 31 65. Roberts RJ, Vincze T, Posfai J, Macelis D. REBASE--a database for DNA restriction and
- 32 modification: enzymes, genes and genomes. *Nucleic Acids Res.* **43**, D298-299 (2015).
- 33
- 34 66. Enright AJ, Van Dongen S, Ouzounis CA. An efficient algorithm for large-scale detection of
- 35 protein families. *Nucleic Acids Res.* **30**, 1575-1584 (2002).
- 36
- 37 67. Katoh K, Standley DM. MAFFT: iterative refinement and additional methods. *Methods Mol.*
- 38 *Biol.* **1079**, 131-146 (2014).
- 39
- 40 68. Gouy M, Guindon S, Gascuel O. SeaView version 4: A multiplatform graphical user interface
- 41 for sequence alignment and phylogenetic tree building. *Mol. Biol. Evol.* **27**, 221-224 (2010).
- 42
- 43 69. Finn RD, Clements J, Eddy SR. HMMER web server: interactive sequence similarity
- 44 searching. *Nucleic Acids Res.* **39**, W29-37 (2011).
- 45
- 46 70. Darling AE, Mau B, Perna NT. progressiveMauve: multiple genome alignment with gene gain,
- 47 loss and rearrangement. *PLoS ONE* **5**, e11147 (2010).
- 48
- 49 71. Rissman AI, Mau B, Biehl BS, Darling AE, Glasner JD, Perna NT. Reordering contigs of draft
- 50 genomes using the Mauve aligner. *Bioinformatics* **25**, 2071-2073 (2009).
- 51
- 52 72. Danecek P, *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156-2158 (2011).
- 53
- 54 73. Novichkov PS, *et al.* RegPrecise 3.0--a resource for genome-scale exploration of
- 55 transcriptional regulation in bacteria. *BMC Genomics* **14**, 745 (2013).
- 56
- 57 74. Bailey TL, Gribskov M. Combining evidence using p-values: application to sequence
- 58 homology searches. *Bioinformatics* **14**, 48-54 (1998).
- 59

- 1 75. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114-2120 (2014).
- 2
- 3
- 4 76. Kopylova E, Noe L, Touzet H. SortMeRNA: fast and accurate filtering of ribosomal RNAs in metatranscriptomic data. *Bioinformatics* **28**, 3211-3217 (2012).
- 5
- 6
- 7 77. Quast C, *et al.* The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res.* **41**, D590-596 (2013).
- 8
- 9
- 10 78. Griffiths-Jones S, Bateman A, Marshall M, Khanna A, Eddy SR. Rfam: an RNA family database. *Nucleic Acids Res.* **31**, 439-441 (2003).
- 11
- 12
- 13 79. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**, 1754-1760 (2009).
- 14
- 15
- 16 80. Li H, *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-2079 (2009).
- 17
- 18
- 19 81. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930 (2014).
- 20
- 21
- 22 82. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).
- 23
- 24
- 25 83. Huang DW, *et al.* DAVID Bioinformatics Resources: expanded annotation database and novel algorithms to better extract biology from large gene lists. *Nucleic Acids Res.* **35**, W169-175 (2007).
- 26
- 27
- 28
- 29 84. Conesa A, Gotz S, Garcia-Gomez JM, Terol J, Talon M, Robles M. Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics* **21**, 3674-3676 (2005).
- 30
- 31
- 32
- 33 85. Mi H, Muruganujan A, Ebert D, Huang X, Thomas PD. PANTHER version 14: more genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools. *Nucleic Acids Res.* **47**, D419-D426 (2019).
- 34
- 35
- 36
- 37 86. Wang M, Zhao Y, Zhang B. Efficient test and visualization of multi-set intersections. *Sci. Rep.* **5**, 16923 (2015).
- 38
- 39
- 40 87. Bordeleau E, Fortier LC, Malouin F, Burrus V. c-di-GMP turn-over in *Clostridium difficile* is controlled by a plethora of diguanylate cyclases and phosphodiesterases. *PLoS Genet.* **7**, e1002039 (2011).
- 41
- 42
- 43
- 44 88. Burton BM, Marquis KA, Sullivan NL, Rapoport TA, Rudner DZ. The ATPase SpoIIIE transports DNA across fused septal membranes during sporulation in *Bacillus subtilis*. *Cell* **131**, 1301-1312 (2007).
- 45
- 46
- 47
- 48

1 Acknowledgements

2 We acknowledge Dr. Richard J. Roberts (New England Biolabs, Inc. USA) for his expertise
 3 and generous help with the prediction of R-M systems and orphan MTases in *C. difficile*
 4 genomes using REBASE Tools, his critical reading, and for providing helpful
 5 comments/suggestions. He was originally an author of this manuscript, however, as a
 6 staunch supporter of the open access movement, he will not author a paper that is not open
 7 access. We also acknowledge Dr. Eduardo P.C. Rocha (Institut Pasteur, Paris, France) for
 8 critical reading and for providing helpful comments/suggestions. The work was primarily
 9 funded by R01 GM114472 (G.F.) from the National Institutes of Health and Icahn Institute for
 10 Genomics and Multiscale Biology. In addition, the work was funded by NIH grants R01
 11 AI119145 (H.v.B and A.B.), R01 AI22232 (A.S.) and R01 AI107029 (R.T.) a Hirschl
 12 Research Scholar award from the Irma T. Hirschl/Monique Weill-Caulier Trust (G.F.), a Pew
 13 Scholar in the Biomedical Sciences grant from the Pew Charitable (A.S.). G.F. is a Nash
 14 Family Research Scholar. A.S. holds an Investigators in the Pathogenesis of Infectious
 15 Disease Award from the Burroughs Wellcome Fund. This work was also supported in part
 16 through the computational resources and staff expertise provided by the Department of
 17 Scientific Computing at the Icahn School of Medicine at Mount Sinai.

18

19 Author Contributions

20 G.F. conceived the hypothesis. A.S. and G.F. supervised the project. P.H.O. and G.F.
 21 designed the computational methods. P.H.O., R.T., A.S., and G.F. designed the
 22 experiments. P.H.O. performed most of the computational analyses and developed most of
 23 the scripts supporting the analyses. J.W.R. performed the growth curves, microscopy
 24 analyses (fluorescence and phase-contrast), analyses of cell length and sporulation stage,
 25 isolated some of the RNA and processed it for qRT-PCR studies; qRT-PCR analyses of
 26 sporulation genes. A.S. constructed the deletion and catalytic $\Delta camA$ mutants, performed

1 complementation, isolated and processed the RNA for several of the RNA analyses and
2 performed many of the sporulation phenotypic assays. E.M.G. and D.T. performed the
3 animal infection experiment and analyzed the data under the supervision of R.T. A.Kim and
4 G.F. performed methylation motif discovery and refinement. O.S. and E.A.M. performed
5 qRT-PCR controls for RNA-seq analyses. O.S., E.A.M., G.D., M.L., C.B., N.Z., D.A., I.O.,
6 G.P., C.H., S.H., R.S., H.v.B. and A.S. contributed to the other experiments. G.D., I.O., and
7 R.S. designed and conducted SMRT sequencing. P.H.O., J.W.R., E.M.G., D.T., A.Kim.,
8 O.S., T.P., S.Z., E.A.M., M.T., S.B., A.A., A.B., R.J.B., R.T., E.E.S., R.S., H.B., A.Kasarskis.,
9 R.T., A.S. and G.F. analyzed the data. P.H.O., R.T., A.S., and G.F. wrote the manuscript
10 with additional information inputs from other co-authors.

11

1 Figure Legends

2 **Fig. 1.** Methyomes of the 36 *C. difficile* strains. (a) Phylogenetic tree of the 36 *C. difficile*
3 strains colored by clade (hypervirulent, human and animal (HA) associated) and MLST
4 sequence type (ST). Heatmap depicting the landscape of methylated motifs per genome,
5 and their average interpulse duration (IPD) ratio. Asterisks refer to new motifs not previously
6 listed in the reference database REBASE. Methylated bases are underlined. The CAAAAA
7 motif was consistently methylated across isolates. Barplot indicates the number and types of
8 active MTases detected per genome. In Type IIC systems, MTase and REase are encoded
9 in the same polypeptide. (b) Representation of the *C. difficile* methylome. Shown are the
10 positions of all methylation motif sites in the reference genome of *C. difficile* 630, colored
11 according to MTase type. Also shown are the average motif occurrences per genome
12 (across the 36 isolates). (c) % of MTases detected according to type. (d) % MTases
13 pertaining to complete R-M systems or without cognate REase (solitary). (e) Breakdown of
14 MTases by location: Integrative Mobile Elements (IMEs), Integrative Conjugative Elements
15 (ICEs), prophages, and other (within the chromosome). No hits were obtained in plasmids.
16 (f) Immediate genomic context of *camA*. The example shown (including coordinates) refers
17 to the reference genome of *C. difficile* 630. + / – signs correspond to the sense and
18 antisense strands respectively. Vertical bars correspond to the distribution of the CAAAAA
19 motif. *CD2754*: phosphodiesterase with a GGDEF domain (PF00990) and a cache domain
20 (PF02743); *ptsI* and *ptsH* belong to a phosphotransferase (PTS) system; *CD2757*: patatin-
21 like phospholipase (PF01734); *CD2758* (*camA*): Type II MTase; *CD2759*: Rrf2-type
22 transcriptional regulator; *CD2760*: phosphodiesterase with a GGDEF domain (PF00990) and
23 a conserved EAL domain (PF00563)⁸⁷; *CD2761*: N-acetylmuramoyl-L-alanine amidase;
24 *CD2762*: undecaprenyl diphosphate synthase.

25
26 **Fig. 2.** CamA modulates sporulation levels in *C. difficile*. (a) Spore purification efficiencies
27 obtained from sporulating cells harvested 3 days after plating. The spore yield (arbitrary
28 units, a.u.), was determined by measuring the optical density at 600 nm of the resulting

1 spore preparations and correcting for the volume of water in which spores were re-
2 suspended. (b) Phase-contrast microscopy after 20 h of sporulation induction. The $\Delta spo0A$
3 strain was used as a negative control because it does not initiate sporulation². Immature
4 phase-dark forespores are marked in pink, and mature phase-bright forespores and free
5 spores are shown in yellow and blue, respectively. Scale bar represents 5 μ m. (c)
6 Morphological analysis of wild-type and $\Delta camA$ cells using fluorescent stains comparing 9
7 and 11 h following sporulation induction. The polar septum formed during asymmetric
8 division is visible using the FM4-64 membrane stain, while the chromosome that is pumped
9 into the forespore after polar septum formation can be seen as a bright foci using the
10 Hoechst 33342 nucleoid stain^{34,88}. FM4-64 staining also allows the engulfing membranes to
11 be visualized. As the mother cell-derived membrane fully encircles the forespore-derived
12 membrane, the FM4-64 signal becomes more intense around the forespore. When these
13 membranes undergo fission, the forespore becomes fully suspended in the mother cell
14 cytosol, and the FM4-64 and Hoechst stains are excluded. Yellow arrows show cells that are
15 undergoing asymmetric division (indicated by a flat polar septum); orange arrows show cells
16 that are in the process of engulfment (indicated by a curved polar septum); and blue arrows
17 show cells that have completed engulfment (indicated by bright membrane staining fully
18 surrounding the forespore). Barplots indicate the number of sporulating cells at different
19 stages of spore assembly in both wild-type and $\Delta camA$ cells. A minimum of 1000 cells per
20 replicate, per strain, and a minimum of two fields were scored for each strain at each
21 timepoint. * $P \leq 0.05$, ** $P < 10^{-2}$, unpaired Student's t-test.

22

23 **Fig. 3.** Abundance, distribution, and conservation of CAAAAA motif sites. (a) Distribution of
24 CAAAAA motif sites in both strands of the reference *C. difficile* 630 genome and
25 corresponding genomic signal obtained by MSR. Letters (A-E) represent regions with
26 particularly high abundance of CAAAAA motifs at scales above 20, i.e., typically above the
27 single gene level (median gene size in *C. difficile* = 0.78 kb) (see Supplementary Table 6d
28 for genes contained in each of the regions). Relation between MSR scale and segment

length is also shown. The significant fold-change (SFC) corresponds to the fold-change ($\log_2$ ratio) between observed and randomly expected overlap statistically significant at $P = 10^{-6}$. Heatmap layers correspond to the number of orthologous conserved (no SNPs/indels, green-shaded) and orthologous variable (with SNPs/indels) CA₅ motif positions. (b) Whole genome alignment of 37 *C. difficile* genomes (36 isolates + *C. difficile* 630 as reference) was performed using Mauve. We defined an orthologous occurrence of the CAAAAA motif (black triangles) if an exact match to the motif was present in each of the 37 genomes (conserved, blue-shaded regions), or if at least one motif (and a maximum of $n-1$, being n the number of genomes) contained positional polymorphisms (maximum of two SNPs or indels per motif) (variable, green-shaded regions). Non-orthologous occurrences of CAAAAA are indicated as orange-shaded regions. The results are shown in Fig. 3a in the form of heatmaps. Numbering in scheme is based on mapping location. (c) DAVID enrichment analysis of the genes containing intragenic and regulatory (100 bp upstream the start codon) orthologous variable CAAAAA motif sites. We considered a Fisher's exact test enrichment statistics, a Benjamini-Hochberg corrected P -value cutoff of 0.05, and a false discovery rate (FDR) < 0.05.

Fig. 4. Distribution of non-methylated CAAAAA motif sites, and overlap with transcription factor binding sites (TFBS) and transcription start sites (TSS). (a) Number of *C. difficile* isolates for which non-methylated CAAAAA motif sites were detected at a given chromosome position (coordinates are relative to the reference genome of *C. difficile* 630). Peak colors correspond to orthologous (conserved and variant) and non-orthologous CAAAAA positions. Some of the major peaks of non-methylated CAAAAA positions were found to overlap with TFBS (e.g., CodY, XylR) and TSS. (b) Genetic regions for which overlap was observed between highly conserved non-methylated CAAAAA motif sites (red ovals) and TFs (CodY and XylR, shown in blue). Other examples of conserved non-methylated CAAAAA motif sites are illustrated in Supplementary Fig. 7b. (c) % CAAAAA motif sites (non-methylated and methylated) overlapping CodY and XylR for each *C. difficile*

1 isolate. (d) Example of a chromosomal region in which non-methylated CAAAAA motifs
2 overlap a TSS (shown as arrow). (e) % CAAAAA motifs (non-methylated (NM) and
3 methylated (M)) overlapping TSSs for each *C. difficile* isolate. (f) % mutated reads (SNPs +
4 indels) in CAAAAA and K(G or T)AAAAA motifs for *M. tuberculosis* (MT), *E. coli* (EC), *C.*
5 *difficile* (CD) and *H. pylori* (HP). AAAAAA was not considered as control motif as it would
6 theoretically be more error-prone. *** $P < 10^{-3}$, Mann-Whitney-Wilcoxon test.

7
8 **Fig. 5.** Gene expression analysis. (a) Heatmap of 161 genes in three replicates of *C. difficile*
9 630 compared to equal number of replicates of *C. difficile* 630 Δ *camA* and that are enriched
10 for the GO terms shown in boxes and detailed in Supplementary Table 8c. The Z score
11 reflects the degree of down- (Z score < 0) or up- (Z score > 0) regulation, computed by
12 subtracting the mean of the log-transformed expression values and dividing by the standard
13 deviation for each gene over all samples scored. (b) Schematic illustrating the sequence of
14 sporulation sigma factor gene transcription and protein activation coupled to morphological
15 changes during sporulation. (c) Comparison of relative transcript levels in wild type and
16 Δ *camA* as determined by qRT-PCR for sporulation sigma factor genes and representative
17 genes in the regulons of sporulation-specific sigma factors at 9 and 11 h after sporulation
18 induction. It should be noted that the primers for *sigK* amplify a region prior to the *sigK*
19 excision site⁴⁹. Statistical significance was determined by one-way ANOVA and Tukey's test
20 for multiple comparisons (* $P \leq 0.05$, ** $P < 10^{-2}$, *** $P < 10^{-3}$, **** $P < 10^{-4}$).

21
22 **Fig. 6.** *In vivo* and additional functional impacts of the Δ *camA* mutation. (a) Kinetics of
23 infection in mice ($n=12$) following inoculation with a sub-lethal amount (10^5 spores) of wild
24 type (WT) *C. difficile* 630 Δ *erm*, MTase mutant Δ *camA*, and complement Δ *camA*-C. Dotted
25 line indicates the limit of detection. Log₁₀-transformed data from each time point were
26 analyzed by ANOVA. * $P < 0.05$, **** $P < 10^{-4}$. (b) Kaplan-Meier survival curves for
27 clindamycin-treated golden Syrian hamsters ($n=6$) infected with 10^3 spores of either wild

1 type (WT) *C. difficile* 630 Δ erm, MTase mutant Δ camA, and complement Δ camA-C. (c)
2 Representative phase-contrast images of vegetative wild type, Δ camA, Δ camA-C, and
3 Δ camA/N165A, grown in BHIS broth (scale bar 5 μ m). (d) Comparison of cell length. A
4 minimum of 400 cells were measured for each strain at mid-log and stationary phase, and a
5 minimum of 1,600 cells were analyzed during sporulation at the indicated timepoints. All cell
6 length measurements were conducted on three independent biological replicates. Statistical
7 significance was determined by one-way ANOVA and Tukey's test for multiple comparisons.
8 (e) Significance of overlap between multiple datasets of DE genes. Comparisons were
9 performed between DE genes called in this study for each time point (blue-shaded) and
10 those from Maldarelli *et al.*⁵⁹ (green-shaded). The latter corresponds to *C. difficile* DE genes
11 in conditions favoring biofilm formation compared to growth on a plate or planktonic form.
12 Color intensities of the outermost layer represent the *P*-value significance of the
13 intersections (3,896 genes were used as background). The height of the corresponding bars
14 is proportional to the number of common genes in the intersection (indicated at the top of the
15 bars for pairwise comparisons between the different studies). (f) Biofilm production as
16 measured by crystal violet staining absorbance at 570 nm. Error bars correspond to
17 standard error of the mean of three independent experiments, with each strain assayed in
18 quadruplicate in each experiment. * $P < 0.05$, ** $P < 10^{-2}$, *** $P < 10^{-3}$, unpaired Student's t-
19 test.

20