Infection of *Borrelia burgdorferi* sensu lato of small mammals in Yunnan Province, China

Zhi-hai He^{1#}, Bao-gui Jiang^{2#}, Zi-hou Gao¹, Zong-ti Shao¹, Yun Zhang¹, Zheng-xiang Liu¹, Yu-qiong Li¹, En-nian Pu¹, Li Tang¹, Ming-guo Yao¹, Na Jia², Michael E. von Fricken³, Jia-fu Jiang^{2*}, Wu-chun Cao^{2*} and Chun-hong Du^{1*}

¹ Yunnan Institute of Endemic Diseases Control and Prevention, Yunnan, P.R. China

² State Key Laboratory of Pathogen and Biosecurity, Beijing Institute of Microbiology and Epidemiology, Beijing, P. R. China

³ George Mason University, Dept. of Global and Community Health, Fairfax, VA, USA

* Correspondence:

Jia-fu Jiang, 15810174122@139.com;

Wu-chun Cao, caowc@bmi.ac.cn;

Chun-hong Du, duchunhong006@hotmail.com

Short Title: BBSL in small mammals from China

#These authors have contributed equally to this work.

Abstract

Background: Lyme disease is caused by *Borrelia burgdorferi* sensu lato (BBSL) which is

usually found in wild and domestic mammals worldwide. Human cases of *B. burgdorferi*

infections have been identified in China, but little direct surveillance of potential rodent

reservoirs has been performed in Yunnan Province, Southwestern China. Yunnan Province is

a tropical area with a diverse topographic range and sustains a high biodiversity of small

mammals that could potentially play an important role in the transmission of a variety of *B.*

burgdorferi sensu lato species.

Methods: 3659 small mammals were captured in 159 sample sites located in 23 counties

in Yunnan Province and screened for BBSL infection by nested PCR based on 5S-23S rRNA

intergenic spacer gene of BBSL. Univariate and multivariate forward stepwise logistic

regression analysis was used to assess the association between infections and related risk

factors.

Results: Infection with BBSL was confirmed in 3.99% (146/3659) of small mammals.

Significant differences in prevalence rates of BBSL were observed at varying landscape types and altitudes. Small mammals in forested areas had higher prevalence rates than other landscape types as did small mammals found at altitudes greater than 2500 meters. The 5S-23S rRNA intergenic spacer revealed that there were 5 genotype of BBSL, including *B. afzelii*, *B. burgdorferi sensu stricto*, *B. japonica*, *B. garinii* and *B. valaisiana*, which demonstrate the genetic diversity and regional distribution.

Conclusions: There exists a wide distribution and genetic diversity of endemic BBSL in Southwestern China, warranting further investigations and monitoring of clinical disease in individuals presenting with symptoms of Lyme disease in these areas.

Author summary

Lyme disease is caused by *Borrelia burgdorferi sensu lato* (BBSL) which is usually found in wild and domestic mammals worldwide. Human cases of *Borrelia burgdorferi sensu lato* infections have been identified in China, but little direct surveillance of potential rodent reservoirs has been performed in Southwestern China. This study documents potential small mammal reservoir hosts collected from a large of sample sites from different landscape types and altitudes, with PCR and sequencing identifying the wide distribution and genetic diversity of endemic *Borrelia burgdorferi sensu lato* in Southwestern China. This was the first report that *B. japonica* was detected in *Apodemus draco* and *Niviventer excelsior* in China. This study adds to body of literature on *Borrelia burgdorferi sensu lato* in China. This work will provide insight regarding small mammals to target for surveillance and we access the association between gender, developmental stage of rodents, environmental landscape and altitude to better prevent human exposure.

Introduction

Lyme borreliosis (LB) is the most commonly reported vector-borne disease across Europe, North America and Asia[1-4]. The causative agents of LB fall within the species complex *B.*

burgdorferi sensu lato (BBSL), and is responsible for a wide spectrum of clinical symptoms. Anti-Borrelia antibodies in rats and humans have been reported in 9 counties and 4 counties of Yunnan, respectively. While there have been documented reports of human cases of Lyme disease in southwestern China[4], the only information on the prevalence of BBSL in rodent reservoirs came from one study, where a majority of rodents were trapped indoors[5]. Yunnan Province is of particular interest given its wide topographic range and high level of small mammal biodiversity, many of which may potential reservoirs for BBSL. We performed a systematic field investigation on the prevalence of BBSL infections in a large quantity of rodents sampled in 23 counties in Yunnan Province, and then analyzed the distribution and genetic diversity of BBSL, as well as the association between infections and suspected risk factors. This study aims to evaluate the role that small mammals play in the transmission of BBSL across Yunnan Province.

Materials and methods

Ethics statement

The research protocol for trapping wild small animals and collecting samples was approved by the Animal Subjects Research Review Boards at the Yunnan Institute of Endemic Diseases Control and Prevention (2013-003), in accordance with the medical research regulations of China and the Regulation of the People's Republic of China for the Implementation of the Protection of Terrestrial Wildlife.

Collection of small mammal samples

From 2011 to 2016, small mammals were captured using animal snap traps set at agricultural, forested, and residential areas at 159 sample sites from 23 counties ranging from 530 to 4300m in Yunnan Province (Table 1). Two hundred snap traps per sample site were placed for three consecutive nights and checked daily. Mammal species were identified according to external morphology, fur color, measurements and visible characters of dentition. Each animal's sex, developmental stage, and location was recorded at the time of sample processing. After identification of species, spleen tissues were removed from the animals and

stored in liquid nitrogen until tested. For unidentified species in the field, the craniums were brought to the laboratory for further identification.

DNA extraction and PCR analysis

DNA was extracted from spleen tissue using the DNA blood and tissue kit (Tiangen Biotechnology, Beijing, China) according to the manufacturer's instruction. A nested PCR for the 5S-23S rRNA intergenic spacer gene of BBSL was done as previously described[6]. The PCR-positive amplicons were directly sequenced with an automated DNA sequencer (ABI PRISM 373; Perkin-Elmer, Norwalk, CT). Sequence analysis was carried out using a FASTA search on the Genbank database, with phylogenetic trees constructed using MEGA software, version 6.06[7]. The 5S-23S rRNA intergenic spacer gene of BBSL obtained in this study were deposited in Genbank under accession numbers MK333406-MK33427 and KP677523.1 respectively.

Statistical analysis

Univariate analysis was used to access the association between gender, developmental stage of rodents, environmental landscape, altitude, and testing positive for BBSL using a chi-square test. All variables with a *P*-value of <0.05 from univariate analysis were entered into a multivariate forward stepwise logistic regression analysis. All analyses were conducted using SPSS (version 17.0, SPSS Inc. Chicago, IL).

Results

A total of 3659 small mammals belonging to 57 species, 29 genera and 10 families from 5 orders were collected (Table 1). The *Apodemus draco* was the most common species (15.82%, 579/3659), followed by *Rattus tanezumi* (15.66%, 573/3659). A total of 146 (3.99%) rodentstested positive for BBSL, with *Ochotona gloveri* (33.33%, 1/3), *Sorex cylindricauda* (14.28%, 7/49), *Soriculusleucops* (14.94%,13/87), and *Rattus tuekkestanicus* (14.28%, 1/7) actively infected with BBSL (Table 2). The positive mammals originated from 14 out of 23 sample counties, including Deqin, Weixi, Yulong, Gongshan, Fugong, Jinggu, Tengchong,

Yongde, Menghai, Yunxian, Shiping, Mile, Yiliang and Yunlong (Table 1), with Gongshan (S1) having the highest prevalence (8.58%), followed by Deqin (S2, 7.85%), and Yiliang (S16, 6.38%). The prevalence of BBSL in small mammals in forested landscapes, agricultural landscapes and residential landscapes were 5.19%, 3.14% and 0.63%, respectively. There was significant difference in prevalence of BBSL in small mammals at the altitude classes of <1500 meters, 1500-2500 meters, >2500 meters with 0.80%, 2.92% and 5.86%, respectively ($\chi^2=43.089$, $p=0.001$), and between different landscapes ($\chi^2=14.945$, $p=0.001$) as depicted in Table 3. The multivariate logistic regression analysis also revealed that samples found at altitudes greater than 1500 meters and in agricultural landscapes were more likely to be infected with BBSL (Table 4).

Table 1. Prevalence of BBSL in small mammals from different survey sites.

Counties	Sampling site	No. of tested	No. of positive for BBSL (%)	B.a	B.b	B.g	B.j	B.v
Gongshan	S1	711	61 (8.58)	39	22	0	0	0
Deqin	S2	662	52 (7.85)	20	1	30	0	1
Shangri-la	S3	106	0 (0)	0	0	0	0	0
Fugong	S4	134	1 (0.75)	0	1	0	0	0
Weixi	S5	239	8 (3.35)	6	2	0	0	0
Yulong	S6	224	1 (0.45)	1	0	0	0	0
Jianchuan	S7	112	0 (0)	0	0	0	0	0
Lushui	S8	114	0 (0)	0	0	0	0	0
Yunlong	S9	177	2 (1.13)	0	0	0	2	0
Tengchong	S10	39	1 (2.56)	0	0	0	0	1
Yingjiang	S11	38	0 (0)	0	0	0	0	0
Yongde	S12	215	3 (1.40)	2	0	0	0	1
Yunxian	S13	68	1 (1.47)	0	0	0	0	1
Jinggu	S14	76	1 (1.32)	1	0	0	0	0
Ninger	S15	88	0 (0)	0	0	0	0	0
Yiliang	S16	94	6 (6.38)	6	0	0	0	0
Shiping	S17	128	6 (4.69)	5	1	0	0	0
Mile	S18	110	2 (1.82)	2	0	0	0	0
Mengzi	S19	22	0 (0)	0	0	0	0	0
Jinping	S20	27	0 (0)	0	0	0	0	0

Wenshan	S21	17	0 (0)	0	0	0	0	0
Menghai	S22	177	1 (0.56)	1	0	0	0	0
Mengla	S23	81	0 (0)	0	0	0	0	0
Total		3659	146 (3.99)	83	27	30	2	4

Table 2. Prevalence of *Borrelia burgdorferi* sensu lato in small mammals of different species.

Orders	Families	Genera	Species	No.of tested	No. of positive (%)	B.a	B.b	B.g	B.v	B.j
Rodetia	Muridae	<i>Rattus</i>	<i>Rattus tanezumi</i>	573	4 (0.70)	3	0	0	1	0
			<i>Rattus nitidus</i>	69	0 (0)	0	0	0	0	0
			<i>Rattus tuekkestanicus</i>	7	1 (14.29)	1	0	0	0	0
			<i>Rattus norvegicus</i>	16	2 (12.50)	1	1	0	0	0
			<i>Rattus brunneusculus</i>	94	2 (2.13)	1	0	0	1	0
		<i>Apodemus</i>	<i>Apodemuslatronum</i>	166	9 (5.42)	1	1	7	0	0
			<i>Apodemuschevrieri</i>	402	20 (4.98)	16	1	3	0	0
			<i>Apodemusdraco</i>	579	19 (3.28)	10	5	3	0	1
		<i>Mus</i>	<i>Mus caroli</i>	75	3 (4.00)	3	0	0	0	0
			<i>Mus pahari</i>	91	6 (6.59)	6	0	0	0	0
			<i>Mus musculus</i>	12	0 (0)	0	0	0	0	0
		<i>Niviventer</i>	<i>Niviventerandersoni</i>	57	3 (5.26)	0	0	2	1	0
			<i>Niviventercoxingi</i>	2	0 (0)	0	0	0	0	0
			<i>Niviventer brahma</i>	1	0 (0)	0	0	0	0	0
			<i>Niviventerreha</i>	32	2 (6.25)	0	2	0	0	0
			<i>Niviventerconfucianus</i>	144	14 (9.72)	6	1	7	0	0
			<i>Niviventer excelsior</i>	29	1 (3.45)	0	0	0	0	1
			<i>Niviventerfulvescens</i>	7	0 (0)	0	0	0	0	0
		<i>Vernaya</i>	<i>Vernaya fulva</i>	7	0 (0)	0	0	0	0	0
		<i>Micromys</i>	<i>Micromysminutus</i>	1	0 (0)	0	0	0	0	0
		<i>Bandicota</i>	<i>Bandicotaindica</i>	2	0 (0)	0	0	0	0	0
		<i>Berylmys</i>	<i>Berylmysbowersi</i>	7	0 (0)	0	0	0	0	0
		<i>Leopoldamys</i>	<i>Leopoldamysedwardsi</i>	6	0 (0)	0	0	0	0	0
	Cricetidae	<i>Eothenomys</i>	<i>Eothenomysmiletus</i>	113	0 (0)	0	0	0	0	0
			<i>Eothenomyseleusis</i>	160	12 (7.50)	8	4	0	0	0

Insectivora	Sciuridae		<i>Eothenomyscachinus</i>	38	4 (10.53)	3	1	0	0	0
			<i>Eothenomys custos</i>	95	2 (2.11)	1	1	0	0	0
			<i>Eothenomysproditor</i>	7	0 (0)	0	0	0	0	0
			<i>Eothenomysolitor</i>	7	0 (0)	0	0	0	0	0
		<i>Pitymys</i>	<i>Pitymysleucurus</i>	42	1 (2.38)	0	0	1	0	0
		<i>Volemys</i>	<i>Volemysclarkei</i>	35	2 (5.71)	1	0	1	0	0
		<i>Dremomys</i>	<i>Dremomyspernyi</i>	26	3 (11.54)	0	0	3	0	0
		<i>Marmota</i>	<i>Marmota himalayana</i>	14	0 (0)	0	0	0	0	0
		<i>Tamiops</i>	<i>Tamiopsswinhoei</i>	7	0 (0)	0	0	0	0	0
	Dipodidae	<i>Eozapus</i>	<i>Eozapussetchuanus</i>	2	0 (0)	0	0	0	0	0
	Soricidae	<i>Crocidura</i>	<i>Crociduralasiura</i>	1	0 (0)	0	0	0	0	0
			<i>Crociduraattenuata</i>	41	2 (4.88)	1	0	0	1	0
			<i>Crocidurahorsfieldi</i>	2	0 (0)	0	0	0	0	0
			<i>Crocidurasuaveolens</i>	3	0 (0)	0	0	0	0	0
			<i>Crociduradracula</i>	62	1 (1.61)	1	0	0	0	0
			<i>Crocidurarussula</i>	33	0 (0)	0	0	0	0	0
		<i>Soriculus</i>	<i>Soriculuscaudatus</i>	46	1 (2.17)	1	0	0	0	0
			<i>Soriculusleucops</i>	87	13 (14.94)	6	7	0	0	0
		<i>Sorex</i>	<i>Sorexalpinus</i>	25	1 (4.00)	1	0	0	0	0
			<i>Sorex cylindricauda</i>	49	7 (14.28)	6	1	0	0	0
			<i>Sorex nigresscens</i>	5	0 (0)	0	0	0	0	0
		<i>Anourosorex</i>	<i>Anourosorex squamipes</i>	114	2 (1.75)	0	2	0	0	0
		<i>Suneus</i>	<i>Suneus murinus</i>	59	1 (1.69)	1	0	0	0	0
	Erinaceidae	<i>Hylomys</i>	<i>Hylomys suillus</i>	14	0 (0)	0	0	0	0	0
		<i>Neotetracus</i>	<i>Neotetracus sinensis</i>	19	0 (0)	0	0	0	0	0
	Talpidae	<i>Scaptonyx</i>	<i>Scaptonyx fusicaudus</i>	5	0 (0)	0	0	0	0	0
		<i>Nasillus</i>	<i>Nasillus gracilis</i>	31	1 (3.23)	1	0	0	0	0

Lagomorpha	Ochotonidae	<i>Ochotona</i>	<i>Ochotona thibetana</i>	92	6 (6.52)	3	0	3	0	0
			<i>Ochotona gloveri</i>	3	1 (33.33)	1	0	0	0	0
Scandentia	Tupaiaidae	<i>Tupaia</i>	<i>Tupaia belangeri</i>	40	0 (0)	0	0	0	0	0
Carnivora	Mustelidae	<i>Mustela</i>	<i>Mustela sibirica</i>	2	0 (0)	0	0	0	0	0
		<i>Meles</i>	<i>Meles meles</i>	1	0 (0)	0	0	0	0	0
Total				3659	146 (3.99)	83	27	30	4	2

Table 3. Risk factors related to *Borrelia burgdorferi* sensu lato based on univariate analyses.

variable	simple size	BBSL Infection		
	constituent ratio (%)	positive rate (%)	χ^2	P
altitude (m)				
~1500	868/3659 (23.72%)	7/868 (0.81%)		
1500~2500	823/3659 (22.49%)	24/823 (2.92%)	43.089	0.001
2500~	1968/3659 (53.79%)	115/1968 (5.84%)		
gender				
male	1753/3659 (47.91%)	81/1753 (4.62%)	3.492	0.062
female	1906/3659 (52.09%)	65/1906 (3.41%)		
age				
adult	3337/3659 (91.20%)	133/3337 (3.99%)	0.002	0.964
pubertal	322/3659 (8.80%)	13/322 (4.04%)		
landscape				
residential	158/3659 (4.32%)	1/158 (0.63%)		
agricultural	1786/3659 (48.81%)	56/1786 (3.14%)	14.945	0.001
forest	1715/3659 (46.87%)	89/1715 (5.19%)		
Total		146/3659 (3.99%)		

Table 4. Risk factors related to BBSL based on multivariate logistic regression.

Variable	OR (95% CI)	p
Altitude (m)		
<1500	1	
1500-2500	4.524 (1.979~10.339)	<0.01
>2500	24.489 (11.351~52.833)	<0.01
Landscape category		
residential	1	
forest	5.617 (0.769~41.052)	0.089
agricultural	8.412 (1.150~61.528)	0.036
forest	1	
residential	0.178 (0.024~1.301)	0.089
agricultural	1.497 (1.153~1.944)	0.002

Sequencing was successful for all 146 positives amplicons samples. The comparative analysis with the BLAST program revealed that 83 samples were *B. afzelii*, 27 were *B. burgdorferi* sensu stricto, 30 were *B. garinii*, four were *B. valaisiana*, and two were *B. japonica*. Deqin county had a distribution of four *Borrelia* spp. except *B. japonica* which was only found in Yunlong county (Table 1). Additionally, four of five *Borrelia* spp. were detected in *Apodemus draco* (Table 2). The nucleotide sequences of the *B. afzelii* sequences

were closely related to the sequence from a patient in China (JX888444.1). All *B. burgdorferi* sensu stricto sequence were identical to the sequence from strain BRE-13 sequenced from a patient's CSF in France (KY594010.1). *B. garinii* sequences in this study showed 99% identity with the strain YN12/2012 from *Canis familiaris* in Yunnan Province. *Borrelia japonica* sequences showed 99% identity with strain Cow611C from a tick in Japan (L30125.1). The *B. valaisiana* sequences were similar to the strain KM2 from *Ixodes granulatus* ticks in Taiwan, China (98%, HM100110.1) and the strain CKA2a from *Apodemus agrarius* in Zhejiang, China (99%, AB022124.1). Phylogenetic analyses based on different representative sequences in this study revealed that all detected *Borrelia* fell within five separate clades belonging to five different types of BBSL including *B. afzelii*, *B. burgdorferi* sensu stricto, *B. garinii*, *B. japonica* and *B. valaisiana* (Fig 1).

Fig 1. Maximum Likelihood phylogenetic tree based on a comparison of *Borrelia burgdorferi* sensu lato 5S-23S rRNA intergenic spacer gene sequences obtained from Yunnan small mammals with *Borrelia burgdorferi* sensu lato reference strains. The number on each branch shows the percent occurrence in 1000 bootstrap replicates. Black circles stood for novel sequences identified in this study.

Discussion

Human cases of LB have been confirmed in almost every province found on mainland China including Yunnan Province. However, most of patients only had serological evidence and were not confirmed for specific genotypes. BBSL has been reported in small mammals trapped in the provinces Qinghai, Hunan, Shanxi, Liaoning, Sichuan, Fujian, Zhejiang, Gansu, Guangdong, Jilin and Yunnan [8-16], suggesting that small mammals are likely the main reservoir hosts in China. This study presents a large sample size extending over a wide geographic area, which provides insight into the prevalence, spatial distribution and genetic diversity of BBSL in small mammals collected in Yunnan Province.

We documented BBSL infection in 30 species of small mammal, among which, 20 species had not been previously documented. These species may be infected occasionally, whether they serve as reservoir hosts need a further study. The *Rattus tanezumii*

(573/3659,15.66%) was the predominant species trapped in households in Yunnan. *Apodemus draco* (579/3659,15.82%) and *A. chevrieri*(402/3659,10.99%) were the predominant hosts species in Yunnan, which was consistent with results from Europe where *Apodemus* are considered a major reservoir of *Borrelia* [17]. BBSL was detected in *Apodemus draco* and in *A. chevrieri* in Yunnan, with *A. draco* capable of carrying four *Borrelia* spp. The *Ochotona gloveri*, *Soriculus leucops* and *Rattus tuekkestanicus* also had a much higher prevalence(>14%) with much larger sample sizes in this study than in other provinces in China [12,18-22]. *Rattus norvegicus* is the prominent household species in Yunnan, which had a high prevalence(12.50%) and was detected positive for pathogenic genotypes (*B. afzelii* and *B. burgdorferi* sensu stricto). We also found that the uncommon species *Sorex cylindricauda* in this study tested positive for BBSL DNA, requiring further investigation to fully understand their role in maintaining or amplifying infections in nature.

Our findings indicated that prevalence rates in rodents are ranked highest to lowest by landscape type as follows: forest landscape > agricultural landscape > residential landscape, which is likely related to tick vector density and preferred habitat. This reiterates the need for individuals traveling into potential tick habitats, like the forest, to take proper protective measures to limit tick bite exposure. Sampling locations in this survey contained a broad range of altitudes from 500 meters to 4500 meters. Among the three altitude classes, small mammals with the highest prevalence of BBSL were found above 2500m. It was reported that *Ixodes ricinus* distribution in Sumava National Park extended toward higher altitudes, probably in relation to warming climates[23]. The roles temperature and humidity play in tick reproduction and reservoir preferences requires further investigation within these altitude ranges. Additionally, there are no reported human cases at these heights, which might reflect lower populations living in these areas.

Our study found five genospecies of BBSL in small mammals in Yunnan Province, four

of them except for *B. japonica*, have previously been associated with LB [24-25]. There exists a wide distribution and genetic diversity of BBSL in Yunnan, compared to only 1-2 genospecies of BBSL in most provinces in China, such as Qinghai, Zhejiang, Guizhou and Guangxi. According to the sequence analysis carried out in this study, most of the *B. afzelii* sequences shared 99% identity with clinical isolates from patients in northeastern China [26]. Most of the *B. burgdorferi* sensu stricto sequences were identical to the sequence from a human case reported in France (KY594010.1). At this time, there have been no confirmed patients with registered sequence of Lyme disease spirochetes in Yunnan province, requiring further investigation in the near future. The sequence of *B. valaisiana* obtained from small mammals cluster into two clades, one cluster within the sequence from Guizhou and Zhejiang province, the other three cluster fell within close proximity to sequences from Europe. Birds are major reservoirs for *B. valaisiana* in Europe, however the transmission cycle maintaining *B. valaisiana* in Yunan may be different from other areas, requiring additional study. *B. japonica* have only been found in Yunlong county, with this representing the first report documenting *B. japonica* in *Apodemus draco* and *Niviventer excelsior* in China. *B. garinii* is the most common genospecies in China, followed by *B. afzelii* [27]. However, we found that *B. afzelii* was the main genospecies detected in Yunnan, which is consistent with previous reports [4]. *B. burgdorferi* sensu stricto has been detected in *Sika deer* from Jilin and in *Caprolagu ssinensis* from Hunan, and detected in small mammals in Yunnan within the more populated counties of Gongshan, Deqin, and Weixi (S1, S2, S5) found in northwestern Yunnan. These findings reflect that Yunnan Province is of particular interest given its diverse topographic range and high level of biodiversity in small mammals that are potential reservoirs for BBSL.

In conclusion, Yunnan Province is an important natural foci of BBSL in China, and given the absence of reported human cases within this region, efforts to expand clinical surveillance

are needed immediately.

Acknowledgements

We thank Professor Zheng-da Gong for identification of rodent species.

References

1. Schwartz AM, Hinckley AF, Mead PS, et al. Surveillance for Lyme Disease - United States, 2008-2015. *MMWR Surveill Summ* 2017;66(22):1-12. <http://doi.org/10.15585/mmwr.ss6622a1> PMID: 29120995
2. Leeflang MM, Ang CW, Berkhout J, et al. The diagnostic accuracy of serological tests for Lyme borreliosis in Europe: a systematic review and meta-analysis. *BMC Infect Dis* 2016;16:140. <http://doi.org/10.1186/s12879-016-1468-4> PMID: 27013465
3. Toshiyuki M. Terrestrial distribution of the Lyme borreliosis agent *Borrelia burgdorferi* sensu lato in East Asia. *Jpn J Infect Dis* 2004;57(6):229-235. PMID: 15623946
4. Hao Q, Hou XX, Geng Z, et al. Distribution of *Borrelia burgdorferi* Sensu Lato in China. *J Clin Microbiol* 2011;49(2):647-650. <http://doi.org/10.1128/JCM.00725-10> PMID: 21106783
5. Du CH, Yin JX, Zuo SQ, et al. Pathogen infection of small mammals from households in Western Yunnan Province. *Chin J Zoonoses* 2016;32(7):623-631 (in Chinese). <http://doi.org/10.3969/j.issn.1002-2694.2016.07.007>
6. Postic D, Assous MV, Grimont PA, et al. Diversity of *Borrelia burgdorferi* sensu lato evidenced by restriction fragment length polymorphism of rrf (5S)-rrl (23S) intergenic spacer amplicons. *Int J Syst Bacteriol* 1994;44(4):743-752. <http://doi.org/10.1099/00207713-44-4-743> PMID: 7981102
7. Tamura K, Peterson D, Peterson N, et al. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 2011;28(10):2731-2739. <http://doi.org/10.1093/molbev/msr121> PMID: 21546353
8. Wu XB, Na RH, Wei SS, et al. Distribution of tick-borne diseases in China. *Parasit Vectors* 2013;6:119. <http://doi.org/10.1186/1756-3305-6-119> PMID: 23617899
9. Wan KL, Zhang ZF, Li YL, et al. Lyme disease spirochete isolated from the kidney of *Rattus coxingi* in Sichuan province. *Chin J Vector Biol Control* 1991;2(6):384-385 (in Chinese).
10. Pan L, Yu ES. Study of Lyme Disease in Fujian Province. *Chin J Public Health* 1992;11(5):271-272 (in Chinese).
11. Shi SZ, Liu ZJ, Sun Y, et al. Investigation on small animals infection of *Borrelia burgdorferi* in Diebu forest. *Chin J Hyg Insect Eq* 2004;10(4):220-221 (in Chinese).
12. Huang HN, Ding Z, He J, et al. Investigation on *Borrelia burgdorferi* infection in ticks and animals from a forest area of Jinlin province. *Chin J Zoonoses* 2006;22(8):785-788 (in Chinese).
13. Zhang DC, Hong Y, Zhong ZQ, et al. Lyme disease spirochete isolated from Rodents and Ticks in Liaoning province. *Chin J Vector Biol Control* 1992;3(1):19 (in Chinese).
14. Hou XX, Geng Z, Hao Q, et al. Rats, the primary reservoir hosts of *Borrelia burgdorferi*, in six representative provinces, China. *Chin J Zoonoses* 2010;26(11):1034-1036 (in Chinese).
15. Meng Z, Jiang LP, Li ZL, et al. Infection with Lyme Spirochetes Detected in Rodents of Zhejiang Middle Area. *Chin J Health Lab Technol* 2010;20(10):2503-2505 (in Chinese).
16. Chen WS, Luo HM, He JF, et al. An Survey on Host and Vector of Lyme Disease in

- Guangdong Province. Chin J Vector Biol Control 2000;11(3):190-193 (in Chinese).
17. Sala V, Faveri ED. Epidemiology of Lyme Disease in Domestic and Wild Animals. Open Dermatol J 2016;10(suppl):15-26. <http://doi.org/10.2174/1874372201610010015>
18. Zhang F, Wang W, Li L. Borrelia burgdorferi Infection and Genotypes in Ticks and Rodents Collected in Qinghai Province. J Prev Med Chin PLA 2016;34(6):799-802 (in Chinese).
19. Wang DM, Hao Q, Cai XH, et al. Study on ribotyping of Lyme borreliosis spirochete in Guizhou province. Chin J Epidemiol 2003;24(12):1129-1131 (in Chinese).
20. Zhang JJ, Zhang F, Li L, et al. Borrelia burgdorferi infection and their genotypes in ticks and rodents collected from Maijishan region, Gansu Province, China. Chin J Zoonoses 2015;31(4):357-360 (in Chinese).
<http://doi.org/10.3969/cjz.j.issn.1002-2694.2015.04.014>
21. Chu CY, He J, Zhao QM, et al. Molecular epidemiological studies on Borrelia burgdorferi in rodents collected in the forest area of several provinces and autonomous regions of China. Chin J Zoonoses 2006;22(9):817-820 (in Chinese).
22. Wulantuya, Gaowa, Yin XH, et al. Survey on Borrelia bacteria in rodents from Bayannur, Inner Mongolia Autonomous Region. Chin J Vector Control 2018;29(3):239-241 (in Chinese).
23. Daniel M, Danielová V, Kříž B, et al. Shift of the Tick Ixodes ricinus and Tick-Borne Encephalitis to Higher Altitudes in Central Europe. Eur J Clin Microbiol Infect Dis 2003;22(5):327-328. <http://doi.org/10.1007/s10096-003-0918-2> PMID: 12736793
24. Rudenko N, Golovchenko M, Grubhoffer L, et al. Updates on Borrelia burgdorferi sensu lato complex with respect to public health. Ticks Tick Borne Dis. 2011;2(3):123-128. <http://doi.org/10.1016/j.ttbdis.2011.04.002> PMID: 21890064
25. Diza E, Papa A, Vezyriet E, et al. Borrelia valaisiana in cerebrospinal fluid. Emerg Infect Dis 2004;10(9):1692-1693. <http://doi.org/10.3201/eid1009.030349> PMID: 15503409
26. Ni XB, Jia N, Jiang BG, et al. Lyme borreliosis caused by diverse genospecies of Borrelia burgdorferi sensu lato in northeastern China. Clin Microbiol Infect 2014;20(8): 808-814. <http://doi.org/10.1111/1469-0691.12532> PMID: 24438159
27. Fang LQ, Liu K, Li XL, et al. Emerging tick-borne infections in mainland China: an increasing public health threat. Lancet Infect Dis 2015;15(12):1467-1479. [http://doi.org/10.1016/S1473-3099\(15\)00177-2](http://doi.org/10.1016/S1473-3099(15)00177-2) PMID: 26453241

Figure 1