

cis dominantly explains regulatory divergence between two indica rice genotypes; drought further enhances regulatory differences

NELZO C. EREFUL^{1,2*}, LI-YU LIU³, SHU-MIN KAO^{4,5}, ERIC TSAI³,
ANTONIO LAURENA⁶, MICHAEL THOMSON⁷, ANDY GREENLAND¹,
WAYNE POWELL⁸, IAN MACKAY^{1,8}, and HEI LEUNG²

¹The John Bingham Laboratory, National Institute of Agricultural Botany (NIAB), Huntingdon Road, Cambridge, CB3 0LE UK

²International Rice Research Institute (IRRI), Los Baños, Laguna 4031 Philippines

³Department of Agronomy, National Taiwan University (NTU), Taipei City 100, Taiwan

⁴Center for Plant Systems Biology, VIB, Ghent B-9052, Belgium

⁵Department of Plant Biotechnology and Bioinformatics, Ghent University, Ghent B-9052, Belgium

⁶Philippine Genome Center – Agriculture, University of the Philippines, Los Baños, Laguna 4031 Philippines

⁷Texas A &M, Department of Soil and Crop Sciences 2474 TAMU, College Station, TX, 77843-2474, USA

⁸SRUC, Peter Wilson Building, West Mains Road, Edinburgh, EH9 3JG UK

*Corresponding author: nelzo.ereful@niab.com

Tel: +44(0)1223 342484 Fax: +44(0)1223 342221

ORCID ID: orcid.org/0000-0003-4460-743X

Abstract

Background. cis and/or trans regulatory divergence within or between related taxa on a genome-wide scale has been largely unexamined in crops, more so, the effect of stress on cis/trans architecture. In this study, the indica genotypes IR64, an elite drought-susceptible lowland variety, and Apo (IR55423-01 or NSIC RC9), a moderate drought-tolerant upland genotype together with their hybrid (IR64 × Apo) were exposed to non- and water-stress conditions. Evidence of cis and/or trans regulatory differences was tested between these two indica rice genotypes.

Results. By sequencing (RNA-seq) the parents and their hybrid, we are able to map genes diverging in cis and/or trans factors between the two genotypes. Under non-stress conditions, cis dominantly explains (11.2%) regulatory differences, followed by trans (8.9%). Further analysis showed that water-limiting conditions largely affect trans and cis + trans factors. Between the two inbred lines, Apo appears to exhibit higher expression fold change of genes enriched in “response to stress” and “photosynthesis” under non- and water-stress conditions. On the molecular level, cis and/or trans regulatory divergence explains their genotypic differences and differential drought response.

Conclusions. Parent–hybrid RNA-seq has the potential to identify genes diverging in cis and/or trans factors even between intra-sub-specifically related genotypes. By comparing cis/trans landscape under stressed and unstressed conditions, this approach has the ability to assess the impact of drought on gene expression. Computational analysis and association of several drought-yield QTL markers with cis- and/or trans-diverging genes provide converging evidences suggestive of a potential approach to identify trait-associated candidate genes using hybrids and their parents alone.

Key words: hybrid, RNA-seq, regulatory divergence, rice, drought, hybrid RNA-seq, cis, trans

Introduction

Allele-specific expression (ASE) imbalance or simply allelic imbalance (AI) happens when one of the two alleles in a heterozygote is expressed more than the other. It has been a subject of investigations in several organisms including crops (rice, barley and maize), model animals (e.g. mouse, *Arabidopsis*, *Drosophila*) and humans. In a hybrid, both alleles are exposed to the same nuclear environment, thus asymmetric expression of the hybrid alleles is attributed to cis regulatory polymorphism. On the other hand, total expression differences between two genotypes which cannot be explained by cis-regulatory differences is ascribed to trans (Wittkopp et al. 2004; McManus et al. 2010). Hence, expression differences between the parental genes that disappear in the hybrid are caused by trans effects (Tirosh et al. 2009).

The relative contribution of both regulatory and coding sequences in generating phenotypic variation and thus evolutionary differences and distinct ecological adaptation has been a subject of recent reports. Regulatory sequences, in particular, have been associated with evolutionary and adaptive divergence in humans (Buil et al. 2015), *Drosophila* (Wittkopp et al. 2004; Wittkopp et al. 2008; McManus et al. 2010; Gompel et al. 2005), yeast (Tirosh et al. 2009), *Arabidopsis* (He et al. 2016), tomato (Guerrero et al. 2016) and stickleback (Verta and Jones, 2019). Some of these reports reveal that it is the changes in cis-acting regulatory sequences that contribute predominantly to phenotypic divergence (Carroll et al. 2008; Stern et al. 2008; Wittkopp et al. 2008; He et al. 2016). However, other investigations have claimed that there is an equal contribution of both cis and trans regulatory factors to evolutionary divergence (Schmitz et al. 2016; Guerrero et al. 2016).

Despite the increasing information of the participation of local and distal regulatory sequences to the expression divergence of organisms, regulatory differences in rice, arguably the world's most important crop, is poorly understood. Moreover, little is known on the effect of stress as a selective agent on gene regulation. A report on rice demonstrated that regulatory divergence is associated with ecological speciation and adaptive alterations (Guo et al. 2015).

A 2015 study shows that modern Green Revolution (GR) rice varieties are typically drought-sensitive due to tight linkage between the loci involved in drought tolerance and plant height (Vikram et al. 2015). Released 50 years ago, IR8, the GR rice is a high-yielding semi-dwarf rice variety, and is derived from a cross between Peta and Dee-geo-woo-gen (DGWG), both of which are *indica* (Hargrove and Coffman, 2006). With a coefficient of parentage (COP) of 0.131 (see pedigrees, S1; from International Rice Information System or IRIS), IR64 and Apo, the materials used in this study, have descended from IR8. Despite their common ancestry, these genotypes show differential response to drought conditions and contrasting ecological adaptation – IR64 thrives in a lowland ecosystem; Apo, in upland. Understanding their regulatory differences may therefore provide insights on the molecular basis of their varying drought response and contrasting adaptation.

Majority of previous studies explored regulatory divergence between or within two or more interspecifically related organisms (e.g., *Drosophila*, *Arabidopsis*, yeast). In this study, two intra-sub-specifically related rice genotypes (both *indica*) were tested for cis- and trans-regulatory differences on a genome-wide scale. This may shed hints on the resolving power of mRNA-seq to dissect regulatory divergence between highly genetically related inbred lines. Such strategy may enable the identification of loci with divergent regulatory regions and potentially causative polymorphism between closely related genotypes.

Our investigation is limited to one-way specific cross (IR64 × Apo) to provide preliminary information on the regulatory differences between the two rice genotypes grown

under normal (well-watered) and water-stress conditions. Such environmental perturbation may further drive regulatory differences.

Results and Discussion

In a companion paper (Ereful et al. 2016), we used mRNA-seq to describe allele-specific expression (ASE) imbalance in IR64×Apo hybrid exposed to both water-stressed and unstressed conditions. Here, we further tested our datasets for evidence of cis and/or trans regulatory differences. Intra-sub-specific comparison of two rice genotypes with contrasting tolerance against water-limiting conditions allows the detection of regulatory differences at candidate loci putatively involved in their distinct ecological adaptation and divergent response against drought. (In this paper, “cis and/or trans” means cis, trans, and their interactions which include cis + trans, cis × trans, and compensatory).

Using paired-end mRNA-seq, we sequenced IR64, Apo and their hybrid exposed to two treatments in two replicates (3 genotypes × 2 treatments × 2 replicates). To be consistent with our previous paper and allow comparison of expression data, we used the *O. sativa* ssp. *japonica* (cv. Nipponbare) MSUv7 cDNA to map the parental reads. Variant calls (SNPs and Indels) between the two inbred parental lines were identified to discern which parent-specific allele belongs to which parental genotype in the F1. We then created a pseudo-molecule to map the heterozygote reads and analyzed for cis/trans regulatory divergence (see Materials and Method). In the hybrid, since both genotype-specific alleles are exposed to the same trans-acting factors, asymmetric expression is attributed to cis differences or allele-specific epigenetic changes (Cowles et al. 2002; Wittkopp et al. 2008).

Using this approach, 9141 genes were found to have one or more mapped reads across genotype–line–treatment combinations (Table S1). Only genes with a total parental read

count of at least 20 in each treatment were considered for further analysis ($IR64 + Apo \geq 20$). We computed the expression ratios (\log_2 -transformed) of genes between the two parental genotypes (Parental, $P = \log_2(IR64/Apo)$) and between the two parent-specific alleles in the F1 heterozygotes (Cis, $C = \log_2(IR64_{F1}/Apo_{F1})$) (see Materials and Methods); trans differences are computed as, $T = P - C$. By plotting the \log_2 -transformed expression ratios of the parents (P) against the hybrid (C) we were able to map isoforms diverging in cis and/or trans regulatory factors and estimate their contributions to the total divergence of the two genotypes under non- and water-stress conditions.

If cis-regulatory differences fully explain divergence between the two genotypes, points (i.e. isoforms) will lie closely on the $y=x$ curve (tested by binomial/Fisher's exact tests, $FDR < 0.5\%$; see Materials and Method for complete statistical tests). On the other hand if trans-regulatory divergence explains their differences, points (i.e. isoforms) will lie at $y=0$ curve (exact tests, $FDR < 0.5\%$) (Wittkopp et al. 2004; Wittkopp et al. 2008; McManus et al. 2010). Additionally, cis + trans (synergistic), cis \times trans (antagonistic), and compensatory effects may also explain their differences. Genes exhibiting cis \times trans difference lies closely on the $y = -x$ curve; compensating on the $x = 0$ curve; for cis + trans, see Fig. 1.

Regulatory divergence under non-stress conditions

Under non-stress conditions, 3705 of the 9141 isoforms were found to have a minimum total mapped read of 20 or more between the parents ($IR64 + Apo \geq 20$). Of these, 415 (11.2%) and 328 (8.9%) of the 3705 isoforms exhibit cis- and trans-regulatory divergence, respectively, between the parents (Table 1; Fig. 1; see Table S2 for complete list of isoforms diverging in cis and/or trans regulatory factors under non-stress treatment). Apparently, cis dominantly explains the regulatory differences over the other categories between IR64 and Apo (exact tests, $FDR < 0.5\%$). ("Conserved" and "ambiguous" categories were omitted from


this comparison). These are shown in red points (i.e. isoforms) lying within or close to the $y=x$ curve (Fig. 1A). GO enrichment analysis of isoforms diverging in cis using AgriGO (Tian et al. 2017) showed significant enrichment of genes associated with “response to stress” (biological process, $FDR < 0.05$; 70 genes; Fig. S1) and “nucleolus” (cellular component, $FDR < 0.05$; 15 genes; Fig. S2). The former suggests that their contrasting response against stress is explained by cis divergence.

On the other hand, GO analysis of isoforms diverging in trans under non-stress conditions showed significant enrichment of genes associated with “photosynthesis” and “generation of precursor metabolites and energy” (biological process, $FDR < 0.05$; Fig. S3) and “thylakoid” and “plastid” (cellular components, $FDR < 0.05$; see Fig. S4 for complete diagram). Analysis of the \log_2 -transformed expression ratios between the two parental genotypes showed that Apo exhibited higher expression fold change (FC) over IR64 in all of the 21 genes enriched in photosynthesis, 18 of which showed at least $2\times$ FC. This photosynthetic efficiency of Apo in terms of expression is ascribed to trans and may further account for its tolerance against drought.


Collectively, cis and trans largely explain divergence of the two genotypes, although compensatory (5.1%), cis + trans (synergistic, 2.0%) and cis \times trans (antagonistic, 3.6%) also contribute to these differences, albeit modestly. GO enrichment analysis of genes diverging in cis + trans showed significant association with “response to stress” and “response to stimulus” (biological process; $FDR < 0.05$) most of which were significantly expressed in the parent Apo (76% or 22 of the 29 genes significantly expressed at $FC \geq 2\times$). Interestingly, one of the genes diverging in cis + trans is the transcription factor ‘no apical meristem’ (OsNAM_{12.1}; LOC_Os12g29330), identified to be a mainstay in the drought-response region in the Vandana/Way Rarem cross (Dixit et al. 2015).

Table 1. Summary of the number of isoforms exhibiting cis and/or trans regulatory divergence between IR64 and Apo under non- and water-stress conditions. χ^2 statistics (with P-values; test for 2×2 contingency or two independent proportions with Yates' continuity correction) indicates the significant difference of the number of isoforms classified in each regulatory category between the two treatments.

Regulatory differences	Non-stress No. of genes (%)	Water-stress No. of genes (%)	χ^2 statistics (P-value)
cis	415 (11.2)	456 (12.6)	3.19 (0.074)
cis + trans (synergistic)	75 (2.0)	194 (5.4)	56.45 (< 0.001)
cis × trans	135 (3.6)	118 (3.3)	0.72 (0.397)
compensatory	187 (5.1)	188 (5.2)	0.05 (0.828)
trans	328 (8.9)	551 (15.2)	69.34 (< 0.001)
Ambiguous	696 (18.8)	752 (20.7)	4.315 (0.038)
Conserved	1869 (50.4)	1366 (37.7)	120.529 (< 0.001)
Total	3705	3626	



A



B

Fig. 1. The “scattering” effect of drought stress on cis and/or trans regulatory architecture between two indica genotypes: IR64 and Apo under (A) non- and (B) water-stress conditions. Visual inspection of the two graphs shows more dispersed points of B compared to A. X-axis indicates expression ratios (in \log_2) between the parental genotypes, $\log_2(\text{IR64}/\text{Apo})$; Y-axis, between the two parent-specific alleles in the hybrid, $\log_2(\text{IR64}/\text{Apo})\text{F1}$.

Cis/trans architecture largely explains genotypic differences. Overall differences between the two parental genotypes were determined using DESeq2 (genotypic differential expression or GDE) to capture loci varying between IR64 and Apo independent of the treatments. Using the 9141 isoforms (described above), GDE between the parents showed 146 genotypically differentially expressed genes (at $\text{FC} \geq 2$ or $|\log_2\text{FC}| \geq 1$, $\text{FDR} < 0.05$; Table S3). Of these GDE genes, 89 (or 61%) exhibited cis divergence between the genotypes; 40, cis + trans (27.3%); 9, trans (6%). Taking all these proportions together, 94% (or 138 of the 146) of the genotypic variations are explained by cis and/or trans regulatory differences between the genotypes. Hence, on the molecular level, cis/trans regulatory landscape explains GDE.

Apparently, local regulatory variations explain mostly of the genotypic differences. Reports showed that natural selection reduces the level of variability of local genomic regions leaving molecular patterns of selection such as increased genetic divergence around adaptive loci (Nielsen et al. 2005; Verta and Jones, 2019; Wachowiak et al. 2018). We further tested this hypothesis by co-localizing cis-diverging genes to drought-yield QTL marker regions (shown below; also see Ereful et al. 2016).

Modern selection and recent breeding procedures have likely generated these differences in which most cis/trans regulatory factors and their interactions appear to favor Apo. These features possibly explain their varying ecological adaptations to lowland (IR64) and upland (Apo) ecosystems and the susceptibility of IR64 and tolerance of Apo to water-

limiting conditions. Our finding is consistent with previous reports which suggests that cis-regulatory changes explain adaptive and ecological divergence between two species of rice (*O. rufipogon* and *O. nivara*; Guo et al. 2015) and among species of *Arabidopsis* (*A. lyrata*, *A. halleri* and *A. thaliana*; He et al. 2016). Furthermore, the finding that cis effects can explain interspecies divergence is common in nature as was previously reported between two subspecies of rice (indica and japonica; Campbell et al. 2019), in *Drosophila* (Wittkopp et al. 2004; Wittkopp et al. 2008; Osada et al. 2017), coffee (Combes et al. 2015), and most recently stickleback (Verta and Jones, 2019).

Regulatory differences under water-limiting conditions

To examine how water-stress treatment changes the regulatory landscape, all lines (parental and hybrid) were exposed to water-limiting conditions. Using the same pipeline as the non-stress treatment, 3626 of the 9141 isoforms were found to have a minimum total mapped read of 20 or more between the parents ($IR64 + Apo \geq 20$) under water-stress conditions. Their cis/trans regulatory landscape under drought stress is plotted in Fig. 1B. Initial comparative visualization of the two figures (Fig. 1A vs B) reflects the impact of stress as a mechanism to propel cis/trans regulatory divergence. Such change in the molecular landscape is suggestive of the effect of drought as an agent of selection for gene regulation.

Of the 3626 isoforms, 456 (or 12.6% of the 3626) and 551 (15.2%) of genes exhibit cis- and trans-regulatory divergence, respectively, which could explain differences between the two genotypes under stress conditions (Table 1; Fig. 1B; see Table S4 for complete list of genes). These results suggest that under water-stress conditions, trans-acting regulatory variations dominantly explain their differences, superseding cis. The χ^2 test confirms that it is the trans and cis+ trans that are significantly affected by the abiotic stress (Table 1), although trans is more significantly affected ($\chi^2=69.34$) than the latter ($\chi^2=56.45$; $P<0.001$). Similar to

a previous finding in *Arabidopsis* (Cubillos et al. 2014), *cis* is relatively stable under water perturbations ($\chi^2=3.19$; $P=0.074$).

The finding that trans regulation differs between environmental regimes agrees with previous studies in yeast (Tirosh et al. 2009) and *Arabidopsis* (Cubillos et al. 2014). We suspect that, as some trans factors are mobile elements (expression; e.g. transcription factors and non-coding RNAs) and can diffuse outside the nucleus, they have been primarily affected by the abiotic stress. Proteins such as TFs have been known to be directly exposed to evolutionary and selection pressures. Additionally, the sensitivity of trans to changes in environmental conditions may be ascribed to its broader mutational target size (sequence) compared to *cis* mutations (Wittkopp et al. 2004; Metzger et al. 2016; Metzger et al. 2017). Similar to non-stress conditions, trans effects under drought conditions showed GO enrichment in “photosynthesis” (Fig. S5), “thylakoid” and “plastid” (Fig. S6). Apo demonstrates a higher expression fold change over IR64 across all genes enriched in “photosynthesis” ($FC \geq 2$, $FDR < 0.05$; 20 genes). Apo appears to photosynthesize more efficiently even under water-limiting regime.

Previous studies revealed that divergence between related taxa could decrease if *cis* and trans act on opposite direction (*cis* × trans, compensating) or could increase if they act on the same direction (*cis* + trans) (Tirosh et al. 2009; Verta and Jones, 2019). Given such a model, IR64 and Apo are predicted to show increased divergence under drought since *cis* + trans was found to be significantly affected by the stress; compensating and antagonistic interactions were not significantly affected. This aligns with a previous paper that disruptive environments or diversifying selection promotes enhancing interactions between *cis* and trans (Shi et al. 2012).

cis/trans architecture explains differential drought response. Using 3-way DE (referred here as drought DE, DDE), expression variations that arise due to the interactions between

the parental genotypes and environment can be detected ($G \times E$) (see Materials and Methods). Results showed that 32 isoforms respond to drought stress across genotypes (Table S5), 13 of which exhibit trans regulatory differences; 12, cis + trans; 3, cis and; 2, cis \times trans. This means that 94% (or 30 of the 32 isoforms) of the drought-responsive genes is explained by cis/trans architecture under water stress. Hence, on the molecular level, their DDE is explained by cis/trans regulatory differences under drought.

Test of ASE in the hybrid

In the previous test above, we showed the cis and/or trans landscape of the two parental genotypes and their hybrid under two contrasting conditions. As allelic imbalance is ascribed to cis differences in a heterozygote state, we further assess the effect of water-limiting conditions on asymmetric expression independent of the parents. This entails plotting the calculated log-transformed expression ratios of the genotype-specific alleles of the hybrid under non-stress or control (x-axis) and water-stress (y-axis) (with binomial exact test at $FDR < 5\%$, against the null hypothesis of having no ASE). This gives information on which isoforms exhibit allelic imbalance under non- and/or water-stress conditions in the hybrid. Results are depicted in Fig. 2 and are summarized in Table S6.

Of the 3058 genes that satisfied our criteria (see Materials and Method), 647 isoforms or 21.2% exhibit AI under both conditions (binomial exact test, $FDR < 5\%$; red dots in Fig. 2); 375 isoforms or 12.3% (yellow dots), under non-stress; 475 or 15.5% (blue), under stress conditions. In summary, almost half of the genes (48.4%) in the hybrids exhibited ASE imbalance across the two conditions. In this study, asymmetric expression was found to be more prevalent under water-limiting compared to non-stress conditions.

IR64-specific allele is favorably expressed in the hybrid over Apo-specific allele under non-stress conditions with 166 and 131 isoforms, respectively, exhibiting ASE

imbalance or cis divergence (under non-stress or control conditions; includes CIS control only; excludes CIS control/stress) (at $FC \geq 2$; binomial exact test, $FDR < 5\%$; Table S6).

However, under drought conditions, Apo-specific allele is favorably expressed over IR64-specific allele with 187 and 159 isoforms, respectively, exhibiting cis divergence (under stress; includes CIS stress only; excludes CIS control/stress) (at $FC \geq 2$; binomial exact test, $FDR < 5\%$; Table S6). There is an apparent genome-wide switching of preferential expression from IR64- to Apo-specific allele from non- to water-stress conditions, respectively, indicating that the Apo-derived alleles were favored under stress in the hybrid.




Fig. 2. Relative expression ratios between IR64- and Apo-specific alleles in the F1 ($\text{Log}_2\text{IR64}_{\text{F1}}/\text{Apo}_{\text{F1}}$) under non- (x-axis) and water-stress (y-axis) conditions. Genes are classified based on their CIS category under non- and water-stress conditions. Inset bar chart: relative proportion (in percentage) of each CIS category including genes with no ASE classification.

Interestingly, 11 genes exhibit allele preference switching in the hybrid, shown as red dots lying closely in the $y = -x$ axis in Fig. 2. These are isoforms which switch from IR64- to Apo-specific allele (or vice-versa) between well-watered and water-stressed conditions. This finding suggests that preferential expression (and possibly dominance) in the heterozygote is condition-dependent. This is an interesting area of further research and needs additional confirmation.

Cis-diverging genes co-localize with drought-yield QTL regions. Previous papers indicated that cis-diverging genes play major roles in interspecific variations and adaptive divergence. In our companion paper, several asymmetrically expressed genes were identified to co-localize with QTL regions known to respond under drought conditions (shown in Fig. 3). These results suggest that several cis-diverging genes are adjacent to regions underlying drought-yield QTLs.

Notably, the drought-response gene LOC_Os12g29330, in particular, co-localizes with RM511, with an estimated distance of 9 kb. It is highly expressed in IR64 as indicated by the positive ratios in both parental (IR64/Apo) and hybrid genotypes ($F1_{IR64}/F1_{Apo}$) in both non- and water-stress conditions (Table S2 and S4, respectively). However, while IR64 expresses this TF, it is drought-susceptible. Rice SNP-Seek database (Alexandrov et al. 2015; Mansueto, et al. 2016) and IR64 assembled draft scaffolds (Cold Spring Harbor Lab (CSHL); Schatz et al. 2014) confirm the structural differences of the 2K upstream region of this locus between the two genotypes: alignments show sequence variations (SNPs and InDels) when pairwise-aligned (Fig. S7; Fig. S8). A multiple intraQTL in the qDTY_{12.1} region centered on the NAM gene (OsNAM_{12.1}) was suggested to provide a concerted drought response (Dixit et al. 2015). In our study, all genes in the qDTY_{12.1} region but one (LOC_Os12g29220, nodulin Mt3, OsMtN3_{12.1}) co-localizing with the NAM gene were not expressed (Table S1).




Fig. 3. Several genes harboring cis divergence were found to co-localize with QTL marker regions (boxed in blue) identified to participate in drought-yield response in the same bi-parental population background (IR64/Apo) under drought conditions (Venuprasad et al. 2009; Ereful et al. 2016). Genes appended with “Trans” exhibited no ASE (as described in the text) but were found to diverge in trans. Markers and genes are said to co-localize if their estimated distance is less than 240kb, the estimated equivalent of 1 cM. (CEN, centromere. Map was generated using ICIMapping (Meng et al. 2015). Positions of markers were estimated based on Cornell and IRMI SSR map at www.archive.grameme.org. Figures not drawn to scale).

In summary, GO enrichment analysis, ASE test, differential expression and results of co-localization analysis consistently suggest that cis/trans architecture may explain the contrasting response of the two genotypes against drought. Taken all these together, these converging evidences underscore the association of cis-diverging regulatory regions with drought response. This leads us to speculate whether it is possible to identify contrasting or diverging adaptive loci, hence, candidate genes associated with important traits using hybrid and their parental lines without generating mapping population. However, such hypothesis needs further investigation.

Finally, mapping cis and/or trans regulatory differences within and between related taxa has been reported in several organisms especially model organisms but is largely unexamined in crops partly due to its highly computational approach. Here, we provided preliminary results suggesting the possibility of assessing cis and/or trans regulatory differences in an important crop using parent – hybrid expression analysis. However, such platform is yet to be fully exploited for crop breeding.

Materials and Methods

In a companion paper, we have described our approach of assessing allelic imbalance in rice hybrid including the description of the datasets (Ereful et al. 2016). In this current work, we report evidence of regulatory divergence of the two genotypes which has not been described before.

Dry-down experiment

Seeds from both parental and hybrid genotypes were initially grown in petri plates under room temperature. After germination, seedlings were transferred to small pots under controlled conditions inside the phytotron, at the International Rice Research Institute (IRRI). Leaf samples were collected to test hybrid status using SSR markers (RM269, RM 511 and RM 80). Hybrid which exhibited clear polymorphism between the parents using these markers were selected for further experiment. Twenty-four (24) seedlings with confirmed polymorphism were transplanted in large pots.

Plant samples were exposed to either non- and water-stress conditions. For water-stressed samples, the fraction of transpirable soil water (FTSW) dry-down approach was applied as previously described (Cal et al. 2013; Serraj et al. 2014; Sinclair and Ludlow 1986). Water-limiting conditions was implemented by initiating a soil dry-down protocol starting 10 days before heading until the plants reached 0.5 FTSW.

RNA extraction

At the end of the dry-down treatment, leaf samples from each plant were collected between 9 and 11 AM. Samples were immediately frozen in liquid nitrogen and stored at –80°C. RNA was extracted using the TRIzol method according to the instructions provided by the supplier (Invitrogen, San Diego, Calif., USA). RNA-seq libraries were prepared as described in Illumina's standard protocol for RNA-seq using the parental (IR64 and Apo) and F1 RNA samples from each treatment (non- and water-stress).

Bioinformatics pipeline

Quality checking. Libraries were sequenced on Illumina GAIIx, generating 38- and 90-base paired end (PE) reads, for the first and second sequencing protocols, respectively. Each sequencing protocol also corresponds to first and second biological replicate.

Quality checking was performed using FASTQC; trimming using FASTX-Toolkit (<http://hannonlab.cshl.edu/>). We checked the 75% percentile of the sequencing qualities for each base of the PE reads in each sample replicate. Bases with 75% percentile of the sequencing qualities ≤ 28 were removed for all reads in the replicate.

Note: Analysis pipeline is shown in S2.

Creating a consensus pseudo-reference. Our bioinformatics approach of estimating ASE levels in the hybrid employs the construction of an *in silico* transcriptome pseudo-reference sequence. This strategy is similar to Shen et al. (2012) and overcame read-mapping biases which may distort ASE estimation (Degner et al. 2009). Use of reference-base mapping (i.e. Nipponbare MSU v7) to align the heterozygote may cause bias (Degner et al. 2009); that is, higher mapping rates towards the allele in the reference sequence. Thus, we preferred the creation of a pseudo-molecule or an *in silico* transcriptome reference to map the reads.

The *O. sativa ssp. japonica* (cv. Nipponbare) MSU v7 cDNA (<http://rice.plantbiology.msu.edu/>) was indexed using bowtie2 (Langmead and Salzberg, 2012). Using the same tool, reads from IR64 and Apo were separately mapped against the indexed reference sequence (no-mixed; no discordant). Alignment options (rdg, rfg, score-min) were set at lenient parameters to allow higher percentage alignment since the materials (IR64 and Apo) are indica and the reference is a japonica.

Using SAMtools (Li et al. 2009), variants (SNPs and Indels) were identified between the MSU7 reference cDNA sequence and the sorted alignment files of both genotypes (commands: sort and mpileup). Custom PERL scripts were created to call SNPs and InDels

common to Apo and IR64 sequences (*indica*), but different from the Nipponbare reference sequence. SNP reads must have at least 5× coverage and the SNP proportion must be more than 0.8. For InDels, read coverage must also be at least 5× but InDel proportion should be at least 0.5. We then developed another PERL script which incorporates these variants (SNP and Indels common to IR64 and Apo but different from the MSUv7 cDNA) to modify the reference sequence.

We further mapped 93-11 *indica* RNAseq reads from BGI (<http://rise2.genomics.org.cn/>) against the pseudo-reference and MSU reference transcriptome sequence and found an increase percentage alignment of 1.09%.

Note: All PERL scripts are available upon request.

SNP calling. Bowtie2 was used to map Apo, IR64 and the F1 genotypes to the pseudo-reference. SAMtools (options: sort and mpileup) was used to sort and parse the mapping results. To find the SNPs between Apo and IR64 sequences, we created a separate PERL script. The criteria, however, should be SNP reads must have at least 3× read coverage and SNP proportion must be more than 0.8. We used these SNPs to discern which read belongs to which genotype in the F1. These variant calls allow us to distinguish and quantify the two genotype-specific SNP-reads in the heterozygote. Reads with multiple SNPs may cause over-estimation of reads. To avoid such scenario, we used a read-wise approach by counting the number of reads with SNPs regardless of their frequency within a read. A separate custom PERL script was created to carry out this step.

Gene expression estimation and normalization procedure. We counted the number of reads from the parents and the genotype-specific alleles in the F1 using eXpress (Roberts and Pachter 2013). All statistical analyses in this study were performed using the software R (v 3.5.1 CRAN).

Normalization was performed using DESeq2 (Love et al. 2014). We preferred DESeq2 scaling factor for the normalization procedure as it is, along with Trimmed Means of M-values (Robinson and Oshlack, 2010), more robust to varying library sizes (Dillies et al. 2013). Normalized count data were rounded off to the nearest integer for downstream statistical tests such as binomial exact test.

Average of the normalized read counts of the two replicates for each sample was calculated. Only genes with a total average normalized read count of 20 of the parental samples ($IR64 + Apo \geq 20$) were considered for further analysis as recommended (Bell et al. 2013; McManus et al. 2010; Shi et al. 2012). In cases where the isoform has zero read count in either of the parents ($IR64 = 0$ or $Apo = 0$) or in either of the parent-specific allele in the hybrid ($F1_{IR64=0}$ or $F1_{Apo=0}$), we adjusted 0 to 1 in order to calculate ratios needing positive integers (e.g. \log_2 transformation). The parental read sum however should be at least 20.

Cis/trans regulatory differences assignment. Normalized read counts of the parental and the hybrid datasets were analyzed for evidence of regulatory divergence using binomial exact tests followed by FDR analysis (Storey and Tibshirani, 2003) at two significant thresholds of 0.5% and 5% FDR. We did not observe any changes in the general trends between the two FDR levels, e.g. dominance of cis followed by trans under non-stress; and dominance of trans followed by cis under water-stress conditions, both of which were consistent at both FDR levels. The most conservative analysis (i.e. $FDR < 0.5\%$) was finally considered in the Results and Discussions.

Genes significantly expressed in either parental or hybrid genotypes were further analyzed for trans effects by comparing the genotype-specific mRNA abundance in the parents and in the hybrid samples using Fisher's exact tests followed by FDR analysis at a significant threshold of 0.5%. Regulatory divergence types are classified according to the categories described below. Such analysis and regulatory divergence assignments are adapted

from previous studies (Landry et al. 2007; McManus et al. 2010; Wittkopp et al. 2008; Bell et al. 2013; Combes et al. 2015).

Cis only: significant DE in both P and C. No significant T ($FDR_P < 0.5\%$, $FDR_C < 0.5\%$, $FDR_T \geq 0.5\%$);

Trans: Significant DE in P and T, but not C ($FDR_P < 0.5\%$, $FDR_C \geq 0.5\%$, $FDR_T < 0.5\%$);

Cis + trans (synergistic): significant DE in P, C and T ($FDR_P < 0.5\%$, $FDR_C < 0.5\%$, $FDR_T < 0.5\%$). The signs of the \log_2FC of these genes in the parental and hybrid are the same. Cis- and trans-regulatory differences favor expression of the same alleles;

Cis \times trans (antagonistic): significant DE in P, C and T ($FDR_P < 0.5\%$, $FDR_C < 0.5\%$, $FDR_T < 0.5\%$). The signs of the \log_2FC of these genes in the parental and hybrid are different. Cis and trans differences favor expression of opposite alleles;

Compensatory: Significant DE in C but not P. Significant T ($FDR_P \geq 0.5\%$, $FDR_C < 0.5\%$, $FDR_T < 0.5\%$);

Ambiguous: statistical test exhibits significant DE in only one of the three tests and has no clear biological interpretations (e.g. $FDR_P \geq 0.5\%$, $FDR_C \geq 0.5\%$, $FDR_T < 0.5\%$).

Conserved: No significant DE in P and C. No significant T ($FDR_P \geq 0.5\%$, $FDR_C \geq 0.5\%$, $FDR_T \geq 0.5\%$);

Both “conserved” and “ambiguous” genes were omitted from discussions.

We then performed chi-square test (at $P < 0.05$) in our data count between the two contrasting treatments to determine regulatory categories responding significantly with respect to environmental changes. The analysis was performed using 2-sample test for equality of proportions with Yates’ continuity correction. The number of isoforms in each regulatory category from each condition constitute the first vector; the total number of isoforms in each condition detected, the second vector.

DESeq2 (Love et al. 2014) multi-factor design was used to test for GDE and DDE. We used the raw data count of 9141 genes expressed with at least one mapped read across genotype – line – treatment to test for DE (Table S2). Only the parental read counts (not the hybrid) were included for this analysis.

GDE was estimated between the genotypes independent of the conditions: (IR64 non-stress, IR64 stressed) vs (Apo non-stress, Apo stressed). Raw read counts were used as the input data count with “IR64” and “Apo” as genotypes. Size factors and dispersions were estimated by DESeq2. On the other hand, differences that arise between the two genotypes under drought conditions relative to non-stress (3-way DE or DDE) were also estimated using the model: (IR64_con, Apo_con) vs (IR64_stress) vs (Apo_stress). Using the function “relevel” of DESeq2, we declared “Apo” and “control (non-stress)” as first level parent and condition, respectively. Results were compared to the genes exhibiting cis/trans regulatory divergence under non- and water-stress conditions, respectively, to find any overlaps.

We used AgriGO (Tian et al. 2017) to perform Gene Ontology (GO) enrichment analysis located at <http://systemsbiology.cau.edu.cn/agriGOv2/>, implementing Singular Enrichment Analysis (SEA) with *O. sativa* MSU v7 as the reference background.

Binomial exact test between the two genotype-specific alleles in the hybrids

A binomial exact test between the two parent-specific alleles in the heterozygote under non- and water-stress conditions was performed using R. This was followed by FDR analysis at a significant threshold of 0.5%. Such analysis was adapted from Cubillos et al. (2014). An isoform should have a total read count (average normalized read counts of parent-specific alleles) of 20 or more ($F1_{Apo} + F1_{IR64} \geq 20$) in one or both conditions. Isoforms with a

total read count of 0 in one or both conditions are discarded. Their log-transformed ratios could not be calculated and binomial exact test could not be performed.

Pairwise-comparison between Apo and IR64 draft genomes for LOC_Os12g29330

We used the published draft genomes of both Apo and IR64 available from Rice SNP-Seek database and assembled IR64 draft scaffolds. To confirm cis regulatory divergence between the two genotypes, *in silico* promoter sequence analysis was performed. Sequence comparison of the promoter sequences between the genotypes was initially visualized using JBrowse located on SNP-Seek (<http://snp-seek.irri.org/jbrowse>). As gaps may be ascribed to absence of sequence coverage, both IR64 and Apo datasets were downloaded from the site to confirm their structural variations. We further confirmed whether there are differences between the genotypes by pairwise-aligning genes and their promoter regions from Apo (Rice SNP-Seek database) against IR64 assembled draft scaffolds (CSHL).

BLAST alignments between IR64 (SNP-Seek) and IR64 (CSHL) using several cDNA query sequences show slight disagreement between the two. We preferred the latter for our succeeding analysis since IR64 (CSHL) – Apo (SNP-Seek) alignment exhibited more conservative structural variations; hence, higher alignment score than IR64 (SNP-Seek) – Apo (SNP-Seek). This is to avoid making conclusions of having promoter structural variations between the two genotypes should we used IR64 (SNPSeek) – Apo (SNPSeek).

Co-localization procedure

Markers which have been previously reported to be involved in drought – yield response were taken from previous papers (Ereful et al. 2016; Venuprassad et al. 2009). Asymmetrically expressed genes which exhibited cis divergence (Fig. 2; Table S6) and the SSR

markers were anchored in the indica genome (Ensembl, <http://plants.ensembl.org>) to estimate their proximity.

Data Availability

All sequencing data from this work are available at NCBI Sequence Read Archive with a submission entry: SUB1568816 with BioProject ID PRJNA338445.

Acknowledgement. This work is supported in part by a grant from Biotechnology and Biological Sciences Research Council (BBSRC: BB/F004265/1.), National Institute of Agricultural Botany (NIAB), and the International Rice Research Institute (IRRI). We thank Jack Lagare and Connie Lotho of the IRRI for providing figures of rice pedigree trees. We also thank Patricia Wittkopp and Joe Coolon for additional help in the scripting during the analysis.

References

- Alexandrov N, et al. (2015) SNP-Seek database of SNPs derived from 3000 rice genomes. *Nucl Acids Res* 43(D1):D1023-D1027. doi: 10.1093/nar/gku1039
- Bell GD, Kane NC, Rieseberg LH, Adams KL (2013). RNA-seq analysis of allele-specific expression, hybrid effects, and regulatory divergence in hybrids compared with their parents from natural populations. *Gen Biol Evol.* 5(7):1309 – 23. doi: 10.1093/gbe/evt072
- Buil A, Brown AA, Lappalainen T, Viñuela A, Davies MN, Zheng HF, Richards JB, Glass D, Small KS, Durbin R, Spector TD, Dermitzakis ET (2015) Gene-gene and gene-environment interactions detected by transcriptome sequence analysis in twins. *Nat Genet* 47(1):88–92
- Cal AJ, Liu D, Mauleon R, Hsing YC, Serraj R (2013) Transcriptome profiling of leaf elongation zone under drought in contrasting rice cultivars. *PLoS* 8(1) doi:10.1371/journal.pone.0054537

- Combes MC, Hueber Y, Dereeper A, Rialle S, Herrera JC, Lashermes P (2015) Regulatory Divergence between Parental Alleles Determines Gene Expression Patterns in Hybrids. *Genome Biol Evol* 7(4):1110–1121. doi:10.1093/gbe/evv057
- Cowles CR, Hirschhorn JN, Altshuler D and Lander ES (2002) Detection of regulatory variation in mouse genes. *Nat Genet.* 32:432–437
- Cubillos FA, Stegle O, Grondin C, Canut M, Tisné S, Gy I, Loudet O (2014) Extensive cis-regulatory variation robust to environmental perturbation in *Arabidopsis*. *The Plant Cell.* 26: 4298–4310. ORCID ID: 0000-0003-3717-0137 (O.L.).
- Degner JF, Marioni JC, Pai AA, Pickrell JK, Nkadori E, Gilad Y, and Pritchard JK (2009) Effect of read-mapping biases on detecting allele-specific expression from RNA-sequencing data. *Bioinformatics* 25(24): 3207–3212. doi:10.1093/bioinformatics/btp579
- Dillies MA, Rau A, Aubert J et al. (2012) A comprehensive evaluation of normalization methods for Illumina high-throughput RNA sequencing data analysis. *Briefings in Bioinformatics* 14(6):671–683. doi:10.1093/bib/bbs046
- Dixit S et al. (2015) Action of multiple intra-QTL genes concerted around a co-localized transcription factor underpins a large effect QTL. *Nat Sci Rep.* doi:10.1038/srep15183
- Ereful NC, Liu LY, Tsai E, Kao SM, Dixit S, Mauleon R, Malabanan K, Thomson M, Laurena A, Lee D, Mackay I, Greenland A, Powell W, Leung H (2016) Analysis of Allelic Imbalance in Rice Hybrids Under Water Stress and Association of Asymmetrically Expressed Genes with Drought-Response QTLs. *Rice* 9:50. DOI: 10.1186/s12284-016-0123-4
- Gompel N, Prud'homme B, Wittkopp PJ, Kassner VA, Carroll SB (2005) Chance caught on the wing: cis-regulatory evolution and the origin of pigment patterns in *Drosophila*. *Nature* 433: 481–487. doi:10.1038/nature03235
- Guerrero RF, Posto AL, Moyle LC, Hahn MW (2016) Genome-wide patterns of regulatory divergence revealed by introgression lines. *Evolution* 70(3):696 – 706
- Guo J, Liu R, Huang L, Zheng XM, Liu PL, Du YS, Cai Z, Zhou L, Wei XH, Zhang FM, Ge S (2015) Widespread and Adaptive Alterations in Genome-Wide Gene Expression Associated with Ecological Divergence of Two *Oryza* Species. *Mol Biol Evol.* 33(1):62–78. doi:10.1093/molbev/msv196
- Hargrove T, Coffman, WR. (2006) Breeding History. In: Barclay A, editor. *Rice Today*. International Rice Research Institute (Philippines): 5(4):35-38

- He F, Arce A, Schmitz G, Beyer A, Koornneef M, Novikova P, de Meaux J (2016) The footprint of polygenic adaptation on stress-responsive cis-regulatory divergence in the *Arabidopsis* genus. *Mol Biol Evol* 33(8):2088-101. doi: 10.1093/molbev/msw096
- Landry CR, Lemos B, Rifkin SA, Dickinson WJ, Hartl DL (2007) Genetic properties influencing the evolvability of gene expression. *Science* 317:18–21
- Langmead B, Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. *Nat Methods* 9:357–359. doi:10.1038/nmeth.1923
- Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup (2009) The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25(16): 2078–2079. doi: 10.1093/bioinformatics/btp352
- Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology* 15: 550. doi: 10.1186/s13059-014-0550-8
- Mansueto, et al. (2016) SNP-Seek II: A resource for allele mining and analysis of big genomic data in *Oryza sativa*. *Current Plant Biology* 7-8:16-25. doi:10.1016/j.cpb.2016.12.003
- McManus CJ, Coolon JD, Duff MO, Eipper-Mains J, Graveley BR, Wittkopp PJ (2010) Regulatory divergence in *Drosophila* revealed by mRNA-seq. *Genome Research* 20:816–825
- Meng L, Li H, Zhang L, Wang J (2015) QTL IciMapping: Integrated software for genetic linkage map construction and quantitative trait locus mapping in biparental populations. *The Crop Journal* 3(3): 269-283. <https://doi.org/10.1016/j.cj.2015.01.001>
- Metzger BPH, et al (2016) Contrasting frequencies and effects of cis- and trans-regulatory mutations affecting gene expression. *Mol Biol Evol* 33:1131–1146
- Metzger BPH, Wittkopp PJ, Coolon JD (2017) Evolutionary Dynamics of Regulatory Changes Underlying Gene Expression Divergence among *Saccharomyces* Species. *Genome Biol Evol* 9(4): 843–854.. doi: 10.1093/gbe/evx035
- Nielsen R, Williamson S, Kim Y, Hubisz MJ, Clark AG, Bustamante C (2005) Genomic scans for selective sweeps using SNP data. *Genome Research* 15:1566–1575. DOI: <https://doi.org/10.1101/gr.4252305>, PMID: 16251466
- Osada, N, Miyagi R, and Takahashi A (2017) Cis- and trans-regulatory effects on gene expression in a natural population of *Drosophila melanogaster*. *Genetics* 206(4): 2139-2148; doi.org/10.1534/genetics.117.201459

- Roberts A, Pachter L (2013) Streaming fragment assignment for real-time analysis of sequencing experiments. *Nat Methods* 10(1):71–73. doi:10.1038/nmeth.2251
- Robinson MD, Oshlack A (2010) A scaling normalization method for differential expression analysis of RNA-seq data. *Genome Biology* 11:R25. doi.org/10.1186/gb-2010-11-3-r25
- Schatz MC, Maron LG, Stein JC, et al. (2014) Whole genome *de novo* assemblies of three divergent strains of rice, *Oryza sativa*, document novel gene space of aus and indica. *Genome Biology*.15:506. <http://genomebiology.com/2014/15/12/506>
- Schmitz JF, Zimmer F, Bornberg-Bauer E (2016) Mechanisms of transcription factor evolution in Metazoa. *Nucleic Acids Res* 44(13): 6287–6297. doi: 10.1093/nar/gkw492
- Serraj R, Dongcheng L, Hong H, Sellamuthu R, Impa S, Cairns J, Dimayuga G, Torres R (2014) Novel approaches for integration of physiology, genomics and breeding for drought resistance improvement in rice., <http://www.intlcss.org/>
- Shen Y, Catchen J, Garcia T, et al (2012) Identification of transcriptome SNPs between *Xiphophorus* lines and species for assessing allele specific gene expression within F1 interspecies hybrids. *Comp Biochem Physiol C Toxicol Pharmacol*. 155(1): 102–108. doi:10.1016/j.cbpc. 2011.03.012
- Shi X, Ng DWK, Zhang C, Comai L, Ye W and Chen ZJ (2012) Cis- and trans-regulatory divergence between progenitor species determines gene-expression novelty in *Arabidopsis* allopolyploids. *Nat Comm* 3:950. DoI: 10.1038/ncomms195
- Sinclair T, Ludlow M (1986) Influence of soil water supply on the plant water balance of four tropical grain legumes. *Aust J Plant Physiol* 13:329. doi:10.1071/pp9860329
- Stern DL, Orgogozo V (2008) The loci of evolution: how predictable is genetic evolution? *Evolution* 62: 2155–2177
- Storey JD, Tibshirani R. (2003) Statistical significance for genomewide studies. *Proc Natl Acad Sci* 100: 9440–9445
- Tian T, Liu Y, Yan H, You Q, Yi X, Du Z, Xu W, Su Z (2017) agriGO v2.0: a GO analysis toolkit for the agricultural community, 2017 update. *Nucleic Acids Res* gkx382. doi: 10.1093/nar/gkx382
- Tirosh I, Reikhav S, Levy AA, Barkai N (2009) A yeast hybrid provides insight into the evolution of gene expression regulation. *Science* 324:659-662
- Venuprasad R, Bool ME, Dalid C, Bernier J, Kumar A, Atlin GN (2009) Genetic loci responding to two cycles of divergent selection for grain yield under drought stress in a rice breeding population. *Euphytica* 167:261–269

Verta JP and Jones FC (2019) Predominance of cis-regulatory changes in parallel expression divergence of sticklebacks. eLife DOI: <https://doi.org/10.7554/eLife.43785>

Vikram P, Swamy BP, Dixit S, Singh R, Singh BP, Miro B, Kohli A, Henry A, Singh NK, Kumar A (2015) Drought susceptibility of modern rice varieties: an effect of linkage of drought tolerance with undesirable traits. Nat Sci Rep 5:14799. doi: 10.1038/srep14799

Wachowiak W, Zaborowska J, Łabiszak B, Perry A, Zucca GM, González-Martínez, SC, and Cavers S (2018) Molecular signatures of divergence and selection in closely related pine taxa. Tree Genetics & Genomes. 14: 83. <https://doi.org/10.1007/s11295-018-1296-3>

Wittkopp PJ, Haerum BK, Clark AG (2004) Evolutionary changes in cis and trans gene regulation. Nature 430:85-88. doi:10.1038/nature02698.

Wittkopp PJ, Haerum BK, Clark AG (2008) Regulatory changes underlying expression differences within and between *Drosophila* species. Nature Genet. 40(3):346-350

Supplementary materials

S1. Pedigrees of IR64 and Apo, taken from IRIS.

S2. Analysis pipeline for determining regulatory divergence and ASE.

Table S1. List of isoforms and their corresponding raw read counts and average normalized read counts which have a minimum mapped read of one or more for at least one column across genotype–line–treatment combinations

Table S2. Isoforms identified to have at least 20 mapped reads in either of the parental genotypes under non-stress conditions and their regulatory divergence classification

Table S3. Results of GDE showing isoforms genotypically differentially expressed ($FC \geq 2$; $FDR < 0.05$)

Table S4. Isoforms identified to have at least 20 mapped reads in either of the parental genotypes under water-stress conditions and their regulatory divergence classification.

Table S5. Results of 3-way differential expression or drought DE (DDE).

Table S6. Results of ASE binomial exact test ($FDR < 0.5\%$). Isoforms were classified as cis-control, cis-stress, or cis (control/stress).

Fig. S1. Graphical results of GO enrichment analysis (biological process) using cis-diverging genes under non-stress conditions as input sequence.

Fig. S2. Graphical results of GO enrichment analysis (cellular component) using cis-diverging genes under non-stress conditions as input sequence.

Fig. S3. Graphical results of GO enrichment analysis (biological process) using trans-diverging genes under non-stress conditions as input sequence.

Fig. S4. Graphical results of GO enrichment analysis (cellular component) using trans-diverging genes under non-stress conditions as input sequence.

Fig. S5. Graphical results of GO enrichment analysis (biological process) using trans-diverging genes under water-stress conditions as input sequence.

Fig. S6. Graphical results of GO enrichment analysis (cellular component) using trans-diverging genes under water-stress conditions as input sequence.

Fig. S7. Pairwise-alignment results using BLAST between IR64 (CSHL) and Apo (SNP-seek) of the promoter region for the gene, No apical meristem (LOC_Os12g29330)

Fig. S8. Graphical representation of the pairwise alignment of the promoter sequence for the gene, No apical meristem (LOC_Os12g29330) using SNP-seek genome browser.