Yeast Bax Inhibitor (Bxi1p/Ybh3p) is a Calcium Channel in *E. coli*

James Mullin, John Kalhorn, Nicholas Mello, Amanda Raffa, Alexander Strakosha,
and Nicanor Austriaco, O.P.*

Department of Biology

Providence College

Providence, Rhode Island

*CORRESPONDING AUTHOR:

1 Cunningham Square, Providence, RI 02918, U.S.A.

Email: naustria@providence.edu

Abstract

Human Bax Inhibitor-1 (HsBI-1/TMBIM6) is the founding member of the evolutionary conserved TMBIM superfamily of proteins that share sequence homology within the transmembrane Bax inhibitor-containing motif (TMBIM). Mechanistically, BI-1/TMBIM6 and all the other mammalian TMBIM proteins appear to be involved in the maintenance of calcium homeostasis, and the crystal structure of a bacterial TMBIM protein, BsYetJ, suggests that the protein is a pH-sensitive calcium leak. The budding yeast, *Saccharomyces cerevisiae*, has a single TMBIM family member (YNL305C) named Bxi1p/Ybh3p. To determine the function of Bxi1p/Ybh3p, we overexpressed Bxi1p-EGFP in *E. coli* to determine if it is a calcium channel. We show that bacterial cells expressing Bxi1p-EGFP are more permeable to calcium than controls. Thus, our data suggests that yeast Bax inhibitor (Bxi1p) is a calcium channel in *E. coli*, lending support to our proposal that Bxi1p is a *bona fide* member of the TMBIM family of proteins. Further, we use our bacterial system to show that gadolinium is an inhibitor of Bxi1p *in vivo*, suggesting a path forward to identifying other small-molecular inhibitors of this clinically-important and highly conserved superfamily of proteins. Finally, parallel experiments revealed that the human Bax Inhibitor-1 (HsBI-1/TMBIM6) is also a calcium channel in bacteria that can be inhibited by gadolinium.

Introduction

Human Bax Inhibitor-1 (HsBI-1/TMBIM6) is the founding member of the evolutionary conserved TMBIM superfamily of proteins that share sequence homology within the transmembrane Bax inhibitor-containing motif (TMBIM) (Xu & Reed, 1998; Reimers et al., 2006; Hu, Smith & Goldberger, 2009; Carrara et al., 2012; Rojas-Rivera & Hetz, 2015; Gamboa-Tuz et al., 2018). The human genome encodes six members of the superfamily (TMBIM1-6) that are homologous to other BI-1 like proteins in vertebrates, plants, yeast, bacteria and viruses. TMBIM1-3 are localized to the Golgi apparatus, TMBIM4-6 are found in the endoplasmic reticulum, and TMBIM5 is a mitochondrial protein (Lisak et al., 2015).

Phenotypically, BI-1/TMBIM6 is an ER-localized, anti-apoptotic protein that was first identified in a screen for human proteins that could inhibit Bax-mediated cell death in yeast (Xu & Reed, 1998). Mammalian cells stably overexpressing BI-1/TMBIM6 are protected against ER-stress induced apoptosis (Chae et al., 2004; Bailly-Maitre et al., 2006). Moreover, mice lacking BI-1/TMBIM6 are more sensitive to stroke-induced cerebral damage and tunicamycin-induced kidney toxicity (Chae et al., 2004). Clinically, BI-1 is known to be overexpressed in breast cancer, glioma, lung, and prostate carcinoma (van 't Veer et al., 2002; Schmits et al., 2002; Grzmil et al., 2003, 2006; Lu et al., 2015). Strikingly, downregulation of BI-1 in prostate cancer cells by RNAi leads to cell death (Grzmil et al., 2003). The other mammalian TMBIM proteins are also cytoprotective against different triggers known to induce cell death (Rojas-Rivera & Hetz, 2015).

Mechanistically, BI-1/TMBIM6 and all the other mammalian TMBIM proteins appear to be involved in the maintenance of calcium homeostasis (Lisak et al., 2015; Liu, 2017). Knocking out TMBIM6 in hepatocytes leads to higher Ca^{2+} content in the ER of hepatocytes (Chae et al., 2004), while overexpressing the gene leads to reduced ER Ca^{2+} content (Westphalen et al., 2005).

Similarly, stably overexpressing each HA-tagged TMBIM1-6 protein family member in HT22 cells reduced ER Ca^{2+} content, and all the TMBIM proteins except TMBIM5 also reduced the basal concentration of calcium in the cytosol (Lisak et al., 2015). TMBIM6 regulates Ca^{2+} flux in a pH-dependent manner (Ahn et al., 2009, 2010; Kiviluoto et al., 2013).

Structurally, the role of the TMBIM family of proteins in calcium homeostasis has been confirmed by the solution of the crystal structure of a bacterial TMBIM protein, BsYetJ, from *Bacillus subtilis* that suggests that the protein is a pH-sensitive calcium leak (Chang et al., 2014). It has a seven-transmembrane-helix fold structure that has either a closed or an open channel conformation depending upon the pH of its environment. It also has a di-aspartyl pH sensor in its C-terminal pore domain (Asp171-Asp195) that corresponds to two aspartate residues in B1/TMBIM6 (Asp188-Asp213) (Chang et al., 2014). Biochemical characterization of BsYetJ proteoliposomes at various pHs revealed that the pH-sensitive calcium-leak activity is intrinsic to the protein (Chang et al., 2014).

The budding yeast, *Saccharomyces cerevisiae*, has a single TMBIM family member (YNL305C) named Bxi1p/Ybh3p (Chae et al., 2003; Cebulski et al., 2011; Büttner et al., 2011). The protein is homologous to the mammalian TMBIM1-6 family members and contains the conserved di-aspartyl pH sensor in its C-terminal pore domain (Asp255-Asp278) that has been identified as the latch responsible for opening and closing the calcium leak. In a previous study, we showed that Bxi1p-GFP is localized to the ER, and that mutant yeast cells deleted of *BXI1* are more susceptible to a range of pharmacological and environmental triggers that induce cell death, especially pharmacological triggers associated with the unfolded protein response (Cebulski et al., 2011). This pro-survival function for Bxi1p has been confirmed by two other laboratories (Chae et al., 2003; Teng et al., 2011), and is consistent with the anti-apoptotic

function associated with the TMBIM superfamily. However, a subsequent paper from the Madeo Laboratory at the University of Graz, published days after our original publication, suggested that the protein encoded by the ORF – which they called Ybh3p for yeast BH3-only protein – is a pro-apoptotic member of the BH3-only family of proteins that translocates from the vacuole to the mitochondria to trigger BH3-domain dependent apoptosis (Büttner et al., 2011).

To begin to experimentally resolve the apparent discrepancy between our data that suggests that Bxi1p/Ybh3p is an anti-apoptotic member of the TMBIM superfamily, and the data that proposed instead that Bxi1p/Ybh3p is a pro-apoptotic member of the BH3-only family of proteins, we overexpressed Bxi1p-GFP in *E. coli* to interrogate its putative calcium channel function. We show that bacterial cells expressing Bxi1p-GFP are more permeable to calcium than controls. Our data suggests that yeast Bax inhibitor (Bxi1p) is a calcium channel in *E. coli*, lending support to our proposal that Bxi1p is a *bona fide* member of the TMBIM family of proteins. We also use our bacterial system to show that gadolinium is an inhibitor of Bxi1p, suggesting a path forward to identifying other small-molecular inhibitors of this clinically-important and highly conserved superfamily of proteins.

99 Materials and Methods

100 Bacterial Strain, Plasmids, and Growth Conditions: All experiments were done with DH5α *E. coli*
 101 cells obtained from New England Biolabs. Bacterial plasmids overexpressing yBxi1p-EGFP and
 102 the hTMBIM6-EGFP fusion were constructed and verified by VectorBuilder. The vector ID for
 103 plasmid, pBAD-EGFP(ns):4XGS:{yBXI1}, which overexpresses wildtype yBxi1p-EGFP in media
 104 containing arabinose is VB160522-1016qch; and the vector ID for plasmid, pBAD-
 105 EGFP(ns):4XGS:{hTMBIM6}, which overexpresses hTMBIM6-EGFP is VB170103-1020maq.
 106 These vector IDs can be used to retrieve detailed information and plasmid maps from
 107 vectorbuilder.com. Cells were transformed and cultured using standard bacterial protocols and
 108 media (Ausubel et al., 2002). Unless noted otherwise, all drugs and reagents were purchased from
 109 SIGMA-Aldrich.

110
 111 Bxi1p-GFP Induction Assays: DH5α *E. coli* cells transformed with a bacterial plasmid
 112 overexpressing either yBxi1p-EGFP or hTMBIM6-EGFP were inoculated in 20mL of LB
 113 containing 100μg/mL ampicillin and allowed to grow to an OD₆₀₀ of 0.6 at 37°C. Expression of
 114 the Bxi1p-EGFP fusion protein was then initiated by the addition of 200μL of 20% arabinose.
 115 Expression levels of the yBxi1p-EGFP or hTMBIM6-EGFP fusion proteins were then visualized,
 116 1, 3, 6, 12, 18, and 24 hours after induction with a Zeiss LSM700 confocal microscope and
 117 quantified with an Accuri C6 Flow Cytometer. For all measurements with the Accuri C6 Flow
 118 Cytometer, bacterial cells were first diluted in 10mL water to an OD₆₀₀ of 0.003.

119
 120 Calcium Permeability Assays: DH5α *E. coli* cells transformed with a bacterial plasmid
 121 overexpressing either yBxi1p-EGFP or hTMBIM6-EGFP were inoculated in 20mL of LB

containing 100µg/mL ampicillin and cultured overnight at 37°C in a floor shaker set to 250 RPM. The following day, 1mL of cells were re-inoculated into 10mLs of fresh LB amp and allowed to grow to an OD₆₀₀ of 0.60. Both induced (with 2% arabinose) and uninduced cultures then were incubated at 37°C, 250 RPM for six hours. Next, the cells were spun down at 3,000g for 10mins, washed twice with 10mL Buffer A (50mM Tris pH 7.5; 100mM KCl; 1mM MgCl₂), and resuspended in 20mL Buffer B (120mM Tris pH 8.0; 0.2mM EDTA) where they were incubated at 37°C, 250 RPM for ten minutes. The cells were then spun down at 3,000g for 10 mins and washed twice with 10mL of buffer A. Finally, the cells were re-suspended in 2.5mL Buffer A containing 10 µM Fura-2AM and incubated at 37°C, 250 RPM for two hours.

Following Fura-2AM incubation, cells were pelleted at 3,000g for 10mins and washed twice with 10mL aliquots of buffer A. Following the washes, the cells were re-suspended in 4mL Buffer A, and plated in 250 µL aliquots in a 96-well black Costar plate. Cells were excited at the standard wavelength for Fura-2AM (510 nm) and measured for wavelength emission of the bound (340nm) and unbound (380 nm) state using a Biotek Cytation 3 Cell Imaging Reader. Emissions were assessed every two minutes for six minutes. Following the initial six minutes, 5 µL of 500 mM CaCl₂ was added and run at the same conditions for an additional 10 minutes. Calcium concentration was represented by the ratio of R₃₄₀/R₃₈₀ of Fura-2 as has been typically done in previous studies (Hudson et al., 1998; Luo et al., 2019). All experiments were done in triplicate. Statistical significance was determined with the Student's t-test, using Graph Pad Prism 6. By default, one asterisk is p<0.05; two asterisks is p<0.01; three asterisks is p<0.001; and four asterisks is p<0.0001.

Gadolinium Assays: To determine whether or not gadolinium inhibited calcium channel activity, the calcium permeability assay described above was repeated with an additional step: Following the Fura-2AM incubation and the emission baseline measurements, but before the addition of external calcium, 10 μ L of 500 mM GdCl₃ was added to the cells in the 96-well plates for 10 minutes to determine if the larger cation had any effect on the cell's permeability to Ca²⁺.

To determine whether or not gadolinium had any effect on bacterial cell growth, DH5 α *E. coli* cells transformed with the appropriate vector were streaked onto LB-ampicillin plates with varying concentrations of gadolinium (GdCl₃). The plates were incubated at 37°C for two days and imaged with a SynGene G:BOX system.

Results

Yeast Bax Inhibitor (Bxi1p/Ybh3p) is a Calcium Channel in *E. coli*

To determine if yeast Bax inhibitor (Bxi1p) has calcium channel function in *E. coli*, we overexpressed a Bxi1p-EGFP fusion protein in bacterial cells using the arabinose-inducible araBAD promoter (Guzman et al., 1995). A similar experiment had been done with the *Bacillus subtilis* homolog, BsYetJ, in bacteria (Chang et al., 2014).

Expression levels of the Bxi1p-EGFP fusion protein were visualized 1, 3, 6, 12, 18, and 24 hours after induction on a Zeiss LSM700 confocal microscope and were quantified with an Accuri C6 Flow Cytometer to measure GFP fluorescence. As shown in Figures 1A and 1B, bacterial cells grown in media containing arabinose for six hours showed significant induction ($p < 0.0001$) of the Bxi1p-EGFP fusion protein. This time point was chosen for all further experiments. We then loaded these DH5 α cells with Fura-2AM, a membrane-permeable, fluorescent calcium indicator that has been used to determine the intracellular calcium dynamics of bacterial cells (Gangola & Rosen, 1987; Chang et al., 2014). Upon addition of external calcium, intracellular calcium concentration as indicated by the R_{340}/R_{380} ratio of Fura-2 increased more rapidly in bacterial cells induced with arabinose as compared with uninduced controls (Figure 1C). This data suggests that Bxi1p-EGFP is a functional TMBIM superfamily member with channel activity that can increase the permeability of bacterial cells to calcium.

Yeast Bax Inhibitor's Channel Function in *E. coli* Can Be Inhibited by Gadolinium.

To see if we could identify small-molecule inhibitors of the TMBIM superfamily of calcium channels using our *in vivo* bacterial system overexpressing Bxi1p-EGFP, we tested several ions for their ability to block calcium permeability in our cells, reasoning that cations

larger than calcium may be able to inhibit Ca^{2+} binding to Bxi1p. To do this, we repeated our overexpression experiments with an additional incubation step: Before the addition of external calcium, the larger cations were added to wells containing Fura-2AM loaded bacterial cells to see if they had any effect on the cell's permeability to Ca^{2+} .

Of several cations tested, we discovered that the gadolinium ion, Gd^{3+} , which is a potent blocker of calcium channels (Bourne & Trifaró, 1982; Biagi & Enyeart, 1990; Malasics et al., 2010) was able to block the permeability of calcium in our DH5 α cells with both basal and overexpressed levels of Bxi1p-EGFP (Figure 2A). Cells pre-exposed to Gd^{3+} did not manifest the calcium spike as measured by the R_{340}/R_{380} ratio of Fura-2 that was observed in non-treated cells. Importantly, the trivalent cation did not significantly decrease the proteins levels of Bxi1p-EGFP as determined by flow cytometry (Figure 2B), nor did it lead either to slow growth or to bacterial cell death (Figure 2C), suggesting that it is a *bona fide* inhibitor of Bxi1p-EGFP's function.

Human Bax Inhibitor's Channel Function in *E. coli* Can Also Be Inhibited by Gadolinium

To determine if the human homolog of Bax inhibitor (HsBI-1/TMBIM6) could function as a calcium channel in our bacterial *in vivo* assay, we overexpressed a protein HsBI1-EGFP fusion using the same arabinose-inducible araBAD promoter as was used with the yeast Bxi1p-EGFP homolog described above. As shown in Figure 3A, bacterial cells grown in media containing arabinose for six hours showed significant induction($p<0.05$) of the HsBI1-EGFP fusion protein. Upon addition of external calcium, intracellular calcium concentration as indicated by the R_{340}/R_{380} ratio of Fura-2 increased more rapidly in cells induced with arabinose as compared with uninduced controls (Figure 3B). This data suggests that HsBI1-EGFP is also a calcium channel when expressed in *E. coli*. This is not unexpected given the significant sequence

199 homology between human Bax inhibitor (hBI-1/TMBIM6) and its bacterial homolog, BsYetJ
200 (Chang et al., 2014).

201 Finally, we discovered that Gd^{3+} is also able to block the permeability of calcium in
202 bacterial cells with both basal and overexpressed levels of HsBI1-EGFP (Figure 4A). Once again,
203 the trivalent cation did not decrease the levels of Bxi1p-EGFP as measured by flow cytometry
204 (Figure 4B), nor did it lead either to slow growth or to bacterial cell death (Figure 4C), suggesting
205 that it is a *bona fide* inhibitor of HsBI1-EGFP's function.

Discussion

The budding yeast, *Saccharomyces cerevisiae*, has a single TMBIM family member (YNL305C) named Bxi1p/Ybh3p (Chae et al., 2003; Cebulski et al., 2011; Büttner et al., 2011). However, the precise function of this protein is disputed. In a previous paper, we showed that Bxi1p-GFP is localized to the ER, and that mutant yeast cells deleted of *BXII* are more susceptible to a range of pharmacological and environmental triggers that induce cell death, especially pharmacological triggers associated with the unfolded protein response (Cebulski et al., 2011). This pro-survival function for Bxi1p has been confirmed by two other laboratories (Chae et al., 2003; Teng et al., 2011), and is consistent with the anti-apoptotic function associated with the TMBIM superfamily.

In contrast, another publication has suggested that the protein encoded by YNL305C – which was named Ybh3p for yeast BH3-only protein – is a pro-apoptotic member of the BH3-only family of proteins that is able to bind *in vitro* with Bcl-X_L (Büttner et al., 2011). Moreover, the authors of this paper propose that Ybh3p translocates from the vacuole to the mitochondria to trigger BH3-domain dependent apoptosis. This would suggest that Bxi1p/Ybh3p has a pro-apoptotic function consistent with its proposed membership in the BH3-only family of proteins.

How do we resolve this discrepancy in the reported putative functions of Bxi1p/Ybh3p? It is notable that the inclusion of Bxi1p/Ybh3p in the BH3-only family of proteins has been criticized by those who have proposed that the BH3-domain-like sequence in this yeast ORF is not a *bona fide* BH3 domain (Aouacheria et al., 2013). These critics point out that the candidate yeast BH3 sequence in YNL305C, which was initially identified by visual inspection, is somewhat truncated as it is located at the C-terminus of the Ybh3p protein, and it overlaps one of six transmembrane segments, two unprecedented features among any other known BH3-containing

proteins. They also affirm that in principle, it is perplexing for budding yeast to have a BH-3 only protein, since it is not clear how this protein would have ever encountered human Bcl-X_L in the course of evolutionary history. Nonetheless, they concede that it is possible that the yeast genome may include a not-yet identified protein whose structure so closely resembles the 3D structure of the Bcl-2 family proteins that it could serve as a potential target of this yeast BH3 domain.

In this paper, we report that the overexpression of either Bxi1p/Ybh3p-EGFP and HsBI1-EGFP increases the permeability of bacterial cells to external calcium. This suggests that Bxi1p-EGFP, like its human counterpart, is a functional TMBIM superfamily member with calcium channel activity. It also suggests that Bxi1p-EGFP is not a member of the BH3-only family of proteins, which are known to be globular rather than transmembrane proteins (Glab, Mbogo & Puthalakath, 2017). However, we are still not sure how to reconcile this observation that Bxi1p is a transmembrane channel protein with the published data that suggests that Bxi1p/Ybh3p is a protein that translocates from the vacuole to the mitochondria to trigger cell death in a manner reminiscent of the BH3-only family of proteins (Büttner et al., 2011).

In addition, we show that the gadolinium ion, Gd³⁺, appears to inhibit both Bxi1p-EGFP and HsBI1-EGFP function in *E. coli*. To the best of our knowledge, this is the first known small-molecule inhibitor of a TMBIM superfamily member that works *in vivo*. However, it is not clear if Gd³⁺ blocks the channel's pore by binding to the Ca²⁺ binding site or if it alters the conformation of the protein itself to prevent Ca²⁺ binding and channel function.

Notably, as we were about to submit this manuscript for publication, a paper was published that showed that gadolinium inhibits the binding of ⁴⁵Ca²⁺ to the bacterial TMBIM homolog, BsYetJ, *in vitro* (Guo et al., 2019). This report also revealed that Yb³⁺ and Lu³⁺ were able to inhibit Ca²⁺ to the BsYetJ channel, albeit to a much lesser extent. As the authors acknowledged,

however, it is still not clear if this inhibition of channel function is the result of direct binding of the lanthanides to the Ca^{2+} binding site or of an induced conformational change in the channel's tertiary structure triggered by the larger cations.

Finally, given the observation that RNA interference of Bax inhibitor activity in the prostate cancer cell lines, PC-3, LNCaP, and DU-145, triggered cell death (Grzmil et al., 2003), it would be interesting to see if Gd^{3+} treatment of these cell lines would also trigger apoptosis by inhibiting the channel function of the protein. In principle, this would suggest that small molecule inhibitors of TMBIM function could be therapeutic in nature.

Conclusion

In this paper, we report that the overexpression of either Bxi1p/Ybh3p-EGFP and HsBI1-EGFP increases the permeability of bacterial cells to external calcium and that this activity can be inhibited by gadolinium *in vivo*. It suggests that Bxi1p like its human counterpart is a *bona fide* member of the TMBIM protein family of calcium channels. We are currently using our *in vivo* system to identify other small-molecular inhibitors of this clinically-important and highly conserved superfamily of proteins.

Figure Legends

FIGURE 1: Yeast Bax Inhibitor (Bxi1p/Ybh3p) is a Calcium Channel in *E. coli*. (A and B) We expressed a yeast Bax inhibitor-EGFP fusion protein in DH5 α bacterial cells using the arabinose-inducible araBAD promoter. Bacterial cells grown in LB media containing arabinose for six hours showed significant induction of the Bxi1p-EGFP fusion protein as measured by an Accuri C6 Flow Cytometer. (C) We then loaded these DH5 α cells with Fura-2AM, a membrane-permeable, fluorescent calcium indicator that has been used to determine the intracellular calcium dynamics of bacterial cells. Upon addition of external calcium, intracellular calcium concentration as indicated by the R340/R380 ratio of Fura-2 increased more rapidly in bacterial cells induced with arabinose as compared with uninduced controls. Error bars indicate standard deviations for trials with at least three independent cultures. The differences in expression and calcium levels were deemed statistically significant by the Student's t-test (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$).

FIGURE 2: Yeast Bax Inhibitor's Channel Function in *E. coli* Can Be Inhibited by Gadolinium. (A) Bacterial cells overexpressing the yeast Bax inhibitor, Bxi1p-EGFP fusion protein were grown in LB media with and without arabinose for six hours. Following the six-minute emission baseline measurements, the cells were loaded with 10 μ L of 500 mM GdCl₃. Upon addition of external calcium ten minutes later, the increase in intracellular calcium concentration as indicated by the R340/R380 ratio of Fura-2 was inhibited by Gd³⁺ treatment. Error bars indicate standard deviations for trials with at least three independent cultures. The differences in expression levels were deemed statistically significant by the Student's t-test (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$). (B) To determine if gadolinium inhibited

expression of the Bxi1p-EGFP fusion, levels of EGFP in bacterial cells grown in arabinose with and without gadolinium were compared with an Accuri C6 Flow Cytometer. Error bars indicate standard deviations for trials with at least three independent cultures. The differences in expression levels were deemed statistically significant by the Student's t-test (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$). (C) To determine whether or not gadolinium had any effect on bacterial cell growth, DH5 α *E. coli* cells transformed with the Bxi1p-EGFP expression vector were streaked onto LB-Ampicillin plates with varying concentrations of gadolinium ($GdCl_3$). The plates were incubated at 37°C for two days and imaged.

FIGURE 3: Human Bax Inhibitor (BI-1/TMBIM6) is a Calcium Channel in *E. coli*. (A and B)

We overexpressed a human Bax inhibitor, HsBI1-EGFP fusion protein in DH5 α bacterial cells using the arabinose-inducible araBAD promoter. Bacterial cells grown in LB media containing arabinose for six hours showed significant induction of the BI1-EGFP fusion protein as measured by an Accuri C6 Flow Cytometer. (C) We then loaded these DH5 α cells with Fura-2AM, a membrane-permeable, fluorescent calcium indicator that has been used to determine the intracellular calcium dynamics of bacterial cells. Upon addition of external calcium, intracellular calcium concentration as indicated by the R340/R380 ratio of Fura-2 increased more rapidly in bacterial cells induced with arabinose as compared with uninduced controls. Error bars indicate standard deviations for trials with at least three independent cultures. The differences in expression and calcium levels were deemed statistically significant by the Student's t-test (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$).

FIGURE 4: Human Bax Inhibitor's Channel Function in *E.coli* Can Be Inhibited by Gadolinium. (A) Bacterial cells overexpressing the human Bax inhibitor, HsBI1-EGFP fusion protein were grown in LB media with and without arabinose for six hours. Following the six-minute emission baseline measurements, the cells were loaded with 10 μ L of 500 mM GdCl₃. Upon addition of external calcium ten minutes later, the increase in intracellular calcium concentration as indicated by the R340/R380 ratio of Fura-2 was inhibited by Gd³⁺ treatment. Error bars indicate standard deviations for trials with at least three independent cultures. The differences in calcium levels were deemed statistically significant by the Student's t-test (* p<0.05; ** p<0.01; *** p<0.001; **** p<0.0001). (B) To determine if gadolinium inhibited expression of the HsBI1-EGFP fusion, levels of EGFP in bacterial cells grown in arabinose with and without gadolinium were compared with an Accuri C6 Flow Cytometer. Error bars indicate standard deviations for trials with at least three independent cultures. The differences in expression levels were deemed statistically significant by the Student's t-test (* p<0.05; ** p<0.01; *** p<0.001; **** p<0.0001). (C) To determine whether or not gadolinium had any effect on bacterial cell growth, DH5 α *E. coli* cells transformed with the HsBI1-EGFP expression vector were streaked onto LB-Ampicillin plates with varying concentrations of gadolinium (GdCl₃). The plates were incubated at 37°C for two days and imaged.

BIBLIOGRAPHY

- van 't Veer LJ., Dai H., van de Vijver MJ., He YD., Hart AAM., Mao M., Peterse HL., van der Kooy K., Marton MJ., Witteveen AT., Schreiber GJ., Kerkhoven RM., Roberts C., Linsley PS., Bernards R., Friend SH. 2002. Gene expression profiling predicts clinical outcome of breast cancer. *Nature* 415:530–536. DOI: 10.1038/415530a.
- Ahn T., Yun C-H., Chae H-JHZ., Kim H-R., Chae H-JHZ. 2009. Refolding and reconstitution of human recombinant Bax inhibitor-1 into liposomes from inclusion bodies expressed in *Escherichia coli*. *Protein expression and purification* 66:35–8. DOI: 10.1016/j.pep.2009.01.005.
- Ahn T., Yun C-H., Kim H-R., Chae H-J. 2010. Cardiolipin, phosphatidylserine, and BH4 domain of Bcl-2 family regulate Ca²⁺/H⁺ antiporter activity of human Bax inhibitor-1. *Cell calcium* 47:387–96. DOI: 10.1016/j.ceca.2010.02.003.
- Aouacheria A., Rech de Laval V., Combet C., Hardwick JM. 2013. Evolution of Bcl-2 homology motifs: homology versus homoplasy. *Trends in cell biology* 23:103–11. DOI: 10.1016/j.tcb.2012.10.010.
- Ausubel FM., Brent R., Kingston RE., Moore DD., Seidman JG., Smith JA., Struhl K. (eds.) 2002. *Short Protocols in Molecular Biology*. New York: John Wiley & Sons, Inc.
- Bailly-Maitre B., Fondevila C., Kaldas F., Droin N., Luciano F., Ricci J-E., Croxton R., Krajewska M., Zapata JM., Kupiec-Weglinski JW., Farmer D., Reed JC. 2006. Cytoprotective gene bi-1 is required for intrinsic protection from endoplasmic reticulum stress and ischemia-reperfusion injury. *Proceedings of the National Academy of Sciences of the United States of America* 103:2809–14. DOI: 10.1073/pnas.0506854103.
- Biagi BA., Enyeart JJ. 1990. Gadolinium blocks low- and high-threshold calcium currents in

pituitary cells. *American Journal of Physiology-Cell Physiology* 259:C515–C520. DOI:

10.1152/ajpcell.1990.259.3.C515.

Bourne GW., Trifaró JM. 1982. The gadolinium ion: A potent blocker of calcium channels and

catecholamine release from cultured chromaffin cells. *Neuroscience* 7:1615–1622. DOI:

10.1016/0306-4522(82)90019-7.

Büttner S., Ruli D., Vögtle F-N., Galluzzi L., Moitzi B., Eisenberg T., Kepp O., Habernig L.,

Carmona-Gutierrez D., Rockenfeller P., Laun P., Breitenbach M., Khoury C., Fröhlich K-U.,

Rechberger G., Meisinger C., Kroemer G., Madeo F. 2011. A yeast BH3-only protein

mediates the mitochondrial pathway of apoptosis. *The EMBO journal* 30:2779–92. DOI:

10.1038/emboj.2011.197.

Carrara G., Saraiva N., Gubser C., Johnson BF., Smith GL. 2012. Six-transmembrane topology for

Golgi anti-apoptotic protein (GAAP) and Bax inhibitor 1 (BI-1) provides model for the

transmembrane Bax inhibitor-containing motif (TMBIM) family. *The Journal of biological*

chemistry 287:15896–905. DOI: 10.1074/jbc.M111.336149.

Cebulski J., Malouin J., Pinches N., Cascio V., Austriaco N. 2011. Yeast Bax inhibitor, Bxi1p, is an

ER-localized protein that links the unfolded protein response and programmed cell death in

Saccharomyces cerevisiae. *PloS one* 6:e20882. DOI: 10.1371/journal.pone.0020882.

Chae H-J., Ke N., Kim H-R., Chen S., Godzik A., Dickman M., Reed JC. 2003. Evolutionarily

conserved cytoprotection provided by Bax Inhibitor-1 homologs from animals, plants, and

yeast. *Gene* 323:101–13.

Chae H-J., Kim H-R., Xu C., Bailly-Maitre B., Krajewska M., Krajewski S., Banares S., Cui J.,

Digicaylioglu M., Ke N., Kitada S., Monosov E., Thomas M., Kress CL., Babendure JR.,

Tsien RY., Lipton SA., Reed JC. 2004. BI-1 regulates an apoptosis pathway linked to

endoplasmic reticulum stress. *Molecular cell* 15:355–66. DOI: 10.1016/j.molcel.2004.06.038.

Chang Y., Bruni R., Kloss B., Assur Z., Kloppmann E., Rost B., Hendrickson WA., Liu Q. 2014. Structural basis for a pH-sensitive calcium leak across membranes. *Science (New York, N.Y.)* 344:1131–5. DOI: 10.1126/science.1252043.

Gamboa-Tuz SD., Pereira-Santana A., Zhao T., Schranz ME., Castano E., Rodriguez-Zapata LC. 2018. New insights into the phylogeny of the TMBIM superfamily across the three of life: Comparative genomics and synteny networks reveal independent evolution of the BI and LFG families in plants. 126:266–278. DOI: 10.1016/j.ympev.2018.04.032.

Gangola P., Rosen BP. 1987. Maintenance of intracellular calcium in *Escherichia coli*. *The Journal of biological chemistry* 262:12570–4.

Glab JA., Mbogo GW., Puthalakath H. 2017. BH3-Only Proteins in Health and Disease. *International review of cell and molecular biology* 328:163–196. DOI: 10.1016/bs.ircmb.2016.08.005.

Grzmil M., Kaulfuss S., Thelen P., Hemmerlein B., Schweyer S., Obenauer S., Kang TW., Burfeind P. 2006. Expression and functional analysis of Bax inhibitor-1 in human breast cancer cells. *The Journal of pathology* 208:340–9. DOI: 10.1002/path.1902.

Grzmil M., Thelen P., Hemmerlein B., Schweyer S., Voigt S., Mury D., Burfeind P. 2003. Bax inhibitor-1 is overexpressed in prostate cancer and its specific down-regulation by RNA interference leads to cell death in human prostate carcinoma cells. *The American journal of pathology* 163:543–52. DOI: 10.1016/S0002-9440(10)63682-6.

Guo G., Xu M., Chang Y., Luyten T., Seitaj B., Liu W., Zhu P., Bultynck G., Shi L., Quick M., Liu Q. 2019. Ion and pH Sensitivity of a TMBIM Ca²⁺ Channel. *Structure*. DOI: 10.1016/j.str.2019.03.003.

Guzman LM., Belin D., Carson MJ., Beckwith J. 1995. Tight regulation, modulation, and high-level expression by vectors containing the arabinose PBAD promoter. *Journal of bacteriology* 177:4121–30.

Hu L., Smith TF., Goldberger G. 2009. LFG: A candidate apoptosis regulatory gene family. *Apoptosis*. DOI: 10.1007/s10495-009-0402-2.

Hudson CA., Rojas JD., Sarvazyan N., Wesson DE., Martínez-Zaguilán R. 1998. Interactions between benzylamiloride and fura-2: studies in vitro and in cardiac myocytes. *Archives of biochemistry and biophysics* 356:25–34. DOI: 10.1006/abbi.1998.0749.

Kiviluoto S., Luyten T., Schneider L., Lisak D., Rojas-Rivera D., Welkenhuyzen K., Missaen L., De Smedt H., Parys JB., Hetz C., Methner A., Bultynck G. 2013. Bax Inhibitor-1-mediated Ca^{2+} leak is decreased by cytosolic acidosis. *Cell calcium* 54:186–92. DOI: 10.1016/j.ceca.2013.06.002.

Lisak DA., Schacht T., Enders V., Habicht J., Kiviluoto S., Schneider J., Henke N., Bultynck G., Methner A. 2015. The transmembrane Bax inhibitor motif (TMBIM) containing protein family: Tissue expression, intracellular localization and effects on the ER Ca^{2+} -filling state. *Biochimica et biophysica acta* 1853:2104–14. DOI: 10.1016/j.bbamcr.2015.03.002.

Liu Q. 2017. TMBIM-mediated Ca^{2+} homeostasis and cell death. *Biochimica et biophysica acta* 1864:850–857. DOI: 10.1016/j.bbamcr.2016.12.023.

Lu B., Li Y., Li H., Zhang Y., Xu J., Ren L., Fu S., Zhou Y. 2015. Bax inhibitor-1 is overexpressed in non-small cell lung cancer and promotes its progression and metastasis. *International journal of clinical and experimental pathology* 8:1411–8.

Luo C., Wang H., Liu Q., He W., Yuan L., Xu P. 2019. A genetically encoded ratiometric calcium sensor enables quantitative measurement of the local calcium microdomain in the

endoplasmic reticulum. *Biophysics Reports* 5:31–42. DOI: 10.1007/s41048-019-0082-6.

Malasics A., Boda D., Valiskó M., Henderson D., Gillespie D. 2010. Simulations of calcium channel block by trivalent cations: Gd 3+ competes with permeant ions for the selectivity filter. *Biochimica et Biophysica Acta (BBA) - Biomembranes* 1798:2013–2021. DOI: 10.1016/j.bbamem.2010.08.001.

Reimers K., Choi CY-U., Mau-Thek E., Vogt PM. 2006. Sequence analysis shows that Lifeguard belongs to a new evolutionarily conserved cytoprotective family. *International journal of molecular medicine* 18:729–34.

Rojas-Rivera D., Hetz C. 2015. TMBIM protein family: ancestral regulators of cell death. *Oncogene* 34:269–80. DOI: 10.1038/onc.2014.6.

Schmits R., Cochlovius B., Treitz G., Regitz E., Ketter R., Preuss K-D., Romeike BFM., Pfreundschuh M. 2002. Analysis of the antibody repertoire of astrocytoma patients against antigens expressed by gliomas. *International journal of cancer* 98:73–7.

Teng X., Cheng W-C., Qi B., Yu T-X., Ramachandran K., Boersma MD., Hattier T., Lehmann P V., Pineda FJ., Hardwick JM. 2011. Gene-dependent cell death in yeast. *Cell death & disease* 2:e188. DOI: 10.1038/cddis.2011.72.

Westphalen BC., Wessig J., Leypoldt F., Arnold S., Methner A. 2005. BI-1 protects cells from oxygen glucose deprivation by reducing the calcium content of the endoplasmic reticulum. *Cell death and differentiation* 12:304–6. DOI: 10.1038/sj.cdd.4401547.

Xu Q., Reed JC. 1998. Bax inhibitor-1, a mammalian apoptosis suppressor identified by functional screening in yeast. *Molecular cell* 1:337–46.

FIGURE 1: Yeast Bax Inhibitor (Bxi1p/Ybh3p) is a Calcium Channel in *E. coli*. (A and B) We expressed a yeast Bax inhibitor-EGFP fusion protein in DH5α bacterial cells using the arabinose-inducible araBAD promoter. Bacterial cells grown in LB media containing arabinose for six hours showed significant induction of the Bxi1p-EGFP fusion protein as measured by an Accuri C6 Flow Cytometer. (C) We then loaded these DH5α cells with Fura-2AM, a membrane-permeable, fluorescent calcium indicator that has been used to determine the intracellular calcium dynamics of bacterial cells. Upon addition of external calcium, intracellular calcium concentration as indicated by the R340/R380 ratio of Fura-2 increased more rapidly in bacterial cells induced with arabinose as compared with uninduced controls. Error bars indicate standard deviations for trials with at least three independent cultures. The differences in expression and

calcium levels were deemed statistically significant by the Student's t-test (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$).

FIGURE 2: Yeast Bax Inhibitor's Channel Function in *E. coli* Can Be Inhibited by

Gadolinium. (A) Bacterial cells overexpressing the yeast Bax inhibitor, Bxi1p-EGFP fusion protein were grown in LB media with and without arabinose for six hours. Following the six-minute emission baseline measurements, the cells were loaded with 10 μL of 500 mM $GdCl_3$. Upon addition of external calcium ten minutes later, the increase in intracellular calcium concentration as indicated by the R_{340}/R_{380} ratio of Fura-2 was inhibited by Gd^{3+} treatment. Error bars indicate standard deviations for trials with at least three independent cultures. The differences in expression levels were deemed statistically significant by the Student's t-test (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$). (B) To determine if gadolinium inhibited expression of the Bxi1p-EGFP fusion, levels of EGFP in bacterial cells grown in arabinose with and without gadolinium were compared with an Accuri C6 Flow Cytometer. Error bars indicate

standard deviations for trials with at least three independent cultures. The differences in expression levels were deemed statistically significant by the Student's t-test (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$). (C) To determine whether or not gadolinium had any effect on bacterial cell growth, DH5 α *E. coli* cells transformed with the Bxi1p-EGFP expression vector were streaked onto LB-Ampicillin plates with varying concentrations of gadolinium ($GdCl_3$). The plates were incubated at 37°C for two days and imaged.

FIGURE 3: Human Bax Inhibitor (BI-1/TMBIM6) is a Calcium Channel in *E. coli*. (A and B)

We overexpressed a human Bax inhibitor, HsBI1-EGFP fusion protein in DH5 α bacterial cells using the arabinose-inducible araBAD promoter. Bacterial cells grown in LB media containing arabinose for six hours showed significant induction of the BI1-EGFP fusion protein as measured by an Accuri C6 Flow Cytometer. (C) We then loaded these DH5 α cells with Fura-2AM, a membrane-permeable, fluorescent calcium indicator that has been used to determine the intracellular calcium dynamics of bacterial cells. Upon addition of external calcium, intracellular calcium concentration as indicated by the R_{340}/R_{380} ratio of Fura-2 increased more rapidly in

bacterial cells induced with arabinose as compared with uninduced controls. Error bars indicate standard deviations for trials with at least three independent cultures. The differences in expression and calcium levels were deemed statistically significant by the Student's t-test (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$).

FIGURE 4: Human Bax Inhibitor's Channel Function in *E.coli* Can Be Inhibited by Gadolinium. (A) Bacterial cells overexpressing the human Bax inhibitor, HsBI1-EGFP fusion protein were grown in LB media with and without arabinose for six hours. Following the six-minute emission baseline measurements, the cells were loaded with 10μL of 500 mM GdCl₃. Upon addition of external calcium ten minutes later, the increase in intracellular calcium concentration as indicated by the R₃₄₀/R₃₈₀ ratio of Fura-2 was inhibited by Gd³⁺ treatment. Error bars indicate standard deviations for trials with at least three independent cultures. The differences in calcium levels were deemed statistically significant by the Student's t-test (* p<0.05; ** p<0.01; *** p<0.001; **** p<0.0001). (B) To determine if gadolinium inhibited expression of the HsBI1-EGFP fusion, levels of EGFP in bacterial cells grown in arabinose with and without gadolinium were compared with an Accuri C6 Flow Cytometer. Error bars indicate standard deviations for trials with at least three independent cultures. The differences in expression levels

were deemed statistically significant by the Student's t-test (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$). (C) To determine whether or not gadolinium had any effect on bacterial cell growth, DH5 α E. coli cells transformed with the HsBI1-EGFP expression vector were streaked onto LB-Ampicillin plates with varying concentrations of gadolinium ($GdCl_3$). The plates were incubated at 37°C for two days and imaged.