Running title: Arabidopsis ERF109 regulates auxin transport

Arabidopsis ERF109 regulates auxin transport-related genes in root development

Rui Liu^{#1}, Xiao-Teng Cai^{#2*}, Ping-Xia Zhao¹, Ping Xu³, and Cheng-Bin Xiang^{1*}

¹School of Life Sciences and Division of Molecular & Cell Biophysics, Hefei National Science Center for Physical Sciences at the Microscale, University of Science and Technology of China, The Innovation Academy of Seed Design, Chinese Academy of Sciences, Hefei, Anhui Province 230027, China.

²Advanced Innovation Center for Tree Breeding by Molecular Design, Beijing University of Agriculture.

³Basic Forestry and Proteomics Research Center, Fujian Agriculture and Forestry University, Fuzhou City, Fujian Province 350002, China.

[#]Equally contributed.

*Corresponding authors:

Xiao-Teng Cai

Email: xtcai@mail.ustc.edu.cn

Cheng-Bin Xiang

Email: xiangcb@ustc.edu.cn

ABSTRACT

The transcription factor ERF109 acts as a crosstalk node between jasmonic acid signaling and auxin biosynthesis by directly regulating *YUC2* and *ASA1* during lateral root formation in Arabidopsis. However, whether ERF109 regulates the auxin transport remains unclear. Here we report a mechanism of ERF109-mediated auxin transport in root system. Through root transcriptome comparison between *erf109*, wild type, and *35S:ERF109*, we found that the genes *PIN2* and *PIN4*, encoding the major membrane-based efflux carriers of auxin, were enriched in the overexpression line. In the promoters of these auxin transport genes, GCC-box cis elements were found and potentially bound by ERF109. Moreover, *PID*, encoding a key regulator in polar auxin transport, was found upregulated in *35S:ERF109* and down regulated in *erf109*. Yeast-one-hybrid and chromatin immunoprecipitation assays showed that ERF109 directly bound to the GCC-box of *PIN2*, *PIN4*, and *PID*. Genetic analyses with double mutants confirmed the function of *ERF109* in the regulation of auxin transport in Arabidopsis roots. Taken together, our results show that ERF109 modulates auxin transport by directly regulating *PIN2*, *PIN4* and *PID*. This ERF109-mediated auxin transport likely works together with ERF109-mediated auxin synthesis to establish auxin maxima for lateral root initiation.

KEYWORDS

Arabidopsis root, *ERF109*, *PID*, *PIN2*, *PIN4*, JA signaling, auxin, auxin transport

INTRODUCTION

Plant root system is essential for nutrition absorption and mechanical functions. The elaboration of root architecture is crucial for its adaptation to environmental constraints (Ge *et al.*, 2019; Gutierrez-Alanis *et al.*, 2018; Julkowska *et al.*, 2017; Lavenus *et al.*, 2016; Nibau *et al.*, 2008; Scheres and Laskowski, 2016). The phytohormone auxin is well known as a key regulator in plant root configuration and development (Friml, 2003; Matthes *et al.*, 2019; Moller *et al.*, 2017; Moore *et al.*, 2017; Overvoorde *et al.*, 2010; Stoeckle *et al.*, 2018). The Trp-dependent primary auxin biosynthesis and metabolic pathway in Arabidopsis is well established (Zhao, 2012). Accumulating evidence supports that local auxin biosynthesis and its regulation play essential roles in plant development and responses to environmental cues (Zhao, 2018b). The synthesis, transport, and cascade signaling pathways of auxin orchestrate almost every aspect of root development, including primary root (PR) growth, lateral root (LR) initiation, and root tropic growth (Friml *et al.*, 2002b; Fukaki *et al.*, 2007; Miao *et al.*, 2018; Ruzicka and Hejatko, 2017; Spalding, 2013; Weiste *et al.*, 2017; Zhao, 2018a).

Auxin gradient created by polar auxin transport (PAT) in root system is fundamental for plant organogenesis and development (Band *et al.*, 2014; Benkova *et al.*, 2003; Paponov *et al.*, 2005). The active directional cell-to-cell movement of auxin, which is mediated by membrane-based influx and efflux carriers, controls auxin distribution in root system (Brumos *et al.*, 2018; Grigolon *et al.*, 2015; Overvoorde *et al.*, 2010; Petrasek and Friml, 2009). Collectively, there are three major auxin carriers that have been identified in Arabidopsis: PIN-FORMED (PIN) family (auxin efflux carriers), AUXIN RESISTANT 1/LIKE AUX1 (AUX1/LAX) family (auxin influx carriers), ATP-BINDING CASSETTE B (ABCB) family (Carrier *et al.*, 2008; Paponov *et al.*, 2005; Rutschow *et al.*, 2014; Tromas and Perrot-Rechenmann, 2010; Zhou and Luo, 2018).

The PIN family proteins facilitate auxin efflux from the cytoplasm to the apoplast (Adamowski and Friml, 2015; Blilou *et al.*, 2005; Friml, 2010; Keicher *et al.*, 2017; Prat *et al.*, 2018; Ren and Lin, 2015). PIN family has a total of 8 members. Each member has particular function and tissue-specific location. PIN proteins can be classified into the PM-localized group including PIN1–PIN4 and PIN7 that contributing to cell-to-cell

95 auxin transport, and the ER-localized group consisting of PIN5, PIN6 and PIN8 that
96 largely contribute to the regulation of the cellular auxin homeostasis (Wabnik *et al.*,
97 2011). PIN1 primarily regulates plant meristem tissue and organ development
98 process at the rootward faces of the stele (Huang *et al.*, 2010; O'Connor *et al.*, 2017;
99 Paponov *et al.*, 2005; Xi *et al.*, 2016). PIN2 is observed on the shootward faces of
100 epidermal cells, and regulates root gravitropism through the redistribution of auxin
101 among root tissue (Mendez-Bravo *et al.*, 2019; Muller *et al.*, 1998; Vieten *et al.*,
102 2005; Zhu *et al.*, 2017). PIN3 concentrates on the basal side of vascular cells as well
103 as the lateral side of pericycle cells in the elongation zone, whose main function is to
104 modulate differential growth and regulate tropic growth (Chen *et al.*, 2015; Friml *et*
105 *al.*, 2002b; Petrasek and Friml, 2009; Zhang *et al.*, 2013). PIN4 is observed on outer
106 layer cells and peripheral cells around the quiescent center (QC), and meanwhile has
107 a polar localization in the basal membranes of provascular cells, which regulates both
108 auxin homeostasis and patterning through sink-mediated auxin distribution in root
109 tips (Blilou *et al.*, 2005; Friml *et al.*, 2002a; Xi *et al.*, 2018). PIN7 is detected in the
110 meristem and elongation zone, and located at lateral and basal membranes of
111 provascular cells, which mainly mediates auxin efflux from the QC area and regulates
112 root tropism and early embryonic development (Band *et al.*, 2014; Blilou *et al.*, 2005;
113 Ki *et al.*, 2016; Ruiz Rosquete *et al.*, 2018). The ER-localized PIN5, PIN6 and PIN8 do
114 not transport auxin from cell to cell but regulates intracellular auxin homeostasis
115 and development (Cazzonelli *et al.*, 2013; Ding *et al.*, 2012; Mravec *et al.*, 2009).

116 AUX1/LAX membrane proteins were shown to play a significant role in the
117 regulation of PAT. There are four members (AUX1, LAX1, LAX2 and LAX3) in
118 Arabidopsis genome, which facilitate auxin influx from the apoplast to the cytoplasm
119 (Robert *et al.*, 2015; Rutschow *et al.*, 2014; Swarup and Peret, 2012). AUX1 mainly
120 regulates root hair development and gravitropism (Dindas *et al.*, 2018; Street *et al.*,
121 2016; Wakeel *et al.*, 2018). Many researches have implicated that AUX1 and LAX3 can
122 regulate lateral root development and apical hook formation (Jones *et al.*, 2009;
123 Porco *et al.*, 2016; Swarup *et al.*, 2005; Swarup and Peret, 2012; Wakeel *et al.*, 2018).

124 Members of the ABCB family are also involved in auxin transportation (Cho and
125 Cho, 2013; Kaneda *et al.*, 2011; Zhang *et al.*, 2018). Some other classes of membrane
126 protein, such as potassium and nitrate transporters, have been reported to play a

127 significant role in facilitating auxin movement as well (Krouk *et al.*, 2010;
128 Vicenteagullo *et al.*, 2004).

129 The PIN family proteins can be specifically regulated by a group of kinases, which
130 includes serine/threonine kinase PINOID (PID), PID2 and WAVY ROOT GROWTH
131 (WAG) proteins (Barbosa *et al.*, 2018; Grones *et al.*, 2018; Haga *et al.*, 2014; Weller *et al.*, 2017). Loss-of-function mutants of the *PID* display apical organogenesis defects
132 and the decline of auxin transport, which is similar to the *pin1* mutant (Benjamins *et al.*, 2001; Bennett *et al.*, 1995). The overexpression of *PID* causes the reduction of
133 lateral auxin transport due to the preferential stimulation of downward directed PAT
134 to the root apex by PID, which leads to the loss of auxin of root apex and peripheral
135 cells, and subsequently results in the reduction of lateral roots, shortened hypocotyl
136 and root, and the loss of gravitropism and phototropism (Benjamins *et al.*, 2001;
137 Haga and Sakai, 2015; Saini *et al.*, 2017a; Sukumar *et al.*, 2009; Willige and Chory,
138 2015). Recent studies have revealed that PID is a crucial contributor of the regulation
139 of apical-basal PIN polarity (Friml *et al.*, 2004; Kleine-Vehn *et al.*, 2009; Saini *et al.*,
140 2017b; Wang *et al.*, 2019; Zourelidou *et al.*, 2014). The mechanisms of PID-mediated
141 PIN polarity alteration have been unraveled (Christensen *et al.*, 2000; Dory *et al.*,
142 2018; Geldner *et al.*, 2003; Kleine-Vehn *et al.*, 2009; Michniewicz *et al.*, 2007).

145 We previously reported a drought resistance mutant *edt1* that displayed an
146 improved root system with more lateral roots, which was caused by an activated
147 expression of the homeodomain-START transcription factor EDT1/HDG11 (Yu *et al.*,
148 2008). We later found that the phytohormone jasmonic acid (JA) was increased in
149 *edt1* roots, consistent with root transcriptome data that show a number of JA
150 biosynthesis-related genes are directly upregulated by EDT1 (Cai *et al.*, 2015b).
151 Moreover, we identified the JA-responsive transcription factor ETHYLENE RESPONSE
152 FACTOR 109 (ERF109) from the same transcriptome analysis. ERF109 is a plant
153 specific transcription factor from the ERF family and acts as an important crosstalk
154 node integrating JA signaling into auxin biosynthesis (Cai *et al.*, 2014). *ERF109* is
155 responsive to JA signaling and promotes auxin biosynthesis by binding to the
156 promoters of *YUC2* and *ASA1*, and therefore affects lateral root formation. As a
157 typical ERF family protein, ERF109 contains one APETALA2 (AP2) domain and can
158 directly bind to the GCC-box (5'-GCCGCC-3') in the promoters of its target genes to

regulate their expressions (Barczak *et al.*, 2016; Cai *et al.*, 2014; Fujimoto *et al.*, 2000; Matsuo *et al.*, 2015). Meanwhile, ERF109 is responsive to a variety of biotic and abiotic redox stimulation and thus also named as Redox-responsive Transcription Factor 1 (RRTF1) (Khandelwal *et al.*, 2008; Matsuo *et al.*, 2015; Matsuo and Oelmüller, 2015; Wang *et al.*, 2008). Recent reports show that ERF109 is also involved in the activation of root stem cells under JA signal (Zhou *et al.*, 2019) and plant regeneration promoted by jasmonate-mediated wound signaling (Zhang *et al.*, 2019).

We previously noticed the reduction of auxin level in root tip of loss-of-function mutant *erf109*. However, *ERF109* is not expressed in primary root tip under normal conditions (Cai *et al.*, 2014), which strongly implicates that ERF109 may also regulate auxin transport. Here we report that ERF109 can also up regulate the transcript level of auxin transport-related *PIN2*, *PIN4*, and *PID*. Yeast-one-hybrid and ChIP-qPCR assays demonstrated that ERF109 can directly bind to GCC-boxes of *PID*, *PIN2* and *PIN4* to up regulate their expression, which is supported by genetic analyses. Our results demonstrate that ERF109 modulates lateral root formation by regulating auxin transport in addition to auxin biosynthesis we previously reported.

RESULTS

ERF109 affects the expression of auxin transport genes

To investigate whether ERF109 regulates auxin transport, we first confirmed the expression of *ERF109* and the phenotype of the *erf109* mutant, *ERF109* overexpression line with the wild type (Figure 1A and B). Then we measured the transcript level of auxin transport-related genes in the root tissue by quantitative RT-PCR. The results showed that the transcript level of *PIN1* was decreased in *erf109* mutant, but not increased in *ERF109* overexpression line significantly (Figure 1C). The transcription of *PIN2* and *PIN4* was significantly increased in *35S:ERF109*, but did not show obvious reduction in *erf109* mutant (Figure 1D and F). There is no significant change in the transcript levels of *PIN3*, *PIN7* and *AUX1* in *erf109* and *35S:ERF109* compared with that in wild type (Figure 1E, G and H).

We also observed the protein level and localization pattern of auxin transporters by using *GFP* reporter lines in *35S:ERF109* and wild type background. Both *PIN2:GFP*

190 and *PIN4:GFP* reporter showed higher GFP signals in the *35S:ERF109* line compared
191 with that in the wild type, but the localization pattern of PIN2 and PIN4 was
192 unchanged (Figure 2). Other auxin transport proteins had no significant change in the
193 expression level or localization pattern as shown by the reporter lines
194 *PIN1pro:PIN1:GFP*, *PIN3pro:PIN3:GFP*, *PIN7pro:PIN7:GFP*, *AUX1pro:AUX1:GFP* in the
195 wild type and *35S:ERF109* background (Supplemental Figure S1).

196 **ERF109 directly binds to the GCC-boxes in the promoter of *PIN2* and *PIN4***

197 Considering that ERF109 is a transcription factor, we speculate that ERF109 may
198 transcriptionally regulate the expression of *PIN2* and *PIN4*. To identify whether these
199 auxin transport genes are potential targets of ERF109, we searched the GCC-boxes in
200 the promoters and transcribed region of these auxin transport genes and found that
201 *PIN2* and *PIN4* contained GCC-box. *PIN2* has one GCC-box in the promoter and two
202 GCC-boxes in its transcribed region. *PIN4* contains two GCC-boxes in the transcribed
203 region (Figure 3A).

204 We performed yeast-one-hybrid assay to determine whether ERF109 could
205 directly bind to the GCC-boxes of *PIN2* and *PIN4*. 30bp DNA fragments contained
206 GCC-box in the middle were chosen from both the promoter and transcribed region
207 of *PIN2* and transcribed region of *PIN4*. The fragments A from *PIN2* and *PIN4*
208 displayed high affinities to ERF109 (Figure 3A and B). The results show that ERF109 is
209 able to bind to the GCC-boxes of *PIN2* and *PIN4* in yeast cells.

210 To confirm whether the interaction between ERF109 and the GCC-boxes of *PIN2*
211 and *PIN4* takes place *in planta*, chromatin immunoprecipitation (ChIP) assays were
212 performed using HA-ERF109 transgenic plants with anti-HA antibodies. These DNA
213 fragments enriched by anti-HA antibodies can be detected by PCR (Figure 3C).
214 Further analysis with quantitative PCR also confirmed the enrichment of GCC-box
215 fragments of *PIN2* and *PIN4* by ERF109 (Figure 3D and E). As the ChIP-PCR and ChIP-
216 qPCR results shown in Figure 3C-E, the *PIN2* and *PIN4* fragments containing the GCC-
217 boxes were significantly enriched by ERF109, which is consistent with yeast-one-
218 hybrid assay results. Thus, the specific binding of ERF109 to the GCC-boxes of *PIN2*
219 and *PIN4* was confirmed in Arabidopsis.

220 ***PID* is another target of ERF109**

221 Since PIN family auxin transporters can be phosphorylated by a group of kinases,

including serine/threonine kinase PINOID (PID), PID2 and WAVY ROOT GROWTH (WAG) proteins, we examined the transcript level of these genes and found that the expression of *PID* was down regulated in *erf109* (*erf109-ko*) and up regulated in *35S:ERF109* (*ERF109-OE*) compared with that in the wild type (Figure 4A). *PID* has a GCC-box in its promoter and two GCC-boxes in transcribed region (Figure 4B). To find out whether these GCC boxes can be bound by ERF109, we conducted yeast-one-hybrid assays and found that ERF109 was indeed able to bind to the *PID* B cis element (Figure 4C). To confirm whether this specific binding occurs in planta, we performed ChIP assays and found that the *PID* B fragment containing the GCC-box was significantly enriched by ERF109 in both ChIP-PCR and ChIP-qPCR (Figure 4D and E).

Genetic analyses of ERF109 function in *pin2*, *pin4* and *pid* mutant background

To detect whether *PIN2*, *PIN4* and *PID* function as target genes of ERF109, *35S:ERF109 pid*, *35S:ERF109 pin2* and *35S:ERF109 pin4* were generated by crosses (Supplemental Figure S2) and their root phenotypes were observed. Genetic assay results showed that the primary root length and lateral root number of *35S:ERF109 pin2*, *35S:ERF109 pin4* and *35S:ERF109 pid* were intermediate between male parent and female parent (Figure 5). These results demonstrate that *PIN2*, *PIN4* and *PID* are each partially responsible for the root phenotype of *35S:ERF109*, indicating that *PIN2*, *PIN4*, and *PID* are targets of ERF109.

DISCUSSION

Plant root system is characterized by its flexibility and adaptability, which allows the optimal adjustment according to environmental constraints (Moore *et al.*, 2017; Overvoorde *et al.*, 2010; Vieten *et al.*, 2005). Both auxin biosynthesis and transport are involved in the regulation of Arabidopsis root development (Du and Scheres, 2018; Morffy and Strader, 2018; Xu *et al.*, 2017). JA has been long recognized as a stress hormone, and can alter root architecture dramatically, but our knowledge about the underlying mechanisms remains limited (Ku *et al.*, 2018; Wasternack, 2007; Wasternack and Hause, 2013). Previous studies have showed that JA can be induced by a variety of environmental stimuli (Carvalhais *et al.*, 2015; de Ollas *et al.*,

253 2013; Gundlach *et al.*, 1992; Wasternack, 2007). The transcription factor
254 EDT1/HDG11 up-regulates JA biosynthesis and thus improves Arabidopsis lateral root
255 formation (Cai *et al.*, 2015a; Yu *et al.*, 2008). ERF109 is responsive to JA signaling and
256 integrates JA signal into auxin biosynthesis (Cai *et al.*, 2014). In this study, we further
257 unraveled the molecular mechanisms by which JA-responsive ERF109 regulates auxin
258 transport in addition to auxin biosynthesis by directly binding to the GCC boxes of
259 *PIN2*, *PIN4*, and *PID* and activating their expression. The increased *PID* stimulates *PIN*
260 activities. Collectively, auxin transport is enhanced.

ERF109 affects auxin transport by directly regulating *PID*, *PIN2* and *PIN4*.

262 Polar auxin transport is mainly mediated by *PIN* efflux carriers family and *AUX1*
263 influx carriers family (Band *et al.*, 2014; Peret *et al.*, 2012; Petrasek and Friml, 2009;
264 Vieten *et al.*, 2005). The transcript level of *PIN2* and *PIN4* increased about 2-fold in
265 *35S:ERF109* compared with that of wide type (Figure 1D, F). Given that ERF109 can
266 directly bind to the GCC-boxes *cis*-element, it was found that both *PIN2* and *PIN4*
267 contained GCC-box (Figure 3A). Through the Y1H and ChIP assays, ERF109 can bind to
268 the GCC-box *cis*-element of *PIN2* and *PIN4* *in vitro* and *in vivo* (Figure 3B-E). Further
269 genetic assay also confirmed that *PIN2* and *PIN4* were responsible for the phenotype
270 of *35S:ERF109* (Figure 5). However, there is no significant down-regulated expression
271 in *erf109* mutant (Figure 1D and F). Considering that the *ERF109* and *PIN2*, 4 have
272 different expression pattern and localization (Cai *et al.*, 2014; Paponov *et al.*, 2005;
273 Wisniewska *et al.*, 2006), *PIN2* and *PIN4* can be only altered under ectopic expression
274 of ERF109. Notably, we found a significant decline of the expression of *PIN1* in the
275 *erf109* line (Figure 1C). Due to that ERF109 mediate auxin biosynthesis in shoot and
276 root tissues (Cai *et al.*, 2014) and *PIN* genes can be regulated by auxin itself (Prat *et*
277 *al.*, 2018; Vieten *et al.*, 2005), it is possible that the down-regulated expression of
278 *PIN1* is caused by the decline of auxin level in *erf109* seedlings.

279 *PID* is a positive regulator of auxin efflux (Benjamins R, et al. 2001, Lee SH, et al.
280 2006, Zourelidou M, et al. 2014). It is notable that the *ERF109* and *PID* have similar
281 expression pattern. Both of them are expressed in neonatal vascular tissue, root tip
282 meristem, root primordial and weakly expressed in plant root tips (Benjamins R, et al.
283 2001, Lee SH, et al. 2006). *PID* was significantly down-regulated in the *erf109*
284 seedlings and up-regulated in *35S:ERF109* lines (Figure 4A). The GCC-box from the

PID displayed a high affinity to ERF109 (Figure 4B-E) and subsequent genetic experiments confirmed that *PID* is a downstream target of ERF109 (Figure 5A, D and G). Taken together, ERF109 can regulate *PID* by directly binding to the GCC *cis*-element in the transcribed region. It is conceivable that an activated expression of *PID* will enhance the phosphorylation of PIN2 and PIN4, leading to enhanced auxin transport.

291 **A working model for ERF109-regulated auxin homeostasis in roots**

292 Based on our results, we propose a working model for the JA-responsive *ERF109*
293 to regulate auxin homeostasis in roots (Figure 6). In this model, *ERF109* responds to
294 JA signaling. As a transcription factor, ERF109 directly binds to the GCC-boxes in the
295 promoters of *ASA1* and *YUC2* to promote auxin biosynthesis as we previously
296 showed (Cai *et al.*, 2014). *ASA1* is a key gene required for auxin biosynthesis under JA
297 signal in lateral root development (Sun *et al.*, 2009). On the other hand, ERF109 can
298 also directly bind to the GCC-box of *PID*, *PIN2* and *PIN4* to increase their transcript
299 levels, thus altering auxin transport. These results suggest that both auxin synthesis
300 and transport may work together to generate the auxin gradient in lateral root
301 formation in response to JA.

302

303 **MATERIALS AND METHODS**

304

305 **Materials and growth conditions**

306 *Arabidopsis thaliana* ecotype Columbia-0 (Col-0) was used as wild type.
307 Homozygous loss-function *erf109* mutant, *35S:ERF109* and *35S:HA-ERF109* lines were
308 previously described (Cai *et al.*, 2014). *PINpro:PIN:GFP* lines and *AUX1pro:AUX1:GFP*
309 were used as female parents in genetic analysis, while Col-0 and *35S:ERF109* were
310 used as male parents. The *pid* (CS9421), *pin2* (CS8058), *pin4-3* mutants were ordered
311 from Arabidopsis Biological Resource Center (ABRC).
312 Surface sterilization of Arabidopsis seeds was conducted with 10% bleach for 10 min
313 and seeds were washed five times in sterile environment subsequently. After keeping
314 in darkness for 3 days at 4°C, sterilized seeds were transferred and germinated on
315 vertical 1/2 Murashige and Skoog (MS) medium agar plates (1% (w/v) sucrose) and

grown in 22°C under 16-h light/8-h dark cycles.

Gene expression analysis

Gene expression analysis was conducted by RT-PCR and quantitative RT-PCR. Total RNA was isolated with TRIZOL reagent (Invitrogen) from 10-day-old Col-0, *erf109* and *35S:ERF109* seedlings. Then, cDNA was generated through Reverse transcription reaction with Prime Script RT reagent Kit (TaKaRa). After PCR amplification of cDNA, PCR products were examined on a 2% agarose gel stained with ethidium bromide (EB). Quantitative RT-PCR was carried out with the SYBR Premix Ex Taq II (TaKaRa) on Applied Biosystems Step One Real-Time PCR system. Experiment data was normalized to *Ubiquitin5* (*At3g62250*). The specific primers used in the examination of transcript levels of target genes were listed in Supplemental Table S1. The results of PCR were also examined on a 2% agarose gel stained with EB.

Confocal microscopy analysis

The GFP observation of root tissues with ZEISS710 confocal laser scanning microscope was described previously (Cai *et al.*, 2014). The GFP fluorescence intensity was quantified by Open Source software Image J (<https://imagej.nih.gov/ij/>) (Collins, 2007).

Bioinformatic analysis of GCC-box *cis*-elements

The GCC-box *cis*-elements were searched using Arabidopsis *cis*-regulatory element database (AtcisDB, <http://arabidopsis.med.ohio-state.edu/AtcisDB/>) (Yilmaz *et al.*, 2011). GCC-box *cis*-elements were searched in the promoter and transcription region of auxin transport related genes subsequently using TAIR - the Arabidopsis information resource (<http://www.arabidopsis.org>) for confirmation and supplement of the AtcisDB results.

Yeast-one-hybrid assay

Yeast-one-hybrid assay (Y1H) was previously described (Cai *et al.*, 2014). The effector plasmid pAD-ERF109 was generated by plasmid pAD-GAL4-2.1 (AD vector) with *ERF109*- coding region sequence cloned into it, which was used for the expression of fusion protein for DNA binding. The 30bp DNA fragments containing GCC-boxes from *PID*, *PIN2*, and *PIN4* were synthesized (Sangon of Shanghai, China) (Supplemental Table S1) with cohesive ends at each side. Then, these DNA fragments were annealed and ligated into reporter plasmid pHIS2 (BD vector) respectively. The

plasmid pHIS2-3*GCC-boxes was also generated by pHIS2 vector with three tandem repeat of GCC-box element. The empty vector of pAD and pHIS2 were used as negative control. Combinations of different AD and BD vectors were co-transformed into Y187 yeast strains. Then, Y187 strains were grown on SD/-Trp-Leu medium and transferred to SD/-Trp-Leu-His medium with 10 mM 3-aminotriazole (3-AT, Sigma) in gradient dilutions (1:10, 1:100 and 1:1,000). The growth state of yeast on SD/-Trp-Leu-His medium indicated the interaction between ERF109 and the corresponding cis-element.

ChIP assay

Chromatin immunoprecipitation (ChIP) assay was previously described (Cai *et al.*, 2014). The nucleus was isolated from 10-day-old wild type and *35S:HA-ERF109* seedlings and the chromatin was sonicated to fragments with various sizes (250 bp-1 kb). The DNA fragments bonded by HA-ERF109 were then enriched by anti-HA antibodies (1:100 for ChIP assay, HA-Tag, Mouse mAb, M20003, Abmart, Shanghai, China) from whole DNA fragment pool. After incubation with Protein A agarose beads (Millipore, USA), enriched DNA fragments used as the templates for PCR were eluted out and purified with phenol/chloroform (1:1, v/v). The enriched DNA fragments can be detected by PCR and quantified by quantitative-PCR with specific primers which were showed in Supplemental Table S1.

ACCESSION NUMBER

Sequence data from this article can be found in the Arabidopsis Genome Initiative or GenBank/EMBL databases under the following accession numbers: *ERF109*, At4g34410; *PID*, At2g34650; *PIN1*, AT1G73590; *PIN2*, At5g57090; *PIN3*, AT1G70940; *PIN4*, At2g01420; *PIN7*, AT1G23080; *AUX1*, AT2G38120; *EDT1/HDG11*, At1g73360; *UBQ5*, At3g62250.

SUPPORTING INFORMATION

Supplemental Figure S1. Expression level of auxin transport-related genes in *35S:ERF109* and wild type.

Supplemental Figure S2. Verification of *ERF109* expression level in the lines for genetic assay (*35S:ERF109 pid*, *35S:ERF109 pin2*, *35S:ERF109 pin4*).

380 Supplemental Table S1. Primers used in this study.

381

382 **ACKNOWLEDGEMENTS**

383 This work was supported by grants from the National Nature Science Foundation of
384 China (31572183) and the Ministry of Science and Technology of China
385 (2016YFD0100701, 2018ZX08009-11B, 2016ZX08005-004-003, 2016ZX08001003).

386 We thank Dr. Chuanyou Li, Institute of Genetics and Developmental Biology, CAS for
387 providing PIN-GFP marker lines

388

389 **AUTHORS CONTRIBUTION**

390 LR, XTC designed the research, performed the research, data analysis and
391 interpretation.

392 PXZ, PX performed the research, data analysis and interpretation.

393 LR wrote the manuscript and XTC revised the manuscript.

394 CBX designed the research, performed interpretation, edited the manuscript, and
395 supervised the project

396

397

REFERENCE

- Adamowski M, Friml J. 2015. PIN-dependent auxin transport: action, regulation, and evolution. *Plant Cell* **27**, 20-32.
- Band LR, Wells DM, Fozard JA, Ghetui T, French AP, Pound MP, Wilson MH, Yu L, Li W, Hijazi HI, Oh J, Pearce SP, Perez-Amador MA, Yun J, Kramer E, Alonso JM, Godin C, Vernoux T, Hodgman TC, Pridmore TP, Swarup R, King JR, Bennett MJ. 2014. Systems analysis of auxin transport in the Arabidopsis root apex. *Plant Cell* **26**, 862-875.
- Barbosa ICR, Hammes UZ, Schwechheimer C. 2018. Activation and Polarity Control of PIN-FORMED Auxin Transporters by Phosphorylation. *Trends Plant Sci* **23**, 523-538.
- Barczak A, Karpińska B, Czajkowska A, Filipecki M. 2016. Molecular insight into ERF109 function in light regulated retrograde signaling. *New Biotechnology* **33**, S164-S164.
- Benjamins R, Quint A, Weijers D, Hooykaas P, Offringa R. 2001. The PINOID protein kinase regulates organ development in Arabidopsis by enhancing polar auxin transport. *Development* **128**, 4057-4067.
- Benkova E, Michniewicz M, Sauer M, Teichmann T, Seifertova D, Jurgens G, Friml J. 2003. Local, efflux-dependent auxin gradients as a common module for plant organ formation. *Cell* **115**, 591-602.
- Bennett SRM, Alvarez J, Bossinger G, Smyth DR. 1995. Morphogenesis in Pinoid Mutants of Arabidopsis-Thaliana. *Plant Journal* **8**, 505-520.
- Blilou I, Xu J, Wildwater M, Willemssen V, Paponov I, Friml J, Heidstra R, Aida M, Palme K, Scheres B. 2005. The PIN auxin efflux facilitator network controls growth and patterning in Arabidopsis roots. *Nature* **433**, 39-44.
- Brumos J, Robles LM, Yun J, Vu TC, Jackson S, Alonso JM, Stepanova AN. 2018. Local Auxin Biosynthesis Is a Key Regulator of Plant Development. *Dev Cell* **47**, 306-318 e305.
- Cai XT, Xu P, Wang Y, Xiang CB. 2015a. Activated expression of AtEDT1/HDG11 promotes lateral root formation in Arabidopsis mutant edt1 by upregulating jasmonate biosynthesis. *J Integr Plant Biol*.
- Cai XT, Xu P, Wang Y, Xiang CB. 2015b. Activated expression of AtEDT1/HDG11 promotes lateral root formation in Arabidopsis mutant edt1 by upregulating jasmonate biosynthesis. *J Integr Plant Biol* **57**, 1017-1030.
- Cai XT, Xu P, Zhao PX, Liu R, Yu LH, Xiang CB. 2014. Arabidopsis ERF109 mediates cross-talk between jasmonic acid and auxin biosynthesis during lateral root formation. *Nat Commun* **5**, 5833.
- Carrier DJ, Bakar NT, Swarup R, Callaghan R, Napier RM, Bennett MJ, Kerr ID. 2008. The binding of auxin to the Arabidopsis auxin influx transporter AUX1. *Plant Physiol* **148**, 529-535.
- Carvalhais LC, Dennis PG, Badri DV, Kidd BN, Vivanco JM, Schenk PM. 2015. Linking Jasmonic Acid Signaling, Root Exudates, and Rhizosphere Microbiomes. *Molecular plant-microbe interactions : MPMI* **28**, 1049.
- Cazonelli CI, Vanstraelen M, Simon S, Yin K, Carron-Arthur A, Nisar N, Tarle G, Cuttriss AJ, Searle IR, Benkova E, Mathesius U, Masle J, Friml J, Pogson BJ. 2013. Role of the Arabidopsis PIN6 auxin transporter in auxin homeostasis and auxin-mediated development. *PLoS One* **8**, e70069.
- Chen Q, Liu Y, Maere S, Lee E, Van Isterdael G, Xie Z, Xuan W, Lucas J, Vassileva V, Kitakura S, Marhavy P, Wabnik K, Geldner N, Benkova E, Le J, Fukaki H, Grotewold E, Li C, Friml J, Sack F, Beeckman T, Vanneste S. 2015. A coherent transcriptional feed-forward motif model for mediating auxin-sensitive PIN3 expression during lateral root development. *Nat Commun* **6**, 8821.
- Cho M, Cho HT. 2013. The function of ABCB transporters in auxin transport. *Plant Signal Behav* **8**, e22990.
- Christensen SK, Dagenais N, Chory J, Weigel D. 2000. Regulation of auxin response by the protein kinase PINOID. *Cell* **100**, 469-478.
- Collins TJ. 2007. ImageJ for microscopy. *Biotechniques* **43**, 25.
- de Ollas C, Hernando B, Arbona V, Gomez-Cadenas A. 2013. Jasmonic acid transient accumulation is needed for abscisic acid increase in citrus roots under drought stress conditions. *Physiol Plant* **147**, 296-306.
- Dindas J, Scherzer S, Roelfsema MRG, von Meyer K, Muller HM, Al-Rasheid KAS, Palme K, Dietrich P, Becker D, Bennett MJ, Hedrich R. 2018. AUX1-mediated root hair auxin influx governs SCF(TIR1/AFB)-type Ca(2+) signaling. *Nat Commun* **9**, 1174.
- Ding Z, Wang B, Moreno I, Duplakova N, Simon S, Carraro N, Reemmer J, Pencik A, Chen X, Tejos R, Skupa P, Pollmann S, Mravec J, Petrasek J, Zazimalova E, Honys D, Rolcik J, Murphy A, Orellana A, Geisler M, Friml J. 2012. ER-localized auxin transporter PIN8 regulates auxin homeostasis and male gametophyte development in Arabidopsis. *Nat Commun* **3**, 941.

- 454 **Dory M, Hatzimasoura E, Kallai BM, Nagy SK, Jager K, Darula Z, Nadai TV, Meszaros T, Lopez-Juez E,**
- 455 **Barnabas B, Palme K, Bogre L, Ditengou FA, Doczi R.** 2018. Coevolving MAPK and PID phosphosites
- 456 indicate an ancient environmental control of PIN auxin transporters in land plants. *FEBS Lett* **592**, 89-
- 457 102.
- 458 **Du Y, Scheres B.** 2018. Lateral root formation and the multiple roles of auxin. *J Exp Bot* **69**, 155-167.
- 459 **Friml J.** 2003. Auxin transport - shaping the plant. *Curr Opin Plant Biol* **6**, 7-12.
- 460 **Friml J.** 2010. Subcellular trafficking of PIN auxin efflux carriers in auxin transport. *Eur J Cell Biol* **89**,
- 461 231-235.
- 462 **Friml J, Benkova E, Blilou I, Wisniewska J, Hamann T, Ljung K, Woody S, Sandberg G, Scheres B,**
- 463 **Jurgens G, Palme K.** 2002a. AtPIN4 mediates sink-driven auxin gradients and root patterning in
- 464 *Arabidopsis*. *Cell* **108**, 661-673.
- 465 **Friml J, Wisniewska J, Benkova E, Mendgen K, Palme K.** 2002b. Lateral relocation of auxin efflux
- 466 regulator PIN3 mediates tropism in *Arabidopsis*. *Nature* **415**, 806-809.
- 467 **Friml J, Yang X, Michniewicz M, Weijers D, Quint A, Tietz O, Benjamins R, Ouwerkerk PB, Ljung K,**
- 468 **Sandberg G, Hooykaas PJ, Palme K, Offringa R.** 2004. A PINOID-dependent binary switch in apical-
- 469 basal PIN polar targeting directs auxin efflux. *Science* **306**, 862-865.
- 470 **Fujimoto SY, Ohta M, Usui A, Shinshi H, Ohme-Takagi M.** 2000. *Arabidopsis* ethylene-responsive
- 471 element binding factors act as transcriptional activators or repressors of GCC box-mediated gene
- 472 expression. *Plant Cell* **12**, 393-404.
- 473 **Fukaki H, Okushima Y, Tasaka M.** 2007. Auxin-mediated lateral root formation in higher plants. *Int Rev*
- 474 *Cytol* **256**, 111-137.
- 475 **Ge Y, Fang X, Liu W, Sheng L, Xu L.** 2019. Adventitious lateral rooting: the plasticity of root system
- 476 architecture. *Physiol Plant* **165**, 39-43.
- 477 **Geldner N, Anders N, Wolters H, Keicher J, Kornberger W, Muller P, Delbarre A, Ueda T, Nakano A,**
- 478 **Jurgens G.** 2003. The *Arabidopsis* GNOM ARF-GEF mediates endosomal recycling, auxin transport, and
- 479 auxin-dependent plant growth. *Cell* **112**, 219-230.
- 480 **Grigolon S, Sollich P, Martin OC.** 2015. Modelling the emergence of polarity patterns for the
- 481 intercellular transport of auxin in plants. *J R Soc Interface* **12**.
- 482 **Grones P, Abas M, Hajny J, Jones A, Waidmann S, Kleine-Vehn J, Friml J.** 2018. PID/WAG-mediated
- 483 phosphorylation of the *Arabidopsis* PIN3 auxin transporter mediates polarity switches during
- 484 gravitropism. *Sci Rep* **8**, 10279.
- 485 **Gundlach H, Müller MJ, Kutchan TM, Zenk MH.** 1992. Jasmonic acid is a signal inducer in elicitor-
- 486 induced plant cell cultures. *Proc Natl Acad Sci U S A. Proceedings of the National Academy of Sciences*
- 487 **89**, 2389-2393.
- 488 **Gutierrez-Alanis D, Ojeda-Rivera JO, Yong-Villalobos L, Cardenas-Torres L, Herrera-Estrella L.** 2018.
- 489 Adaptation to Phosphate Scarcity: Tips from *Arabidopsis* Roots. *Trends Plant Sci* **23**, 721-730.
- 490 **Haga K, Hayashi K, Sakai T.** 2014. PINOID AGC kinases are necessary for phytochrome-mediated
- 491 enhancement of hypocotyl phototropism in *Arabidopsis*. *Plant Physiol* **166**, 1535-1545.
- 492 **Haga K, Sakai T.** 2015. PINOID functions in root phototropism as a negative regulator. *Plant Signaling*
- 493 *& Behavior* **10**, e998545.
- 494 **Huang F, Zago MK, Abas L, van Marion A, Galvan-Ampudia CS, Offringa R.** 2010. Phosphorylation of
- 495 conserved PIN motifs directs *Arabidopsis* PIN1 polarity and auxin transport. *Plant Cell* **22**, 1129-1142.
- 496 **Jones AR, Kramer EM, Kirsten K, Ranjan S, Bennett MJ, Lazarus CM, Ottoline LHM, Grierson CS.** 2009.
- 497 Auxin transport through non-hair cells sustains root-hair development. *Nature Cell Biology* **11**, 78-84.
- 498 **Julkowska MM, Koevoets IT, Mol S, Hoefsloot H, Feron R, Tester MA, Keurentjes JJB, Korte A, Haring**
- 499 **MA, de Boer GJ, Testerink C.** 2017. Genetic Components of Root Architecture Remodeling in Response
- 500 to Salt Stress. *Plant Cell* **29**, 3198-3213.
- 501 **Kaneda M, Schuetz M, Lin BS, Chanis C, Hamberger B, Western TL, Ehling J, Samuels AL.** 2011. ABC
- 502 transporters coordinately expressed during lignification of *Arabidopsis* stems include a set of ABCBs
- 503 associated with auxin transport. *Journal of Experimental Botany* **62**, 2063.
- 504 **Keicher J, Jaspert N, Weckermann K, Moller C, Throm C, Kintzi A, Oecking C.** 2017. *Arabidopsis* 14-3-3
- 505 epsilon members contribute to polarity of PIN auxin carrier and auxin transport-related development.
- 506 *Elife* **6**.
- 507 **Khandelwal A, Elvitigala T, Ghosh B, Quatrano RS.** 2008. *Arabidopsis* transcriptome reveals control
- 508 circuits regulating redox homeostasis and the role of an AP2 transcription factor. *Plant Physiol* **148**,
- 509 2050-2058.
- 510 **Ki D, Sasayama D, Cho HT.** 2016. The M3 Phosphorylation Site Is Required for Trafficking and

Biological Roles of PIN-FORMED1, 2, and 7 in Arabidopsis. *Front Plant Sci* **7**, 1479.

Kleine-Vehn J, Huang F, Naramoto S, Zhang J, Michniewicz M, Offringa R, Friml J. 2009. PIN auxin efflux carrier polarity is regulated by PINOID kinase-mediated recruitment into GNOM-independent trafficking in Arabidopsis. *Plant Cell* **21**, 3839-3849.

Krouk G, Lacombe B, Bielach A, Perrinewalker F, Malinska K, Mounier E, Hoyerova K, Tillard P, Leon S, Ljung K. 2010. Nitrate-regulated auxin transport by NRT1.1 defines a mechanism for nutrient sensing in plants. *Developmental Cell* **18**, 927.

Ku YS, Sintaha M, Cheung MY, Lam HM. 2018. Plant Hormone Signaling Crosstalks between Biotic and Abiotic Stress Responses. *Int J Mol Sci* **19**.

Lavenus J, Guyomarch S, Laplace L. 2016. PIN Transcriptional Regulation Shapes Root System Architecture. *Trends Plant Sci* **21**, 175-177.

Matsuo M, Johnson JM, Hieno A, Tokizawa M, Nomoto M, Tada Y, Godfrey R, Obokata J, Sherameti I, Yamamoto YY, Bohmer FD, Oelmüller R. 2015. High REDOX RESPONSIVE TRANSCRIPTION FACTOR1 Levels Result in Accumulation of Reactive Oxygen Species in Arabidopsis thaliana Shoots and Roots. *Mol Plant* **8**, 1253-1273.

Matsuo M, Oelmüller R. 2015. REDOX RESPONSIVE TRANSCRIPTION FACTOR1 is involved in age-dependent and systemic stress signaling. *Plant Signaling & Behavior* **10**, e1051279.

Matthes MS, Best NB, Robil JM, Malcomber S, Gallavotti A, McSteen P. 2019. Auxin EvoDevo: Conservation and Diversification of Genes Regulating Auxin Biosynthesis, Transport, and Signaling. *Mol Plant* **12**, 298-320.

Mendez-Bravo A, Ruiz-Herrera LF, Cruz-Ramirez A, Guzman P, Martinez-Trujillo M, Ortiz-Castro R, Lopez-Bucio J. 2019. CONSTITUTIVE TRIPLE RESPONSE1 and PIN2 act in a coordinate manner to support the indeterminate root growth and meristem cell proliferating activity in Arabidopsis seedlings. *Plant Sci* **280**, 175-186.

Miao ZQ, Zhao PX, Mao JL, Yu LH, Yuan Y, Tang H, Liu ZB, Xiang CB. 2018. HOMEBOX PROTEIN52 Mediates the Crosstalk between Ethylene and Auxin Signaling during Primary Root Elongation by Modulating Auxin Transport-Related Gene Expression. *Plant Cell* **30**, 2761-2778.

Michniewicz M, Zago MK, Abas L, Weijers D, Schweighofer A, Meskiene I, Heisler MG, Ohno C, Zhang J, Huang F, Schwab R, Weigel D, Meyerowitz EM, Luschnig C, Offringa R, Friml J. 2007. Antagonistic regulation of PIN phosphorylation by PP2A and PINOID directs auxin flux. *Cell* **130**, 1044-1056.

Moller BK, Xuan W, Beeckman T. 2017. Dynamic control of lateral root positioning. *Curr Opin Plant Biol* **35**, 1-7.

Moore S, Liu J, Zhang X, Lindsey K. 2017. A recovery principle provides insight into auxin pattern control in the Arabidopsis root. *Sci Rep* **7**, 43004.

Morffy NJ, Strader LC. 2018. Locally Sourced: Auxin Biosynthesis and Transport in the Root Meristem. *Dev Cell* **47**, 262-264.

Mravec J, Skupa P, Bailly A, Hoyerova K, Krecek P, Bielach A, Petrasek J, Zhang J, Gaykova V, Stierhof YD, Dobrev PI, Schwarzerova K, Rolcik J, Seifertova D, Luschnig C, Benkova E, Zazimalova E, Geisler M, Friml J. 2009. Subcellular homeostasis of phytohormone auxin is mediated by the ER-localized PIN5 transporter. *Nature* **459**, 1136-1140.

Muller A, Guan C, Galweiler L, Tänzler P, Huijser P, Marchant A, Parry G, Bennett M, Wisman E, Palme K. 1998. AtPIN2 defines a locus of Arabidopsis for root gravitropism control. *Embo Journal* **17**, 6903-6911.

Nibau C, Gibbs DJ, Coates JC. 2008. Branching out in new directions: the control of root architecture by lateral root formation. *New Phytol* **179**, 595-614.

O'Connor DL, Elton S, Ticchiarelli F, Hsia MM, Vogel JP, Leyser O. 2017. Cross-species functional diversity within the PIN auxin efflux protein family. *Elife* **6**.

Overvoorde P, Fukaki H, Beeckman T. 2010. Auxin control of root development. *Cold Spring Harb Perspect Biol* **2**, a001537.

Paponov IA, Teale WD, Trebar M, Blilou I, Palme K. 2005. The PIN auxin efflux facilitators: evolutionary and functional perspectives. *Trends Plant Sci* **10**, 170-177.

Peret B, Swarup K, Ferguson A, Seth M, Yang Y, Dhondt S, James N, Casimiro I, Perry P, Syed A, Yang H, Reemmer J, Venison E, Howells C, Perez-Amador MA, Yun J, Alonso J, Beemster GT, Laplace L, Murphy A, Bennett MJ, Nielsen E, Swarup R. 2012. AUX/LAX genes encode a family of auxin influx transporters that perform distinct functions during Arabidopsis development. *Plant Cell* **24**, 2874-2885.

Petrasek J, Friml J. 2009. Auxin transport routes in plant development. *Development* **136**, 2675-2688.

568 **Porco S, Larrieu A, Du Y, Gaudinier A, Goh T, Swarup K, Swarup R, Kuempers B, Bishopp A, Lavenus J,**
569 **Casimiro I, Hill K, Benkova E, Fukaki H, Brady SM, Scheres B, Peret B, Bennett MJ.** 2016. Lateral root
570 emergence in Arabidopsis is dependent on transcription factor LBD29 regulation of auxin influx carrier
571 LAX3. *Development* **143**, 3340-3349.
572 **Prat T, Hajny J, Grunewald W, Vasileva M, Molnar G, Tejos R, Schmid M, Sauer M, Friml J.** 2018.
573 WRKY23 is a component of the transcriptional network mediating auxin feedback on PIN polarity.
574 *PLoS Genet* **14**, e1007177.
575 **Ren H, Lin D.** 2015. ROP GTPase regulation of auxin transport in arabidopsis. *Mol Plant* **8**, 193-195.
576 **Robert HS, Grunewald W, Sauer M, Cannoot B, Soriano M, Swarup R, Weijers D, Bennett M,**
577 **Boutillier K, Friml J.** 2015. Plant embryogenesis requires AUX/LAX-mediated auxin influx. *Development*
578 **142**, 702-711.
579 **Ruiz Rosquete M, Waidmann S, Kleine-Vehn J.** 2018. PIN7 Auxin Carrier Has a Preferential Role in
580 Terminating Radial Root Expansion in Arabidopsis thaliana. *Int J Mol Sci* **19**.
581 **Rutschow HL, Baskin TI, Kramer EM.** 2014. The carrier AUXIN RESISTANT (AUX1) dominates auxin flux
582 into Arabidopsis protoplasts. *New Phytol* **204**, 536-544.
583 **Ruzicka K, Hejatk J.** 2017. Auxin transport and conjugation caught together. *J Exp Bot* **68**, 4409-4412.
584 **Saini K, Abdelgawad H, Markakis MN, Schoenaers S, Asard H, Prinsen E, Beemster GTS, Vissenberg K.**
585 2017a. Perturbation of Auxin Homeostasis and Signaling by PINOID Overexpression Induces Stress
586 Responses in Arabidopsis. *Front Plant Sci* **8**, 1308.
587 **Saini K, Markakis MN, Zdanio M, Balcerowicz DM, Beeckman T, De Veylder L, Prinsen E, Beemster**
588 **GTS, Vissenberg K.** 2017b. Alteration in Auxin Homeostasis and Signaling by Overexpression Of PINOID
589 Kinase Causes Leaf Growth Defects in Arabidopsis thaliana. *Front Plant Sci* **8**, 1009.
590 **Scheres B, Laskowski M.** 2016. Root patterning: it takes two to tangle. *J Exp Bot* **67**, 1201-1203.
591 **Spalding EP.** 2013. Diverting the downhill flow of auxin to steer growth during tropisms. *Am J Bot* **100**,
592 203-214.
593 **Stoeckle D, Thellmann M, Vermeer JE.** 2018. Breakout-lateral root emergence in Arabidopsis thaliana.
594 *Curr Opin Plant Biol* **41**, 67-72.
595 **Street IH, Mathews DE, Yamburkenko MV, Sorooshzadeh A, John RT, Swarup R, Bennett MJ, Kieber**
596 **JJ, Schaller GE.** 2016. Cytokinin acts through the auxin influx carrier AUX1 to regulate cell elongation in
597 the root. *Development* **143**, 3982-3993.
598 **Sukumar P, Edwards KS, Rahman A, Delong A, Muday GK.** 2009. PINOID kinase regulates root
599 gravitropism through modulation of PIN2-dependent basipetal auxin transport in Arabidopsis. *Plant*
600 *Physiol* **150**, 722-735.
601 **Sun J, Xu Y, Ye S, Jiang H, Chen Q, Liu F, Zhou W, Chen R, Li X, Tietz O, Wu X, Cohen JD, Palme K, Li C.**
602 2009. Arabidopsis ASA1 is important for jasmonate-mediated regulation of auxin biosynthesis and
603 transport during lateral root formation. *Plant Cell* **21**, 1495-1511.
604 **Swarup R, Kramer EM, Perry P, Knox K, Leyser HM, Haseloff J, Beemster GT, Bhalerao R, Bennett MJ.**
605 2005. Root gravitropism requires lateral root cap and epidermal cells for transport and response to a
606 mobile auxin signal. *Nat Cell Biol* **7**, 1057-1065.
607 **Swarup R, Peret B.** 2012. AUX/LAX family of auxin influx carriers-an overview. *Front Plant Sci* **3**, 225.
608 **Tomas A, Perrot-Rechenmann C.** 2010. Recent progress in auxin biology. *C R Biol* **333**, 297-306.
609 **Vicenteagullo F, Rigas S, Desbrosses G, Dolan L, Hatzopoulos P, Grabov A.** 2004. Potassium carrier
610 TRH1 is required for auxin transport in Arabidopsis roots. *Plant Journal for Cell & Molecular Biology* **40**,
611 523-535.
612 **Vieten A, Vanneste S, Wisniewska J, Benkova E, Benjamins R, Beeckman T, Luschnig C, Friml J.** 2005.
613 Functional redundancy of PIN proteins is accompanied by auxin-dependent cross-regulation of PIN
614 expression. *Development* **132**, 4521-4531.
615 **Wabnick K, Kleine-Vehn J, Govaerts W, Friml J.** 2011. Prototype cell-to-cell auxin transport mechanism
616 by intracellular auxin compartmentalization. *Trends Plant Sci* **16**, 468-475.
617 **Wakeel A, Ali I, Upreti S, Azizullah A, Liu B, Khan AR, Huang L, Wu M, Gan Y.** 2018. Ethylene mediates
618 dichromate-induced inhibition of primary root growth by altering AUX1 expression and auxin
619 accumulation in Arabidopsis thaliana. *Plant Cell Environ* **41**, 1453-1467.
620 **Wang P, Shen L, Guo J, Jing W, Qu Y, Li W, Bi R, Xuan W, Zhang Q, Zhang W.** 2019. Phosphatidic Acid
621 Directly Regulates PINOID-Dependent Phosphorylation and Activation of the PIN-FORMED2 Auxin
622 Efflux Transporter in Response to Salt Stress. *Plant Cell* **31**, 250-271.
623 **Wang Z, Cao G, Wang X, Miao J, Liu X, Chen Z, Qu LJ, Gu H.** 2008. Identification and characterization
624 of COI1-dependent transcription factor genes involved in JA-mediated response to wounding in

Arabidopsis plants. *Plant Cell Rep* **27**, 125-135.

Wasternack C. 2007. Jasmonates: an update on biosynthesis, signal transduction and action in plant stress response, growth and development. *Ann Bot* **100**, 681-697.

Wasternack C, Hause B. 2013. Jasmonates: biosynthesis, perception, signal transduction and action in plant stress response, growth and development. An update to the 2007 review in *Annals of Botany*. *Ann Bot* **111**, 1021-1058.

Weiste C, Pedrotti L, Selvanayagam J, Muralidhara P, Froschel C, Novak O, Ljung K, Hanson J, Droge-Laser W. 2017. The Arabidopsis bZIP11 transcription factor links low-energy signalling to auxin-mediated control of primary root growth. *PLoS Genet* **13**, e1006607.

Weller B, Zourelidou M, Frank L, Barbosa IC, Fastner A, Richter S, Jurgens G, Hammes UZ, Schwechheimer C. 2017. Dynamic PIN-FORMED auxin efflux carrier phosphorylation at the plasma membrane controls auxin efflux-dependent growth. *Proc Natl Acad Sci U S A* **114**, E887-E896.

Willige BC, Chory J. 2015. A current perspective on the role of AGCVIII kinases in PIN-mediated apical hook development. *Frontiers in Plant Science* **6**, 767.

Wisniewska J, Xu J, Seifertova D, Brewer PB, Ruzicka K, Blilou I, Rouquie D, Benkova E, Scheres B, Friml J. 2006. Polar PIN localization directs auxin flow in plants. *Science* **312**, 883.

Xi D, Chen X, Wang Y, Zhong R, He J, Shen J, Ming F. 2018. Arabidopsis ANAC092 regulates auxin-mediated root development by binding to the ARF8 and PIN4 promoters. *J Integr Plant Biol*.

Xi W, Gong X, Yang Q, Yu H, Liou YC. 2016. Pin1At regulates PIN1 polar localization and root gravitropism. *Nat Commun* **7**, 10430.

Xu D, Miao J, Yumoto E, Yokota T, Asahina M, Watahiki M. 2017. YUCCA9-Mediated Auxin Biosynthesis and Polar Auxin Transport Synergistically Regulate Regeneration of Root Systems Following Root Cutting. *Plant Cell Physiol* **58**, 1710-1723.

Yilmaz A, Mejia-Guerra MK, Kurz K, Liang X, Welch L, Grotewold E. 2011. AGRIS: the Arabidopsis Gene Regulatory Information Server, an update. *Nucleic Acids Res* **39**, D1118-1122.

Yu H, Chen X, Hong YY, Wang Y, Xu P, Ke SD, Liu HY, Zhu JK, Oliver DJ, Xiang CB. 2008. Activated expression of an Arabidopsis HD-START protein confers drought tolerance with improved root system and reduced stomatal density. *Plant Cell* **20**, 1134-1151.

Zhang G, Zhao F, Chen L, Pan Y, Sun L, Bao N, Zhang T, Cui C-X, Qiu Z, Zhang Y, Yang L, Xu L. 2019. Jasmonate-mediated wound signaling promotes plant regeneration. *Nat. Plants* **In press**.

Zhang KX, Xu HH, Yuan TT, Zhang L, Lu YT. 2013. Blue-light-induced PIN3 polarization for root negative phototropic response in Arabidopsis. *Plant J* **76**, 308-321.

Zhang Y, Nasser V, Pisanty O, Omary M, Wulff N, Di Donato M, Tal I, Hauser F, Hao P, Roth O, Fromm H, Schroeder JI, Geisler M, Nour-Eldin HH, Shani E. 2018. A transportome-scale amiRNA-based screen identifies redundant roles of Arabidopsis ABCB6 and ABCB20 in auxin transport. *Nat Commun* **9**, 4204.

Zhao Y. 2018a. Essential Roles of Local Auxin Biosynthesis in Plant Development and in Adaptation to Environmental Changes. *Annu Rev Plant Biol* **69**, 417-435.

Zhao YD. 2012. Auxin Biosynthesis: A Simple Two-Step Pathway Converts Tryptophan to Indole-3-Acetic Acid in Plants. *Molecular Plant* **5**, 334-338.

Zhao YD. 2018b. Essential Roles of Local Auxin Biosynthesis in Plant Development and in Adaptation to Environmental Changes. *Annual Review of Plant Biology*, Vol 69 **69**, 417-435.

Zhou JJ, Luo J. 2018. The PIN-FORMED Auxin Efflux Carriers in Plants. *Int J Mol Sci* **19**.

Zhou W, Lozano-Torres JL, Blilou I, Zhang X, Zhai Q, Smant G, Li C, Scheres B. 2019. A Jasmonate Signaling Network Activates Root Stem Cells and Promotes Regeneration. *Cell*.

Zhu R, Dong X, Hao W, Gao W, Zhang W, Xia S, Liu T, Shang Z. 2017. Heterotrimeric G Protein-Regulated Ca(2+) Influx and PIN2 Asymmetric Distribution Are Involved in Arabidopsis thaliana Roots' Avoidance Response to Extracellular ATP. *Front Plant Sci* **8**, 1522.

Zourelidou M, Absmanner B, Weller B, Barbosa IC, Willige BC, Fastner A, Streit V, Port SA, Colcombet J, de la Fuente van Bentem S, Hirt H, Kuster B, Schulze WX, Hammes UZ, Schwechheimer C. 2014. Auxin efflux by PIN-FORMED proteins is activated by two different protein kinases, D6 PROTEIN KINASE and PINOID. *Elife* **3**.

FIGURE LEGENDS

Figure 1. ERF109 regulates the expression of auxin transport genes

A. Confirming the expression level of *ERF109* by RT-PCR. RNA samples isolated from 10-d-old seedlings of *erf109*, *35S:ERF109* and wild type plants were analyzed by quantitative RT-PCR.

B. Confirming the phenotype of 7-day-old *35S:ERF109* plants. *35S:ERF109* seedlings had shorter primary root and more lateral roots than the wild type. Scale bar = 1 cm.

C-H. Quantitative RT-PCR analysis of the transcript levels of auxin transport genes. Transcript level of *PIN1* (C), *PIN2* (D), *PIN3* (E), *PIN4* (F), *PIN7* (G), and *AUX1* (H) in the wild type, *erf109*, and *35S:ERF109* seedlings. RNA was isolated from the two-week-old seedlings. Values are mean $\pm$ SD (n=3 experiments, *P<0.05, **P<0.01). Asterisks indicate Student's t-test significant differences.

Figure 2. Fluorescence intensity of PIN2-GFP and PIN4-GFP was elevated in ERF109 over-expression line

PIN2- and PIN4-GFP reporters were introduced in the wild type and *35S:ERF109* by genetic crosses [*35S:ERF109* $\times$ *PIN2pro:PIN2:GFP*] (ERF109 PIN2, A), wild type $\times$ *PIN2pro:PIN2:GFP* (WT PIN2, B), *35S:ERF109* $\times$ *PIN4pro:PIN4:GFP* (ERF109 PIN4, C), and wild type $\times$ *PIN4pro:PIN4:GFP* (WT PIN4, D)]. The GFP signals were observed using 7-day-old seedlings on a confocal microscope. Bar=50 μ M. Three independent lines were analyzed for wild type and *35S:ERF109* background, and consistent results were obtained. GFP intensity of PIN2 (E) and PIN4 (F) was quantified by Image J (<https://imagej.nih.gov/ij/>). Values are mean $\pm$ SD (n=3 experiments, **P<0.01). Asterisks indicate Student's t-test significant differences.

Figure 3. ERF109 directly regulates the transcription of *PIN2* and *PIN4*.

A. The schematic view of the locations of GCC-boxes (inverted red triangles) in the promoters and transcription regions of *PIN2* and *PIN4* as well as the primers used for ChIP-PCR (blue lines for fragments contained GCC-box and green lines for fragments without GCC-box worked as negative control).

709 **B.** Yeast-one-hybrid assay for ERF109 binding to the GCC-box of *PIN2* and *PIN4*. 30bp
710 fragments contained GCC-box were chosen for BD vector constructions. A serial yeast
711 dilutions (1:1, 1:10, 1:100 and 1:1000) were grown on SD medium lacking Leu and
712 Trp (SD -Leu-Trp) and transferred to SD medium lacking Leu, Trp and His + 3-AT (SD -
713 Leu-Trp-His) respectively. The empty AD (pGADT7) and BD (pHis2) vectors were
714 used as negative controls.

715 **C.** ChIP-PCR assay for the binding between ERF109 and GCC-boxes of *PIN2* and *PIN4*.
716 DNA fragments with HA-ERF109 were precipitated from input DNA with anti-HA
717 antibodies or with no antibody. The enrichment of DNA fragments was determined
718 by PCR. The regions of *UBQ*, *PIN2* and *PIN4* that do not contain GCC-box were used
719 as negative controls.

720 **D-E.** Quantitative PCR analysis for ChIP assay. The enrichments of *PIN2* (D) and *PIN4*
721 (E) fragments contained GCC-box were confirmed by Quantitative-PCR. Values are
722 mean $\pm$ SD (n=3 experiments, ***P<0.001). Asterisks indicate Student's t-test
723 significant differences.

724

725 **Figure 4. *PID* is another direct target of ERF109**

726 **A.** Expression levels of *PID* by quantitative RT-PCR. RNA samples were isolated from
727 7-d-old seedlings of *erf109* knockout mutant (*erf109 ko*), wild type (WT), and the
728 overexpression line *35S:ERF109* (*ERF109-OE*). Values are mean $\pm$ SD (n=3
729 experiments, **P<0.01, ***P<0.001). Asterisks indicate Student's t-test significant
730 differences.

731 **B.** The schematic view of the location of GCC-box *cis*-element (inverted red triangle)
732 in the promoter and transcribed region of *PID* as well as the primers used for PCR
733 (blue lines for fragments contained GCC-box and green lines for fragments without
734 GCC-box worked as negative control).

735 **C.** Yeast-one-hybrid assay for ERF109 binding to the GCC-box *cis*-element of *PID*. The
736 assay was performed as same as figure 3B.

737 **D.** ChIP-PCR assay was conducted using *35S:HA-ERF109* transgenic seedlings and anti-
738 HA antibodies. The assay was performed as for Figure 3C.

739 **E.** Quantitative PCR analysis for ChIP assay. The enrichment of *PID* promoter

fragment was confirmed by quantitative PCR. Values are mean $\pm$ SD (n=3 experiments, ***P<0.001). Asterisks indicate Student's t-test significant differences.

Figure 5. Genetic analyses of ERF109 in *pid*, *pin2* and *pin4* mutant background

A-C. Root phenotype. Seeds of indicated lines were separately germinated on MS for 12 days before photographs were taken (Scale bar =1cm).

D-F. The primary root length was measured at the indicated time points. Values are mean $\pm$ SD (n=30 seedlings, *P<0.05, **P<0.01). Statistically significant differences were calculated based on the Student's t-tests.

G-I. The numbers of LR (per cm) of different plants grown on MS were counted at the indicated time points. Values are mean $\pm$ SD (n=30 seedlings, *P<0.05, **P<0.01). Statistically significant differences were calculated based on the Student's t-tests.

Figure 6. A working model of ERF109 in auxin homeostasis

JA-responsive ERF109 upregulates auxin biosynthesis by directly binding to GCC-boxes in the promoters of *ASA1* and *YUC2*. ERF109 also directly bind to the GCC-boxes *cis*-elements in *PIN2*, *PIN4*, and *PID*. Elevated PID enhances PINs phosphorylation and stimulates auxin transport. Therefore, ERF109 regulates root development through both auxin biosynthesis and transport.

Figure 1. ERF109 regulates the expression of auxin transport genes

A. Confirming the expression level of ERF109 by RT-PCR. RNA samples isolated from 10-d-old seedlings of erf109, 35S:ERF109 and wild type plants were analyzed by quantitative RT-PCR.

B. Confirming the phenotype of 7-day-old 35S:ERF109 plants. 35S:ERF109 seedlings had shorter primary root and more lateral roots than the wild type. Scale bar = 1 cm.

Figure 2. Fluorescence intensity of PIN2-GFP and PIN4-GFP was elevated in ERF109 over-expression line PIN2- and PIN4-GFP reporters were introduced in the wild type and 35S:ERF109 by genetic crosses [35S:ERF109 ♂ × PIN2pro:PIN2:GFP ♀ (ERF109 PIN2, A), wild type ♂ × PIN2pro:PIN2:GFP ♀ (WT PIN2, B), 35S:ERF109 ♂ × PIN4pro:PIN4:GFP ♀ (ERF109 PIN4, C), and wild type ♂ × PIN4pro:PIN4:GFP ♀ (WT PIN4, D)]. The GFP signals were observed using 7-day-old seedlings on a confocal microscope. Bar=50 μ M. Three independent lines were analyzed for wild type and 35S:ERF109 background, and consistent results were obtained. GFP intensity of PIN2 (E) and PIN4 (F) was quantified by Image J (<https://imagej.nih.gov/ij/>). Values are mean $\pm$ SD (n=3 experiments, **P<0.01). Asterisks indicate Student's t-test significant differences.

Figure 3. ERF109 directly regulates the transcription of PIN2 and PIN4.

A. The schematic view of the locations of GCC-boxes (inverted red triangles) in the promoters and transcription regions of PIN2 and PIN4 as well as the primers used for ChIP-PCR (blue lines for fragments contained GCC-box and green lines for fragments without GCC-box worked as negative control).

B. Yeast-one-hybrid assay for ERF109 binding to the GCC-box of PIN2 and PIN4. 30bp fragments contained GCC-box were chosen for BD vector constructions. A serial yeast dilutions (1:1, 1:10, 1:100 and 1:1000) were grown on SD medium lacking Leu and Trp (SD -Leu-Trp) and transferred to SD medium lacking Leu, Trp and His + 3-AT (SD -Leu-Trp-His) respectively. The empty AD (pGADT7) and BD (pHis2) vectors were used as negative controls.

C. ChIP-PCR assay for the binding between ERF109 and GCC-boxes of PIN2 and PIN4. DNA fragments with HA-ERF109 were precipitated from input DNA with anti-HA antibodies or with no antibody. The enrichment of DNA fragments was determined by PCR. The regions of UBQ, PIN2 and PIN4 that do not contain GCC-box were used as negative controls.

D-E. Quantitative PCR analysis for ChIP assay. The enrichments of PIN2 (D) and PIN4 (E) fragments contained GCC-box were confirmed by

Figure 4. PID is another direct target of ERF109

A. Expression levels of PID by quantitative RT-PCR. RNA samples were isolated from 7-d-old seedlings of *erf109* knockout mutant (*erf109* ko), wild type (WT), and the overexpression line 35S:ERF109 (ERF109-OE). Values are mean $\pm$ SD (n=3 experiments, **P<0.01, ***P<0.001). Asterisks indicate Student's t-test significant differences.

B. The schematic view of the location of GCC-box cis-element (inverted red triangle) in the promoter and transcribed region of PID as well as the primers used for PCR (blue lines for fragments contained GCC-box and green lines for fragments without GCC-box worked as negative control).

C. Yeast-one-hybrid assay for ERF109 binding to the GCC-box cis-element of PID. The assay was performed as same as figure 3B.

D. ChIP-PCR assay was conducted using 35S:HA-ERF109 transgenic seedlings and anti-HA antibodies (Abmart Shanghai). The assay was

Figure 5. Genetic analyses of ERF109 in *pid*, *pin2* and *pin4* mutant background

A-C. Root phenotype. Seeds of indicated lines were separately germinated on MS for 12 days before photographs were taken (Scale bar =1cm).

D-F. The primary root length was measured at the indicated time points. Values are mean $\pm$ SD (n=30 seedlings, *P<0.05, **P<0.01). Statistically significant differences were calculated based on the Student's t-tests.

G-I. The numbers of LR (per cm) of different plants grown on MS were counted at the indicated time points. Values are mean $\pm$ SD (n=30 seedlings, *P<0.05, **P<0.01). Statistically significant differences were calculated based on the Student's t-tests.

Figure 6. A working model of ERF109 in auxin homeostasis
JA-responsive ERF109 upregulates auxin biosynthesis by directly binding to GCC-boxes in the promoters of ASA1 and YUC2. ERF109 also directly bind to the GCC-boxes cis-elements in PIN2, PIN4, and PID. Elevated PID enhances PINs phosphorylation and stimulates auxin transport. Therefore, ERF109 regulates root development through both auxin biosynthesis and transport.