Patterns of phenotypic plasticity among populations of three Mediterranean pine species and implications for evolutionary responses to climate change

List of Authors

Natalia Vizcaíno-Palomar^{1*}, Bruno Fady², Ricardo Alía³, Annie Raffin⁴, Sven Mutke³, Marta Benito Garzón¹

Institutional affiliations

¹ BIOGECO, INRA, Univ. Bordeaux, 33615 Pessac, France

² INRA Unité de Recherches Ecologie des Forêts Méditerranéennes (URFM), Avignon, France

³ INIA, Forest Research Centre & iuFOR UVa-INIA, Ctra La Coruña km 7.5, 28040 Madrid – Spain

⁴ INRA Unité Expérimentale Forêt Pierroton (UEFP), 33610 Cestas, France

* corresponding email's author: natalia.vizcaino.palomar@gmail.com

Contact Information: natalia.vizcaino.palomar@gmail.com.

[ORCID ID 0000-0002-3481-7567](https://orcid.org/0000-0002-3481-7567)

Abstract

Aim: Under rapid environmental change, phenotypic plasticity, if adaptive, could increase the odds for organisms to persist. Environmental variation over time is an important source of phenotypic plasticity. Likewise, phenotypic plasticity can vary with age in many organisms. However, little is known on phenotypic plasticity variation across species' ranges. Our aims are: (i) to assess whether populations' phenotypic plasticity is related to the inter-annual climate variation under which populations have evolved during the last century; (ii) to compare phenotypic plasticity among developmental classes; and (iii) to predict phenotypic plasticity across' species ranges.

Location: Europe and North-Africa.

Time period: 1901-2014.

Major taxa studied: *Pinus nigra*, *P. pinaster* and *P. pinea*.

Methods: We used 372 646 individual tree height measurements at three developmental classes from a wide network of 38 common gardens in Europe and North Africa with provenances covering the distribution range of the species. With this data, we: i) build linear mixed-effect models of tree height as a function of tree age, population and climate; ii) estimate populations' reaction norms from the fitted models; iii) calculate populations' phenotypic plasticity indexes; iv) build models of populations' phenotypic plasticity indexes as a function of inter-annual climate variation during the last century.

Results: We found that i) most populations that have evolved under high inter-annual climate variation, in either maximum or minimum values in temperature or precipitation, exhibited high values of plasticity in tree height; ii) phenotypic plasticity for tree height was higher in young trees than in older ones, iii) phenotypic plasticity did not follow any particular geographical pattern across species' ranges.

Main conclusions: Phenotypic plasticity across the three Mediterranean pines' ranges is related with the climate variation experienced over time and calls into question whether this plasticity could be adaptive and hence beneficial to cope with climate change in the short-term.

Keywords: Acclimation, Black pine, Climate change, Developmental stage, Inter-annual climate variation, Maritime pine, Mixed-effect models, Stone pine.

Introduction

Climate change is reshuffling species distribution ranges from marine to terrestrial systems, altering current ecosystems functioning and structure through disruption of species interactions at temporal and/or spatial scales (Lenoir *et al.*, 2008; Poloczanska *et al.*, 2013). To survive under new climates, organisms can move to more favorable environments (Chen *et al.*, 2011; Sunday *et al.*, 2011), or persist *in-situ* by changes in their genetic composition or adjusting to environmental changes using phenotypic plasticity (West-Eberhard, 2003; Pulido & Berthold, 2004). Evolutionary responses to climate change will imply changes in allele frequencies that need many generations to arise (Bradshaw & Holzapfel, 2001; Reale *et al.*, 2003; Franks *et al.*, 2007), whereas plastic responses can occur without changes in the genetic structure (Sultan, 2000; Valladares *et al.*, 2014) within one generation (or even longer when including trans-generational effects, Donelson *et al.*, (2018)). Thus, phenotypic plasticity can provide a rapid response, whereas evolutionary responses need longer time depending on the lifespan of organisms. For the particular case of trees, with very long generation times and large gene flow among populations, genetic adaptation occurs at long time scales (Savolainen *et al.*, 2007). For example, evolutionary adjustments to match new climates could need more than 1500 years in *Pinus sylvestris* (Rehfeldt *et al.*, 2002). Therefore, plasticity is often the main mechanism for tree populations to respond *in-situ* to rapid climate change (Benito Garzón *et al.*, 2019).

Environmental variation, either spatial or temporal, may promote differentiation in phenotypic plasticity among populations (Vizcaíno-Palomar *et al.*, 2016). In this context, some studies have shown that more plastic genotypes are promoted under greater heterogeneity (Lind & Johansson, 2007; Canale & Henry, 2010; Baythavong, 2011; Lázaro-Nogal *et al.*, 2015). However, phenotypic plasticity may not be always advantageous, and sometimes it can be detrimental. For example, high values of plasticity can be associated with low values of fitness-related traits as survival, biomass, or reproduction (e.g. Sánchez-Gómez *et al.*, (2006); Molina-Montenegro & Naya, (2012)). Likewise, changes in plasticity can occur during the lifespan of organisms due to morphological and physiological adjustments to the environment (Evans, 1972; Coleman *et al.*, 1994; Mitchell & Bakker, 2014). Hence, we could expect to find differences in phenotypic plasticity for fitness-related traits between early and mature stages of development (Mediavilla & Escudero, 2004). For instance, we can expect high plasticity in seedlings that present small root systems located in the shallow soil layers with great variation in soil moisture in contrast to mature trees with deep root systems reaching to more stable layers of soil moisture over the year. Hence, greater plasticity at the recruitment stage could be favorable for plant establishment in the community. Taken altogether, we could expect that phenotypic plasticity can vary across the species' distribution ranges and within-species lifespan.

The complex topography and orography of the Mediterranean basin, with its inter and intra-seasonal climate variation, and its recent story of species' expansions from refugia after the Last Glacial Maximum could have promoted differentiation in phenotypic plasticity among populations across species' ranges (Médail & Diadema, 2009). As a result, Mediterranean pine species present patchy distributions with differentiated patterns of genetic diversity and local adaptation, reviewed in Fady, (2012). For instance, although *P. nigra* has a larger distribution than *P. pinaster*, both present moderate-high population differentiation in neutral genetic diversity (Soto *et al.*, 2010) and in quantitative traits, such as tree height, diameter, height-diameter allometry, survival, etc. (Varelides *et al.*, 2001; Taïbi *et al.*, 2016; Vizcaíno-Palomar *et al.*, 2016). On the contrary, *P. pinea* presents very low levels in genetic diversity across its range (Vendramin *et al.*, 2008), as well as low differentiation for quantitative traits, such as in tree height (Mutke *et al.*, 2010, 2013; Sánchez-Gómez *et al.*, 2011).

Assessing populations' phenotypic plasticity responses across the species' ranges requires the use of phenotypic data measured from multiple common gardens, ideally a minimum of three (Arnold *et al.*, 2019), installed across large environmental gradients in which a suite of populations from varied origins are planted. These experimental designs permit to fit populations' non-linear phenotypic responses to the environment, known as 'reaction norm' curves (Gavrilets & Scheiner, 1993; Schlichting & Pigliucci, 1998), from which quantifying populations' phenotypic plasticity is straightforward (Arnold *et al.*, 2019). Furthermore, populations' phenotypic responses can be used to quantify populations' phenotypic plasticity using indices (Valladares *et al.*, 2006).

In this study, we used tree height, a fitness-related trait (King, 1990; Savolainen *et al.*, 2007), measured in a wide network of common gardens established in Europe and North Africa for *Pinus nigra*, *P. pinaster* and *P. pinea* (Vizcaíno-Palomar *et al.*, 2019). We fitted linear mixed-effect models of tree height to: i) assess whether populations' phenotypic plasticity is related to the inter-annual climate variation under which populations have evolved during the last century; ii) compare phenotypic plasticity among developmental classes; iii) predict phenotypic plasticity across species ranges.

Material and methods

Provenance trials, species and phenotypic data

We used tree height recorded in common garden networks for three pine species: *Pinus nigra* Arn., *P. pinaster* Aiton and *P. pinea* L. (see Figure S1 in Supporting Information). For *Pinus nigra*, we used 192 221 measurements of individual tree height recorded in 15 trials distributed across three countries (France, Germany and Spain) where 78 populations (provenances) from origins covering the entire range of the species were planted. Trials were planted between years 1968 and 2009 and tree heights were measured between 2 and 18 year-old. For *P. pinaster* we used 123 801 measurements of individual tree height recorded in 14 trials established across three countries (France, Morocco and Spain) and 182 populations covering the entire range of the species. Trials were installed between years 1966 and 1992 and tree heights were measured between 2 and 34 year-old. For *P. pinea*, we used 56 624 measurements of individual tree height recorded in 9 trials established in France and Spain, where a total 55 populations covering the entire range of the species were planted. Trials were established between years 1993 and 1997, and tree heights were measured between 2 and 22 year-old. Further description of these databases can be found in Vizcaíno-Palomar *et al.*, (2019).

To analyse the effect of age on phenotypic plasticity we defined three developmental classes (DC. 1, DC. 2 and DC. 3) covering the range of ages of each species. In all species, DC.1 included information for 4 year-old trees, DC.2 included trees of 8, 13, and 9 year-old, and DC.3 included information for 14, 24, and 22 year-old trees, for *P. nigra*, *P. pinaster* and *P. pinea*, respectively.

Climate data

We used the EuMedClim dataset that provides annual measurements between 1901 and 2014, at 30 arc-seconds (~ 1km) of resolution (<http://gentree.data.inra.fr/climate/datasets/>; Fréjaville & Benito Garzón, (2018)). We used a total of 21 climatic variables related with either annual or seasonal parameters of climate in terms of precipitation and temperature (Table S1). From this database, we computed the following climate-related variables and indices:

i) Long-term climate effect on trees' height population (clim_p) was calculated as the average climate at the population origin between the beginning of the 20th century (1901) and the year before the trees were planted in the trials. This 'long-term' effect reflects the climate that occurred when the planted seeds were generated, and it can be related to population effects.

ii) Short-term climate effect on trees' height population (clim_t) was calculated as the average climate at the trial of the last 3 years including the year when measurements were taken. This definition assures that plastic responses are measured under equal periods of time in all trees

for the three species, easing comparisons. This ‘short-term’ effect was defined to reflect the plastic response of tree height to recent climate.

iii) Inter-annual climate variation indices during the 20th at the population origin. We computed the standard deviation (sd) of seven climate variables selected to reflect the past climate variation encountered by the tree populations since the beginning of 20th century (1901) and the year before the trees were planted. Specifically, we computed the standard deviation (sd) of the mean annual temperature (sd bio1), sd of the mean diurnal temperature range (sd bio2), sd of the maximum temperature of the warmest month (sd bio5), sd of the minimum temperature of the coldest month (sd bio6), sd of the annual precipitation (sd bio12), sd of the precipitation of the wettest month (sd bio13) and sd of the precipitation of the driest month (sd bio14).

All climate-related variables and indices were standardized for further analyses.

Statistical analyses

We used linear mixed-effect models to account for the following effects: tree age, genetics (approached by the climate at the population origin, $clim_p$) and plasticity (approached by the climate at the trial, $clim_t$) on tree height measured across the networks of provenance tests for the three species. Afterwards, we predict populations’ phenotypic responses across the climatic range covered by the trials to compute phenotypic plasticity indices to summarize these curves. Our approach allowed us to obtain linear or non-linear curve responses as these are very common in nature (Arnold *et al.*, 2019). Then, phenotypic plasticity values can be estimated with more accurate.

1. Linear mixed-effect models of tree height responses accounting for age and climate

For each species, we selected one climate variable for the population (hereafter $clim_p$) and another for the trial (hereafter $clim_t$). This selection was based upon the complementary use of linear mixed-effects models and on principal components analyses (PCA) of the climate variables (see Appendix S1 for a detailed description). For *P. nigra*, we selected mean annual temperature (bio1) for $clim_t$ and annual water availability (WAI) for $clim_p$. For *P. pinaster*, annual potential evapotranspiration (PET) for $clim_t$ and winter precipitation (prec.djf) for $clim_p$. And for *P. pinea*, maximum temperature of the warmest month (bio5) for $clim_t$ and summer precipitation (prec.jja) for $clim_p$ (see Appendix S2, Table S2 and Figure S2).

Linear mixed-effect models were fitted to quantify the effects of tree age, $clim_p$ and $clim_t$ on tree height. The model equation takes the form (Eq. 1):

$$H = \sum_{i=1}^p \alpha_i X_i + \varepsilon_{ran} + \varepsilon_{res} \quad \text{Eq. 1}$$

where H is tree height, α_i is the set of p parameters associated with the main and interactive fixed effects of X_i (tree age, clim_p , clim_t), ε_{ran} is the variance component associated with the random terms, and ε_{res} is the residual distributed error, usually following a Gaussian distribution (see Results section).

The saturated model for the fixed part, $\sum_{i=1}^p \alpha_i X_i$, contained the linear and quadratic terms for each explanatory variable and all the potential pair-wise and three variable interactions (i.e. $\text{Age} \times \text{clim}_p$, $\text{Age} \times \text{clim}_t$, $\text{clim}_p \times \text{clim}_t$, $\text{Age} \times \text{clim}_p \times \text{clim}_t$). The random part of the model allowed us to consider three dimensions of common gardens experiments: a) the hierarchical nature of the data derived from the experimental design (i.e. populations nested within blocks, and blocks nested within trials), b) the temporal correlation between repeated measurements within tree individuals (i.e. individual tree), and c) the potential sources of variation not included in the fixed effects (such as soil, variation occurring at smaller spatial scales than blocks, etc.). All the variables were examined for outliers and departures from normality and the linearity of the relationships of each predictor with the response variable was checked (through residual plots for each predictor variable in the final model).

We selected the best-supported model starting from a saturated model following a hierarchical backward selection procedure (Burnham & Anderson, 2002; Zuur *et al.*, 2009). For the random part of the model, we selected the structure with the lowest AIC value (round 1). For the fixed part, we used the Akaike Information Criterion (AIC) (Akaike, 1992), following the rule that net increments of lower than two units of AIC associated with the elimination of any parameter in the full model determined the exclusion of the parameter from the final model. We started by testing the three-variable interaction (round 2), followed by the two-variable interaction (round 3), main effects (round 4), and linear effects (round 5).

Differences in AIC between models allowed us quantifying the relative importance of each predictor variable. The random effects were tested using restricted maximum likelihood of the parameter (REML), and fixed effects using maximum-likelihood (ML). Finally, parameter estimates of the best-supported model were obtained using restricted maximum likelihood (REML), which minimizes the likelihood of the residuals from the fixed-effect portions of the model (Zuur *et al.*, 2009). The variance explained by the model was assessed by pseudo- R^2 (Nakagawa & Schielzeth, 2013) that splits the variance into the marginal MR^2 (explained solely by the fixed effects) and conditional CR^2 (explained by fixed and random effects together). This pseudo- R^2 cannot be calculated with all the combinations between the family distributions and link functions (e.g. Gaussian family with identity link “log”) for linear mixed-effect models, therefore the goodness of fit of the models was also assessed by computing the capacity of generalization of the

model (CG). To do this, we calculated the Pearson coefficient, ρ , between a model fitted with the 2/3 parts of the data and independently validated with the remaining 1/3 part of the data. Additionally, to detect collinearity between explanatory variables in the best-supported model we used the variance inflation factor (VIFs) and set maximum value of VIF to 5 which is considered acceptable (Belsley, 1991). Appropriateness of the best model was assessed by plots of predicted vs. observed values. We used the R version 3.2.3 (R Core Team, 2016) run in linux-gnu operating system to perform all the analyses, and the “lme4” package (Bates *et al.*, 2015).

2. Computing populations’ phenotypic responses

Using the best-supported model for each species, we predicted populations’ phenotypic responses curves of tree height across the climatic range covered by the trials, clim_t , for the three developmental classes (DC). Specifically, we fixed tree age using the DC, and the climate of origin of each population (clim_p), and then we predict tree height responses curves along the climate of the trial (clim_t) varying between the 99% percentiles observed in clim_t data.

3. Computing phenotypic plasticity indices

Using the populations’ phenotypic responses curves of tree height and developmental classes (DC.1, DC.2 and DC.3), we computed two phenotypic plasticity indices across the climatic ranges covered by the trials (reviewed in Valladares *et al.*, (2006)).

1) Phenotypic plasticity index (PP) computed as follows:

$$\text{PP} = (\text{PR}_M - \text{PR}_m) / \text{PR}_M \quad \text{Eq. 2}$$

where PR_M is the highest phenotypic value across the population’s phenotypic response and across the climatic range studied. PR_m is the lowest phenotypic value across the climatic range studied. This index ranges between values of zero and one. The closer the values are to zero the less plastic the population is; and the opposite with values close to one.

2) Coefficient of variation of the phenotypic response (CV) computed as follow:

$$\text{CV} = \text{sd}(\text{PR}) / \text{mean}(\text{PR}) \quad \text{Eq. 3}$$

where PR is the phenotypic value at each point of the climatic range studied across the population’s phenotypic response fitted. sd is the standard deviation. This index reflects well the range of phenotype variation across the studied range. This index ranges between zero and one; the smaller the CV is, the smaller the plasticity is; and the opposite, the greater the CV is, the greater the plasticity is.

4. The developmental class effect on populations’ tree height plasticity indices

We tested if plasticity in tree height changed with the developmental class within species. To this aim, we performed analyses of variance of the two phenotypic plasticity indices, PP and CV, for the developmental classes and post-hoc pairwise comparisons of Tukey HSD (Honestly Significant Difference).

5. The inter-annual climate variation during the 20th century effect on tree height plasticity indices

For each species and each developmental class, we tested if inter-annual climate variation at the populations' origin could explain the current degree of phenotypic plasticity measured by the two phenotypic plasticity indices (PP and CV). To this end, we fitted linear fixed-effect models between the phenotypic plasticity index (PP or CV, as the response variable) and the set of inter-annual climate variation indices (sd bio1, sd bio2, sd bio5, sd bio6, sd bio12, sd bio13 and sd bio14, as explanatory variables) for each developmental class (Eq. 4). Collinearity in the models was controlled by including climate variation indices whose co-variation measured with the Pearson's correlation coefficient was below |0.7|.

$$PI_j = \sum_{i=1}^p \alpha_i X_i + \varepsilon_{res} \quad \text{Eq. 4}$$

where PI_j is the phenotypic plasticity index at the developmental class j ($j= 1, 2$ or 3), α_i is the set of p parameters associated with the effects of X_i (sd bio _{i}) and ε_{res} is the residual error. Models were fitted with a Gaussian distribution of errors and identity link function. A step-wise procedure (direction= "backward") was implemented to choose the best-supported model. Appropriateness of the models were assessed by plots of residuals vs. fitted values, qq-plots and the Cook's distance that identify outliers in the data that could over-influence the model fitting, if necessary they were removed from the analysis.

Results

The model: Tree height responses accounting for tree age and climate

The model that included all the factors tested (tree age, clim_p and, clim_t) with the linear and quadratic effects, and three and two-pairwise interactions, was the best-supported one for the three species (Table 1). The final model for *P. nigra* included age, mean annual temperature at the trial (bio1_t) and annual water availability at the populations' origin (WAI_p); *P. pinaster* included age, annual potential evapotranspiration at the trial (PET_t) and winter precipitation at the populations' origin (prec.djf_p); and for *P. pinea* included age, maximum temperature of the warmest month at the trial (bio5_t) and summer precipitation at the populations' origin (prec.jja_p) (Table 2, Table S3 and Fig. S3). All models produced unbiased estimates of tree height and high capacity of generalization, as well as high marginal and conditional explained variance, $\text{CG}/\text{MR}^2/\text{CR}^2$, with 0.79/not available/not available, 0.72/0.83/0.96 and 0.80/0.69/0.97) for *P. nigra*, *P. pinaster* and *P. pinea*, respectively (Table 2).

Main drivers of tree height triggering populations' phenotypic responses

Overall, tree age made the largest contribution to tree height, followed in order of importance by the climate at the trial and at the populations' origin: clim_t and clim_p , respectively (Table 1; see ΔAIC comparisons). The mean annual temperature of the trial presented a positive effect on tree height in *P. nigra* (Fig.1a and Fig.S4a), but in the other two species, at a certain evaporative demand (either expressed in mm by annual potential evapotranspiration or degrees Celsius by the maximum temperature of the warmest month), the temperature had a negative effect on tree height, see populations' phenotypic responses in Fig.1 and Fig.S4 for *P. nigra* and *P. pinaster*.). The most important interaction was age with clim_t , except in *P. pinaster* that was age with clim_p (Tables 1 & 2). Finally, the interaction term between $\text{clim}_t \times \text{clim}_p$ overall contributed the least to tree height, although in *P. pinaster* this contribution was higher than in the other species (Table 1 and populations' phenotypic responses in Fig.1. and Fig.S4.).

The developmental class effect on populations' tree height plasticity indices

Overall, phenotypic plasticity indices decreased significantly across developmental classes, i.e. young trees are the most plastic ones (Fig. 2, Tables S4 and S5). The greatest values of plasticity were found for *P. pinea* and the least for *P. nigra* for all developmental classes (Fig. 2). Finally, intraspecific phenotypic plasticity variation was the greatest in *P. pinaster* and the lowest in *P. pinea* (Fig. 2).

The inter-annual climate variation during the 20th century effect on tree height plasticity indices

For the three species, we did not include the standard deviation of annual precipitation (sd bio12) in the fixed-effect models because it was highly correlated with the standard deviation of the precipitation of the wettest month (sd bio13): Pearson' correlation coefficients of 0.76, 0.82 and 0.91 for *P. nigra*, *P. pinaster* and *P. pinea*, respectively. Moreover, we removed some populations whose Cook's distances were above 1 and over-influenced the fitted models (Appendix S3). Also, the model fitted for early adults in *P. pinea* using the CV index was not included in the results as it did not accomplished the linearity assumptions.

Overall, inter-annual temperature and precipitation variation during the 20th century in the standard deviation (sd) of the maximum temperature of the warmest month (sd bio5), sd of the precipitation of the wettest month (sd bio13) and sd of the precipitation of the driest month (sd bio14) were positively correlated with phenotypic plasticity indices (Table 3); while inter-annual variation in sd of the average annual temperature (sd bio1) was negatively correlated with phenotypic plasticity indices, with the exception of the developmental class 3 in *P. nigra* and *P. pinea* (Table 3). The variance explained by the phenotypic plasticity indices models (P.I.) was high, and ranged between 0.69 and 0.70 for *P. nigra*, 0.80 and 0.84 for *P. pinaster*, and 0.71 and 0.76 for *P. pinea* (Table 3).

The results of the two indices were similar (Table 3). For illustrative purposes we plot the PP and CV indices along the sd bio5 variable that was statistically significant in all the models tested. Overall, we found that populations that experienced higher inter-annual climate variation during the 20th century (sd bio5) presented higher plasticity in tree height for the three developmental classes, except in the developmental class 3 for *P. nigra* and *P. pinea* (Fig.3 and Fig.S5). Spatial differences among populations of PP and CV were also similar, with more contrasted differences among populations for *P. pinaster* (Fig. 1 and Fig. S5).

Discussion

The use of range-wide multi-year tree height measurements compiled on common gardens allowed us to quantify the effect of tree age, population differentiation and plasticity across the distribution ranges of three Mediterranean pines. Overall, our results show that: (i) A significant part of phenotypic plasticity in tree height was explained by the inter-annual climate variation during the 20th century under which tree populations evolved; (ii) Younger trees were more plastic in tree height than older trees; (iii) Although populations' responses to climate were largely driven by phenotypic plasticity we did not find a geographical pattern of phenotypic plasticity across the species' ranges.

Main climatic drivers of tree height triggering populations' phenotypic responses

Plastic responses in tree height were mainly driven by temperature-related variables (annual mean temperature, annual potential evapotranspiration and maximum temperature of the warmest month), and in general, rising temperatures led to higher trees up to a certain value. This result is in agreement with previous studies suggesting that higher heights in pines are found at warmer sites, if drought is not limiting (Vizcaíno-Palomar *et al.*, 2016). This could be explained because warm temperatures, up to a certain threshold, allow trees to have higher photosynthetic capacity, resulting in a higher rate of carbon assimilation (Way & Oren, 2010) but going beyond that threshold, it can imply the opposite effect. Accordingly, in *P. pinaster* and *P. pinea* tree height decreases when the evaporative demand is too high due to stomatal closure and reduction of the photosynthetic activity (Pasho *et al.*, 2012; Mazza *et al.*, 2014). However, we did not find that threshold in *P. nigra*. This species is found at higher altitudes than *P. pinaster* and *P. pinea*, in mountainous areas where high temperatures can be counterbalanced by the altitude effect and hence allow for increments of tree height growth. Moreover, it can be explained by the fact that the range of climate covered by the trials do not cover the complete population's phenotypic response, explaining the lack of a maximum tree height as it have been found in the other two species. Altogether, these results suggest that high temperatures linked with water stress are the main climatic drivers limiting tree height in the three pine species studied.

The main driver of population differentiation (population effect) in tree height was precipitation (for *P. pinaster* and *P. pinea*) and annual water availability (for *P. nigra*). This points out to the selective role of water availability across the distribution range of these mostly Mediterranean trees (Pigott & Pigott, 1993). Our findings suggest that evolutionary processes in tree height were mostly driven by water availability (Aranda *et al.*, 2009; Sánchez-Gómez *et al.*, 2011), although local adaptation is driven by minimum winter temperatures for *P. nigra* (Kreyling *et al.*, 2012), and by mean annual temperature for *Pinus pinea* (Mutke *et al.*, 2010). The highest

differences in tree height among populations were found for *P. nigra* and *P. pinaster* (Fig. 1a. and Fig. 1e.). In general, populations originating from the extremes of the climatic gradient, either under high or low values in rainfall or water availability, underperformed compared to populations originating from intermediate climates, but these differences are more marked in *P. pinaster*. For example, *P. pinaster* populations from the south of the distribution are better adapted to drought: they invest higher biomass to root and less to stem development than populations from northern parts of the distribution (Aranda *et al.*, 2009). In *P. nigra*, differences in tree height due to genetic effects have also been recorded (Thiel *et al.*, 2012), whereas *P. pinea* shows low genetic variation among populations (Fig.1i.). This is in agreement with previous studies reporting little genetic variation in morphological and physiological quantitative traits -e.g. photosynthesis, biomass partitioning, SLA, etc. (Court-Picon *et al.*, 2004; Mutke *et al.*, 2010; Sánchez-Gómez *et al.*, 2011) but null in Chambel *et al.*, (2007).

The developmental class effect on populations' tree height plasticity indices

Young pine trees were more plastic than early adults (Fig. 2). This result suggests that the capacity to respond plastically changes along the life cycle of trees. Phenotypic plasticity differ among species, among populations, among traits (Valladares *et al.*, 2002; Bradshaw, 2006), and here we show that it also varies with age. The first stages of recruitment are critical for plant establishment, and hence greater capacity of plasticity in tree height can be advantageous to avoid competition and reach light. In addition, small changes in the environment can be more noticeable for seedlings than to saplings or adult trees that are already well established with their root systems installed into deeper layers of the soil compared to seedlings. For instance, soil moisture variation is higher in the shallow layers of the soil than in deeper ones, hence promoting greater phenotypic plasticity. As a consequence, phenotypic plasticity variation across developmental classes can impact into many ecological processes, such as population and community dynamics, the community assembly and ecosystem functioning.

The inter-annual climate variation during the 20th century effect on tree height plasticity indices

Our findings suggest that plasticity in tree height is a trait that is under selection driven by climate variability (Table 3, Fig. 3 and Fig. S5). This finding is consistent for the three Mediterranean species regardless the origin of the populations. Populations that evolved under high inter-annual climate variation, in either maximum or minimum values in temperature or precipitation of climate variables associated with extreme values (standard deviation of the maximum temperature of the warmest month, sd of the precipitation of the wettest month and sd of the precipitation of the driest

month) during the 20th century, have great capacity to respond plastically in tree height to changes in climate (Fig. 3), regardless their position at the core or at the margin of the distribution range (maps in Fig. 1 and Fig.S4). These results are in agreement with previous studies in plant species where plastic responses were associated with climate variation in *Convolvulus chilensis* and *Senna candolleana* (Gianoli & González-Teuber, 2005; Lázaro-Nogal *et al.*, 2015). Although local adaptation (population effect in common garden data) seems to clearly follow a geographical pattern in European trees (including the ones studied here) (Frejaville *et al.*), our results show that plasticity geographical patterns are more complex (Valladares *et al.*, 2014).

Implications of phenotypic plasticity for evolutionary responses to climate change

Our findings are important in the context of climate change because plastic genotypes would likely increase their odds to persist at the short-time if plasticity is adaptive, and can also be advantageous if plastic genotypes are subject to further evolution that promotes the necessary genetic changes to reach the new optimum and get adapted to the new environment (Pigliucci, 2005; Richards *et al.*, 2006). Among the three studied species, *P. pinaster* presents high values of plasticity in tree height combined with a high differentiation among populations (Fig. 1e. and Fig. 2), suggesting good chances to respond to climate change in the short term by phenotypic plasticity and keeping evolutionary potential to adapt in the long term. *P. pinea* presents the highest phenotypic plasticity values out of the three studied species, but combined with low differentiation among populations and low genetic diversity (Fig. 1i. and Fig. 2), which makes plasticity virtually the unique way for this species to respond to changes in the environment. However, we cannot rule out that plasticity for tree height is related to higher fitness and our results call into question whether higher plasticity could be adaptive and hence beneficial to cope with climate change in the short-term.

Limitations

Although the network of provenance tests used in the present study cover relatively well the species distribution ranges, our results of phenotypic plasticity in tree height are confined to the climatic gradients covered by these common gardens. For example, in *P. nigra* phenotypic plasticity values could have been underestimated because the maximum tree height is not reached within the mean annual temperature range studied. Improved results regarding phenotypic plasticity could be obtained by establishing new common gardens both within the species' distribution to complete the range and outside the climatic range of the species.

Conclusions

Under a climate change context, the potential of the three species to persist *in-situ* largely rely on their plastic responses, regardless of their genetic diversity. The predominance of the plastic effect over the genetic one highlights that at the short-term, species' strategy to keep pace with climate change will likely rely on eco-physiological adjustments to environmental changes rather than on evolutionary responses. However, our current understanding of plasticity makes difficult to ascertain if plasticity will be adaptive and in case, which will be the real limits of plasticity. Therefore, to allow species persistence in the long-term, genetic variation within populations is essential to respond by evolutionary processes to environmental changes. Likewise, our results call into question whether future climate change variation would promote plasticity in the near future as our results showed that happened in the 20th century.

References

- Akaike, H. (1992) *Information theory and an extension of the maximum likelihood principle. Breakthroughs in statistics* (ed. by S Kotz and N Johnson), pp. 610–624. Springer-Verlag, London, UK.
- Aranda, I., Alía, R., Ortega, U., Dantas, Â.K. & Majada, J. (2009) Intra-specific variability in biomass partitioning and carbon isotopic discrimination under moderate drought stress in seedlings from four *Pinus pinaster* populations. *Tree Genetics and Genomes*, **6**, 169–178.
- Arnold, P.A., Kruuk, L.E.B. & Nicotra, A.B. (2019) How to analyse plant phenotypic plasticity in response to a changing climate. *New Phytologist*, **222**, 1235–1241.
- Bates, D., Maechler, M., Bolker, B. & Walker, S. (2015) Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, **67**, 1–48.
- Baythavong, B.S. (2011) Linking the Spatial Scale of Environmental Variation and the Evolution of Phenotypic Plasticity: Selection Favors Adaptive Plasticity in Fine-Grained Environments. *The American Naturalist*, **178**, 75–87.
- Belsley, D.A. (1991) A Guide to using the collinearity diagnostics. *Computer Science in Economics and Management*, **4**, 33–50.
- Benito Garzón, M., Robson, T.M. & Hampe, A. (2019) Δ Trait SDMs: species distribution models that account for local adaptation and phenotypic plasticity. *New Phytologist*, **222**, 1757–1765.
- Bradshaw, A.D. (2006) Unravelling phenotypic plasticity? why should we bother? *New Phytologist*, **170**, 644–648.
- Bradshaw, W. & Holzapfel, C. (2001) Genetic shift in photoperiodic response correlated with global warming. *Proceedings of the National Academy of Sciences*, **98**, 14509–14511.
- Burnham, K.P. & Anderson, D.R. (2002) *Model selection and multimodel inference: A practical Information-Theoretic approach*, (ed. by 2nd edn.) Springer-Verlag, New York.
- Canale, C. & Henry, P. (2010) Adaptive phenotypic plasticity and resilience of vertebrates to increasing climatic unpredictability. *Climate Research*, **43**, 135–147.
- Chambel, M.R., Climent, J. & Alía, R. (2007) Divergence among species and populations of Mediterranean pines in biomass allocation of seedlings grown under two watering regimes. *Annals of Forest Science*, **64**, 87–97.
- Chen, I.C., Hill, J.K., Ohlemüller, R., Roy, D.B. & Thomas, C.D. (2011) Rapid range shifts of species associated with high levels of climate warming. *Science (New York, N.Y.)*, **333**, 1024–6.
- Coleman, J.S., McConnaughay, K.D. & Ackerly, D.D. (1994) Interpreting phenotypic variation in plants. *Trends in ecology & evolution*, **9**, 187–91.
- Court-Picon, M., Gadbin-Henry, C., Guibal, F. & Roux, M. (2004) Dendrometry and morphometry

- of *Pinus pinea* L. in Lower Provence (France): adaptability and variability of provenances. *Forest Ecology and Management*, **194**, 319–333.
- Donelson, J.M., Salinas, S., Munday, P.L. & Shama, L.N.S. (2018) Transgenerational plasticity and climate change experiments: Where do we go from here? *Global Change Biology*, **24**, 13–34.
- Evans, C.G. (1972) *The Qualitative Analysis of Plant Growth*, Blackwell Scientific Publications, Oxford, UK.
- Fady, B. (2012) Biogeography of neutral genes and recent evolutionary history of pines in the Mediterranean Basin. *Annals of Forest Science*, **69**, 421–428.
- Franks, S.J., Sim, S. & Weis, A.E. (2007) Rapid evolution of flowering time by an annual plant in response to a climate fluctuation. *Proceedings of the National Academy of Sciences*, **104**, 1278–1282.
- Fréjaville, T. & Benito Garzón, M. (2018) The EuMedClim Database: Yearly Climate Data (1901–2014) of 1 km Resolution Grids for Europe and the Mediterranean Basin. *Frontiers in Ecology and Evolution*, **6**, 31.
- Frejaville, T., Vizcaíno-Palomar, N., Fady, B., Kremer, A. & Benito Garzón, M. Climate adaptation lags are higher in populations at the range margins than at the distribution core of European trees. *under review*.
- Gavrilets, S. & Scheiner, S.M. (1993) The genetics of phenotypic plasticity. V. Evolution of reaction norm shape. *Journal of Evolutionary Biology*, **6**, 31–48.
- Gianoli, E. & González-Teuber, M. (2005) Environmental Heterogeneity and Population Differentiation in Plasticity to Drought in *Convolvulus Chilensis* (Convolvulaceae). *Evolutionary Ecology*, **19**, 603–613.
- King, D.A. (1990) The adaptive significance of tree height. *The American Naturalist*, **135**, 809–828.
- Kreyling, J., Wiesenberg, G.L.B., Thiel, D., Wohlfart, C., Huber, G., Walter, J., Jentsch, A., Konnert, M. & Beierkuhnlein, C. (2012) Cold hardiness of *Pinus nigra* Arnold as influenced by geographic origin, warming, and extreme summer drought. *Environmental and Experimental Botany*, **78**, 99–108.
- Lázaro-Nogal, A., Matesanz, S., Godoy, A., Pérez-Trautman, F., Gianoli, E. & Valladares, F. (2015) Environmental heterogeneity leads to higher plasticity in dry-edge populations of a semi-arid Chilean shrub: insights into climate change responses. *Journal of Ecology*, **103**, 338–350.
- Lenoir, J., Gegout, J.C., Marquet, P.A., de Ruffray, P. & Brisse, H. (2008) A Significant Upward Shift in Plant Species Optimum Elevation During the 20th Century. *Science*, **320**, 1768–1771.
- Lind, M.I. & Johansson, F. (2007) The degree of adaptive phenotypic plasticity is correlated with

- the spatial environmental heterogeneity experienced by island populations of *Rana temporaria*.
Journal of Evolutionary Biology, **20**, 1288–1297.
- Mazza, G., Cutini, A., Manetti, M.C. & Chiara Manetti, M. (2014) Site-specific growth responses to
climate drivers of *Pinus pinea* L. tree rings in Italian coastal stands. *Annals of Forest Science*,
71.
- Médail, F. & Diadema, K. (2009) Glacial refugia influence plant diversity patterns in the
Mediterranean Basin. *Journal of Biogeography*, **36**, 1333–1345.
- Mediavilla, S. & Escudero, A. (2004) Stomatal responses to drought of mature trees and seedlings
of two co-occurring Mediterranean oaks. *Forest Ecology and Management*, **187**, 281–294.
- Mitchell, R.M. & Bakker, J.D. (2014) Intraspecific Trait Variation Driven by Plasticity and
Ontogeny in *Hypochaeris radicata*. *PLoS ONE*, **9**, e109870.
- Molina-Montenegro, M.A. & Naya, D.E. (2012) Latitudinal Patterns in Phenotypic Plasticity and
Fitness-Related Traits: Assessing the Climatic Variability Hypothesis (CVH) with an Invasive
Plant Species. *PLoS ONE*, **7**.
- Mutke, S., Gordo, J., Chambel, M.R., Prada, M.A., Álvarez, D., Iglesias, S. & Gil, L. (2010)
Phenotypic plasticity is stronger than adaptative differentiation among Mediterranean stone
pine provenances. *Forest Systems*, **19**, 354.
- Mutke, S., Gordo, J., Khouja, M. & Fady, B. (2013) Low genetic and high environmental diversity
at adaptive traits in *Pinus pinea* from provenance tests in France and Spain. *Options
Méditerranéennes A*, **105**, 73–79.
- Nakagawa, S. & Schielzeth, H. (2013) A general and simple method for obtaining R^2 from
generalized linear mixed-effects models. *Methods in Ecology and Evolution*, **4**, 133–142.
- Pasho, E., Camarero, J.J., Luis, M. & Vicente-Serrano, S.M. (2012) Factors driving growth
responses to drought in Mediterranean forests. *European Journal of Forest Research*, **131**,
1797–1807.
- Pigliucci, M. (2005) Evolution of phenotypic plasticity: where are we going now? *Trends in
Ecology & Evolution*, **20**, 481–486.
- Pigott, C.D. & Pigott, S. (1993) Water as a determinant of the distribution of trees at the boundary
of the Mediterranean zone. *Journal of Ecology*, **81**, 557–566.
- Poloczanska, E.S., Brown, C.J., Sydeman, W.J., Kiessling, W., Schoeman, D.S., Moore, P.J.,
Brander, K., Bruno, J.F., Buckley, L.B., Burrows, M.T., Duarte, C.M., Halpern, B.S., Holding,
J., Kappel, C. V., O'Connor, M.I., Pandolfi, J.M., Parmesan, C., Schwing, F., Thompson, S.A.
& Richardson, A.J. (2013) Global imprint of climate change on marine life. *Nature Climate
Change*, **3**, 919–925.
- Pulido, F. & Berthold, P. (2004) Microevolutionary Response to Climatic Change. *Advances in*

- 551 *Ecological Research*, **35**, 151–183.
- 552 R Core Team (2016) A language and environment for statistical computing.
- 553 Reale, D., McAdam, A.G., Boutin, S. & Berteaux, D. (2003) Genetic and plastic responses of a
- 554 northern mammal to climate change. *Proceedings of the Royal Society B: Biological Sciences*,
- 555 **270**, 591–596.
- 556 Rehfeldt, G.E., Tchebakova, N.M., Parfenova, Y.I., Wykoff, W.R., Kuzmina, N.A. & Milyutin, L.I.
- 557 (2002) Intraspecific responses to climate in *Pinus sylvestris*. *Global Change Biology*, **8**, 912–
- 558 929.
- 559 Richards, C.L., Bossdorf, O., Muth, N.Z., Gurevitch, J. & Pigliucci, M. (2006) Jack of all trades,
- 560 master of some? On the role of phenotypic plasticity in plant invasions. *Ecology Letters*, **9**,
- 561 981–993.
- 562 Sánchez-Gómez, D., Valladares, F. & Zavala, M.A. (2006) Performance of seedlings of
- 563 Mediterranean woody species under experimental gradients of irradiance and water
- 564 availability: trade-offs and evidence for niche differentiation. *The New phytologist*, **170**, 795–
- 565 806.
- 566 Sánchez-Gómez, D., Velasco-Conde, T., Cano-Martín, F.J., Ángeles Guevara, M., Teresa Cervera,
- 567 M. & Aranda, I. (2011) Inter-clonal variation in functional traits in response to drought for a
- 568 genetically homogeneous Mediterranean conifer. *Environmental and Experimental Botany*, **70**,
- 569 104–109.
- 570 Savolainen, O., Pyhäjärvi, T. & Knürr, T. (2007) Gene flow and local adaptation in trees. *Annual*
- 571 *Review of Ecology, Evolution, and Systematics*, **38**, 595–619.
- 572 Schlichting, C.D. & Pigliucci, M. (1998) *Phenotypic Evolution. A Reaction Norm Perspective*,
- 573 Sinauer Associates Inc, Sunderland, Massachusetts.
- 574 Soto, A., Robledo-Arnuncio, J.J., González-Martínez, S.C., Smouse, P.E. & Alía, R. (2010)
- 575 Climatic niche and neutral genetic diversity of the six Iberian pine species: a retrospective and
- 576 prospective view. *Molecular ecology*, **19**, 1396–409.
- 577 Sultan, S.E. (2000) Phenotypic plasticity for plant development, function and life history. *Trends in*
- 578 *Plant Science*, **5**, 537–542.
- 579 Sunday, J.M., Bates, A. & Dulvy, N. (2011) Global analysis of thermal tolerance and latitude in
- 580 ectotherms. *Proceedings of the Royal Society B: Biological Sciences*, **278**, 1823–1830.
- 581 Taïbi, K., del Campo, A.D., Aguado, A. & Mulet, J.M. (2016) Early establishment response of
- 582 different *Pinus nigra* ssp. *salzmanii* seed sources on contrasting environments: Implications for
- 583 future reforestation programs and assisted population migration. *Journal of Environmental*
- 584 *Management*, **171**, 184–194.
- 585 Thiel, D., Nagy, L., Beierkuhnlein, C., Huber, G., Jentsch, A., Konnert, M. & Kreyling, J. (2012)

Uniform drought and warming responses in *Pinus nigra* provenances despite specific overall performances. *Forest Ecology and Management*, **270**, 200–208.

Valladares, F., Balaguer, L., Martínez-Ferri, E., Perez-Corona, E. & Manrique, E. (2002) Plasticity, instability and canalization: Is the phenotypic variation in seedlings of sclerophyll oaks consistent with the environmental unpredictability of Mediterranean ecosystems? *New Phytologist*, **156**, 457–467.

Valladares, F., Matesanz, S., Guilhaumon, F., Araújo, M.B., Balaguer, L., Benito Garzón, M., Cornwell, W., Gianoli, E., van Kleunen, M., Naya, D.E., Nicotra, A.B., Poorter, H. & Zavala, M.A. (2014) The effects of phenotypic plasticity and local adaptation on forecasts of species range shifts under climate change. *Ecology Letters*, **17**, 1351–1364.

Valladares, F., Sánchez-Gómez, D. & Zavala, M.A. (2006) Quantitative estimation of phenotypic plasticity: Bridging the gap between the evolutionary concept and its ecological applications. *Journal of Ecology*, **94**, 1103–1116.

Varelides, C., Brofas, G. & Varelides, Y. (2001) Provenance variation in *Pinus nigra* at three sites in Northern Greece. **58**, 893–900.

Vendramin, G.G., Fady, B., González-Martínez, S.C., Hu, F.S., Scotti, I., Sebastiani, F., Soto, Á. & Petit, R.J. (2008) Genetically depauperate but widespread: the case of an emblematic mediterranean pine. *Evolution*, **62**, 680–688.

Vizcaíno-Palomar, N., Benito Garzón, M., Alía, R., Giovannelli, G., Huber, G., Mutke, S., Pastuszka, P., Raffin, A., Sbay, H., Seho, M., Vauthier, D. & Fady, B. (2019) Geographic variation of tree height of three pine species (*Pinus nigra* Arn., *P. pinaster* Aiton and *P. pinea* L.) gathered from common gardens in Europe and North-Africa. *Annals of Forest Science*, **accepted**.

Vizcaíno-Palomar, N., Ibáñez, I., González-Martínez, S.C., Zavala, M.A. & Alía, R. (2016) Adaptation and plasticity in aboveground allometry variation of four pine species along environmental gradients. *Ecology and Evolution*, **6**, 7561–7573.

Way, D.A. & Oren, R. (2010) Differential responses to changes in growth temperature between trees from different functional groups and biomes: a review and synthesis of data. *Tree Physiology*, **30**, 669–688.

West-Eberhard, M.J. (2003) *Developmental Plasticity and Evolution*, Oxford University Press, incl., New York.

Zuur, A.F., Ieno, E.N., Walker, N., Saveliev, A.A. & Smith, G.M. (2009) *Mixed Effects Models and Extensions in Ecology with R*, Springer.

619 **Data Accessibility Statement**

620 All phenotypic data used in this study are available on ZENODO with DOIs

621 10.5281/zenodo.3250704, 10.5281/zenodo.3250698 and 10.5281/zenodo.3250707 for *P. nigra*, *P.*

622 *pinaster* and *P. pinea*, respectively (Vizcaíno-Palomar *et al.*, 2019).

Table 1 Random and fixed effects selection of tree height model in response to age, clim_t and clim_p using the Akaike Information Criterion (AIC) for the three pine species.

<u><i>Pinus nigra</i></u>	df	AIC	ΔAIC	<u><i>Pinus pinaster</i></u>	df	AIC	ΔAIC
<i>Random effects selection</i>							
# round 1: testing random structure							
full model just fixed effects (fixed)	12	2239318		fixed	12	175302	
fixed + block effect	15	2002183		fixed + block effect	15	80711	
fixed + tree ind	13	1917911		fixed + tree ind	13	143270	
fixed + block effect + tree ind	16	1888388		fixed + block effect + tree ind	16	77968	
<i>Fixed effects selection</i>							
# round 2: three way interactions							
full model	16	1888388	0	full model	16	77862	0
no age×bio1 _t ×WAI _p	15	1888727	339	age×PET _t ×prec.djf _p	15	77885	23
# round 3: two way interactions							
full model	16	1888388	0	full model	16	77862	0
no age×bio1 _t	15	1897136	8748	no age×PET _t	15	78141	279
no age×WAI _p	15	1888506	117	no age×prec.djf _p	15	78772	910
no bio1 _t ×WAI _p	15	1888401	13	no PET _t ×prec.djf _p	15	77959	97
# round 4: main effects							
full model	16	1888388	0	full model	16	77862	0
no age	11	2605168	716780	no age	11	367581	289719
no bio1 _t	11	1916110	27722	no PET _t	11	123183	45321
no WAI _p	11	1888975	587	no prec.djf _p	11	79360	1498
# round 5: linear effects							
full model	16	1888388	0	full model	16	77862	0
no quadratic effect age	15	2083109	194721	no quadratic effect age	15	181836	103974
no quadratic effect bio1 _t	15	1889004	616	no quadratic effect PET _t	15	85070	7208
no quadratic effect WAI _p	15	1888398	10	no quadratic effect prec.djf _p	14	80797	2935

<i>Pinus pinea</i>	df	AIC	ΔAIC
<i>Random effects selection</i>			
# round 1: assessing random structure			
fixed	12	53113	
rand_bl	15	19110	
rand_ind	13	37939	
rand_both	16	16898	
<i>Fixed effects selection</i>			
# round 2: three way interactions			
full model	16	16790	0
age bio5 _t ×prec.jja _p	15	16797	7
# round 3: two way interactions			
full model	16	16790	0
no age×bio5 _t	15	23396	6606
no age×prec.jja _p	15	16807	17
no bio5 _t ×prec.jja _p	15	16794	4
# round 4: main effects			
full model	16	16790	0
no age	11	154070	137280
no bio5 _t	11	29128	12337
no prec.jja _p	11	16829	39
# round 5: linear effects			
full model	16	16790	0
no quadratic effect age	15	41981	25190
no quadratic effect PET _t	15	28835	12044
no quadratic effect prec.djf _p	14	19025	2234

626 df are the degrees of freedom, AIC is the Akaike Information Criterion value, Δ AIC is the difference between the full model and the models tested in
627 each round. In random effects selection, the smaller the AIC is, the better supported the random structure part is. In fixed effects selection, if Δ AIC
628 comparing two models is smaller than 2 units then we remove that effect from the full model.

Table 2 Detailed description of the best-supported model for each pine species analyzed including the family distribution and link function, the variance and standard deviation for the random effects (Var and SD respectively); and the estimated parameter, standard error and t-values for the fixed-effects (Estimate, SE and t-values, respectively).

Pinus nigra

Family distribution and link function:	Gaussian distribution error and identity link "log"		
<i>Random effects</i>		Var	SD
Id_tree	(Intercept)	20.613	4.54
Provenance/Block/Trial	(Intercept)	9.567	3.09
Block/Trial	(Intercept)	2.222	1.49
Trial	(Intercept)	63.428	7.96
Residual		282.506	16.81
<i>Fixed effects</i>	Estimate	SE	t value
(Intercept)	4.710	0.120	39.35
age	1.166	0.001	1250.89
bio1 _t	0.318	0.002	128.67
bio1 _t ²	-0.035	0.001	-24.88
age ²	-0.212	0.000	-527.92
WAI _p	0.021	0.006	3.59
WAI _p ²	-0.015	0.004	-3.45
age × bio1 _t	-0.048	0.001	-93.39
age × WAI _p	0.002	0.001	3.73
bio1 _t × WAI _p	0.008	0.002	3.84
age × bio1 _t × WAI _p	-0.010	0.001	-18.43
CG	0.79		
MR ²	-		
CR ²	-		

Pinus pinaster

Family distribution and link function:	Gaussian distribution error and log (H)		
<i>Random effects</i>		Var	SD
Id_tree	(Intercept)	0.01	0.11
Provenance/Block/Trial	(Intercept)	0.041	0.20
Block/Trial	(Intercept)	0.004	0.07
Trial	(Intercept)	0.271	0.52
Residual		0.093	0.31
<i>Fixed effects</i>	Estimate	SE	t value
(Intercept)	6.150	0.140	44.022
age	1.422	0.002	923.084
PET _t	-0.812	0.006	-127.091
PET _t ²	-0.379	0.004	-86.587
prec.djf _p	0.043	0.005	8.278
prec.djf _p ²	-0.044	0.003	-14.141
age ²	-0.366	0.001	-418.081
age × PET _t	0.028	0.002	16.764
age × prec.djf _p	-0.040	0.001	-30.268
PET _t × prec.djf _p	-0.034	0.003	-9.983
age × PET _t × prec.djf _p	0.008	0.002	5.030
CG	0.72		
MR ²	0.83		
CR ²	0.96		

Pinus pinea

Family distribution and link function:	Gaussian distribution error and log (H)		
<i>Random effects</i>		Var	SD
Id_tree	(Intercept)	0.014	0.12
Provenance/Block/Trial	(Intercept)	0.007	0.09
Block/Trial	(Intercept)	0.017	0.13
Trial	(Intercept)	0.601	0.78
Residual		0.062	0.25
<i>Fixed effects</i>	Estimate	SE	t value
(Intercept)	5.504	0.259	21.300
age	1.321	0.002	719.800
bio5 _t	0.168	0.012	14.600
bio5 _t ²	-0.719	0.006	-118.000
prec.jja _p	0.007	0.002	3.000
prec.jja _p ²	-0.004	0.001	-4.800
age ²	-0.278	0.002	-183.900
age × bio5 _t	0.185	0.002	84.800
age × prec.jja _p	0.005	0.001	4.300
bio5 _t × prec.jja _p	-0.006	0.002	-2.400
age × bio5 _t × prec.jja _p	0.004	0.001	3.000
CG	0.80		
MR ²	0.69		
CR ²	0.97		

Table 3. Results from the linear fixed-effect models between phenotypic plasticity indices (P.I.) and the standard deviation climate variables (sd bio). Three sub-tables are presented for each pine species, a), b) and c). Each sub-table shows the results for the two indices and the three developmental classes (DC) analyzed. Developmental Class 1: green, DC. 2: orange and DC. 3: blue

a) *P. nigra*

DC. 1	Variable	Estimate	effect	p-value	R ²	DC. 2	Variable	Estimate	effect	p-value	R ²	DC. 3	Variable	Estimate	effect	p-value	R ²
PP	sd bio 1	-5.11×10 ⁻³	-	***	0.69	PP	sd bio 1	-4.42×10 ⁻³	-	***	0.70	PP	sd bio 1	2.95×10 ⁻³	+	***	0.70
	sd bio5	6.46×10 ⁻³	+	***			sd bio5	4.26×10 ⁻³	+	***			sd bio5	-2.78×10 ⁻³	-	***	
	sd bio 13	5.68×10 ⁻³	+	***			sd bio6	1.11×10 ⁻³	+	n.s.			sd bio6	-7.36×10 ⁻⁴	-	n.s.	
	sd bio 14	2.94×10 ⁻³	+	***			sd bio 13	4.06×10 ⁻³	+	***			sd bio 13	-2.69×10 ⁻³	-	***	
CV	sd bio 1	-7.43×10 ⁻³	-	***	0.70	CV	sd bio 1	-5.35×10 ⁻³	-	***	0.70	CV	sd bio 1	2.81×10 ⁻³	+	***	0.70
	sd bio5	7.09×10 ⁻³	+	***			sd bio5	5.14×10 ⁻³	+	***			sd bio5	-2.66×10 ⁻³	-	***	
	sd bio6	1.86×10 ⁻³	+	n.s.			sd bio6	1.30×10 ⁻³	+	n.s.			sd bio6	-7.30×10 ⁻⁴	-	n.s.	
	sd bio 13	6.84×10 ⁻³	+	***			sd bio 13	4.90×10 ⁻³	+	***			sd bio 13	-2.54×10 ⁻³	-	***	
	sd bio 14	4.04×10 ⁻³	+	***			sd bio 14	2.89×10 ⁻³	+	***			sd bio 14	-1.51×10 ⁻³	-	***	

Variable refers to the explanatory variables of the model (standard deviation of the inter-annual climate variation in the 20th century). Estimate refers to the estimated value for that variable. Effect makes reference whether the parameter estimate has a positive (+) or negative (-) impact on the phenotypic plasticity index. p-value refers whether the Estimate is statistically significant with a p-value lower than 0.001 ***, 0.01 **, 0.05 *, 0.1 □ and 1 n.s. R² is the explained variance of the model.

642	b) <i>P. pinaster</i>															
643																
C. 1	Variable	Estimate	effect	p-value	R ²	DC. 2	Variable	Estimate	effect	p-value	R ²	DC. 3	Variable	Estimate	effect	p-value
P	sd bio 1	-4.34×10 ⁻³	-	***	0.84	PP					0.80	PP				
	sd bio 2	-7.23×10 ⁻⁴	-	□			sd bio 2	-1.20×10 ⁻³	-	**			sd bio 2	-8.59×10 ⁻⁴	-	**
	sd bio 5	3.83×10 ⁻³	+	***			sd bio 5	1.70×10 ⁻³	+	***			sd bio 5	1.20×10 ⁻³	+	***
	sd bio 6	-3.86×10 ⁻³	-	***			sd bio 6	-1.57×10 ⁻³	-	***			sd bio 6	-1.16×10 ⁻³	-	***
	sd bio 13	7.92×10 ⁻³	+	***			sd bio 13	8.69×10 ⁻³	+	***			sd bio 13	6.33×10 ⁻³	+	***
V	sd bio 14	2.91×10 ⁻³	+	***	0.84	CV	sd bio 14	7.65×10 ⁻⁴	+	□	0.80	CV	sd bio 14	5.82×10 ⁻⁴	+	□
	sd bio 1	-4.30×10 ⁻³	-	***			sd bio 2	-1.39×10 ⁻⁴	-	**			sd bio 2	-8.39×10 ⁻⁴	-	**
	sd bio 2	-7.74×10 ⁻⁴	-	n.s.			sd bio 5	2.14×10 ⁻³	+	***			sd bio 5	1.33×10 ⁻³	+	***
	sd bio 5	4.43×10 ⁻³	+	***			sd bio 6	-2.25×10 ⁻³	-	***			sd bio 6	-1.42×10 ⁻³	-	***
	sd bio 6	-5.10×10 ⁻³	-	***			sd bio 13	1.06×10 ⁻²	+	***			sd bio 13	6.52×10 ⁻³	+	***
	sd bio 13	1.01×10 ⁻²	+	***	0.84		sd bio 14	8.40×10 ⁻⁴	+	□	0.80		sd bio 14	4.86×10 ⁻⁴	+	n.s.
	sd bio 14	3.45×10 ⁻³	+	***												

644
645 Variable refers to the explanatory variables of the model (standard deviation of the inter-annual climate variation in the 20th century). Estimate refers to the estimated
646 value for that variable. Effect makes reference whether the parameter estimate has a positive (+) or negative (-) impact on the phenotypic plasticity index. p-value
647 refers whether the Estimate is statistically significant with a p-value lower than 0.001 ***, 0.01 **, 0.05 *, 0.1 □ and 1 n.s. R² is the explained variance of the model.

648

649 c) *P. pinea*

650

DC. 1	Variable	Estimate	effect	p-value	R ²	DC. 2	Variable	Estimate	effect	p-value	R ²	DC. 3	Variable	Estimate	effect	p-value	R ²
PP	sd bio1	-2.26×10 ⁻⁵	-	n.s.		PP	sd bio1	-2.47×10 ⁻⁵	-	n.s.		PP	sd bio1	4.09×10 ⁻⁵	+	□	
	sd bio 5	1.08×10 ⁻⁴	+	***			sd bio 5	9.69×10 ⁻⁵	+	***			sd bio 5	-1.44×10 ⁻⁴	-	***	
	sd bio 6	1.02×10 ⁻⁴	+	***			sd bio 6	8.99×10 ⁻⁵	+	***			sd bio 6	-1.26×10 ⁻⁴	-	***	
	sd bio 14	9.99×10 ⁻⁵	+	***	0.76		sd bio 14	9.07×10 ⁻⁵	+	***	0.71		sd bio 14	-1.23×10 ⁻⁴	-	***	0.7
CV	sd bio1	-6.40×10 ⁻⁵	-	□		CV	sd bio1	-4.11×10 ⁻⁵	-	□							
	sd bio 5	2.43×10 ⁻⁴	+	***			sd bio 5	1.50×10 ⁻⁴	+	***							
	sd bio 6	2.07×10 ⁻⁴	+	***			sd bio 6	1.37×10 ⁻⁴	+	***							
	sd bio 14	2.21×10 ⁻⁴	+	***	0.73		sd bio 14	1.40×10 ⁻⁴	+	***	0.75						

651

652 Variable refers to the explanatory variables of the model (standard deviation of the inter-annual climate variation in the 20th century). Estimate refers to the estimated

653 value for that variable. Effect makes reference whether the parameter estimate has a positive (+) or negative (-) impact on the phenotypic plasticity index. p-value

654 refers whether the Estimate is statistically significant with a p-value lower than 0.001 ***, 0.01 **, 0.05 *, 0.1 □ and 1 n.s. R² is the explained variance of the model.

Figures

Figure 1. Populations' phenotypic tree height responses across climate, particularized for three populations' origin (High, Average and Low in terms of $clim_p$ values) and for the three developmental classes, DC, (Developmental Class 1: green, DC. 2: orange and DC. 3: blue) for a) *P. nigra*, e) *P. pinaster* and i) *P. pinea*. Values of the phenotypic plasticity index (PP) for the three developmental classes across the species natural distribution ranges are shown. DC. 1: b), f) and j), DC. 2: c), g) and k); and DC. 3: d), h) and i).

Figure 2. Phenotypic plasticity values for the three species and for the three developmental classes, DC, (Developmental Class 1: green, DC. 2: orange and DC. 3: blue) for the two indices computed (PP and CV). An inset graph is included in the CV index as the values of *P. nigra* and *P. pinaster* are significantly smaller compared with those obtained in *P. pinea*.

673

Phenotypic plasticity index (PP)

674

675 **Figure 3.** Phenotypic plasticity predictions of the PP index across the standard deviation of
676 the maximum temperature of the warmest month (sd bio5) for the three developmental
677 classes and pines species.

678 **Appendix S1.** Detailed description for climate variable selection.

679 We used two analysis to select the short-term climate related to the trial ($clim_t$) and the long-
680 term climate related to the climate at the population origin ($clim_p$): i) linear mixed-effect
681 models and ii) principal component analyses.

682 i) Linear Mixed-Effect Models

683 To select $clim_t$, we ran 21 linear mixed-effects models to analyse the response of tree height
684 to each of the 21 climate variables to the climate of the trials (Table S1). To select $clim_p$, we
685 ran 21 linear mixed-effects models to analyse the response of tree height to each of the 21
686 climate variables to the climate at the populations' origin. Random effects included
687 populations nested into blocks, and those nested within trials, and trees nested within
688 population, block and trial, to control for differences among sites and populations, and to
689 control for repeated measurements of the same trees, respectively. Fixed effects included tree
690 age and the climate variable, including the linear and the quadratic forms, and the linear
691 interaction between them. All climate variables were standardized (the mean was subtracted
692 from each value and divided by the standard deviation). The model equation takes the form:

$$693 \quad H = \sum_{i=1}^p \alpha_i X_i + \varepsilon_{ran} + \varepsilon_{res} \quad \text{Eq. 1}$$

694 where H is tree height, α_i is the set of p parameters associated with the main and interactive
695 effects of X_i climate (either $clim_t$ or $clim_p$) and age variables, ε_{ran} is the variance component
696 associated with the random terms, and ε_{res} is the residual distributed error.

697 We selected the model in which the climate variable met three requisites: 1) the
698 sign of the estimated quadratic coefficient must be negative in order to get concave responses
699 (maximum tree height is expected at intermediate climatic values, decreasing towards the
700 extremes) 2) high absolute values of the estimated coefficients as an approach of the size
701 effect of the climate variable on tree height, 3) being biological meaningful variable. For
702 example, potential evapotranspiration climate variable can provide us with more information
703 than just a variable of annual precipitation. Usually, curve responses between $clim_t$ and tree
704 height are more likely to be concaves than tree height with $clim_p$ (personal observation from
705 preliminary analysis). Therefore for $clim_t$ we focused on the absolute highest value of the
706 negative estimated parameter in the quadratic term when statistically significant, while for
707 $clim_p$, we focused on whether the linear and quadratic terms were statistically significant and
708 then we chose among those variables with the absolute highest negative parameter in the
709 quadratic term. If only the linear parameter was significant, then we chose the variable with
710 the highest estimated parameter.

711 ii) Principal Component Analysis

712 We ran two independent principal component analyses (PCA) in R (R Core Team, 2016) for
 713 $clim_t$ and $clim_p$ variables for each species. These analyses help selecting the variable which is
 714 highly representative of the climate variation covered by the data. We generally choose the
 715 climate variable that belongs to the first PCA axis, which captures a higher variance of the
 716 data. However, sometimes we needed to choose the climate variable from the second PCA
 717 axis if the climate variable is highly correlated with the first PCA axis.

718

719 These two complementary analyses were defined to facilitate climate selection, but this was
 720 not always straightforward.

721 **Appendix S2.** Description of the results from the linear mixed-effect models and principal
722 component analyses to select the climate variables of clim_p and clim_t .

723 *P. nigra*. Mean annual temperature, bio1, was selected for clim_t . It showed the highest size
724 effect among the total of the climate variables tested in the linear mixed-effect models (Table
725 S2). Moreover, mean annual temperature was highly correlated with the first PCA axis (0.76)
726 and showed a moderate-high contribution to the first axis (6.37% being the maximum 9.36%)
727 (Figure S2). Annual water availability, WAI, was selected for clim_p . Although none of the
728 tested climate variables in the linear mixed-effect models were statistically significant, the
729 principal component analysis facilitated to choose it (Table S2). Thus, annual water
730 availability was highly correlated with the first PCA axis (-0.93) and it showed the highest
731 contribution to the first PCA axis, 8.33% (Figure S2).

732 *P. pinaster*. Annual potential evapotranspiration, PET, was selected for clim_t .
733 Although precipitation of the driest month could have been selected for modeling, we chose
734 annual potential evapotranspiration as it was statistically significant in the linear mixed-effect
735 models and it integrates temperature and precipitation values which make this variables more
736 informative (Table S2). Annual potential evapotranspiration was highly correlated with the
737 first PCA axis (0.94), and it showed the second highest contribution (7.57%) to this axis after
738 summer daily mean temperature (Figure S2). Winter precipitation, prec.djf , was selected for
739 clim_p . It was statistically significant in the linear mixed-effect models, and the two estimated
740 coefficients were biologically meaningful and their size effects were the highest (Table S2).
741 The variable of winter precipitation was moderate-to-highly correlated with the second PCA
742 axis (0.76), with a moderate-to-high contribution to the axis (8.21%, being the maximum
743 contribution of 12.55%) (Figure S2). The second axis of the PCA explained nearly the same
744 amount of variability, 33.38%, compared to the variance explained by the first axis of
745 45.19%.

746 *P. pinea*. Maximum temperature of the warmest month, bio5, was selected to
747 represent clim_t . It showed the highest size effect on the estimated quadratic term in the linear
748 mixed-effect models. Moreover, maximum temperature of the warmest month was highly
749 correlated with the first PCA axis (-0.91) and it showed a moderate-to-high high contribution
750 to the first axis of the PCA (6.88%, being the maximum 7.86%) (Figure S2). Summer
751 precipitation was selected to represent clim_p . Both of the terms estimated, the linear and the
752 quadratic terms, were statistically significant. The linear and quadratic terms were
753 statistically significant. Moreover, summer precipitation was moderately correlated with the

754 second PCA axis (-0.59) and it showed a moderate contribution to the axis (4.91%, being the
755 maximum 11.51%) (Figure S2). The second axis of the PCA explained nearly the same
756 amount of variability, 33.3% compared to 38.83%.

757

758 **Appendix S3.** Populations' with a Cook's distance value above one.

759 The Cook's distance measures the effect of deleting a given population due to the presence of
760 large residuals that can influence the accuracy of the model. We removed those populations
761 from the linear-fixed effect models. Specifically, in *P. nigra* we deleted the population of
762 Parapluberg in the three developmental classes (DC.1, DC. 2 and DC.3). In *P. pinaster*, we
763 deleted four populations (the population of Val Freda in DC. 3, and the populations of Ain
764 Baccouche, Tabarka and Valencia in DC. 1). In *P. pinea*, we deleted the population of Artvin
765 in DC1, DC2 and DC3.

766 **Supporting Information**

767 **Table S1.** Climatic variables used from the EuMedClim (Fréjaville & Benito Garzón, 2018).

768 **Table S2.** Results from linear mixed-effect models to select the climate variables of $clim_p$
769 and $clim_t$. Bold letters indicate the selected variables. Complementary results from the
770 Principal Component Analyses are shown in Figure S2

771 **Table S3.** Variance inflator factors (VIF) of the best-supported model for each pine species
772 analyzed.

773 **Table S4.** Mean and standard deviation values for each phenotypic plasticity index. Analysis
774 of the variance and post-hoc analyses adjusted by Tukey HSD were performed to test
775 differences among developmental classes (DC).

776 **Table S5.** Results from the analysis of the variance to test phenotypic plasticity variation
777 across developmental classes (DC) and for each pine species.

778

779

780 **Figure S1.** Maps showing the trials, provenances and natural distribution of the three species.
781 Red triangles represent the common gardens (trials) and light blue circles the provenances.
782 The light green area represents the natural distribution of the species according to
783 EUFORGEN (<http://www.euforgen.org/>). Top left: *Pinus nigra*, top right: *Pinus pinaster*,
784 bottom left: *Pinus pinea*. Adapted from Vizcaíno-Palomar *et al.*, (2019).

785 **Figure S2.** Plots of principal component analyses (PCA) for the short-term climate ($clim_t$)
786 and the long-term climate ($clim_p$). These results are complementary to the linear mixed-effect
787 model results shown in Appendix S2.

788 **Figure S3.** Plots of residuals of the best-supported model for tree height. Figures show the
789 residuals across age, $clim_t$ and $clim_p$ in standardized values. a) *P. nigra*, b) *P. pinaster*, and c)
790 *P. pinea*.

791 **Figure S4.** Populations' phenotypic tree height responses across $clim_t$ particularized for three
792 populations' origin (High, Average and Low in terms of $clim_p$ values) and for the three
793 developmental classes, DC, (Developmental Class 1: green, DC. 2: orange and DC. 3: blue)
794 for a) *P. nigra*, e) *P. pinaster* and i) *P. pinea*. Values of the coefficient of variation index (CV)
795 for the three developmental classes across the species natural distribution ranges are shown.
796 DC. 1: b), f) and j), DC. 2: c), g) and k); and DC. 3: d), h) and i).

797 **Figure S5.** Phenotypic plasticity predictions of the CV index across the standard deviation of
798 the maximum temperature of the warmest month (sd bio5) for the three developmental
799 classes and pines species.
800

801 **Table S1.** Climatic variables used from the EuMedClim (Fréjaville & Benito Garzón, 2018).

Annual climate variables		
Number	Variable name	Description
1	bio1	annual daily mean temperature (° C)
2	bio2	mean diurnal temperature range (max-min, ° C)
3	bio5	maximum temperature of the warmest month, (° C)
4	bio6	minimum temperature of the coldest month (° C)
5	bio12	annual precipitation (mm)
6	bio13	precipitation of the wettest month (mm)
7	bio14	precipitation of the driest month (mm)
8	PET	annual potential evapotranspiration [PET] (mm)
9	WAI	water availability [bio12- PET] (mm)
Seasonal climate variables		
Number	Variable name	Description
10	tmean.djf	winter daily mean temperature (° C)
11	tmean.mam	spring daily mean temperature (° C)
12	tmean.jja	summer daily mean temperature (° C)
13	tmean.son	autumn daily mean temperature (° C)
14	prec.djf	winter precipitation (mm)
15	prec.mam	spring precipitation (mm)
16	prec.jja	summer precipitation (mm)
17	prec.son	autumn precipitation (mm)
18	PET.min	PET of the wettest month (mm)
19	PET.max	PET of the driest month (mm)
20	WAI.min	WAI of the driest month (mm)
21	WAI.max	WAI of the wettest month (mm)

802

803 **Table S2.** Results from linear mixed-effect models to select the climate variables of clim_p
804 and clim_t. Bold letters indicate the selected variables. Complementary results from the
805 Principal Component Analyses are shown in Figure S2
806

<i>Pinus nigra</i>	Short term climate effects		Long term climate effects				
Variable	coef^2	t value	Variable	coef	t value	coef^2	t value
bio1_t	-0.24	-112.12	prec.jja _p	0.00	-0.01	-0.04	-0.28
tmean.djf _t	-0.20	-140.38	PET _p	-0.02	-0.12	-0.03	-0.19
PET.min _t	-0.13	-45.51	WAI.min _p	-0.02	-0.08	-0.02	-0.14
bio6 _t	-0.10	-127.72	bio5 _p	-0.02	-0.11	-0.02	-0.18
prec.jja _t	-0.08	-84.68	PET.max _p	-0.01	-0.07	-0.02	-0.10
prec.son _t	-0.07	-180.30	WAI_p	-0.01	-0.03	-0.01	-0.08
bio14 _t	-0.03	-149.63	bio2 _p	-0.02	-0.09	-0.01	-0.06
tmean.son _t	-0.03	-19.31	prec.djf _p	-0.05	-0.20	-0.01	-0.08
bio13 _t	-0.01	-29.53	tmean.jja _p	-0.02	-0.09	-0.01	-0.12
pet.max _t	-0.01	-11.58	prec.son _p	0.00	0.00	-0.01	-0.04
WAI.min _t	-0.01	-9.42	WAI.max _p	-0.07	-0.27	-0.01	-0.05
tmean.jja _t	0.01	10.06	bio13 _p	-0.05	-0.19	-0.01	-0.05
prec.mam _t	0.01	22.98	tmean.son _p	0.01	0.04	-0.01	-0.06
bio5 _t	0.02	28.96	bio1 _p	0.01	0.07	0.00	-0.05
bio12 _t	0.02	65.66	tmean.djf _p	0.06	0.31	0.00	-0.01
WAI _t	0.03	72.89	tmean.mam _p	-0.01	-0.05	0.00	0.00
PET _t	0.04	28.05	bio6 _p	0.05	0.26	0.00	0.00
prec.djf _t	0.05	100.84	prec.mam _p	-0.01	-0.04	0.00	0.00
tmean.mam _t	0.05	41.13	bio12 _p	-0.05	-0.20	0.01	0.03
bio2 _t	0.13	92.50	PET.min _p	0.04	0.16	0.01	0.04
WAI.max _t	0.13	211.40	bio14 _p	-0.04	-0.18	0.03	0.12

807

808

<i>Pinus pinaster</i>	Short term climate effects		Long term climate effects				
Variable	coef^2	t value	Variable	coef	t value	coef^2	t value
bio14 _t	-79.13	-77.05	prec.djf_p	92.89	5.83	-116.90	-15.43
PET_t	-62.09	-37.09	bio13 _p	124.55	10.02	-95.65	-13.56
WAI _t	-40.50	-34.15	bio2 _p	58.20	5.07	-75.43	-11.05
WAI.max _t	-28.87	-52.10	PET.max _p	101.64	7.73	-84.34	-9.88

<i>Pinus pinaster</i>	Short term climate effects		Long term climate effects				
Variable	coef^2	t value	Variable	coef	t value	coef^2	t value
tmean.son _t	-25.77	-21.76	prec.mam _p	75.40	6.56	-37.47	-6.85
bio2 _t	-17.74	-27.77	prec.jja _p	163.64	14.18	-42.90	-6.05
WAI.min _t	-15.79	-24.15	bio6 _p	5.58	0.47	-36.23	-5.04
bio5 _t	-11.70	-6.69	WAI.max _p	83.98	7.58	-26.93	-3.66
bio12 _t	-7.56	-9.46	prec.son _p	35.45	3.26	-33.29	-3.21
prec.djf _t	-4.04	-14.02	bio14 _p	123.28	8.70	-14.10	-1.89
PET.min _t	3.78	1.71	WAI _p	52.54	5.85	-12.40	-1.55
bio13 _t	5.18	15.19	bio12 _p	54.79	5.45	-10.01	-1.25
bio6 _t	10.16	13.75	PET _p	-39.41	-4.07	-6.07	-0.97
prec.son _t	11.66	49.06	tmean.son _p	22.65	1.78	-3.60	-0.47
PET.max _t	12.06	21.90	PET.min _p	-70.27	-5.89	-0.67	-0.11
tmean.djf _t	13.18	10.17	tmean.djf _p	29.28	2.29	-0.40	-0.05
prec.mam _t	14.62	27.30	WAI.min _p	64.09	7.41	2.43	0.32
prec.jja _t	18.26	15.69	bio5 _p	321.98	15.39	28.08	2.16
tmean.jja _t	62.56	31.59	bio1 _p	49.67	3.57	21.93	2.87
tmean.mam _t	89.81	71.02	tmean.mam _p	63.97	4.65	24.23	3.20
bio1 _t	142.09	56.23	tmean.jja _p	333.48	13.47	148.08	9.88

809

<i>Pinus pinea</i>	Short term climate effects		Long term climate effects				
Variable	coef^2	t value	Variable	coef	t value	coef^2	t value
bio5_t	-181.07	-152.89	pet.mean _p	0.38	0.21	-2.23	-2.43
prec.son _t	-99.92	-76.75	bio5 _p	-1.61	-0.80	-2.07	-1.91
pet.min _t	-54.20	-29.40	prec.jja_p	4.67	1.96	-1.88	-2.39
tmean.jja _t	-49.02	-67.42	PET.max _p	0.68	0.33	-1.34	-0.96
bio1 _t	-21.73	-24.04	WAI.min _p	-0.43	-0.20	-1.33	-1.03
bio6 _t	-17.76	-35.58	bio14 _p	2.99	1.09	-1.14	-1.64
bio14 _t	-11.35	-26.11	tmean.jja _p	-4.86	-2.47	-1.08	-0.75
tmean.son _t	-9.99	-16.22	tmean.mam _p	-5.34	-2.91	-0.66	-0.46
tmean.mam _t	-9.29	-16.19	tmean.djf _p	-5.26	-2.87	-0.36	-0.19
PET.max _t	-7.60	-12.53	PET.min _p	-2.05	-0.97	-0.34	-0.44
WAI.min _t	2.66	9.36	bio6 _p	-5.23	-2.68	0.21	0.10
prec.mam _t	5.11	13.96	bio2 _p	3.25	1.52	0.29	0.16
bio13 _t	7.00	10.69	bio1 _p	-5.87	-3.20	0.54	0.31
prec.djf _t	12.97	45.43	WAI _p	-5.80	-2.90	1.80	1.28
WAI.max _t	22.84	38.69	prec.son _p	-9.19	-3.92	2.59	2.26

<i>Pinus pinea</i>	Short term climate effects		Long term climate effects				
Variable	coef ²	t value	Variable	coef	t value	coef ²	t value
prec.jja _t	25.26	84.43	tmean.son _p	-7.19	-4.05	3.71	2.13
tmean.djft	35.07	47.88	prec.mam _p	-7.98	-3.36	3.84	2.59
bio2 _t	38.08	20.62	bio12 _p	-9.18	-4.64	4.60	3.16
PET _t	57.09	26.45	bio13 _p	-13.95	-6.45	8.76	5.41
bio12 _t	74.68	88.44	ppet.max _p	-14.19	-6.45	9.41	5.28
WAI _t	81.80	66.39	prec.djfp	-15.37	-6.54	9.61	5.70

810

811 **Table S3.** Variance inflator factors (VIF) of the best-supported model for each pine species
812 analyzed.

<i>Pinus nigra</i>		<i>Pinus pinaster</i>		<i>Pinus pinea</i>	
Variable	VIF	Variable	VIF	Variable	VIF
age	5	age	2	age	2
bio1 _t	1	PET _t	2	bio5 _t	2
bio1 _t ²	2	PET _t ²	2	bio5 _t ²	4
age ²	5	prec.djf _p	2	prec.jja _p	2
WAI _p	1	prec.djf _p ²	2	prec.jja _p ²	2
WAI _p ²	1	age ²	2	age ²	2
age × bio1 _t	3	age × PET _t	1	age × bio5 _t	4
age × WAI _p	3	age × prec.djf _p	1	age × prec.jja _p	1
bio1 _t × WAI _p	1	PET _t × prec.djf _p	1	bio5 _t × prec.jja _p	1
age × bio1 _t × WAI _p	3	age × PET _t × prec.djf _p	1	age × bio5 _t × prec.jja _p	1

813 **Table S4.** Mean and standard deviation values for each phenotypic plasticity index are shown.
814 Analysis of variance and post-hoc analyses adjusted by Tukey HSD were performed to test
815 differences among developmental classes (DC).
816

		DC. 1		DC. 2		DC. 3	
	P.I	mean	standard deviation	mean	standard deviation	mean	standard deviation
<i>P. nigra</i>	PP	0.75a	0.012	0.73b	0.009	0.65c	0.006
	CV	0.48a	0.015	0.45b	0.011	0.36c	0.006
<i>P. pinaster</i>	PP	0.86a	0.011	0.85b	0.001	0.84c	0.007
	CV	0.52a	0.015	0.51b	0.012	0.50c	0.007
<i>P. pinea</i>	PP	0.98a	0.000	0.98b	0.000	0.95c	0.000
	CV	0.82a	0.000	0.82b	0.000	0.80c	0.000

817 a, b and c (from the highest to the lowest) mean that mean values of phenotypic plasticity values are
818 statistically different among developmental stages for each index analyzed.
819

820 **Table S5.** Results from the analysis of the variance to test phenotypic plasticity variation across
821 developmental classes (DC) and for each pine species.
822

<i>Pinus nigra</i>	df	Sum Sq	F	p value	
PP index (DC)	2	0.478	2865.9	< 2.2e-16	***
Residuals	234	0.020			
CV index (DC)	2	0.569	2244.6	< 2.2e-16	***
Residuals	234	0.030			
<i>Pinus pinaster</i>	df	Sum Sq	F	p value	
PP index (DC)	2	0.036	209.59	< 2.2e-16	***
Residuals	552	0.048			
CV index (DC)	2	0.048	183.01	< 2.2e-16	***
Residuals	552	0.073			
<i>Pinus pinea</i>	df	Sum Sq	F	p value	
PP index (DC)	2	0	212713	< 2.2e-16	***
Residuals	162	0			
CV index (DC)	2	0.017	45936	< 2.2e-16	***

Residuals	162	0			
-----------	-----	---	--	--	--

823 df: degrees of freedom, Sum Sq means sum of squares, *F*-test, *** for p values < 0.001

824

Figure S1. Maps showing the trials, provenances and natural distribution of the three species. Red triangles represent the common gardens (trials) and light blue circles the provenances. The light green area represents the natural distribution of the species according to EUFORGEN (<http://www.euforgen.org/>). Top left: *Pinus nigra*, top right: *Pinus pinaster*, bottom left: *Pinus pinea*. Adapted from Vizcaíno-Palomar *et al.*, (2019).

Figure S2. Plots of principal component analyses (PCA) for the short-term climate (clim_t) and the long-term climate (clim_p). These results are complementary to the linear mixed-effect model results shown in Appendix S2.

838 a)

839 b)

840 c)

841
842 **Figure S3.** Plots of residuals of the best-supported model for tree height. Figures show the residuals
843 across age, clim_t and clim_p in standardized values. a) *P. nigra*, b) *P. pinaster*, and c) *P. pinea*.

Figure S4. Populations' phenotypic tree height responses across $clim_t$ particularized for three populations' origin (High, Average and Low in terms of $clim_p$ values) and for the three developmental classes, DC, (Developmental Class 1: green, DC. 2: orange and DC. 3: blue) for a) *P. nigra*, e) *P. pinaster* and i) *P. pinea*. Values of the coefficient of variation index (CV) for the three developmental classes across the species natural distribution ranges are shown. DC. 1: b), f) and j), DC. 2: c), g) and k); and DC. 3: d), h) and i).

Coefficient of variation index (CV)

Figure S5. Phenotypic plasticity predictions of the CV index across the standard deviation of the maximum temperature of the warmest month (sd bio5) for the three developmental classes and pines species.

References

- Fréjaville, T. & Benito Garzón, M. (2018) The EuMedClim Database: Yearly Climate Data (1901–2014) of 1 km Resolution Grids for Europe and the Mediterranean Basin. *Frontiers in Ecology and Evolution*, **6**, 31.
- Vizcaíno-Palomar, N., Benito Garzón, M., Alía, R., Giovannelli, G., Huber, G., Mutke, S., Pastuszka, P., Raffin, A., Sbay, H., Seho, M., Vauthier, D. & Fady, B. (2019) Geographic variation of tree height of three pine species (*Pinus nigra* Arn., *P. pinaster* Aiton and *P. pinea* L.) gathered from common gardens in Europe and North-Africa. *Annals of Forest Science*, **accepted**.