Case studies and mathematical models of ecological speciation. V. Adaptive divergence of whitefish in Fennoscandia.

Sunday 4th August, 2019

Xavier Thibert-Plante¹, Kim Præbel², Kjartan Østbye^{3,4}, Kimmo K. Kahilainen⁵, Per-Arne Amundsen⁶,
and Sergey Gavrilets⁷

¹Previous affiliations: Department of Ecology and Environmental Science, and IceLab, Umeå University, Umeå, Sweden. Phone: +1 (865) 974-9334. Fax: +1 (865) 974-9300.

²Norwegian College of Fishery Science, UiT The Arctic University of Norway, N-9037 Tromsø, Norway.

³Inland Norway University of Applied Sciences, Department of Forestry and Wildlife Management, Campus Evenstad, Anne Evenstadsvei 80, NO-2480 Koppang, Norway.

⁴Centre for Ecological and Evolutionary Synthesis (CEES), Department of Biosciences, University of Oslo, PO Box 1066 Blindern, NO-0316 Oslo, Norway.

⁵Inland Norway University of Applied Sciences, Department of Forestry and Wildlife Management, Campus Evenstad, Anne Evenstads vei 80, NO-2480 Koppang, Norway.

⁶Department of Arctic and Marine Biology, Faculty of Biosciences, Fisheries and Economics, UiT The Arctic University of Norway, N-9037 Tromsø, Norway.

⁷Department of Ecology and Evolutionary Biology, Department of Mathematics, National Institute for Mathematical and Biological Synthesis (NIMBioS), Center for the Dynamics of Social Complexity (DySoC), University of Tennessee, Knoxville, TN 37996 USA.

24 **Running title:** Diversification in Scandinavian whitefish

Keywords: gene flow; whitefish; speciation; divergence; individual-based simulation; parapatric
26 speciation

Supporting online material can be found at <http://xavier.thibert-plante.com/whitefish>. The
28 material is not published yet, therefore it is password protected. To access the information you will
need the username and password below (case sensitive).

30 User name: Scandinavia2

Password: Evolution45!

32 When the paper gets published, the online material will be archived to DRYAD or a similar service.

Acknowledgments

34 We thank Göran Englund, Jung koo Kang, Abisko Research Station, Polar Research grant, the
Norwegian Research Council (grant no. 186320), Centre for Ecological and Evolutionary Synthe-
36 sis (CEES), University of Oslo, the Office of Naval Research grant W911NF-17-1-0150, and the
National Institute for Mathematical and Biological Synthesis supported through the NSF awards
38 EF-0832858 and DBI-1300426, with additional support from The University of Tennessee. We
have no conflict of interest to declare.

40 Authorship Contributions

XTP, KP, KØ, KKK, PAA, and SG designed the experiment. XTP and SG designed the numerical
42 model. XTP coded and ran the simulations. XTP and SG analyzed the simulation output. XTP, KP,
KØ, KKK, PAA, and SG interpreted the results. XTP, KP, KØ, KKK, PAA, and SG wrote the paper.

Case studies and mathematical models of ecological speciation. V. Adaptive divergence of whitefish in Fennoscandia.

Running title: Diversification in Scandinavian whitefish

Keywords: gene flow; whitefish; speciation; divergence; individual-based simulation; parapatric speciation

Abstract

Modern speciation theory has greatly benefited from a variety of simple mathematical models focusing on the conditions and patterns of speciation and diversification in the presence of gene flow. Unfortunately the application of general theoretical concepts and tools to specific ecological systems remains a challenge. Here we apply modeling tools to better understand adaptive divergence of whitefish during the postglacial period in lakes of northern Fennoscandia. These lakes harbor up to three different morphs associated with the three major lake habitats: littoral, pelagic, and profundal. Using large-scale individual-based simulations, we aim to identify factors required for in situ emergence of the pelagic and profundal morphs in lakes initially colonized by the littoral morph. The importance of some of the factors we identify and study - sufficiently large levels of initial genetic variation, size- and habitat-specific mating, sufficiently large carrying capacity of the new niche - is already well recognized. In addition, our model also points to two other factors that have been largely disregarded in theoretical studies: fitness-dependent dispersal and strong predator pressure in the ancestral niche coupled with the lack of it in the new niche(s). We use our theoretical results to speculate about the process of diversification of whitefish in Fennoscandia and to identify potentially profitable directions for future empirical research.

Introduction

The diversity of species on Earth continues to provide inspiration for scientists studying speciation and the origins and maintenance of biodiversity. What makes these processes extremely complex and difficult to understand is that different evolutionary and ecological factors controlling their dynamics act simultaneously and often have opposing effects. The complexity of speciation processes implies that mathematical modeling can potentially play a very important role in uncovering its general rules and patterns. By now we have an impressive array of models and modeling techniques that shed light on the conditions, probability, waiting time to and duration of speciation, the degree of genetic and phenotypic divergence between the emerging species, and the way different resources (including space) are partitioned between the sister species. Models also explain the effects of different parameters and factors (such as the strength of selection, rates of mutation, recombination, migration, population size, number of loci, distribution of allelic effects, etc.) on the dynamics of speciation.^{1–9}

Most of this work has focused on models of speciation aims for both generality and mathematical simplicity. These models are very useful and insightful in uncovering general rules and patterns of speciation, adaptive radiation, and biological diversification. However, their generality almost necessarily implies that these models are very difficult to apply to any particular systems and species studied by empirical biologists. Therefore it is very important to supplement general models of speciation with those tailored for specific biological cases.

Studying models tailored for particular case studies can be very useful from several perspectives.⁹ First, mathematical models emerging from these projects do lead to a better understanding of the evolutionary dynamics of the studied specific systems. Second, although the relevant models are case-specific, they contribute towards building the general theory of speciation, e.g. by supporting or undermining the generality of particular observations and patterns. Third, the process of building a mathematical model even for a particularly well-studied empirical system usually reveals the lack of biological understanding or crucial empirical data needed to make appropriate modeling assumptions or specify parameters. This can greatly stimulate further empirical work to remove these limitations.

By now a relatively small number of such models have been developed for some of the best

studied systems. These include models aiming to capture the dynamics of non-allopatric speciation of cichlids in a lake^{10–13} and palms on an oceanic island,¹⁴ hybrid speciation in butterflies in Central America,¹⁵ ecomorph formation in marine snails in Sweden,¹⁶ pulmonate snails,¹⁷ and parallel adaptation in threespine stickleback.¹⁸

Here, we continue this work by focusing on a young and well-documented empirical system – the lacustrine European whitefish (*Coregonus lavaretus* (L.)) in Fennoscandian postglacial lakes. Ten to twenty thousand years ago Fennoscandia was covered by a thick ice sheet.¹⁹ The ice sheet retracted 8000 – 10000 years ago,^{20,21} creating a landscape dominated by inter-connected lakes and rivers and thus providing the opportunity for postglacial immigration of cold-water adapted freshwater fish, such as whitefish. After the glacier melting and land uplifting, this region has a high density of lakes, with deep lakes having three major habitats (littoral, pelagic and profundal) for fishes. Almost all lakes in this region contain whitefish, but most lakes have only one littoral morph, some lakes two morphs (littoral and pelagic), and a few large and deep Fennoscandian lakes three morphs.^{22–24} The different morphs of whitefish show habitat specific patterns in their resource use, body size, gill raker number, and life-history traits.^{22,23} Genetic evidence suggests a rapid divergence of these morphs from an ancestral monophyletic lineage, driven by natural selection on gill rakers and body size.^{22,25,26} Besides historical contingency, ecological opportunity mediated by interactions, such as resource competition and predation, may play an important role in diversification of whitefishes with regard to ecomorphology and life-history.^{24,27,28}

What is remarkable about this system, compared to other postglacial lakes with pelagic and littoral morphs, is the presence of up to three distinct morphs in some lakes. This raises the question: could they have diverged within the lake, or have different morphs evolved in different lakes and came into contact later? Our goal here is to use individual-based simulations to better understand conditions for and various factors (abiotic and biotic) controlling the divergence of Fennoscandian whitefish into three principal habitats of subarctic lakes. Our particular focus will be on the effects of selection for local adaptation, gene flow, carrying capacity of the habitat, and predation.

Methods

Model

Environment

We consider finite sexual diploid populations inhabiting isolated lakes. Each lake has three different ecological niches (habitats) with its own whitefish population: littoral, pelagic, and profundal. Each lake can have up to four predator species feeding on whitefish: pike, perch, burbot, and brown trout (Table 1).^{27,29,30} Time is discrete, the unit of time is one year, and the generations are overlapping.

Individual whitefish

Each individual is characterized by three phenotypic traits: size s , the number of gill rakers x , and whether or not it is sexually mature. We discretize size s into four stages: 0, 1, 2 and 3, roughly corresponding to the size of 2cm, 15cm, 20cm, and 30cm in whitefish. Size stage 0 describes newly born individuals.

We assume that immature fish can grow while mature fish stop growing and invest all their available energy into reproduction. If an immature fish survives to the end of the year, it either matures, with a probability m_s depending on its current size s , or grows to the next size class $s + 1$, with probability $1 - m_s$. Newly born individuals can only grow but not mature (i.e., $m_0 = 0$), whereas individuals reaching the largest size 3 always mature (i.e., $m_3 = 1$). Note that in our model maturation always happens at the end of the year reflecting the need to accumulate energy for gonad development e.g.³¹ Figure 1 illustrates these assumptions.

The studies on Coregonus clupeaformis^{32,33} and salmonids³⁴ show that maturation rate and the number of gill rakers are polygenic traits. Correspondingly, we assume that maturation rates m_1 and m_2 are additively controlled by L diallelic loci each. (In numerical simulations $L = 4$.) The corresponding allelic effects are scaled so that m_1 and m_2 stay between zero and one. The number of gill rakers x is also additively controlled by L loci but each locus has multiple discrete alleles. These alleles are subject to step-wise mutation and the effects are scaled so that trait x can take only integer values and mutation changes x only by plus and minus one. We assume that all genes

Figure 1: Life-history of a whitefish. The fish grows in size until it matures. The probabilities of maturation at stage 1 and 2 are m_1 and m_2 , respectively. If a whitefish reaches stage 3, it definitely matures the next year and stops growing.

are physically unlinked. Mutations happens with probability $\mu = 10^{-5}$ per gene per reproduction.

Selection

We assume that the whitefish population is subject to density-dependent viability selection due to intraspecific competition, habitat-specific stabilizing selection on gill raker number, size-dependent mortality due to predation as well as fertility selection due to maturation rates and body size differences.

Condition. The gill raker number x controls the food available to fish in a given environment.

We define the condition $\omega_j(x)$ of a whitefish with x gill rakers in niche j as

$$\omega_j(x) = \exp\left(-\frac{(x - \theta_j)^2}{2\sigma^2}\right). \quad (1)$$

Here θ_j is the optimum gill rakers number in niche j and σ^2 is a parameter measuring the strength of stabilizing selection on x towards this optimum. For whitefish, we set θ at 26, 36, and 18 for the littoral, pelagic, and profundal habitats, respectively. These values are representative for well-established morphs and are close to those observed in whitefish morphs LSR, DR, and SSR discussed above.^{24, 26}

Density-dependent selection. We assume that the population is subject to density-dependent mortality with the survival rate of individuals with x gill rakers and of size s in niche j being

$$v_j(s, x) = \frac{1}{1 + \frac{N_{s,j}}{K_{s,j}\omega_j(x)}}. \quad (2)$$

Here $N_{s,j}$ is the total number of individuals of size s in niche j , and constant parameter $K_{s,j}$ is the population size at which a population of perfectly adapted individuals (i.e., individuals with perfect condition $\omega_j = 1$) would have a survival rate of 0.5 (see the Appendix). We will interpret parameters $K_{s,j}$ as measures of carrying capacity of the corresponding sizes in the corresponding niches. The larger the number of competitors $N_{s,j}$ and the worse is the individual condition ω_j , the smaller is the survival rate v_j . Our formulation follows the Beverton-Holt model³⁵ and implies that density-dependent competition occurs only between individuals of the same size in the same ecological niche. The latter assumption reflects that fact that resource partitioning among whitefish morphs is very strong e.g.^{23, 36} and that intra-morph resource competition is much higher and the most pronounced between individuals of nearest age cohorts and size classes using same resources at the same habitats.^{37, 38}

Predation. Maximum gape of a predator restricts the sizes and shapes of prey that can be eaten. We assume that each predator species in our system is characterized by a maximum prey size g_k . To describe predation, we posit that predator k removes a random proportion π_k of surviving whitefish with sizes smaller than or equal to its gape size g_k . Values π_k are parameters in our model. When whitefish are bigger than the maximum prey size of a predator, they are not predated upon. The maximum prey size of the four predators for whitefish are approximated in Table 1. Note that each predator is found in only a subset of habitats.

Predator	Max prey size	Habitats
Pike	25cm	Littoral
Perch	15cm	Littoral
Burbot	20cm	Littoral, Profundal
Brown trout	15cm	Littoral, Pelagic

Table 1: Habitats and the maximum prey size for different whitefish predators.^{27,29,30}

Dispersal

Individuals can change their ecological niches within the lake. We assume that, each year, after reproduction each fish chooses the ecological niche to go to with probabilities proportional to their overall survival in each niche (accounting for both density-dependent selection and predation). One interpretation of this assumption is that each fish samples different niches before deciding on the one to stay. Fitness-dependent migration has been used in a number of earlier ecological models^{39–42} but has apparently been neglected in speciation modeling. Note that after each dispersal event, the niche population sizes change, so niche “attractiveness” can change as well. In numerical implementation of the model, to avoid a bias due to a sequence of events during the dispersal, individuals are chosen at random (without replacement, so that individuals move, at most, once per time step) to make a dispersal move.

Reproduction

We assume that mating is assortative by the size and the niche. [This modeling choice can also accommodate a scenario where fish do not mate within their physical niches but rather mate at a niche- and size-specific mating ground or mating time.] Each surviving female chooses a mate at random from an appropriate set of males. She then produces a random number of offspring sampled from a Poisson distribution with size-specific means b_1, b_2 and b_3 . All offspring are born in the same niche as their parents.

Life cycle

We assume there is a yearly sequence of events in the life of every fish. It starts with random death of eggs which is followed by density-dependent survival which is followed by predation mortality.

The surviving fish grow or reproduce. Then all fish, except the newborn, have the opportunity to
 204 disperse among the niches within the lake, before this cycle starts again.

Numerical simulations

206 Scenarios

We simulated two scenarios of diversification differing in initial conditions. The Colonization-L
 208 scenario reflects the current view of the post-glacial colonization of lakes by large littoral fishes.²⁴
 In this scenario, the initial population of size 3,000 eggs is introduced in the littoral niche to which
 210 it is adapted. Its modal trait values are: $x = 26$, $m_1 = 0$, and $m_2 = 1$, that is, fish grows to the
 “large” stage 2 only. The Colonization-G scenario is similar as Colonization-L scenario, but all
 212 initial colonizers are assumed to grow bigger, i.e. reach the “giant” stage three ($x = 26$, $m_1 = 0$, and
 $m_2 = 0$).

214 In both scenarios, the initial population harbors genetic and phenotypic variation. The standard
 deviation of the gill raker number is five (which is close to values observed in natural populations).
 216 In the genes controlling the probabilities of maturation, one allele has frequency 95% and another
 has frequency of 5%. After introduction, the population then evolves for 10,000 years. We then
 218 evaluate if and how many different morphs emerge and are maintained in the lake.

Parameter values

220 In numerical simulations, besides the initial conditions, we also varied the predation intensity π .
 Specifically, for each of the four predators we set π at 0%, 30%, and 50%.

222 In our model, fertility parameters $\{b_1, b_2, b_3\}$ are set to $\{4, 16, 64\}$ respectively, following an
 exponential relationship with length.^{43,44} Note that these numbers should be interpreted not as
 224 an actual number of eggs produced by a fish but rather as a number of offspring surviving to the
 moment when they are subject to selection.

226 To estimate carrying capacities $K_{s,j}$ (used in eq. (2)) we used available data assuming that
 extant fish are adapted to their environment, i.e. that they have the optimal number of gill rakers
 228 and reach maturation at a proper size ($m_1 = m_2 = 0$ for the littoral niche, $m_1 = 1$ for the pelagic niche,

Stage	Littoral	Pelagic	Profundal
0 (2cm)	15125	5517	656
1 (15cm)	3025	6000	66
2 (20cm)	756	0	150
3 (30cm)	1000	0	0

Table 2: Equilibrium population densities $N_{s,j}$ for fish of different sizes in different habitats in the baseline model.

and $m_1 = 0, m_2 = 1$ for the profundal niche). We also assumed that extant fish in three niches are reproductively isolated and that fertility is a function of the female size.^{43, 44}

We set the equilibrium densities of the largest morph in each niche to 1,000, 6,000, and 150 individuals for the littoral, pelagic, and profundal niche, respectively. These values are proportional to the observed densities in each niche.^{45, 46}

From these values and empirical data, we estimated the corresponding equilibrium densities for other morphs (Table 2) and the values of corresponding carrying capacity parameters $K_{s,j}$ (Table S1 in the Appendix).

The carrying capacities of different niches vary among different lakes. A particularly important factor affecting the likelihood of diversification is the carrying capacity of the profundal niche which can vary because of the size of the lake, its shape, and the nutrient deposition rate. In this study we used three different carrying capacities for the profundal niche resulting in equilibrium densities equal to one, two, and four times the values given in Table 2. We kept carrying capacities of the other two niches constant.

Overall, the results we present here explore $2 \times 3^4 \times 3 = 486$ different combinations of parameters, assuming assortative mating and fitness-dependent dispersal, for each of which we did 10 independent stochastic runs. We also performed a similar investigation of four related sets of models: 1) with random (rather than size-assortative) mating and random (rather than fitness-dependent) dispersal, 2) with size-assortative mating and random dispersal, and 3) with random mating and fitness-dependent dispersal. We also investigated our model with 4) smaller initial standard deviations in gill raker number and lower variation in the maturation loci. We never observed the emergence of the profundal morph in these models so we do not present the corresponding results.

Evolutionary outcomes

To interpret our numerical results, we say that a given niche has been successfully colonized if in the last year the population of newborn both 1) has gill rakers number adapted to their niche and 2) they mostly grow to a size which characterizes the niche in real lakes.

The former assumption was formalized as the requirement that the average gill raker number x is close to the optimum value, specifically: $\theta_j - \sigma \leq x \leq \theta_j + \sigma$. (Recall that σ is a parameter measuring the strength of stabilizing selection.) The latter assumption was formalized as requirements on the evolved average maturation rates m_1 and m_2 . Specifically,

- if $m_1 > \frac{2}{3}$ (i.e., fish mostly matures at size 1), then we say the fish is pelagic;
- if $m_1 < \frac{1}{3}, m_2 > \frac{2}{3}$ (i.e., fish mostly matures at size 2), then we say the fish is profundal;
- if $m_1 < \frac{1}{3}, m_2 < \frac{1}{3}$ (i.e., fish mostly reaches size 3), then we say the fish is littoral.

Results

Here we present our main results on the conditions under which each niche was colonized and by how many morphs. [Our results for the entire parameter space studied can be found online (<https://bit.ly/2qCFR6P>) (username: Scandinavia2, password: Evolution45! - the password protection will be removed when the paper is published). The online browser will be archived on DRYAD (DOI available later).]

When we look across the entire parameter space, the majority of lakes fall into four different compositions of morphs present: littoral morph only, pelagic morph only, littoral and pelagic morphs, and all three morphs present (Tables 3 and 4). In most case, the system reached a stochastic equilibrium state on the time scale of a few hundred generations (see the link above for some examples). This observation on the speed of diversification is comparable with earlier speciation models (e.g., [gavrilets2004](#), [gavriletsVose2005](#), [gavriletsEtAl2007](#)). Below we will break down the conditions under which each outcome was more likely to be observed.

Relative carrying capacity	1.0	2.0	4.0
None	2	2	1
Monomorphic	479	495	509
Littoral	439	477	501
Pelagic	39	17	8
Profundal	1	1	0
Dimorphic	329	313	295
Littoral Pelagic	329	313	279
Profundal Littoral	0	0	16
Profundal Pelagic	0	0	0
Three morphs	0	0	5

Table 3: Number of simulations with different compositions of morphs present for different relative carrying capacities of the profundal niche in Colonization-L scenario.

Relative carrying capacity	1.0	2.0	4.0
None	0	0	0
Monomorphic	425	484	520
Littoral	422	482	519
Pelagic	3	2	1
Profundal	0	0	0
Dimorphic	385	326	290
Littoral Pelagic	385	326	282
Profundal Littoral	0	0	8
Profundal Pelagic	0	0	0
Three morphs	0	0	0

Table 4: Number of simulations with different compositions of morphs present for different relative carrying capacities of the profundal niche in Colonization-G scenario.

Only littoral morph (no diversification). Survival of the littoral morph with no other morphs emerging was the most common outcome in our simulations (Tables 3, 4 and Figures 2, 3). It typically required the presence of predation by trout. Lakes with only the littoral morph are common in Fennoscandia.²⁴ Their abundance in the geologically younger region is one of the reasons that we hypothesize that the littoral morph is ancestral to Fennoscandia lakes.

Two morphs. The emergence of the pelagic morph and the maintenance of the littoral morph was a frequent outcome in both colonization scenarios (about 40%). This outcome occurs more often when trout predation was absent or small ((Tables 3, 4 and Figures 2, 3)). There were also a few cases (24) where the divergence occurred toward the profundal morph, instead of the

pelagic (Tables 3, 4 and Figures 2c, 3c). All these cases were observed when the profundal environment was large and no burbot predator was present. This outcome is usually not observed in Fennoscandia.

Single pelagic morph. Although lakes with only the pelagic morph present are not observed in Fennoscandia,²⁴ but they were observed in our simulations. This outcome implies the extinction of the littoral morph after it gives rise to the pelagic morph. This outcome was observed in both colonization scenarios under high predation by pike and perch but much more common in the Colonization-G scenario (Tables 3, 4 and Figures 2d, 3d).

Three morphs. The cases with all three morphs present were very rare in our simulations - 5 cases only. They all happened in Colonization-L scenario under large carrying capacity for the profundal niche (Table 3) when predation from pike and perch was strong but that from trout and burbot was weak (Figure 2e). Lakes with all three morphs are rarely observed in Fennoscandia.²⁴

Failed adaptation. Four simulations resulted in populations with intermediate traits according to our metrics. These failed adaptations all involved high predation. Additionally, in one simulation with strong predation, the population went completely extinct.

Overall, both monomorphic and polymorphic lakes emerged from our simulations. The outcomes of Colonization-L and Colonization-G scenarios were similar, but Colonization-L scenario was more prone to diversification. The reason is that Colonization-L scenario started with the gill raker numbers typical for the littoral morph but with the size close to those for the pelagic and profundal morphs. This simplified the emergence of these two morphs. The lakes with all three niches colonized were observed but very rarely.

Discussion

We used individual-based simulations to study the likelihood of within-lake ecological and morphological diversification in Scandinavian whitefish in the post-glacial time frame (about 10,000 years). Our simulations show a common emergence of lakes where, in addition to the originally colonizing littoral morph, either the pelagic or profundal morphs have evolved and become established. The former outcome was much more frequent than the latter. We also observed the presence of lakes with all three morphs present, although very infrequently. However, this result fits with the

observations from the wild which have revealed only a handful of trimorphic systems.

We started our simulations with a small founder population adapted to the littoral niche and no individuals adapted to the two other available niches: pelagic or profundal. Colonizing the pelagic niche required individuals to evolve increased gill raker number and smaller size. Colonizing the profundal niche required individuals to evolve decreased gill raker number and smaller size. The formation of new morphs typically happened on the time-scale of a few hundred generations. This is similar to other models with some initial genetic variation and strong selection.^{7,47,48}

Speciation theory tells us though that the mere existence of “empty” ecological niches does not guarantee that they will be colonized and “filled” by locally adapted organisms, especially if there is a possibility of gene flow from the ancestral niche. Some additional factors must usually be in place to simplify survival in the new environment and reduce the effects of deleterious gene flow preventing local adaptation. Several such factors turned out to be very important in our model as well.

First, populations must have sufficiently large initial genetic variation. In our simulations presented above, we set the initial standard deviation of the gill rakers at 5. With a lower standard deviation (and, correspondingly, lower initial genetic variation), the state with both littoral and pelagic morphs was possible, but the profundal morph never emerged. Second, diversification required mating to be assortative with respect to both size and ecological niche. Third, dispersal had to be fitness-dependent. Both these assumptions result in reduced effects of gene flow from the ancestral niche. With random mating and/or dispersal, empty niches remained largely empty. Colonization of the profundal niche was mainly observed when its carrying capacity was the largest. Biologically, this describes lakes that are large and deep. Large carrying capacity simplified survival in the new niche by effectively reducing within-niche competition. These four assumptions alone were still not enough to ensure diversification. The fourth important factor was the elevated predation pressure in the littoral niche. Colonization of the pelagic niche largely required the absence of brown trout which is the only pelagic predator in model assumptions. Correspondingly, colonization of the profundal niche largely required the absence of burbot which is the only predator that can be found there. By starting to utilize the niche with no predators, the fish gets an immediate additional survival advantage which does not require evolving any genetic adaptations. Moreover, the absence of predators in a niche implies that fish does not have to grow large and

thus could evolve earlier maturation. The latter in turn would lead to faster population growth.

The appearance of the pelagic morph in addition to the littoral one was much more common than that of the profundal morph, which is in accordance with findings from empirical studies.²⁴ This happens because in our model, reflecting the situation in most lakes, the carrying capacity of the pelagic niche was much larger than that of the profundal niche making the pelagic niche much easier to colonize. Diversification was promoted if initial colonizers were littoral fish of “large” (Colonization-L scenario) rather than “giant” size (Colonization-G scenario), which happens because the size of the former is closer to those of the two other morphs (that are smaller than the littoral) and thus much smaller evolutionary change is required. However, body size is very plastic in whitefish and influenced by resource availability and strength of competition (e.g. amundsenEtAl2004, kahilainenEtAl2005). The Colonization-G scenario may nevertheless be relevant as large size likely is beneficial for migration. Following this scenario, “giant” colonizers may initially have experienced untapped resources and fast growth rates, but subsequently turned into the “large” category due to reduced growth rates from increasing fish density and intraspecific resource competition. The modeling results corroborate with a recent empirical study from northern Fennoscandian whitefish indicating that body size and number of gill rakers are both targeted by natural selection.²⁶

Next, we discuss whether the factors identified in our model are present in natural populations of whitefish. Large genetic variation in colonizing individuals may have derived from the ancestral populations in the postglacial refugia, hybridization events during the loss of their pre-glacial habitats, and ongoing hybridization in contemporary times. Hybridization has an important role in historical and contemporary evolution of whitefish.^{49,50} Whitefish is well-known for its high capacity for both fast forward and reverse speciation, which suggest a role of hybridization behind the high genetic variation harbored by the whitefish.^{22,25} In the current modeling approach the minimum variation of five gill rakers of the ancestral morph was needed for the divergence to other morphs. Such variation of gill raker number is typical for generalist whitefish throughout much of the distribution suggesting ecological relevance of the model assumptions. However, the very few lakes with all three morphs in the simulation model overlooks the early individual specialization to all three habitats by the ancestral colonizing morph.⁵¹ Such step is very likely a crucial initial step towards full-fledged ecological speciation described as five model simulations leading to occur-

rence of all three morphs. While the proportion of simulations leading to three morphs is around
 374 0.6%, it could be indeed comparable to natural conditions where this morph is present only at very
 few large lakes having wide profundal habitat coverage and many predators in littoral and pelagic
 376 niches.^{27,52} Furthermore, frequent occurrence of pelagic and littoral morphs (36-40%) of simula-
 tions obviously refers to frequent divergence to habitats providing main energy sources (pelagic
 378 phytoplankton and littoral algae/macrophytes) to the lake food webs. In contrast, the profundal
 habitat is an unproductive habitat and dependent on energy inputs (settling materials) from the
 380 pelagic and littoral. These pelagic and littoral inputs vary among systems due to lake morphology
 and productivity and this complexity makes whitefish divergence to this habitat rare.^{24,52}

Size assortative mating is common in fish.⁵³⁻⁵⁵ Body size and size at sexual maturation of
 whitefish morphs are highly correlated with their resource use. Specifically, specialization to littoral
 384 benthic prey leads to large size and late maturation, pelagic prey to smallest size and fastest mat-
 uration, and finally profundal benthic prey to intermediate body size and late maturation.^{22,23,37,38}
 386 Such differences in body size and maturation would be a logical step towards size assortative
 mating, if similar sized mature individuals among morphs are rarely present at the spawning sites.
 388 At least in the pelagic morph this seems likely as the very high mortality of small pelagic morph
 suggest that very few individuals are able to survive and grow to the size when other morphs
 390 mature.^{37,38} Reproductive isolation may be further strengthened by temporal and spatial differ-
 ences in time and place of spawning.^{56,57} Such differences in spawning may arise via the habitat
 392 specific differences in temperature that regulates both prey resource availability and initiation of
 spawning.^{31,58} In Fennoscandia, the littoral benthic resources are at the highest from early to
 394 mid-summer, pelagic zooplankton at mid to late summer and profundal benthic resources at late
 season.^{31,37,38} The littoral reaches the highest water temperature in summer, but cools down at
 396 the earliest time followed by the pelagic and profundal.⁵⁹ Differences in realized water temperature
 among the whitefish morphs may be a major driver of divergence in Fennoscandian whitefish,[?] as
 398 divergent temperature regimes alter the maturity status of all three morphs. There is also some
 evidence of spatial divergence of spawning sites among the morphs and in many salmonids such
 400 spawning site fidelity maintains population divergence.⁶⁰⁻⁶² In our model, we assumed that in-
 dividuals exhibit certain habitat preferences. In fish, those can take different form such as natal
 402 environment imprinting, condition dependent, density dependent, and predator avoidance.^{63,64}

In the model, the lack of pelagic (brown trout) and profundal (burbot) predation was a needed prerequisite for the raise of both pelagic and profundal whitefish morphs. Such conditions where strong predation occurs solely on littoral habitats of large lakes seems unlikely in contemporary conditions.^{27,29} However, in the large lakes without polymorphic whitefish, brown trout and other predators (pike, burbot, perch) indeed used mostly littoral habitat.⁵² It is likely that the raise of pelagic whitefish morph induce brown trout shift to pelagic habitat use and piscivory.^{30,52} Similar process could be present with regard to burbot, where divergence of profundal morph provides a new forage fish. However, burbot is a dark active predator that frequently use diel bank migration to feed on more abundant littoral prey resources.⁶⁵ Thus, the empirical data suggest that modeling requirements for predation could be met in nature, but such conditions are rare.

Due to computational considerations, our model has some obvious limitations. For example, we assumed that fish laid a relatively small number of eggs. In reality, fish can produce a very large number of eggs; much larger than what we could simulate using an individual-based approach. We expect that with larger number of eggs, selection will be more efficient and divergence may occur faster than what is observed here (cf. gir10). The initial conditions were centered only on scenarios of “giant” and “large” colonizers, but it is very difficult to determine the actual body size of the ancestor starting to diverge. However, increasing intraspecific competition for resources is likely after the colonization of a new lake, which tend to shrink the body size supporting shift from giant to smaller body size. Predation by multiple species had a strong effect on whitefish divergence, but in the wild there are lake systems with whitefish morphs without main predators or very low amount of predation.²⁶ However, the majority of lakes with three morphs have abundant predator populations and predation is likely to have strong influence on life-history divergence of prey. It is also likely that prey and predators co-evolve during the divergence process⁶⁶ but individual-based modeling of multi-species evolutionary processes is inherently difficult. The process of building the model has also revealed important gaps in the available data on whitefish populations and their environments. In particular, having more precise estimates of the population sizes, predation rates, and fecundity of different morphs would increase the power of our model.

In our model, the body size was subject to direct selection by predation and also due to fertility and maturation rate differences. We did not consider explicitly body size adaptation to the ecological niche. However since body size correlates with gill raker number,^{23,36,67} our model partially

captures this effect.

The sizes and depth of lakes in Fennoscandia vary a lot as well as their productivity with evident implications to species divergence.²⁴ In our simulations, we considered only three major lake habitats and assumed pre-determined carrying capacities for each habitat. Lake morphometry and productivity in addition to de-glaciation history likely fine-tunes the divergence processes further,²⁴ but these are far too complex scenarios to model conclusively. Further modelling effort is needed to understand vertical pelagic divergence documented for example to Alpine whitefish⁶⁸ and North American ciscoes.⁶⁹ Also, it is well established that a small number of loci of large effect are more conducive to speciation than a large number of loci with small effects.^{7,48,70} We have chosen a small number of loci to run our simulation to facilitate the process of adaptive radiation. With larger number of loci, we would not expect much diversification to be observed in our model. Given that diversification in whitefish does happen, we expect that the underlying traits are controlled by a small number of loci with large effects.

Overall, our modeling supported the possibility of divergence to three lake habitats during the postglacial time-frame, although such cases were rare and required a large profundal habitat without predators. Divergence to pelagic and littoral morphs was much more frequent and occurred with various levels of predation. The modeling effort indicated how little actually we know about reproductive isolation mechanisms as well as about habitat-specific carrying capacities and essential population dynamic parameters such as age-specific mortality in whitefish. Nevertheless, it would be interesting to apply our modeling approach to a true deep water species, such as Lake Baikal *Cottus* and various members of *Salvelinus*, in very large lakes using species-specific parameters.^{71–73} Recent empirical studies have highlighted the important role of deep water habitats which have been previously neglected in monitoring efforts.^{74,75} In these respects, mathematical modeling could provide an efficient predictive tool for finding new lakes with three morphs, as well as identification of key factors contributing to their disappearance from the postglacial lakes.

References

- ¹ S. A. H. Geritz and É. Kisdi. Adaptive dynamics in diploid, sexual populations and the evolution of reproductive isolation. Proceedings of the Royal Society Biological Sciences Series B,

267(1453):1671–1678, 2000.

² M. Kirkpatrick and V. Ravigné. Speciation by natural and sexual selection: Models and experiments. The American Naturalist, 159(Supplement):S22–S35, 2002.

³ M. R. Servedio and M. A. F. Noor. The role of reinforcement in speciation: Theory and data. Annual Review of Ecology, Evolution, and Systematics, 34(1):339–364, 2003.

⁴ U. Dieckmann, M. Doebeli, J. A. J. Metz, and D. Tautz. Adaptive speciation. Cambridge Univ. Press, Cambridge, U.K., 2004.

⁵ Michael Doebeli. Adaptive Diversification (MPB-48). Princeton University Press, 2011.

⁶ S. Gavrillets. Perspective: Models of speciation: What have we learned in 40 years? Evolution, 57(10):2197–2215, 2003.

⁷ S. Gavrillets. Fitness landscapes and the origin of species. Princeton University Press, Princeton, New Jersey, USA, 2004.

⁸ G. S. van Doorn, P. Edelaar, and F. J. Weissing. On the origin of species by natural and sexual selection. Science, 326(5960):1704–1707, 2009.

⁹ S. Gavrillets. Models of speciation: Where are we now? Journal of Heredity, 105(Special Issue):743–755, 2014.

¹⁰ R. Lande, O. Seehausen, and J. J. M. van Alphen. Mechanisms of rapid sympatric speciation by sex reversal and sexual selection in cichlid fish. Genetica, 112-113:435–443, 2001.

¹¹ S. Gavrillets, A. Vose, M. Barluenga, W. Salzburger, and A. Meyer. Case studies and mathematical models of ecological speciation. 1. Cichlids in a crater lake. Molecular Ecology, 16(14):2893–2909, 2007.

¹² M. Kawata, A. Shoji, S. Kawamura, and O. Seehausen. A genetically explicit model of speciation by sensory drive within a continuous population in aquatic environments. BMC Evolutionary Biology, 7:99, 2007.

¹³ R. Aguilée, A. Lambert, and D. Claessen. Ecological speciation in dynamic landscapes. Journal of Evolutionary Biology, 24(12):2663–2677, DEC 2011.

¹⁴ S. Gavrillets and A. Vose. Case studies and mathematical models of ecological speciation. 2.
488 Palms on an oceanic island. Molecular Ecology, 16(14):2910–2921, 2007.

¹⁵ E. A. Duenez-Guzman, J. Mavárez, M. D. Vose, and S. Gavrillets. Case studies and mathemat-
490 ical models of ecological speciation. 4. Hybrid speciation in butterflies in a jungle. Evolution,
63(10):2611–2626, 2009.

¹⁶ Suzanne Sadedin, Johan Hollander, Marina Panova, Kerstin Johannesson, and Sergey
492 Gavrillets. Case studies and mathematical models of ecological speciation. 3: Ecotype formation
494 in a swedish snail. Molecular Ecology, 18(19):4006–4023, 2009.

¹⁷ Masato Yamamichi and Akira Sasaki. Single-gene speciation with pleiotropy: effects of allele
496 dominance, population size and delayed inheritance. Evolution, 67(7):2011–2023, JUL 2013.

¹⁸ M. Roesti, S. Gavrillets, A. P. Hendry, W. Salzburger, and D. Berner. The genomic signature of
498 parallel adaptation from shared genetic variation. Molecular Ecology, 23(16):3944–3956, 2014.

¹⁹ Jan Mangerud, Martin Jakobsson, Helena Alexanderson, Valery Astakhov, Garry KC Clarke,
500 Mona Henriksen, Christian Hjort, Gerhard Krinner, Juha-Pekka Lunkka, Per Möller, Andrew
Murray, Olga Nikolskaya, Matti Saarnisto, and John Inge Svendsen. Ice-dammed lakes and
502 rerouting of the drainage of northern eurasia during the last glaciation. Quaternary Science
Reviews, 23(11):1313–1332, 2004.

²⁰ J. L. Sollid, S. Andersen, N. Hamre, O. Kjeldsen, O. Salvigsen, S. Sturød, T. Tveitå, and A. Wil-
504 helmsen. Deglaciation of finnmark, north norway. Norsk Geografisk Tidsskrift - Norwegian
506 Journal of Geography, 27(4):233–325, 1973.

²¹ Raimo Kujansuu, Brita Eriksson, and Tuulikki Grönlund. Lake
508 Inarijärvi, northern Finland: sedimentation and late Quaternary evolution. Number 143.
Geological Survey of Finland, 25 p., 1998.

²² K. Østbye, P-A. Amundsen, L. Bernatchez, A. Klemetsen, R. Knudsen, R. Kristoffersen, T. F.
510 Naesje, and K. Hindar. Parallel evolution of ecomorphological traits in the european white-
512 fish coregonus lavaretus (l.) species complex during postglacial times. Molecular Ecology,
15(13):3983–4001, 2006.

²³ Chris Harrod, Jennie Mallela, and Kimmo K. Kahilainen. Phenotype-environment correlations in a putative whitefish adaptive radiation. Journal of Animal Ecology, 79(5):1057–1068, 2010.

²⁴ A. Siwertsson, R. Knudsen, K. Kahilainen, K. Præbel, R. Primicerio, and P.-A. Amundsen. Sympatric diversification as influenced by ecological opportunity and historical contingency in a young species lineage of whitefish. Evolutionary Ecology Research, 12:929–947, 2010.

²⁵ Kim Præbel, Rune Knudsen, Anna Siwertsson, Markku Karhunen, Kimmo K. Kahilainen, Otso Ovaskainen, Kjartan Østbye, Stefano Peruzzi, Svein-Erik Fevolden, and Per-Arne Amundsen. Ecological speciation in postglacial european whitefish: rapid adaptive radiations into the littoral, pelagic, and profundal lake habitats. Ecology and evolution, 3(15):4970–4986, 2013.

²⁶ K. Häkli, K. Østbye, K.K. Kahilainen, P.-A. Amundsen, and K. Præbel. Diversifying selection drives parallel evolution of adaptive trait utility to the main lake habitats along the speciation continuum in European whitefish. Ecology and Evolution, 8:2617–2631, 2018.

²⁷ Kimmo Kahilainen and Hannu Lehtonen. Piscivory and prey selection of four predator species in a whitefish dominated subarctic lake. Journal of Fish Biology, 63(3):659–672, 2003.

²⁸ Per-Arne Amundsen, Rune Knudsen, Anders Klemetsen, and Roar Kristoffersen. Resource competition and interactive segregation between sympatric whitefish morphs. 50(1):301–307, 2004.

²⁹ Per-Arne Amundsen, Thomas Bøhn, Olga A Popova, Frode J Staldvik, Yuri S Reshetnikov, Nikolay A Kashulin, and Anatoly A Lukin. Ontogenetic niche shifts and resource partitioning in a subarctic piscivore fish guild. Hydrobiologia, 497(1-3):109–119, 2003.

³⁰ Hallvard Jensen, Kimmo K. Kahilainen, Per-Arne Amundsen, Karl Øystein Gjelland, Antti Tuomaala, Tommi Malinen, and Thomas Bøhn. Predation by brown trout (Salmo trutta) along a diversifying prey community gradient. Canadian Journal of Fisheries and Aquatic Sciences, 65(9):1831–1841, 2008.

³¹ Brian Hayden, Chris Harrod, and Kimmo K. Kahilainen. Dual fuels: intra-annual variation in the relative importance of benthic and pelagic resources to maintenance, growth and reproduction in a generalist salmonid fish. Journal of Animal Ecology, 83(6):1501–1512, 2014.

³² Pierre-Alexandre Gagnaire, Eric Normandeau, Scott A. Pavey, and Louis Bernatchez. Mapping phenotypic, expression and transmission ratio distortion qtl using rad markers in the lake whitefish (*Coregonus clupeaformis*). *Molecular Ecology*, 22:30363048, 2015.

³³ Anne-Marie Dion-Cote Rogers, Eric Normandeau, Pierre-Alexandre Gagnaire, Anne C. Dalziel, Jobran Chebib, and Louis Bernatchez. Rad-qtl mapping reveals both genome-level parallelism and different genetic architecture underlying the evolution of body shape in lake whitefish (*Coregonus clupeaformis*) species pairs. *G3: GENES, GENOMES, GENETICS*, 5:1481–1491, 2015.

³⁴ Arne Jacobs, Robyn Womack, Mel Chen, Karim Gharbi, and Kathryn R. Elmer. Significant synteny and colocalization of ecologically relevant quantitative trait loci within and across species of salmonid fishes. *Genetics*, 207:741–754, 2017.

³⁵ M. Kot. *Elements of Mathematical Ecology*. Cambridge, Cambridge University Press, 2001.

³⁶ K. Kahilainen and K. Østbye. Morphological differentiation and resource polymorphism in three sympatric whitefish *coregonus lavaretus* (L.) forms in a subarctic lake. *Journal of Fish Biology*, 68(1):63–79, 2006.

³⁷ K. Kahilainen, H. Lehtonen, and K. Könönen. Consequence of habitat segregation to growth rate of two sparsely rakered whitefish (*Coregonus lavaretus* (L.)) forms in a subarctic lake. *Ecology of Freshwater Fish*, 12(4):275–285, 2003.

³⁸ K. Kahilainen, E. Alajärvi, and H. Lehtonen. Planktivory and diet-overlap of densely rakered whitefish (*coregonus lavaretus* (L.)) in a subarctic lake. *Ecology of Freshwater Fish*, 14(1):50–58, 2005.

³⁹ Paul R. Armsworth and Joan E. Roughgarden. The structure of clines with fitness-dependent dispersal. *American Naturalist*, 172(5):648–657, NOV 2008.

⁴⁰ Paul R. Armsworth. Conditional dispersal, clines, and the evolution of dispersiveness. *Theoretical Ecology*, 2(2):105–117, JUN 2009.

⁴¹ Jonathan T. Rowell. The limitation of species range: A consequence of searching along resource gradients. *Theoretical Population Biology*, 75(2-3):216–227, MAR-MAY 2009.

⁵⁶⁸ ⁴² Robert Stephen Cantrell, Chris Cosner, Yuan Lou, and Chao Xie. Random dispersal versus fitness-dependent dispersal. Journal of Differential Equations, 254(7):2905–2941, APR 1 2013.

⁵⁷⁰ ⁴³ J. P. Schneider, P. W. Laarman, and H. Gowing. Length-weight relationships. chapter 17. In Manual of fisheries survey methods II: with periodic updates., volume Fisheries Special Report
⁵⁷² 25. Michigan Department of Natural Resources, Ann Arbor, 2000.

⁵⁷⁴ ⁴⁴ Odd Terje Sandlund, Karl Øystein Gjelland, Thomas Bøhn, Rune Knudsen, and Per-Arne Amundsen. Contrasting population and life history responses of a young morph-pair of european whitefish to the invasion of a specialised coregonid competitor, vendace. PloS one,
⁵⁷⁶ 8(7):e68156, 2013.

⁵⁷⁸ ⁴⁵ K. Kahilainen, T. Malinen, A. Tuomaala, and H. Lehtonen. Diel and seasonal habitat and food segregation of three sympatric coregonus lavaretus forms in a subarctic lake. Journal of Fish Biology, 64(2):418–434, 2004.

⁵⁸⁰ ⁴⁶ Tommi Malinen, Antti Tuomaala, Hannu Lehtonen, and Kimmo K Kahilainen. Hydroacoustic assessment of mono-and polymorphic coregonus density and biomass in subarctic lakes. Ecology of freshwater fish, 23(3):424–437, 2014.
⁵⁸²

⁵⁸⁴ ⁴⁷ S. Gavrillets and A. Vose. Dynamic patterns of adaptive radiation. Proceedings of the National Academy of Sciences of the United States of America, 102(50):18040–18045, 2005.

⁵⁸⁶ ⁴⁸ X. Thibert-Plante and S. Gavrillets. Evolution of mate choice and the so called magic traits in ecological speciation. Ecology Letters, 16(8):1004–1013, 2013.

⁵⁸⁸ ⁴⁹ A. G. Hudson, P. Vonlanthen, and O. Seehausen. Rapid parallel adaptive radiations from a single hybridogenic ancestral populations. Proceedings of the Royal Society London B, 278:58–66, 2011.

⁵⁹⁰ ⁵⁰ S. Bhat, P-A. Amundsen, R. Knudsen, K.. Gjelland, S-E. Fevolden, L. Bernatchez, and K. Præbel. Speciation reversal in European whitefish (Coregonus lavaretus (L.)) caused by competitor
⁵⁹² invasion. PLoS One, 9:e91208, 2014.

⁵¹ A. Siwertsson, Knudsen R., K. Præbel, C. E. Adams, J. Newton, and P-A. Amundsen. Discrete
foraging niches promote ecological, phenotypic, and genetic divergence in sympatric whitefish
(*Coregonus lavaretus*). *Evolutionary Ecology*, 27:547–564, 2013.

⁵² S. M. Thomas, C. Harrod, B. Hayden, T. Malinen, and K. K. Kahilainen. Ecological speciation
in a generalist consumer expands the trophic niche of a dominant predator. *Scientific Reports*,
7:8765, 2017.

⁵³ M. R. Servedio, G. S. Van Doorn, M. Kopp, A. M. Frame, and P. Nosil. Magic traits in speciation:
'magic' but not rare. *Trends in Ecology and Evolution*, 26(8):389–397, 2011.

⁵⁴ Megan L Head, Genevieve M Kozak, and Janette W Boughman. Female mate preferences for
male body size and shape promote sexual isolation in threespine sticklebacks. *Ecology and
evolution*, 3(7):2183–2196, 2013.

⁵⁵ Theresa Rueger, Naomi M Gardiner, and Geoffrey P Jones. Size matters: male and female
mate choice leads to size-assortative pairing in a coral reef cardinalfish. *Behavioral Ecology*,
27:1585–1591, 2016.

⁵⁶ G. Svårdson. Speciation of Scandinavian *Coregonus*. *Reports of the Institute of Freshwater
Research Drottningholm*, 57:1–95, 1979.

⁵⁷ R. S. Taylor and V. L. Friesen. The role of allochrony in speciation. *Molecular Ecology*,
26:3330–3342, 2017.

⁵⁸ Josef Wanzenböck, Barbara Pamminer-Lahnsteiner, Kathrin Winkler, and Steven J. Weiss.
Experimental evaluation of the spawning periods of whitefish (*Coregonus lavaretus* complex) in
Lake Mondsee, Austria. *Advances in Limnology*, 63:89–97, 2012.

⁵⁹ Kimmo Kalevi Kahilainen, William Paul Patterson, Eloni Sonninen, Chris Harrod, and Mikko
Kiljunen. Adaptive radiation along a thermal gradient: Preliminary results of habitat use and
respiration rate divergence among whitefish morphs. *PLOS ONE*, 9(11):e112085, 2014.

⁶⁰ Skúli Skúlason, Sigurur S. Snorrason, and Bjarni Jónsson. Sympatric morphs, populations
and speciation in freshwater fish with emphasis on Arctic charr. In AE Magurran and RM May,
editors, *Evolution of biological diversity*, pages 70–92. Oxford University Press, Oxford, 1999.

⁶¹ A. P. Hendry, J. K. Wenburg, E. C. Bentzen, P. Volk, and T. P. Quinn. Rapid evolution of reproductive isolation in the wild: Evidence from introduced salmon. Science, 290(5491):516–518, 2000.

⁶² K. Østbye, T.F. Næsje, L. Bernatchez, O. T. Sandlund, and K. Hindar. Morphological divergence and origin of sympatric populations of european whitefish (Coregonus lavaretus L.) in Lake Femund, Norway. Journal of Evolutionary Biology, 18:683–70, 2005.

⁶³ P. Edelaar and D. I. Bolnick. Non-random gene flow: an underappreciated force in evolution and ecology. Trends in Ecology and Evolution, 27(12):659–665, 2012.

⁶⁴ D. Berner and X Thibert-Plante. How mechanisms of habitat preference evolve and promote divergence with gene flow. Journal of Evolutionary Biology, 28(9):1641–1655, 2015.

⁶⁵ P. A. Cott, M. . Guzzo, A. J. Chapelsky, S. W. Milne, and P. J. Blanchfield. Diel bank migration of burbot (Lota lota). Hydrobiologia, 757:3–20, 2015.

⁶⁶ J. Brodersen, D. M. Post, and O. Seehausen. Upward adaptive radiation cascades: predator diversification induced by prey diversification. Trends in Ecology and Evolution, 33:59–71, 2018.

⁶⁷ Kimmo K. Kahilainen, Anna Siwertsson, Karl Ø Gjelland, Rune Knudsen, Thomas Bøhn, and Per-Arne Amundsen. The role of gill raker number variability in adaptive radiation of coregonid fish. Evolutionary Ecology, 25(3):573–588, 2011.

⁶⁸ P. Vonlanthen, D. Roy, A.G. Hudson, C.R. Largiader, D. Bittner, and O. Seehausen. Divergence along a steep ecological gradient in lake whitefish (Coregonus sp.). Journal of Evolutionary Biology, 22:498–514, 2009.

⁶⁹ R.L Eshenroder, P. Vecsei, O.T. Gorman, D.L. Yule, T.C. Pratt, N.E. Mandrak, D.B. Bunnell, and A.M. Muir. Ciscoes (Coregonus, subgenus Leucichthys) of the Laurentian Great Lakes and Lake Nipigon. 2016.

⁷⁰ S. Gavrillets and J. B. Losos. Adaptive radiation: Contrasting theory with data. Science, 323(5915):732, 2009.

⁷¹ A. M. Muir, M. J. Hansen, C. R. Bronte, and C. C. Krueger. If arctic charr salvelinus alpinus is
 646 the most diverse vertebrate, what is the lake charr salvelinus namaycush? Fish and Fisheries,
 17:1194–1207, 2016.

⁷² L. D. Radnaeva, D. V. Popov, O. Grahl-Nielsen, I. V. Khanaev, S. V. Bararsadueva, and R. Kakela.
 648 Fatty acids composition in the whitefish muscle of Cottoidei fishes of Lake Baikal reflects their
 650 habitat depth. Environmental Biology of Fishes, 100:1623–1641, 2017.

⁷³ E. V. Esin and G. N. Markvich. Evolution of the charrs, genus Salvelinus (Salmonidae). 1. origins
 652 and expansion of the species. Journal of Ichthyology, 58:187–203, 2018.

⁷⁴ J. Brodersen and O. Seehausen. Why evolutionary biologists should get seriously involved in
 654 ecological monitoring and applied biodiversity assessment programs. Evolutionary Applications,
 7:968–983, 2014.

⁷⁵ C. J. Doenz, D. Bittner, P. Vonlanthen, C. E. Wagner, and O. Seehausen. Rapid buildup of
 656 sympatric species diversity in alpine whitefish. Ecology and Evolution, 8:9398–9412, 2018.

Figure 2: The frequencies of different outcomes in the Colonization-L scenario for different predation rates. 30 simulations for each parameter combination. The numbers are also reflected in the size of the font used. (a) Only littoral morphs. (b) Both littoral and pelagic morphs. (c) Both littoral and profundal morphs. (d) Only pelagic morph. (e) All three morphs.

Figure 3: The frequencies of different outcomes in the Colonization-G scenario for different predation rates. 30 simulations for each parameter combination. The numbers are also reflected in the size of the font used. (a) Only littoral morphs. (b) Both littoral and pelagic morphs. (c) Both littoral and profundal morphs. (d) Only pelagic morph. No cases with all three morphs were observed.

Appendices

Demographic equilibrium

To estimate the carrying capacity parameters $K_{s,j}$ we proceeded in several steps. First, we estimated the birth rate for whitefish of different size using data in.^{43,44} Second, using personal observations on the proportions of mature fish in each niche for trimorphic lakes we came up with realistic values of their equilibrium densities (N) at 1000, 6000 and 150 individuals, for littoral, pelagic and profundal morphs, respectively. These values serve as default values in our simulations.

Figure S1: Demographic analysis of each population. N_s , v_s and b_s are the population density, survival rate, and fertility at size s , respectively. Subscripts u and m specify immature and mature individuals.

Then, using the Leslie matrices approach (see Figure S1) and assuming perfect adaptation and no predation, we find that the equilibrium population densities N_s of fish of different size in

different niches must satisfied the following equalities:

$$\begin{array}{lll}
 \text{Littoral} & \text{Profundal} & \text{Pelagic} \\
 N_0 = \frac{N_{3m}b_3}{3} & N_0 = \frac{N_{2m}b_2}{3} & N_0 = \frac{N_{1m}b_1}{3} \\
 N_1 = N_0v_0 & N_1 = N_0v_0 & N_{1u} = N_0v_0 \\
 N_2 = N_1v_1 & N_{2u} = N_1v_1 & N_{1m} = v_1(N_{1u} + N_{1m}) \\
 N_{3u} = N_2v_2 & N_{2m} = v_2(N_{2u} + N_{2m}) & \\
 N_{3m} = v_3(N_{3u} + N_{3m}) & &
 \end{array} \tag{S1}$$

where b_s is the average number of offspring per female of size s , and subscripts u and m denote immature and mature individuals, respectively. Note that the division by three in all equations for N_0 above is due to the 2:1 male:female sex ratio in whitefish (Author personal observations).

Littoral morph. Exact data needed to estimate all parameters of our model are difficult to come up with. As a result in some cases we have to take an educated guess. Using our unpublished data (Author unpublished), we know that one of the 2,048 sampled littoral fish survived up to 26 years, thus living 23 years at stage three. This gives the yearly survival rate of $v_3 \approx 0.71$. We then can find a relation between the mature and immature population sizes at stage three. With $N_3 = 1000$, we have $N_{3m} = 709$ and $N_{3u} = 291$. From Eq. 2 in the main text, we calculate $K_{s,j}$:

$$K_{s,j} = \frac{N_{s,j}v_{s,j}(x)}{\omega_{s,j}(x)(1 - v_{s,j}(x))} \tag{S2}$$

Assuming adapted individuals ($\omega_{s,j}(x) = 1$), we can now calculate the carrying capacity parameter $K_3 = \frac{v_3N_3}{1-v_3} = 2448$. Assuming that $b_3 = 64$, from Eq. S1 we get the number of stage zero individuals in the littoral environment $N_0 = 15125$.

We now have to set the other three survival rate v_0, v_1 and v_2 . We know their product because it is equal to the ratio of stage 3 and 0 individuals: $v_0v_1v_2 = \frac{N_{3u}}{N_0} = 0.019$. One possible simple combination of surviving rates that appears to be realistic is $(v_0, v_1, v_2) = (0.2, 0.25, 0.4)$. With these values, we have equilibrium densities at $(N_0, N_1, N_2, N_{3u}, N_{3m}) = (15125, 3025, 756, 291, 709)$. Using equation S2, the carrying capacity parameters for sizes 0, 1 and 2 for the littoral morph become $K_0 = 3781, K_1 = 1008, K_2 = 504, K_3 = 2448$.

Stage	Littoral	Pelagic	Profundal
0 (2cm)	0.2/3781	0.338/2817	0.1/73
1 (15cm)	0.25/1008	0.69/13355	0.4/44
2 (20cm)	0.4/504	0/0	0.82/683
3 (30cm)	0.71/2448	0/0	0/0

Table S1: Survival $v_{s,j}$ /carrying capacity $K_{s,j}$ at equilibrium for the different stage in different habitats.

Profundal morph. In order to find parameters for the carrying capacity of the profundal population, we need to make further assumptions: First, we set its fertility at $b_2 = 16$ which fits the known relationship between fish size and the number of eggs produced [SchneiderEtAl2000, sandlundEtAl2013]. In our unpublished data, of the 243 profundal whitefish sampled, the oldest one was 30 years old. Using the same logic as above, we find its survival rate $v_2 = 0.82$ at stage two. We set the equilibrium density of the largest profundal fish to $N_2 = 150$. Using those assumptions we get a profundal niche with the following equilibrium densities $(N_0, N_1, N_{2u}, N_{2m}) = (656, 66, 123, 27)$ and survival $(v_0, v_1, v_2) = (0.10, 0.40, 0.82)$. Using equation S2, the carrying capacity parameters for sizes 0, 1, and 2 for the profundal morph become $K_0 = 73, K_1 = 44, K_3 = 683$.

Pelagic: To extract the survival and carrying capacities of the pelagic population, we again use the relationship between fish fertility and size [SchneiderEtAl2000, sandlundEtAl2013] and get $b_1 = 4$. Of the 913 profundal whitefish sampled, five reached the age of 16 years old. Using the same logic as above, we have a survival rate of $v_1 = 0.69$ at stage one. In this case, we use $N_1 = 6000$, and $v_1 = 0.69$ and extract the other parameters using the Leslie equations above. We now have the equilibrium densities $(N_0, N_{1u}, N_{1m}) = (5517, 1862, 4138)$ and survival $(v_0, v_1) = (0.338, 0.69)$ for the pelagic niche. Using equation S2 results in carrying capacity parameters $K_0 = 2817, K_1 = 13355$ for the pelagic environment.

Table S1 summarizes the estimates of equilibrium survival rates v and K parameters.