Endosomal explosion induced by hypertonicity enhances chitosan transfection

Shupeng Wang^{1,2}, Shaohua Jin¹, Guangzhi Li², Rui Sun², Qinghai Shu^{1,*}, Song Wu^{2,*}

¹ School of Material Science and Engineering, Beijing Institute of Technology, Beijing 100081, China

² Department of Urology, The Third Affiliated Hospital of Shenzhen University, Shenzhen Luohu Hospital Group, Shenzhen 518001, China

* Corresponding author

E-mail: qhshu121@bit.edu.cn (Q. Shu)

wusong@szu.edu.cn (S. Wu)

Abstract

Transferring DNA into cells to regulate cell function is a novel research field in recent decades. Chitosan is a gene vector with the properties of low-cost and safe, but high efficient delivery has remained challenging. We developed a strategy termed EEIH, for endosomal explosion induced by hypertonicity, in which short-time exposure to hypertonic solutions triggers endosomes destabilization like explosions. EEIH can force chitosan/DNA polyplexes to break through the endosomal barriers to approach the nucleus, which results in boosting the transfection efficiency of chitosan in several cell lines. We demonstrate that EEIH is a significant and practical strategy in chitosan transfection system without sophisticated modification of chitosan.

Keywords: Chitosan; Endosome; Hypertonicity; Gene delivery; Transfection

1. Introduction

The introduction of specific genes into the cell nucleus has raised considerable interest in quite a long time, both in research and the clinic. Non-viral gene vector transfection is relatively safe compared with viral delivery. Chitosan is safe and degradable as a non-viral gene vector, but the transfection efficiency is limited[1, 2]. Modification of chitosan with arginine[3], histidine[4], lysine[4], Polyethylenimine [5], polyethylene glycol (PEG)[1], lipid[6], inorganic nanoparticles[7], and branched structure[8, 9] can improve chitosan transfection to a certain degree, but the resulting

materials are still not efficient enough or mass produced, which limits the wide range of applications..

PEGylated gene carriers generally achieve higher gene transfection efficiency[1, 10, 11]. PEG could promote transfection by another method. In 1985, Gopal demonstrated that at the end of incubating cells with DEAE-dextran and DNA, the treatment with a 40% PEG3400 solution for 2-4 min could improve transfection efficiency effectively. Based on this finding, we examined whether brief exposure to PEG200, PEG1000 and PEG6000 could improve transfection efficiency at the end of treatment of cells with chitosan/DNA polyplexes. We found the treatment of PEG of lower molecular weight enhanced transfection more effectively, which suggested high osmotic pressure may play a nonnegligible role.

In this study, we describe a strategy to boost chitosan transfection. We demonstrate that the briefly hypertonic treatment of cells incubated with chitosan/DNA polyplexes offers superior gene transfection efficiency. We term this strategy “EEIH” for endosomal explosion induced by hypertonicity. The mechanism is that the hypertonicity treatment can change the intracellular osmotic pressure, and then endosomes destabilize like explosions, and foreign DNA escapes from endosomes, which lead to foreign DNA enter into the nucleus to boost transfection (Fig.1). Our results highlight two points, the first is that EEIH is easy to operate, chitosan can achieve satisfied transfection efficiency combination with EEIH for used widely, the second is that EEIH force exogenous substance uptaken to escape from endosomes to disperse in the cytosol, which has great promise in gene transfection and drug delivery system.

Fig.1. Schematic illustration showing the endosomal explosion induced by hypertonicity

2. Materials and Methods

2.1 Materials and cell lines

Chitosan (5-20 mPa·s (0.5% in 0.5% Acetic Acid at 20°C), 75-85% deacetylated, product number: C2395) was purchased from TCI (Japan). Glycerol, dimethylsulfoxide (DMSO), PEG200, PEG1000 and PEG6000 were purchased by Sigma (USA). YOYO-1, LysoTracker Red and Hoechst 33342 were purchased by Invitrogen (USA). Sodium chloride (NaCl) was purchased from Sangon Biotech (China). The chemicals were used as received without further purification. Plasmid DNA pCMV-C-EGFP(5010 bp, Cat. No. D2626) was purchased by Beyotime Biotechnology (China).

HEK293T (a human embryonic kidney cell line, ATCC) and HeLa (a human cervical cancer cell line, ATCC) were cultured in Dulbecco's Modified Eagle Medium (DMEM) (Gibco, USA) supplemented with 10% fetal bovine serum and 1% penicillin-streptomycin. 5637 (a bladder cancer cell line, ATCC) was cultured in RPMI-1640 medium (Gibco, USA) supplemented with 10% fetal bovine serum and 1% penicillin-streptomycin.

2.2 Preparation of chitosan/DNA polyplexes

A chitosan solution (200 µg/ml in 5 mM sodium acetate buffer, pH 5.5) was prepared as previously described[1]. An equal volume of the DNA solution (100 µg/ml in molecular biology grade water) and the sterile-filtered (0.22 µm) chitosan solution were mixed and vortexed for 30 s. The chitosan/DNA polyplexes were incubated at room temperature for 20 min before further use.

2.3 In vitro transfection

The transfection media for chitosan (DMEM, 5 mM MES, pH 6.5) was prepared for further use, according to previous studies[12, 13]. The cells were seeded at 7.5×10^4 cells/well in 24-well plates for 24 h before gene transfection, followed by the medium in each well was replaced with 250 µl of the transfection media, Chitosan/DNA polyplexes were added into each well and incubated with

the cells at the dose of 1 μ g DNA/well (4 μ g DNA per ml of transfection media). After 6 h at 37 °C and 5% CO₂, the transfection media was replaced with 250ul 1x PBS containing different concentrations (weight/volume percentage) of compounds, followed by exposed treatment at room temperature for 3 min. Then the cells were washed twice with PBS and then replaced with complete media for another 42 h before transfection efficiency analysis. The expression of EGFP in cells was observed by fluorescent microscopy (Olympus, Japan) and analyzed by flow cytometry (BD, USA).

2.4 Confocal microscopy

To analyze the intracellular trafficking behaviors of chitosan/DNA polyplexes, Hela cells were seeded in 15 mm glass-bottom cell culture dishes (NEST Biotechnology, China) for 24 h, plasmids were labeled with YOYO-1 as described previously[14], the chitosan/DNA polyplexes were formed as described above save for the use of YOYO-1 labeled DNA. Cells were incubated with the polyplexes as mentioned above (at the dose of 4 μ g DNA per ml of transfection media). After 6 h incubation, the endosomes and the cell nuclei were stained with LysoTracker Red and Hoechst33342 according to the protocol, respectively. The dishes were directly observed by confocal laser scanning microscopy (CLSM, Zeiss, Germany) with a 40 $\times$ objective without fixed. After the treatment of the 10% NaCl solution following the same method as above, the changes of the intracellular trafficking behaviors were also directly observed by CLSM.

3. Results and discussion

3.1. The effect of PEG treatment on chitosan transfection

In an early study, Gopal found briefly exposed to a 40% solution of PEG3400 had increased the transfection efficiency of DEAE-dextran, but the mechanism has not illuminated clearly[15]. To validate the effect of the high concentration of PEG on chitosan transfection, we used different molecular weight and concentration of PEG to treat the Hela cells incubated with chitosan/DNA polyplexes. 3-min PEG treatment had a distinct influence on chitosan transfection (Fig.2). It is worth noting that lower molecular weight PEG achieved higher transfection efficiency at the same

concentration, so we speculated that osmolality might influence transfection efficiency.

Fig.2. The effect of PEG treatment on chitosan transfection in HeLa cells. (A) Fluorescence microscopy images of HeLa cells transfected (scale bar, 500 μ m). (B) Positive EGFP cells (%) and mean fluorescence intensity were recorded by flow cytometry. Data are indicated as mean $\pm$ SD (n = 3).

3.2. The effect of glycerol or DMSO treatment on chitosan transfection

To further examine the role of osmolality, we explored the impact of the hypertonic solution induced by glycerol or DMSO on chitosan transfection. As shown in Fig.3, the results of high transfection efficacy were confirmed in HeLa cells, HEK293T cells and 5637 cells.

Based on demonstrated above, we noticed that glycerol, DMSO and low molecular weight PEG were permeable cryoprotectants[16]. We wondered whether the high transfection efficiency was mainly due to the process of permeable compounds diffused into cells.

Fig.3. The effect of different concentration of glycerol or DMSO solutions on the chitosan in HeLa, HEK293T and 5637 cells. Positive EGFP cells (%) and mean fluorescence intensity were recorded by flow cytometry. Data are indicated as mean $\pm$ SD (n = 3).

3.3 The effect of NaCl hypertonicity treatment on chitosan transfection

Next, we tested whether hypertonicity induced by the compound without the ability to penetrate cell membranes could improve chitosan/DNA polyplexes transfection efficiency. We examined the effect of NaCl hypertonicity treatment on chitosan transfection. As shown in Fig.4, 3-min exposed to 10% NaCl solution improves transfection efficacy significantly. Here, we speculated that hypertonicity played the main role in high chitosan transfection rather than the effect of permeable compounds.

Fig.4. The effect of NaCl hypertonicity treatment on chitosan transfection in Hela, HEK293T, and 5637 cells. (A) Fluorescence microscopy images of cells transfected (scale bar, 500 μ m). (B, C) Positive EGFP cells (%) and mean fluorescence intensity were recorded by flow cytometry. Data are indicated as mean $\pm$ SD (n = 3).

3.4 The effect of hypertonicity on intracellular trafficking behaviors of chitosan/DNA polyplexes

To verify the role of hypertonicity on chitosan/DNA polyplexes endocytosed, we compared

intracellular trafficking behaviors of chitosan/DNA polyplexes before and after hypertonicity treatment. The usual method of monitoring intracellular behaviors of gene carriers is to observe cell fixed coverslips by CLSM. For observing the responses of the same living cells to hypertonicity, we used glass-bottom cell culture dishes instead of fixed coverslips. As shown in Fig.5, after 6 h incubation, co-localizations of DNA (labeled with a green fluorescent dye, YOYO-1) within the endosomes (stained by LysoTracker Red) were observed. Next, we treated the cells with 10% NaCl. Surprisingly, the treatment seemed to trigger endosomes to explode. DNA labeled with YOYO-1 dispersed in the cytoplasm in a short time, and showed high co-localization with the nucleus, which indicated that DNA could be transferred into the nucleus to enable gene transcription.

We try to explain the "endosomal explosion-like" phenomenon from the biophysical perspective. The cells in the hypertonic environment tend to dehydrate and shrink, which leads to an increase in intracellular osmotic pressure. What happens to endosomes in the face of intracellular hyperosmotic environment, one possible answer is that endosomes may be sensitive to the change of osmotic pressure, which may shrink and explode like a squeezed balloon in hypertonicity. In a word, EEIH (endosomal explosion induced by hypertonicity) forced DNA to break through the barrier of endosomes to the nucleus.

Fig.5. CLSM images of HeLa cells incubated with chitosan/DNA polyplexes before and after hypertonicity treatment. DNA was labeled with YOYO-1, the endosomes and the nuclei were stained with LysoTracker and Hoechst33342, respectively (scale bar, 10 μ m).

4. Conclusion

Here we present a low-cost, highly efficient and flexibly adaptable method of chitosan transfection. Endosomal escape is a key step in the transfection process. In the hypertonic condition, the EEIH strategy can destabilize endosomes, facilitate chitosan/DNA polyplexes endocytosed to escape from the endosomes into the nucleus. This strategy can be used as a versatile tool in the development of gene and drug delivery, especially for the therapy of hollow organs such as the cervix uteri and the bladder.

Acknowledgements

This work was supported by the National Research Programs from the Ministry of Science and Technology (MOST) of China (No. 2017YFA0105900) and Shenzhen Basic Research Project (JCYJ20160429172247015 and JCYJ20160429093033251).

Competing interests: The authors declare no competing interests

References

- [1] H.-Q. Mao, K. Roy, V.L. Troung-Le, K.A. Janes, K.Y. Lin, Y. Wang, J.T. August, K.W. Leong, Chitosan-DNA nanoparticles as gene carriers: synthesis, characterization and transfection efficiency, *Journal of controlled release* 70(3) (2001) 399-421.
- [2] Y. Cao, Y.F. Tan, Y.S. Wong, M.W.J. Liew, S. Venkatraman, Recent Advances in Chitosan-Based Carriers for Gene Delivery, *Mar Drugs* 17(6) (2019) 381.
- [3] Y. Gao, Z. Xu, S. Chen, W. Gu, L. Chen, Y. Li, Arginine-chitosan/DNA self-assemble nanoparticles for gene delivery: In vitro characteristics and transfection efficiency, *Int J Pharm* 359(1-2) (2008) 241-6.
- [4] J. Chang, X. Xu, H. Li, Y. Jian, G. Wang, B. He, Z. Gu, Components simulation of viral envelope via amino acid modified chitosans for efficient nucleic acid delivery: in vitro and in vivo study, *Advanced Functional Materials* 23(21) (2013) 2691-2699.
- [5] Q.Q. Zhao, J.L. Chen, M. Han, W.Q. Liang, Y. Tabata, J.Q. Gao, Combination of poly(ethylenimine) and chitosan induces high gene transfection efficiency and low cytotoxicity, *J Biosci Bioeng* 105(1) (2008) 65-8.
- [6] E. Baghdan, S.R. Pinnapireddy, B. Strehlow, K.H. Engelhardt, J. Schafer, U. Bakowsky, *Lipid*

coated chitosan-DNA nanoparticles for enhanced gene delivery, *Int J Pharm* 535(1-2) (2018) 473-479.

[7] F.M. Kievit, O. Veiseh, N. Bhattarai, C. Fang, J.W. Gunn, D. Lee, R.G. Ellenbogen, J.M. Olson, M. Zhang, PEI-PEG-chitosan-copolymer-coated iron oxide nanoparticles for safe gene delivery: synthesis, complexation, and transfection, *Advanced functional materials* 19(14) (2009) 2244-2251.

[8] J. Malmo, K.M. Varum, S.P. Strand, Effect of chitosan chain architecture on gene delivery: comparison of self-branched and linear chitosans, *Biomacromolecules* 12(3) (2011) 721-9.

[9] S.P. Strand, M.M. Issa, B.E. Christensen, K.M. Varum, P. Artursson, Tailoring of chitosans for gene delivery: novel self-branched glycosylated chitosan oligomers with improved functional properties, *Biomacromolecules* 9(11) (2008) 3268-76.

[10] H.X. Wang, Z. Song, Y.H. Lao, X. Xu, J. Gong, D. Cheng, S. Chakraborty, J.S. Park, M. Li, D. Huang, L. Yin, J. Cheng, K.W. Leong, Nonviral gene editing via CRISPR/Cas9 delivery by membrane-disruptive and endosomolytic helical polypeptide, *Proc Natl Acad Sci U S A* 115(19) (2018) 4903-4908.

[11] L. Feng, X. Yang, X. Shi, X. Tan, R. Peng, J. Wang, Z. Liu, Polyethylene glycol and polyethylenimine dual-functionalized nano-graphene oxide for photothermally enhanced gene delivery, *Small* 9(11) (2013) 1989-1997.

[12] M. Thibault, M. Astolfi, N. Tran-Khanh, M. Lavertu, V. Darras, A. Merzouki, M.D. Buschmann, Excess polycation mediates efficient chitosan-based gene transfer by promoting lysosomal release of the polyplexes, *Biomaterials* 32(20) (2011) 4639-46.

[13] M. Lavertu, S. Methot, N. Tran-Khanh, M.D. Buschmann, High efficiency gene transfer using chitosan/DNA nanoparticles with specific combinations of molecular weight and degree of deacetylation, *Biomaterials* 27(27) (2006) 4815-4824.

[14] N.P. Gabrielson, J. Cheng, Multiplexed supramolecular self-assembly for non-viral gene delivery, *Biomaterials* 31(34) (2010) 9117-27.

[15] T.V. Gopal, Gene transfer method for transient gene expression, stable transformation, and cotransformation of suspension cell cultures, *Mol Cell Biol* 5(5) (1985) 1188-90.

[16] M. Lee, M. Hong, Cryoprotection of lipid membranes for high-resolution solid-state NMR studies of membrane peptides and proteins at low temperature, *Journal of biomolecular NMR* 59(4) (2014) 263-277.