Metagenomic analysis of soil samples collected from estuarine mangroves of Arabian Sea reveals rich microbiota and high numbers of sulphate reducing bacteria accompanied with methanogen bacteria.

Mandar S. Paingankar[#] and Deepti D. Deobagkar *

Molecular Biology Research Laboratory, Centre for Advanced Studies, Department of Zoology, Savitribai Phule Pune University, Pune - 411 007, India.

[#] Current address: Department of Zoology, Government Science College, Chamorshi Road, Gadchiroli - 442605, Maharashtra, India.

*Corresponding Author:

Deepti D. Deobagkar,
ISRO Chair Professor,
Department of Zoology, Centre for Advance Studies,
Savitribai Phule Pune University, Pune 411007, India.
Contact : +919921184871.

deepti.deobagkar@gmail.com

Abstract

This study reports the analyses of the microbiome of the estuarine soil of mangroves of the Arabian Sea. Mangroves soil samples were collected from 12 locations of Arabian Sea coast of Maharashtra, India. 16S rRNA gene V3–V4 region amplicon sequencing was performed using the Miseq Illumina platform to identify the microbial communities present in the mangroves ecosystem. The metagenomics analysis provided an insight into the abundance, diversity and spatial variations in the mangrove microbial communities in relation to physico-chemical parameters and revealed that Proteobacteria, Flavobacteria and Planctomycetes are abundant in mangroves system. The differences in bacterial abundance, composition and diversity can partly be attributed to the physico-chemical characteristics of the samples, geographical location and anthropogenic activities in the locality. High numbers of sulphate reducing bacteria accompanied with methanogen bacteria were characteristic of Indian mangroves. The results obtained in the current study indicate rich species diversity and add valuable insights about the diversity of microbial communities of the mangroves in Maharashtra along the west coast of India and can provide better information for effective measures for conservation of mangroves. GIS based prediction suggest that the sulphur utilizing communities are under threat from anthropogenic activities and may decline in future if immediate measures are not implemented.

Keywords: Mangroves of west coast, Metagenomics, Microbial ecology, pollution

Introduction

Mangroves system is known to play an important role maintaining the sea level and protecting the coast. These mangrove systems are characterized by structurally and functionally unique salt-tolerant, arboreal, flowering plant forests¹. Taxonomically diverse group of plants and associated microbial communities are known to play fundamental roles in the nutrient cycle, productivity, functioning, and maintenance of the mangrove ecosystem². The mangrove microbial community plays an integral role in nutrients cycling and thus nurturing the sustainable productivity³. Furthermore, fluctuations in temperature, aerobic conditions and salinity make these microbial communities more complex, diverse and unique⁴. Understanding the complex nature of mangrove microbial communities is one of the important aspects in microbes-plant interaction dynamics³. Similarly, microbes-plant interactions in mangroves have great implications in local and global ecological impact⁵⁻⁸. Furthermore these microbes are known to provide interesting biomolecules for human welfare⁹. Therefore exploration and identification of microbial communities in mangroves system have a potential application in human welfare as well as ecological conservation.

The microbial community studies have been carried out by various research groups to understand the role of microbial communities in the complex functioning of mangrove ecosystems¹⁰⁻²¹. In recent years, this ecosystem is rapidly getting destroyed and is threatened due to anthropological activities and modernisation²²⁻²³. In India, mangroves are present on eastern coastline, western coastline and in the Andaman and Nicobar Islands and are spread over an area of 4,921 sq. km. Presence of unique microbial communities in Sundarbans of India and Andaman Nicobar islands have been documented by various workers^{12,13,16,18}. While on western coastline of India, mangroves are concentrated in Gujarat and Maharashtra. The mangroves of

India are under immense pressure of conversion in to agriculture land and discharge of organic and inorganic matter from rapidly developing cities^{9,23}. Mumbai, Thane, Raigad and Ratnagiri area in Maharashtra state are developing rapidly and anthropogenic activities in these are exerting the tremendous pressure on flora, fauna and microbial diversity^{23,24}. Very few studies have been conducted to understand the microbial communities present in mangrove ecosystems of western coastline India^{20,21,24}. Comparative studies on distinct mangrove microbial communities based on metagenomics will provide better insights into microbial community structure in the mangroves of India. Therefore the metagenomics explorative studies on microbial community structure in mangroves of India is essential in acquiring the baseline data on the diversity and functional genomics of microbial communities at higher resolution and in greater depth. The objective of the present study was to identify and to provide baseline data on the microbial community structure present in different mangrove areas along the west coast in Maharashtra State, India using the Illumina MiSeq sequencing platform.

Material and Methods

Ethics Statement

The collection sites of the current study are not part of any reserve forest or National park or privately-owned areas and therefore no specific permits were needed for the field studies. Endangered or protected species were not collected or included in the study.

Sample Collection

Information on mangroves of India and shape files for GIS mapping were retrieved from earlier reports^{22,23}. Sediment samples were collected superficially (0-5 cm depth) during the period of low tide at the GPS coordinates. Anjarle (17.840N, 73.101E and 17.843N, 73.103E),

Kelashi(17.928N,73.074E and 17.931N, 73.077E), Bankot(17.958N, 73.031E and 17.961N,73.033E), Uran(18.896,72N.939E and 18.898N,72.944E), Agarkot(18.546N,72.934E and 18.549N,72.943E), Vashim(18.812N,73.033E and 18.814N,73.026E). These mangrove forests were located in estuarine regions of the Jog, Bharja, Savitri, Kundalika and Patalganga rivers of Maharashtra, India (Supplementary Information Table S1).

Environmental parameters

The average temperature, average rainfall, pH, SO₄, NO₃, PO₄, Mg, Cl₂ and salinity of samples was measured and is depicted in Supplementary information Table S2.

DNA extraction from soil samples

Soil collected from respective samples was processed for DNA isolation using Powersoil DNA isolation kit (MoBio laboratories Inc. Carlsbad, CA) as per manufacturer's instructions. DNA concentration was measured using the Quantus fluorimeter (Promega, USA).

Amplification primers

Regions corresponding to V3 and V4 regions of 16s rRNA were amplified from the DNA samples and were subjected to pyrosequencing using the Miseq Illumina platform.

Sequence analysis

Sequence analysis was carried out by employing the methods described earlier²⁴. The taxonomic assignment of unassembled metagenomic sequences was performed using BLASTX against the SEED and Pfam databases on the MG-RAST server v2.0 (<http://metagenomics.nmpdr.org>) using a cut-off E-value of 1e-10. BLASTX was also used to conduct a similarity search against the NCBI-NR database, and MetaGenome Analyzer software (MEGAN v6.0)²⁵ (<https://github.com/danielhuson/megan-ce>) with the LCA algorithm (maximum number of matches per read: 5, min support: 5, min score: 35, top percent:10) was used to visualize results.

Statistical analysis

Several indices of clonal diversity were estimated using the MEGAN and PAST3 program available from the University of Oslo website link (<http://folk.uio.no/ohammer/past>). A similarity matrix was generated using a probabilistic distance metric. The statistical significance of the relationship between the differences in chemical composition and (i) species diversity, (ii) species alpha diversity indices and (iii) species beta diversity indices was tested by Mantel tests. P values were calculated using 9999 permutations on rows and columns of dissimilarity matrices.

GIS mapping and prediction using DIVA GIS and MAXent

The mangroves sulphur utilizing bacteria data was retrieved from current study and earlier study²⁴. The baseline (1950–2000) temperature and precipitation layers were obtained from WorldClim-Global Climate data repository (www.worldclim.org). Temperature and precipitation layers of the future (year 2020, year 2030) were obtained from repository (www.ccafs-climate.org). The data were resampled at the native WorldClim 30 arcsec (approximately 1x1km) resolution. Statistical modelling was conducted with DIVA GIS and MAXENT (https://biodiversityinformatics.amnh.org/open_source/maxent/) software with the present data and from the environmental variables.

Results

Description of the Community

Sequence clustering resulted in the identification of 1357 (364.16±84.80 per sample) different bacterial species (Supplementary Information Table S3). In general, Proteobacteria and Bacteroidetes were more abundant and other taxon such as Cyanobacteria, Euryarchaeota, Chloroflexi and Verrucomicrobia were present at moderate abundance. The Acidobacteria were

more abundant in Kelashi sample whereas Fusobacteria showed moderate abundance in the Uran sample. The numbers of sequences affiliated with each taxon are depicted in Figure 2A, with a major abundance of Gammaproteobacteria (8.86-20.26 %), Flavobacteriia (7.24-25.76%), Alphaproteobacteria (9.34-24.65%), Deltaproteobacteria (2.88-14.57%) and Actinobacteria (0.17-6.1%) followed by other minor class represented by Planctomycetia (1.2-3.57%). Sulphate utilizer species such as *Desulfobulbus mediterraneus*, *Desulfomicrobium sp.*, *Desulfonatronum thiosulfatophilum*, *Desulfosarcina sp.* were present in most of the samples. These sulphate reducers were accompanied by methanogens at all sites. *Aciditerrimonas ferrireducens*, *Algisphaera agarilytica*, *Ardenticatena maritima*, *Bacteriovorax marinus*, *Bdellovibriobacteriovorus*, *Bellilinea caldifistulae*, *Blastocatella fastidiosa*, *Blastopirellula marina*, *Calothrixdesertica*, *Candidatus Solibacterusitatus*, *Litorilinea aerophila*, *Ornatilinea apprima*, *Owenweeksia hongkongensis*, *Phycisphaera mikurensis* *Thiohalobacterthiocyanaticus* *Trueperaradiovictrix* were present in moderate numbers in all collected samples. *Deferrisomacramini* exclusively present in ASUN samples whereas *Nafulsella turpanensis* *Roseiflexus sp.* RS-1 exclusively present in KLM samples. Species such as *Lewinella nigricans* *Methanomassiliicoccus luminyensis*, *Methanothermococcus kinawensis*, *Robiginitaleabiformata* showed moderate presence in all sampling sites except KLM samples. *Ignavibacterium album* and *Ilumatobacter fluminis* were present in moderate numbers only in ASUN, ASWD, BNM samples whereas *Gramella aestuarii* present significant number only in ADM BNM samples.

Alpha and Beta diversity of bacterial communities in different mangrove soil samples

Alpha and beta diversity analysis revealed rich taxonomic diversity and dominance of few species in mangroves samples (Supplementary Information Table S4; S5). Interestingly, soil samples from KLM1 (D = 0.06438), KLM2 (D = 0.03303), VASH1 (D = 0.03259) and VASH2

($D = 0.03802$) showed higher dominance of fewer bacterial groups. Chao-1 analysis identified (274-729) species in each sample. At species level, high beta diversity was observed in all mangrove samples (Supplementary Information Table S5) wherein 36 species were common in all 12 samples. ASUN1 sample showed maximum number (61) of unique species.

Habitat Type Differences in Communities

In order to examine the community structure and its specific features, PCoA analysis was carried out. Most samples clustered closely together, indicating that the microbial communities residing in these mangroves are similar and represent characteristic and typical community structures. Interestingly, communities in the mangroves facing industrialization threats, KLM1, KLM2 and ADM1, ADM 2 were different from all other (Fig. 3A). Similarly, neighbour joining tree revealed the distinctness of KLM1, KLM2 and ADM1, ADM 2 samples (Fig. 3B). Canonical correlation analysis (CCorA) was employed to find out the correlations between alpha diversity indices and environmental factors. A strong positive correlation was observed between the SO_4 concentration and relative dominance of a few species. A positive correlation was observed in PO_4 and Cl_2 concentration and Berger-Parker index in samples. A negative correlation was observed in NO_3 concentration and dominance in samples (Fig. 4A). Mantel test was used to determine the factors (diversity, chemical parameters, and Whitaker beta diversity) which best predicted the community diversity across the different mangrove areas samples (Fig. 4B). Differences in chemical parameters correlated significantly with diversity across different collection sites ($r = 0.278$; p value = 0.02). Similarly, chemical parameters of soil had significant effect on diversity and beta diversity ($r = 0.717$; p value = 0.0001). These results indicated that soil chemical composition and presence of pollutants is playing an important role and influences the bacterial diversity.

GIS based prediction of distribution of sulphur utilizing bacteria in Maharashtra

Using the current climate layers (68 layers) the distribution of sulphur utilizing bacteria in Maharashtra was predicted (Fig. 5). The prediction suggests that areas such as Dighi, Chiplun, Uran, Jawaharlal Nehru port are suitable for growth of sulphur utilizing bacteria. Based on future climatic conditions, sulphur bacteria prediction map for the year 2020 and 2030 was predicted. The prediction suggests that area suitable for growth of sulphur utilizing bacteria will be decline in 2020 and 2030 (Fig. 5).

Discussion

The mangroves of India are under tremendous pressure of anthropogenic activities and are rapidly declining²³. Maharashtra state of the west coast of India is one of the rapidly industrializing and urbanizing region and discharging a huge amount of industrial and organic waste to Arabian Sea^{26,27}. Jha Fernandes. The draining of industrial and domestic waste the mangroves of Maharashtra is significantly affecting the mangroves and associated flora, fauna and microbes. Therefore there is urgent need to gather the baseline data of microbes associated with mangroves as they play important role in mangroves ecosystem dynamics. In the current study metagenomic profiles of mangrove sediments of west coastline of India was carried out using the miSeq Illumina platform. All the samples showed the high abundance of bacterial reads and a small percentage of sequences from Archaea and unidentified organisms.

A taxonomic analysis of samples collected from Kerala, Tamil Nadu, Sundarbans, Andaman Nicobar islands, Maharashtra and current study clearly indicates the higher abundance of Proteobacteria in the mangroves sediments^{12,16,17,18,20,21,24} (Fig. 2A). The metagenomic analysis identified more than 80% species inhabited in these mangrove sediments (Fig.2B).

201 Among the Proteobacteria detected, Gammaproteobacteria were most abundant followed by
 202 Alphaproteobacteria. The *nifH* gene, the key players in nitrogen fixation, has been detected
 203 predominantly Alphaproteobacteria, Betaproteobacteria, and Gammaproteobacteria³⁰⁻³²(Fig.
 204 2A).Along with Proteobacteria, *Bacteroidetes*, *Acidobacteria*, *Firmicutes*, *Actinobacteria*,
 205 *Nitrospirae*, *Cyanobacteria*, and *Planctomycetes* were also showed their moderate presence in
 206 the current study and Indian mangroves^{12,16,17,18,20,21,24}. A similar patterns were obtained from the
 207 samples belonging to mangroves areas of different parts of the world^{11,14,15,19,20,21,28,29}. As
 208 compared to Indian and Brazilian mangrove samples, Red sea mangrove samples showed
 209 relatively higher number of Achaea^{12,16- 21,24}.Bacteroidetes known for hydrocarbon degradation
 210 and substantial abundance of Bacteroidetes has been previously noted from mangroves
 211 samples^{10,16,20,21,24,33}.The mixed marine and terrestrial conditions in mangroves are suitable for
 212 nitrogen fixer proteobacteria and Bacteroidetes. Current study and samples collected from Kerala
 213 showed the moderate numbers of Actinobacteria^{20,21}. Although Chloroflexi and Verrucomicrobia
 214 detected routinely in metagenomics analysis, little is known about the possible role of these
 215 bacterial phyla. Samples from Indian mangroves showed the presence of high numbers of
 216 sulphate reducing bacteriaaccompanied with methanogen bacteria. Sulphate reducing bacteria,
 217 such as *Desulfobulbus*, *Desulfomicrobium*, *Desulfonatronum*, *Desulfosarcina*were found
 218 accompanied with methanogen genera, including *Methanothermobacter*, *Methanococcus*,
 219 *Methanocaldococcus*, *Methanosarcina*, *Methanococcoides*, *Methanospirillum*^{12,16,17,18,20,21,24}. The
 220 results obtained in current study and earlier studies clearly suggest that sulphate reducers and
 221 methanogens might be using the different metabolic pathways in non-competitive manner to
 222 shape up the microbial metabolism in mangroves ecosystem³⁴⁻³⁸. Sequence reads assign to

aromatic hydrocarbons degraders and an iron and sulphur-reducing mesophilic anaerobe were found significantly lower in Indian mangroves samples.

The alpha and beta diversity indices clearly indicate the presence of high species richness and high diversity in these samples. PCoA and neighbour joining analysis clearly suggest the distinctness of KLM and ADM samples from rest of the samples (Fig. 3A,3B). Relatively less organic and inorganic pollution might be responsible for distinctness of KLM and ADM samples. The Canonical correlation analysis and Mantel test clearly suggest that chemical properties of soil sediments and micro environment in this ecological niche influenced bacterial diversity in these mangrove areas (Fig. 4A;4B). It has been well documented that geographical location, plant species, and/or physico-chemical parameters play an important role in shaping the microbial structure. The observed variation in microbial diversity and composition in studied samples might be attributed to the environmental factors and anthropogenic activities in respective mangrove forests.

GIS based prediction tools are important in examining the potential problems arising from natural and artificial forces on environment as well as microhabitats. Furthermore, GIS is one of the important tools which can be used in ecology to preserve and monitor the makeup of biological life within a given area. However, its use in predicting microbial ecology is limited^{39,40}. In the current study GIS prediction was used to predict the sulphur utilizing bacteria in Maharashtra region. Similarly future climate data was used to predict the trends of sulphur utilizing bacteria in response to climate change. Using current climate condition layers (total 68 layers) the distribution of sulphur utilizing bacteria in Maharashtra was predicted. The prediction suggests that climatic conditions in Raigad district especially near Dighi, Uran and Jawaharlal Nehru port are suitable for growth of sulphur utilizing bacteria. These bacteria are useful in

recycling of various compounds such as sulphates, alkanes and many more. The prediction using future data suggest the decrease in suitable area for growth of these sulphur utilizing bacteria. These observations suggest that anthropogenic activities will upset the delicate ecology of these microbes and may lead to massive effect on flora, fauna and human welfare. Therefore there is urgent need to conserve this ecologically delicate and important niche

In conclusion, present study identified the presence of novel groups of bacteria (Fig.6). The current study generated the baseline data of bacterial diversity and composition in sediments of mangroves of west coast of India. The data obtained in the current study clearly indicate that biogeochemical parameters, anthropogenic activities and micro niche play an important role in structuring the bacterial diversity. GIS based prediction suggests that the mangrove microbial communities may decline in future if immediate conservation measures are not implemented.

Acknowledgement

We thank Prof. Dileep N. Deobagkar for his valuable suggestions.

Funding

The authors would like to acknowledge financial support provided by the Ministry of Earth Sciences (MoES), India and UGC-UPE-II.

Conflict of interests

The authors declare that they have no competing interests.

References

- 269 1. Alongi, D. M., Present state and future of the world's mangrove forests. *Environ. Conserv.*
270 2002, **29**(3), 331–349.
- 271 2. McLusky, D. *The estuarine ecosystem*. Springer Science and Business Media 2013.
- 272 3. Field, C. *et al.*, Mangrove biodiversity and ecosystem function. *Global Ecol. Biogeogr. Lett.*
273 1998, **7**, 3-14.
- 274 4. Duke, N.C. *et al.*, A world without mangroves. *Science*, 2007, **317**, 41.
- 275 5. Gomes, N.C.M., Borges, L.R., Paranhos, R., Pinto, F.N., Mendonça-Hagler, L.C. and Smalla,
276 K. Exploring the diversity of bacterial communities in sediments of urban mangrove forests.
277 *FEMS Microbiol. Ecol.* 2008, **66**(1), 96-109.
- 278 6. Kulkarni, R., Deobagkar, D., Zinjarde, S. Metals in mangrove ecosystems and associated
279 biota: A global perspective *Ecotoxicol. Environ. Safety* 2018, **153**, 215–228
- 280 7. Kandasamy, K. A. Review of studies on Pichavaram mangrove, Southeast India
281 *Hydrobiologia* 2000, **430**(1), 185-205
- 282 8. Ragavan P. Mohan, P. M. Saxena, A. Jayaraj, R. S. C. Ravichandran, K. Saxena, M.
283 Mangrove floristics of the Andaman and Nicobar Islands: critical review and current scenario
284 *Marine Biodiver.* 2016, 1-21 <https://doi.org/10.1007/s12526-016-0581-3>
- 285 9. Upadhyay, V.P., Ranjan, R. and Singh, J.S. Human-mangrove conflicts, The way out. *Curr.*
286 *Sci.* 2002, **83**, 1328–1336.
- 287 10. Gomes, N.C.M. *et al.*, Taking root, enduring effect of rhizosphere bacterial colonization in
288 mangroves. *PLoS ONE* 2010, **5**(11), e14065.
- 289 11. Taketani, R.G., Yoshiura, C.A., Dias, A.C.F., Andreote, F.D. and Tsai, S.M. Diversity and
290 identification of methanogenic archaea and sulphate-reducing bacteria in sediments from a
291 pristine tropical mangrove. *A van Leeuw J. Microb.* 2010, **97**(4), 401–411.

12. Ghosh, A., Dey, N. and Bera, A. Culture independent molecular analysis of bacterial communities in the mangrove sediment of Sundarbans, India. *Saline Syst.* 2010, **6(1)**, 1
<http://dx.doi.org/10.1186/1746-1448-6-1>.
13. Mishra, R.R., Swain, M.R., Dangar, T.K. and Thatoi, H. Diversity and seasonal fluctuation of predominant microbial communities in Bhitarkanika, a tropical mangrove ecosystem in India. *Rev. Biol. Trop.* 2012, **60(2)**, 909-924.
14. Andreote, F.D. *et al.*, The microbiome of Brazilian mangrove sediments as revealed by metagenomics. *PLoS One.* 2012, **7(6)**, e38600.
15. Jiang, X.T., Peng, X., Deng, G.H., Sheng, H.F., Wang, Y. and Zhou, H.W. Illumina sequencing of 16S rRNA tag revealed spatial variations of bacterial communities in a mangrove wetland. *Microb. Ecol.* 2013, **66**, 96–104.
16. Basak, P. *et al.*, Spatiotemporal analysis of bacterial diversity in sediments of Sundarbans using parallel 16S rRNA gene tag sequencing. *Microb. Ecol.* 2014, **69**, 500–511.
17. Prakash, S. *et al.*, Influence of physicochemical and nutritional factors on bacterial diversity in mangrove sediments along the southwest coast of Tamilnadu, India. *Environ Monit. Assess.* 2015, **187**, 562.
18. Basak, P. *et al.*, Bacterial diversity assessment of pristine mangrove microbial community from Dhulibhashani, Sundarbans using 16S rRNA gene tag sequencing. *Genomics Data.* 2016, **7**, 76-78.
19. Alzubaidy, H. *et al.*, Rhizosphere microbiome metagenomics of gray mangroves (*Avicennia marina*) in the Red Sea. *Gene.* 2016, **576**, 626–636
20. Imchen, M. *et al.*, Searching for signatures across microbial communities: Metagenomic analysis of soil samples from mangrove and other ecosystems *Sci. Rep.*, 2017, **7**, 859

21. ImchenM. et al., Comparative mangrove metagenome reveals global prevalence of heavy metals and antibiotic resistome across different ecosystems *Sci. Rep.*, 2018, **8**,11187.
22. Giri, C. *et al.*, Status and distribution of mangrove forests of the world using earth observation satellite data. *Global Ecol. Biogeogr.* 2011, **20**, 154–159.
23. Sahu, S.C., Suresh, H.S., Murthy, I.K. and Ravindranath, N.H. Mangrove Area Assessment in India, Implications of Loss of Mangroves. *J. Earth Sci. Clim. Change.* 2015, **6**, 280. doi, 10.4172/2157-7617.1000280
24. Paingankar, M.S. and Deobagkar, D.D. Pollution and environmental stressors modulate the microbiome in estuarine mangroves: a metagenome analysis. *Curr. Sci.* (In Press)
25. Huson, D.H., Beier, S., Flade, I., Górska, A., El-Hadidi, M., Mitra, S., et al. MEGAN Community Edition - Interactive Exploration and Analysis of Large-Scale Microbiome Sequencing Data. *PLoS Comput. Biol.*, 2016, **12**(6), e1004957. <https://doi.org/10.1371/journal.pcbi.1004957> (<https://github.com/danielhuson/megan-ce>)
26. Jha, S.K., Chavan, S.B., Pandit, G.G., Negi, B.S. and Sadasivan, S. Behaviour and fluxes of trace and toxic elements in creek sediment near Mumbai, India. *Environ. Monit. Assess.* 2002, **76**, 249–262.
27. Fernandes, L., Nayak, G.N., Ilangoan, D. and Borole, D.V. Accumulation of sediment, organic matter and trace metals with space and time, in a creek along Mumbai coast, India. *Estuar. Coast Shelf Sci.* 2011, **91**, 388–399.
28. Lyimo, T.J., Pol, A. and Op den Camp, H.J.M. methane emission, sulphide concentration and redox potential profiles in Mtoni mangrove sediment, Tanzania. *Western Indian Ocean J. Mar. Sci.* 2002, **1**, 71–80.

29. Lyimo, T.J., Pol, A. and Op den Camp, H.J.M. Sulfate reduction and methanogenesis in
sediments of Mtoni mangrove forest, Tanzania. *Ambio*. 2002, **31**, 614–616.
30. Bagwell, C.E., La Rocque, J.R., Smith, G.W., Polson, S.W., Friez. M.J., Longshore, J.W. and
Lovell CR. Molecular diversity of diazotrophs in oligotrophic tropical seagrass bed
communities. *FEMS Microb. Ecol.* 2002, **39**, 113– 119.
31. Bird, C., Martinez, J.M., O'Donnell, A.G. and Wyman, M. Spatial distribution and
transcriptional activity of an uncultured clade of planktonic diazotrophic γ -proteobacteria in
the Arabian Sea. *Appl. Environ. Microbiol.* 2005, **71**, 2079– 2085.
32. Church, M.J., Short, C.M., Jenkins, B.D., Karl, D.M. and Zehr, J.P. Temporal patterns of
nitrogenase gene (*nifH*) expression in the oligotrophic North Pacific Ocean. *Appl. Environ.*
Microbiol. 2005, **71**, 5362– 5370.
33. Kim, S.J. and Kwon, K.K. Bacteroidetes, Handbook of Hydrocarbon and Lipid
Microbiology, 2010, pp 1813–1817.
34. Muyzer, G. and Stams, A.J. The ecology and biotechnology of sulphate-reducing bacteria.
Nat. Rev. Microbiol. 2008, **6(6)**, 441-454.
35. Lyimo, T.J., Pol, A. and Op den Camp, H.J.M. Sulfate reduction and methanogenesis in
sediments of Mtoni mangrove forest, Tanzania. *Ambio*. 2002, **31**, 614–616.
36. Muyzer, G. and Stams, A.J. The ecology and biotechnology of sulphate-reducing bacteria.
Nat. Rev. Microbiol. 2008, **6(6)**, 441-454.
37. Alongi, D.M., Boto, K.G. and Robertson, A.I. Nitrogen and Phosphorus Cycles in Tropical
Mangrove Ecosystems, Am GeophysUniv, Washington DC. 1992, **41**, 251–292
38. Rodríguez, H. and Fraga R. Phosphate solubilizing bacteria and their role in plant growth
promotion. *Biotechnol. Adv.* 1999, **17 (4–5)**, 319–339.

39. Terrat S, Horrigue W, Dequiedt S, Saby NPA, Lelièvre M, Nowak V, et al. Mapping and predictive variations of soil bacterial richness across France. *PLoS ONE* 2017, **12**(10), e0186766.
40. Griffiths RI, Thomson BC, Plassart P, Gweon HS, Stone D, Creamer RE, et al. Mapping and validating predictions of soil bacterial biodiversity using European and national scale datasets. *Appl Soil Ecol* 2016, **97**, 61–68.

Figure Legends

Figure 1: Sampling sites: Mangrove soil samples were collected from different locations from eusturine mangrove along the coast of Arabian Sea.

Fig. 2 Distribution of predominant bacterial class in samples based on 16S rRNA gene sequencing. A) Class distribution Observations are displayed as stacked bar charts for individual mangrove sample (x-axis) against the percent class abundance (y-axis). B) Rarefaction curves for mangrove samples

Figure 3: A) Principal coordinate analysis (PCoA) of the bacterial communities derived from the weighted UniFrac distance matrix. B) Neighbor-joining phylogenetic tree

Figure 4: A Canonical correlation analysis (CCorA) B) Mantel Test

Figure 5: GIS based prediction of sulphur utilizing bacteria in Maharashtra

Figure 6: Microbial diversity at various mangrove sites

Supplementary Information Table S1: Information of sampling sites

Supplementary Information Table S2: Information of chemical parameters of sampling sites

Supplementary Information Table S3: Information of bacterial species

Supplementary Information Table S4: Alpha Diversity Indices

384 **Supplementary Information Table S5:** Global beta diversity indices

385 **Supplementary Information Table S6:** Information of sampling sites used for GIS based

386 prediction

Table 1: Details of soil samples collected from mangroves

Place	District	Sample code	Latitude	Longitude	River
Anjarle	Ratnagiri	ADM1	17.840	73.101	Jog River
Anjarle	Ratnagiri	ADM2	17.843	73.103	Jog River
Kelashi	Ratnagiri	KLM1	17.928	73.074	Bharja River
Kelashi	Ratnagiri	KLM2	17.931	73.077	Bharja River
Bankot	Ratnagiri	BNMN1	17.958	73.031	Savitri River
Bankot	Ratnagiri	BNMN2	17.961	73.033	Savitri River
Uran	Raigad	ASUN1	18.896	72.939	
Uran	Raigad	ASUN2	18.898	72.944	
Agarkot	Raigad	ASWD1	18.546	72.934	Kundalika River
Agarkot	Raigad	ASWD2	18.549	72.943	Kundalika River
Vashim	Raigad	VASH1	18.812	73.033	Patalganga River
Vashim	Raigad	VASH2	18.814	73.026	Patalganga River

Figure 1: Sampling sites: Mangrove soil samples were collected from different locations from eusturine mangrove along the coast of Arabian Sea.

Fig. 2 Distribution of predominant bacterial class in samples based on 16S rRNA gene sequencing. A) Class distribution Observations are displayed as stacked bar charts for individual mangrove sample (x-axis) against the percent class abundance (y-axis). B) Rarefaction curves for mangrove samples

Figure 3: A) Principal coordinate analysis (PCoA) of the bacterial communities derived from the weighted UniFrac distance matrix. B) Neighbor-joining phylogenetic tree

Figure 4: A Canonical correlation analysis (CCorA) B) Mantel Test

Figure 5: GIS based prediction of sulphur utilizing bacteria in Maharashtra

412 **Figure 6: Microbial diversity at various mangrove sites**

413