Predation strategies of the bacterium *Bdellovibrio bacteriovorus* result in bottlenecks, overexploitation, minimal and optimal prey sizes

J. Kimberley Summers^{1*} and Jan-Ulrich Kreft¹

¹Institute of Microbiology and Infection & Centre for Computational Biology & School of Biosciences, University of Birmingham, Edgbaston, Birmingham, B15 2TT, United Kingdom

***Corresponding author**

J. Kimberley Summers

Tel: +44 (0)121 41-45424

E-mail address: jks282@bham.ac.uk

Running title: Predation strategies of *Bdellovibrio*

Abstract

With increasing antimicrobial resistance, alternatives for treating infections or removing resistant bacteria are urgently needed, such as the bacterial predator *Bdellovibrio bacteriovorus* or bacteriophage. Therefore, we need to better understand microbial predator-prey dynamics. We developed mass-action mathematical models of predation for chemostats, which capture the low substrate concentration and slow growth typical for intended application areas of the predators such as wastewater treatment, aquaculture or the gut. Our model predicted a minimal prey size required for predator survival, explaining why *Bdellovibrio* is much smaller than its prey. A too good predator (attack rate too high, mortality too low) overexploited its prey leading to extinction (tragedy of the commons). Surprisingly, a predator taking longer to produce more offspring outcompeted a predator producing fewer offspring more rapidly (rate versus yield trade-off). Predation was only efficient in a narrow region around optimal parameters. Moreover, extreme oscillations under a wide range of conditions led to severe bottlenecks. A bacteriophage outcompeted *Bdellovibrio* due to its higher burst size and faster life cycle. Together, results suggest that *Bdellovibrio* would struggle to survive on a single prey, explaining why it must be a generalist predator and suggesting it is better suited to environments with multiple prey than phage.

27 Introduction

28 Predator-prey relationships are some of the oldest and most important interactions in nature and
 29 occur at every level, from the smallest virus infecting a bacterium, to lions attacking wildebeest.
 30 Predators are often keystone species in natural ecosystems [1]. Investigations into predator-prey
 31 dynamics in natural settings are, however, complicated by an array of confounding factors [2].
 32 Microbes by contrast make attractive models for studying predator-prey interactions in a controlled
 33 environment, with millions of individuals in a drop of liquid that can go through many generations in
 34 days. A completely separate, but increasingly important, reason for the interest in microbial
 35 predators is the antimicrobial resistance crisis, with resistance to even last resort antibiotics such as
 36 colistin rising [3]. There is an urgent need for ‘living antibiotics’ as alternatives to antibiotics, such as
 37 phage and bacterial predators.

38 The best-studied bacterial predator is *Bdellovibrio bacteriovorus*. It is a Gram-negative bacterium
 39 that predaes a wide range of other Gram-negative bacteria [4], regardless of their antimicrobial
 40 resistance or pathogenicity. *Bdellovibrio* alternates between two phases in its lifecycle. In the free-
 41 living attack phase, it does not grow but hunts prey. It swims faster than most bacteria, at speeds of
 42 up to $160 \mu\text{m s}^{-1}$ or ~ 100 body lengths s^{-1} [5]. This requires a high metabolic rate and loss of viability
 43 within 10 hours if prey is not encountered [6]. Once a cell is encountered, its suitability is
 44 investigated for several minutes [7]. If suitable, it enters the prey’s periplasm by creating and
 45 squeezing through a pore in the outer membrane and peptidoglycan layer, shedding its flagellum in
 46 the process [8]. Once inside the periplasm, *Bdellovibrio* kills the prey and lets the cytoplasmic
 47 nutrients leak into the periplasm [9]. It also alters the prey’s peptidoglycan, causing the prey cell to
 48 round up into a “bdelloplast”. In this bdelloplast phase, *Bdellovibrio* uses these nutrients to grow into
 49 a long filament rather than dividing [5]. When the nutrients have been used up, this filament septates
 50 into as many new cells as resources allow for, typically between 3 and 6 new predators per *E. coli*
 51 cell [10]. If it were using normal binary fission, it could only produce 2^n offspring, potentially forsaking
 52 prey resources if they would only suffice for say 5 rather than 8 offspring. The new *Bdellovibrio* form
 53 flagella and lyse the remains of the prey cell, allowing them to burst out in search of fresh prey.

Mathematical models explain how predator-prey interactions can generate stable oscillations [11]. Most models of microbial predator prey interactions focussed on protists or bacteriophage. Only a few models considered predatory bacteria, see Table S1 for an overview of models and the review by Wilkinson [12]. Varon & Zeigler [13] fitted a Lotka-Volterra model to their experimental results, which led them to propose that a minimal prey density is necessary for the survival of the predator. Crowley's [14] model tracked substrate levels and used Monod kinetics for prey growth, it also included a delay between prey attack and the production of new predators to reflect the ~3 h long bdelloplast phase. The dynamics were destabilised by higher nutrient concentrations, especially at low dilution rates, leading to extreme oscillations. Wilkinson [15] developed two *Bdellovibrio* models, one based on a Holling type II functional response, but without specifically modelling the bdelloplast stage. The other model did include a combined predator-prey complex, but used a Holling type I functional response [15]. Both models again showed a destabilising effect of nutrient enrichment, albeit somewhat ameliorated by the presence of a decoy species. Similarly, Hobley et al. [7], Baker et al. [16] and Said et al. [17] included a predator-prey complex and a Holling type I functional response. Two further models have been developed by Hol et al. [18] and Dattner et al. [19], both including spatial structure. These previous models focussed on the effects of dilution rate and substrate inflow on the dynamics of the chemostat systems. However, the effects of prey and predator characteristics such as prey size, attack kinetics and predation efficiency have not been studied.

In this study, we developed a family of models, based on 'ingredients' from previous models, to identify a unique combination of ingredients that generates realistic outcomes from realistic model assumptions and to ask many novel questions. This model predicts that *Bdellovibrio* has to be much smaller than its prey to survive, which is in agreement with empirical evidence. We also found that high attack rates, large prey sizes and low predator mortality, which might be expected to help the predator, are in fact detrimental for predator survival. Instead, these parameters have optimal values that maximize predator density. These optima occurred at the tipping points into oscillations, and were often very narrow relative to the natural variation of these parameters. Moreover, we found that

81 the system was prone to extreme oscillations in bacterial densities, leading to bottlenecks that would
 82 result in stochastic extinction. Additionally, *Bdellovibrio* would easily be outcompeted by a
 83 bacteriophage. Together, these three key predictions (narrow optima, population bottlenecks and
 84 phage superiority) suggest that *Bdellovibrio* would struggle to survive on a single prey species in a
 85 natural environment full of phages, and where conditions are unlikely to be optimal, and if so, not for
 86 long. Our findings are consistent with the fact that *Bdellovibrio* is a generalist predator so we posit
 87 that our model explains why *Bdellovibrio* is a generalist predator and why it has to be as small as it
 88 is.

89 **Model development**

90 We developed models to investigate the effects of a consumer, the predatory bacterium *Bdellovibrio*
 91 *bacteriovorus* or a bacteriophage, on a bacterial population under continuous culture conditions in a
 92 chemostat. A chemostat captures the low substrate concentrations and slow growth rates typical of
 93 the natural environment [20]. Also, the prey on its own will reach a steady state population density in
 94 the chemostat, which facilitates the investigation of oscillations caused by predation. Oscillations
 95 could not be observed in a batch culture model. We developed a family of models to investigate the
 96 effect of changes in the structure of the model on the predator prey dynamics. This structural
 97 sensitivity analysis [21] identified Model 6 as the most realistic in terms of assumptions and
 98 predictions (see SI text, Table S2 and Fig. S4), so we explored Model 6 further and describe it in Fig.
 99 1 and list its parameters and their values in Table 1. Model 6 included a single abiotic resource
 100 (substrate S), single prey species (N) and a single obligate predator (P). The prey species grew by
 101 consuming the substrate according to Monod kinetics. The predator had a Holling type II functional
 102 response (predation rate proportional to predator density but saturating at high predator density), a
 103 bdelloplast (for *Bdellovibrio*) or infected cell (for bacteriophage) stage and mortality. The bdelloplast
 104 (B) is a distinct stage in the *Bdellovibrio* life cycle that usually lasts for 2 to 4 hours. There are distinct
 105 parallels between this bdelloplast stage and a bacteriophage infected cell as the prey cell does not
 106 grow and replicate when infected by a bacteriophage or consumed by *Bdellovibrio* and the predator
 107 or phage does not prey on or infect further prey. We modelled the bdelloplast or infected cell stage
 108 as a separate entity to account for the delay in producing offspring and the combining of the
 109 consumer and its prey. For simplicity, these entities will be referred to as a bdelloplast from now on,
 110 but the principles apply equally to an infected cell.

Results

Model implementation and validation showed the system to be brittle

We first ensured that the numerical results were reliable by testing all MatLab Ordinary Differential Equation (ODE) solvers. Only the Runge-Kutta ode45 solver could correctly handle all test cases including extreme oscillations (Fig. S1). It was also necessary to set tolerances of this solver to very low values and to constrain variables to be non-negative (Fig. S2). Secondly, we used the same test cases to compare simulations using biomass-based units with those using particle-based units, required to obtain numerically stable simulations with bacteriophage parameters. The results were in agreement (SI text and Fig. S3). These unusual difficulties of obtaining correct numerical results highlight that the system tends to undergo extremely rapid changes, followed by periods of stasis, and is very sensitive to parameter settings.

Structural sensitivity analysis identified most appropriate model

To gain a better understanding of the impact of various modelling choices, we compared a number of ordinary differential equation (ODE) and delay differential equation (DDE) models (Table S2, Fig. S4). We first ran the simulation without predators until the prey had reached a steady state (same for all models). (Model 1) Addition of a predator without a bdelloplast stage (or other form of delay between prey killing and predator birth) gave rise to sustained, extreme oscillations. (Model 2) Incorporating an explicit delay of 4 hours, approximately doubled the oscillatory period from ~150 hours to ~300 hours. (Model 3) Adding mortality reduced the oscillatory period to approximately 100 hours. (Models 4-6) Addition of an explicit bdelloplast stage stabilised the system, resulting in a stable steady state of co-existing predator and prey, regardless of the predator's Holling type functional response. (Model 4 vs. 6) With a Holling type I predator functional response (Model 4), the final prey density was lower, and the predator density higher, than with the saturating type II response (Model 6). (Model 5) Constant input of prey to the system; this is akin to growing the prey on its own in one chemostat that feeds into a second chemostat containing predator. This gave a very similar response to the single chemostat with constant input of substrate. Model 6 was chosen as the most biologically appropriate model for all further work, for several reasons. Firstly, the time

required for *Bdellovibrio* to consume the contents of a prey cell, convert these into new *Bdellovibrio* predators, septate and finally lyse the prey cell to release new predators is about four hours [10]. As the prey is killed very shortly after penetration [24], there is a significant delay between prey killing and birth of new predators, which is best modelled by treating the bdelloplast stage as a separate entity. Secondly, *Bdellovibrio* has a high endogenous respiration rate and a correspondingly low life span in the absence of suitable prey [6]. Hence, it is appropriate to include predator mortality in the model. Thirdly, the saturating Holling type II functional response results from the ‘handling time’ of a predator [25]. This would correspond to the minimum time for a successful attack in a prey-saturated environment, i.e., the attachment and penetration time of *Bdellovibrio* of ~10 minutes [26].

Dynamic regimes from steady states to extreme oscillations

Model 6 has 12 parameters and six possible dynamic regimes (Fig. 2a). In order to gain a better understanding of the factors determining which regimes were observed, we swept through a range of inflow substrate concentrations (S_0) and dilution rates and evaluated the steady state and its stability analytically (see model analysis in SI). As expected, at the highest dilution rates and lowest S_0 , all biological species washed out. Reducing the dilution rate or increasing S_0 enabled survival of first the prey alone and then the predator. Further increases in S_0 destabilised the system, resulting first in damped and then sustained oscillations, before finally reaching a linearly unstable state. We ran simulations in each of the regimes and found that they agreed with the outcomes predicted from the model analysis (Fig. 2b-g), apart from giving sustained, extreme oscillations where the analytical results predicted a linearly unstable state that would correspond to washout (Fig. 2d).

Since the oscillations generated by our *Bdellovibrio* model had a much longer period, extremely abrupt rises and falls and a strong asymmetry in wave shapes compared to the typical Lotka-Volterra models for animal populations, we compared our model with a protist predator model by Curds and Bazin [27] that generated similarly extreme but shorter period oscillations. The period of the protist model could be increased to that of the *Bdellovibrio* model just by replacing protist with *Bdellovibrio* kinetic parameters (SI text on Protist model, Fig. S5).

Dimensional analysis was then performed to deduce which combinations of parameters determine the qualitative behaviour of the system, yielding 7 independent parameter combinations. Essentially, rates were relative to the dilution rate, inflow substrate concentration relative to the prey's substrate affinity and predator affinity replaced by its ratio to prey affinity (for details see SI text). The next 7 sections show the effect of these 7 dimensionless parameter combinations that determine qualitative behaviour.

Bdellovibrio has a minimal and optimal attack rate constant

To our knowledge, Varon & Zeigler [13] is the only study of bacterial predation kinetics, using a relative of *Bdellovibrio*, the marine strain BM4, and *Photobacterium leiognathi* as prey. We used this study to calculate our default values for the attack kinetics. Since different predator and prey combinations may have different kinetics, we investigated the effect of varying the attack rate constant, which in the Holling type II functional response corresponds to the catalytic rate constant of an enzyme with Michaelis-Menten type saturation. The ratio of attack rate constant (μ_p) and half-saturation constant ($K_{N,p}$) gives the initial slope of the Holling type II function, which determines the predation rate at low prey densities. We found that the highest attack rate constant was not best for the predator. Instead, a lower attack rate constant was optimal for the abundance of the predator and conversely for the prey (Fig. 3). The position and width of this optimum varied with dilution rate (Fig. 3c, d). Increasing levels of S_0 narrowed the range of μ_p in which the predator could achieve near maximal density. Below the optimal μ_p , there was a sharp drop to predator extinction. The optimal μ_p was also the rate at which the system underwent a Hopf bifurcation [28] from a stable steady state of co-existence into an oscillatory regime (Fig. 3e-g).

Higher prey growth rate does not benefit prey

Since *Bdellovibrio* can prey on a wide range of Gram-negative bacteria with a wide range of maximum specific growth rates (μ_N), not just *E. coli*, we investigated the effects of changing this. At low inflow substrate concentration (S_0), populations were co-existing in a stable steady state where surprisingly prey growth rate had no effect on prey and predator density (Fig. 4b, cf. phase diagram

in Fig. 2a). At high S_0 , populations were co-existing in sustained oscillations and increasing prey growth rate led to decreasing prey abundance and then a sharp drop in prey abundance at the bifurcation point where the system became stable (Fig. 4a, c). Overall, higher prey growth rate did not benefit the prey.

Predators benefit from mortality

Mortality of *Bdellovibrio* is much higher than that of other bacteria. Therefore, we included it in our model and swept death rates from 0 to 0.2 h^{-1} – substantially more than the 0.06 h^{-1} reported for *Bdellovibrio* [6]. Surprisingly, predator death was beneficial for the predator with an optimal death rate just above a critical mortality where oscillations were replaced with stable co-existence (Fig. S6). Further increases in mortality caused predator extinction. Increasing inflow substrate concentration (S_0) and dilution rate (D) narrowed the predator peak, giving a narrow window for predator persistence (Fig. S6).

There is an optimal maturation rate for bdelloplasts

The bdelloplast stage, where the predator grows inside the prey periplasm with a certain specific growth rate (maturation rate) until prey resources are exhausted and offspring is produced, takes ~3 hours with *E. coli*. We found as with most other parameters that there was a minimal maturation rate required for predator survival and an optimal maturation rate (Fig. S7). This optimal rate was just below a critical rate, above which populations oscillated at higher inflow substrate concentration.

Highest affinity of the predator for its prey is not always optimal

The dimensional analysis identified that the system behaviour depends on the affinity of the predator for its prey relative to the prey affinity for its substrate (SI text), so we swept through a range of predator affinities. At low inflow substrate concentrations (S_0), populations did not cycle and the highest affinity (lowest $K_{N,P}$) was optimal (Fig. S8a). At high S_0 , in contrast, too high affinities (too low $K_{N,P}$) resulted in oscillations with reduced average predator levels (Fig. S8b-e). Lowering the affinity (increasing $K_{N,P}$) resulted in a bifurcation to a stable co-existence that benefited the predator at the expense of the prey. The optimal affinity for the predator was just above this critical $K_{N,P}$ (SI text).

Increasing prey productivity benefits predator until an optimum is reached

Substrate inflow determines prey productivity. Increasing prey productivity from 0 at the outset benefits the predator much more than the prey, which remains at very low levels, until a predator maximum is reached. Further increasing substrate inflow led to a drop in predator and rise in prey (SI text, Fig. S9).

Bdelloplast burst size

The last of the 7 dimensionless parameter combinations is the burst size, which increases with prey size. It is not surprising that there was a minimal burst size for predator survival but it was not expected that this minimal burst size (at higher dilution rates or inflow substrate concentrations) was higher than the experimentally determined burst size for *E. coli* of 3.5 (Fig. S10). We also found an optimal burst size, after which predator abundance declined, i.e., too large prey was not optimal. The lower the dilution rate, the lower the minimal and optimal burst size and the broader the optimum such that the optimum was in the range corresponding to typical prey sizes (Fig. S10). Increasing the burst size above the optimal value resulted in a bifurcation to extreme oscillations, corresponding to a sharp rise in prey and drop in predator average densities, but only at low inflow substrate concentration. At higher S_0 , the optimum was very narrow, above values corresponding to typical prey and oscillations did not occur (Fig. S10).

Bacteriophage outcompetes *Bdellovibrio*

We asked whether bacteriophages would outcompete *Bdellovibrio* on single prey populations and if so, under which conditions and why. The bacteriophage was parameterised based on the well-studied T4 phage infecting *E. coli*. T4 caused oscillations and outcompeted *Bdellovibrio* (Fig. 5). Why did the phage win? There are three processes where the two predators differ. Firstly, the attack kinetics, where the phage had a higher attack rate constant (μ_P), but a lower affinity for prey (higher $K_{N,P}$). Secondly, the kinetics of prey consumption, where the phage had a higher burst size ($Y_{P/B}$) and a faster maturation rate (k_P). Thirdly, the phage did not have mortality like *Bdellovibrio*. To find out which advantage(s) allowed the phage to win, we ran competitions where the phage kept the

prey affinity disadvantage and had one or more of the advantages. Increased burst size ($Y_{P/B}$) alone was sufficient for the phage to win (Fig. S11d). A combination of increased attack rate constant (μ_P) and reduced mortality was also sufficient (Fig. S11a, b, e). Increased maturation rate (k_P) was insufficient even in the presence of either an increased attack rate constant (μ_P) or reduced mortality (Fig. S11c, f, g). Simulating the experiments of Williams et al. [29] where a bacterial predator (but no phage) responded upon addition of prey to a mesocosm, suggests that if suitable phage were present, they should have responded better to the prey addition (SI text, Fig. S12).

Trade-off between high rate versus high yield for predators

Since it takes time to convert prey resources into predator biomass, more complete exploitation of prey resources (higher yield or burst size) should come at the cost of a longer maturation time before offspring will emerge (lower maturation rate). The fast but wasteful predator was assumed to convert bdelloplasts into offspring at a higher rate (k_P one third higher), but had a lower burst size ($Y_{P/B}$ halved) to reflect a reduced yield. The high yield strategy outcompeted the high rate strategy (Fig. 6). Note that the phage had both higher burst size and faster maturation rate, considering this, it is not surprising that the phage won (cf. Fig. 5).

Co-existence of two predators on one prey

We assumed that a trade-off between attack rate constant and affinity for prey exists because time has to be invested for finding prey as well as for binding and examining potential prey. We found that this trade-off would allow coexistence of a ‘fast’ predator with a higher attack rate constant and a ‘high affinity’ predator with lower $K_{N,P}$ on a single prey (Fig. S13).

Minimal and optimal prey cell sizes

Bacteria have a large range of sizes [30]. Clearly there is a minimum size for a prey cell to enter and consume. *Bdellovibrio* also needs to produce at least two offspring per prey cell. Cells of *B. bacteriovorus* are around seven times smaller than *E. coli* cells; this small size might enable *Bdellovibrio* to prey on a wider range of cell sizes. While consuming a larger prey cell will produce more offspring, it will take longer. Also, larger prey cells will mean fewer prey are available if prey

growth is limited by the substrate entering the system as in our chemostat case and most environments. It is not obvious whether the higher number of offspring per prey would offset the disadvantages of longer maturation times and fewer cells to hunt. We found that there was both a minimum prey to predator size ratio required for predator persistence and an optimal value for maximal predator biomass (Fig. 7). Fat prey caused the system to display extreme oscillations. The optimal prey size was just below the size causing these oscillations. Increases in S_0 narrowed the optimum towards the minimal prey size (Fig. 7a-c). Increases in dilution rate also narrowed the peak, but increased the optimal prey size (Fig. 7c, d).

Bottlenecks, permanence and paradox of enrichment

When sweeping parameters, it became clear that the system oscillated with long periods of hundreds of hours and extreme amplitudes for many of the conditions tested, with extremely low minimum values ($< 1 \times 10^{-30}$ mg dry mass ml^{-1}). Such bottlenecks would result in the extinction of the predator (or both predator and prey) in any natural system prone to extrinsic fluctuations and governed by stochastic processes. We therefore examined how prey size (previous section) in combination with dilution rate and inflow substrate concentration (cf. Fig. 2a) would affect robust predator persistence (union of regions of stable co-existence and damped oscillations in Fig. 2a). Robust persistence is known as permanence and means that population densities do not approach zero as the boundary is a repeller [31]. The prey size range allowing permanence narrowed with increasing inflow substrate concentration (Fig. S14). This effect of increased productivity (higher influx of resource for prey) destabilising the system is known as the paradox of enrichment [32] and is typical for predator prey systems. Prey also had to be unrealistically large to enable permanence at higher dilution rates and lower productivity.

Tragedy of the commons

Similarly to increasing productivity, the prey size range enabling permanence shrank with increasing attack rate constant of the predator (Fig. S15). This is an example of the tragedy of the commons, where overexploitation of a shared resource known as the commons is to the detriment of the users of the resource, such as overfishing [33]. Here, a too effective predator that consumed its prey faster

295 than it could regrow led to extinction of the predator. Increases in the inflow substrate concentration
 296 (S_0) narrowed the prey size range for permanence further whilst increases in dilution rate expanded
 297 the range (Fig. S15).

298 ***Global parameter sensitivity analysis***

299 To understand how much the system behaviour would change if the parameters were different due
 300 to changes in substrate, prey or predator species, we conducted a global sensitivity analysis (SI text
 301 and Figs. S16, S17). Substrate concentration was only sensitive to the inflow substrate
 302 concentration. Prey density was not sensitive to substrate concentration and prey growth kinetics but
 303 sensitive to predator parameters. Predator densities were sensitive to the attack rate constant but
 304 also to the maximal specific prey growth rate.

Discussion

Our results suggest that *Bdellovibrio* consuming a single prey species can only survive permanently within a very narrow range of conditions. We therefore characterise this predator prey system as 'brittle'. For example, over a wide range of conditions, the system is prone to extreme oscillations with periods of over a hundred hours and bacterial densities dropping below 0.1 pg ml^{-1} . In a deterministic mathematical model, species densities can eventually recover from even the smallest positive number but in a biological system, there has to be at least a single cell left ($\sim 1 \text{ pg}$). More importantly, when a system contains just a few cells over a long time, stochastic fluctuations will almost inevitably result in the loss of those few cells, leading to local extinction. For most system parameters, there was a narrow range of values that allowed the predator to reproduce fast enough to avoid being washed out without triggering these oscillations, and the optimal value to maximise predator numbers occurred near the threshold triggering these oscillations. Increasing nutrient concentrations narrowed this survival range. Previous models of microbial predator prey interactions in chemostats, including those with *Bdellovibrio* as the predator, also showed a tendency to extreme oscillations, especially for higher nutrient concentrations [14, 15, 27, 34]. Only the group of Varon studied this experimentally in chemostats with the *Bdellovibrio* like predator BM4 and *Photobacterium leiognathi* as prey. They did observe oscillations, albeit less extreme ones than in our model [35, 36]. They also found that BM4 could survive in a chemostat with a prey density of $2\text{-}5 \times 10^4 \text{ CFU ml}^{-1}$ [36], substantially less than the $7 \times 10^5 \text{ CFU ml}^{-1}$ predicted by their model [13] and less still than the $4.4 \times 10^6 \text{ CFU ml}^{-1}$ minimum required for survival in our model. In contrast Keya and Alexander [37] found *Bdellovibrio* strain PF13, isolated from soil, would only replicate in the presence of at least $3 \times 10^7 \text{ CFU ml}^{-1}$ of its *Rhizobium* prey.

329 Given the brittle behaviour predicted by the model it is surprising that *Bdellovibrio* is
 330 ubiquitous in non-marine and *Bdellovibrio* like organisms are ubiquitous in marine
 331 environments [38–41]. This suggests that there must be other forces at work stabilising
 332 population numbers. One possibility is that *Bdellovibrio* is a hotspot organism targeting
 333 habitats of high prey density, or structured environments containing areas of high prey
 334 density such as biofilms, and that these hotspots are the sources of *Bdellovibrio* and other
 335 habitats are sinks. Biofilms represent a lump of prey for *Bdellovibrio*. It possesses the lytic
 336 enzymes needed to chew the extracellular matrix that holds cells within the biofilm and can
 337 likely derive valuable nutrients from this [42]. *Bdellovibrio* can predate the metabolically
 338 inactive cells found deeper within biofilms and its gliding motility allows it to move within the
 339 biofilm [43]. *Bdellovibrio* is also highly motile and contains chemotaxis genes that might
 340 enable it to locate these prey rich areas [44]. Indeed, there is ample evidence that
 341 *Bdellovibrio* is a hotspot organism. Higher numbers are found in sewage rather than in rivers
 342 [45, 46], and downstream sewage treatment plants rather than upstream [47] or in sewage
 343 treated rather than untreated soils [48]. The prey rich rhizosphere sustains higher predator
 344 numbers than the bulk soil [49]. Eutrophic lakes support higher numbers than oligotrophic
 345 lakes [50]. *Bdellovibrio* like organisms are more abundant in marine sediments than the
 346 water column and much higher in oyster shell biofilms [51]. *Bdellovibrio* are likewise more
 347 common in trickling filter biofilms than in their inflow [52].

348 However, the model predicts a narrower range of conditions for survival at the higher
 349 nutrient concentrations expected in hotspots. Therefore, it is unlikely that these hotspots
 350 would sustain *Bdellovibrio* if it were preying only on a single prey species. Moreover, our
 351 model also suggests that bacteriophage would outcompete *Bdellovibrio* under all conditions
 352 tested. This is because phage, as specialist predators, have several big advantages.
 353 Indeed, bacteriophage are far more numerous in nature than *Bdellovibrio* [53, 54] (this

comparison is reasonable assuming that *Bdellovibrio* in total can prey on about half of all bacteria that phage in total can prey on). This is despite the fact that phage have several disadvantages that we did not consider. First, half of the sequenced bacterial genomes contain a CRISPR-CAS system providing adaptive immunity against phage [55]; restriction enzymes cutting up phage DNA are also common [56]. Second, bacteria can rapidly evolve resistance to phages [57], in contrast to *Bdellovibrio*, although phenotypic plasticity of prey can afford temporary protection against *Bdellovibrio* [41]. Third, phage require a metabolically active host cell to replicate whereas *Bdellovibrio* can consume dormant prey [4]. Fourth, phage do not benefit from chemotaxis as *Bdellovibrio* likely does. Nevertheless, phage outnumber *Bdellovibrio*, suggesting that these four factors can be overcome by the huge diversity of phage. Overall, our results suggest that *Bdellovibrio* must prey on several prey species to survive. Indeed, all known *Bdellovibrio* and like organisms have a wide prey range [4, 49, 58, 59].

One might expect that prey cells have to be large enough to give rise to two predator offspring. Surprisingly, our model predicts that *Bdellovibrio* needs prey that is at least 7 times larger than itself. Indeed, this is about the difference in size between *Bdellovibrio* and *E. coli* [60] and other typical prey are of similar size. However, the predicted optimal prey size is considerably larger. Maybe *Bdellovibrio* cannot be much smaller than it already is or accessing diverse prey species avoids precarious oscillations.

Our model also predicts that a predator that kills its prey too efficiently (has a too high attack rate, or too little mortality) will drive its prey to extinction and then become extinct itself, resulting in a tragedy of the commons [33]. Indeed, *Bdellovibrio* has a much higher mortality rate than other bacteria. While this may make over-exploitation of prey less likely, the high mortality is likely caused by its high energy expenditure when swimming fast and not feeding at the same time, also its small size prevents storage of large energy reserves [6].

In conclusion, our model results suggest that *Bdellovibrio* and like organisms are unlikely to survive in most natural environments if they were preying only on single prey species. They would also be outcompeted by phage. In line with empirical evidence, *Bdellovibrio* ought to be a generalist predator and would only thrive in prey density hotspots – which it should be able to find by chemotaxis. For application as a living antibiotic to reduce the abundance of pathogens or antimicrobial resistant bacteria in aquaculture or plant and animal agriculture, *Bdellovibrio* would be expected to be more effective where multiple prey species, not only the target species, are naturally available or added artificially.

Acknowledgments

We would like to thank Andrew Lovering (University of Birmingham) and Liz Sockett (University of Nottingham) for helpful discussions of *Bdellovibrio* biology. KS is supported by a Biotechnology and Biological Sciences Research Council (BBSRC) UK funded, Midlands Integrative Biosciences Training Partnership (MIBTP) PhD studentship.

Competing Interests

The authors declare no competing financial interests.

References

1. Paine RT. A note on trophic complexity and community stability. *Am Nat* 1969; **103**: 91–93.
2. Krebs CJ, Boonstra R, Stan B, Sinclair ARE. What drives the 10-year cycle of Snowshoe hares? *Bioscience* 2001; **51**: 25–35.
3. Arcilla MS, van Hattem JM, Matamaros S, Melles DC, Penders J, de Jong MD. Dissemination of the *mcr-1* colistin resistance gene. *Lancet Infect Dis* 2016; **16**: 147–149.
4. Stolp H, Starr MP. *Bdellovibrio bacteriovorus* gen. et sp. n., a predatory, ectoparasitic, and bacteriolytic microorganism. *Antonie Van Leeuwenhoek* 1963; **29**: 217–248.
5. Lambert C, Morehouse KA, Chang C-Y, Sockett RE. *Bdellovibrio*: growth and development during the predatory cycle. *Curr Opin Microbiol* 2006; **9**: 639–644.
6. Hespell RB, Thomashow MF, Rittenberg SC. Changes in cell composition and viability of *Bdellovibrio bacteriovorus* during starvation. *Arch Microbiol* 1974; **97**: 313–327.
7. Hobley L, King JR, Sockett RE. *Bdellovibrio* predation in the presence of decoys: Three-way bacterial interactions revealed by mathematical and experimental analyses. *Appl Environ Microbiol* 2006; **72**: 6757–6765.
8. Starr MP, Baigent NL. Parasitic interaction of *Bdellovibrio bacteriovorus* with other bacteria. *J Bacteriol* 1966; **91**: 2006–2017.
9. Romo AJ, Ruby EG, Saier MHJ. Effect of *Bdellovibrio bacteriovorus* infection on the phosphoenolpyruvate:sugar phosphotransferase system in *Escherichia coli*: evidence for activation of cytoplasmic proteolysis. *Res Microbiol Paris* 1992; **143**: 5–14.
10. Seidler RJ, Starr MP. Factors affecting the intracellular parasitic growth of *Bdellovibrio bacteriovorus* developing within *Escherichia coli*. *J Bacteriol* 1969; **97**: 912–23.
11. Berryman AA. The origins and evolution of predator-prey theory. *Ecology* 1992; **73**:

- 419 1530–1535.
- 420 12. Wilkinson MHF. Mathematical modelling of predatory prokaryotes. In: Jurkevitch E
421 (ed). *Predatory Prokaryotes*. 2006. pp 94–130.
- 422 13. Varon M, Zeigler BP. Bacterial predator-prey interaction at low prey density. *Appl*
423 *Environ Microbiol* 1978; **36**: 11–17.
- 424 14. Crowley PH, Straley SC, Craig RJ, Culin JD, Fu YT, Hayden TL, et al. A model of prey
425 bacteria, predator bacteria, and bacteriophage in continuous culture. *J Theor Biol*
426 1980; **86**: 377–400.
- 427 15. Wilkinson MHF. Predation in the presence of decoys: an inhibitory factor on pathogen
428 control by bacteriophages or *Bdellovibrios* in dense and diverse ecosystems. *J Theor*
429 *Biol* 2001; **208**: 27–36.
- 430 16. Baker M, Negus D, Raghunathan D, Radford PM, Moore C, Clark G, et al. Measuring
431 and modelling the response of *Klebsiella pneumoniae* KPC prey to *Bdellovibrio*
432 *bacteriovorus* predation, in human serum and defined buffer. *Sci Rep* 2017; **7**: 1–18.
- 433 17. Said N, Chatzinotas A, Schmidt M. Have an ion on it: The life-cycle of *Bdellovibrio*
434 *bacteriovorus* viewed by helium-ion microscopy. *Adv Biosyst* 2019; **3**: 1800250.
- 435 18. Hol FJH, Rotem O, Jurkevitch E, Dekker C, Koster DA. Bacterial predator–prey
436 dynamics in microscale patchy landscapes. *Proc R Soc B Biol Sci* 2016; **283**:
437 20152154.
- 438 19. Dattner I, Miller E, Petrenko M, Kadouri DE, Jurkevitch E, Huppert A. Modelling and
439 parameter inference of predator–prey dynamics in heterogeneous environments using
440 the direct integral approach. *J R Soc Interface* 2017; **14**.
- 441 20. Jannasch HW. Estimations of bacterial growth rates in natural waters. *J Bacteriol*
442 1969; **99**: 156–160.
- 443 21. Hellweger FL, Clegg RJ, Clark JR, Plugge CM, Kreft J-U. Advancing microbial

- 444 sciences by individual-based modelling. *Nat Rev Microbiol* 2016; **14**: 461–471.
- 445 22. Kreft J-U, Booth G, Wimpenny JWT. BacSim, a simulator for individual-based
446 modelling of bacterial colony growth. *Microbiology* 1998; **144**: 3275–3287.
- 447 23. Kubitschek HE, Friske JA. Determination of bacterial cell volume with the Coulter
448 Counter. *J Bacteriol* 1986; **168**: 1466–7.
- 449 24. Rittenberg SC, Shilo M. Early host damage in the infection cycle of *Bdellovibrio*
450 *bacteriovorus*. *J Bacteriol* 1970; **102**: 149–60.
- 451 25. Jeschke JM, Kopp M, Tollrian R. Predator functional responses: Discriminating
452 between handling and digesting prey. *Ecol Monogr* 2002; **72**: 95–112.
- 453 26. Hespell RB. Glycolytic and tricarboxylic acid cycle enzyme activities during
454 intraperiplasmic growth of *Bdellovibrio bacteriovorus* on *Escherichia coli*. *J Bacteriol*
455 1976; **128**: 677–80.
- 456 27. Curds CR, Bazin MJ. Protozoan predation in batch and continuous cultures. In: Droop
457 MR, Jannasch HW (eds). *Advances in Aquatic Microbiology*. 1980. Academic Press,
458 pp 115–176.
- 459 28. Hopf E. A mathematical example displaying features of turbulence. *Commun Pure*
460 *Appl Math* 1948; **1**: 303–322.
- 461 29. Williams HN, Lympelopoulou DS, Athar R, Chauhan A, Dickerson TL, Chen H, et al.
462 *Halobacteriovorax*, an underestimated predator on bacteria: potential impact relative
463 to viruses on bacterial mortality. *ISME J* 2016; **10**: 491–499.
- 464 30. Schulz HN, Barker B, Jørgensen BB. Big bacteria. *Annu Rev Microbiol* 2001; **55**: 105–
465 37.
- 466 31. Jansen VAA, Sigmund K. Shaken not stirred: on permanence in ecological
467 communities. *Theor Popul Biol* 1998; **54**: 195–201.
- 468 32. Rosenzweig ML. Paradox of enrichment: destabilization of exploitation ecosystems in

- ecological time. *Science* 1971; **171**: 385–7.
33. Hardin G. The tragedy of the commons. The population problem has no technical solution; it requires a fundamental extension in morality. *Science* 1968; **162**: 1243–8.
34. Nisbet RM, Cunningham A, Gurney WS. Endogenous metabolism and the stability of microbial prey-predator systems. *Biotechnol Bioeng* 1983; **25**: 301–306.
35. Varon M. Selection of predation-resistant bacteria in continuous culture. *Nature* 1979; **277**: 386–388.
36. Varon M, Fine M, Stein A. The maintenance of *Bdellovibrio* at low prey density. *Microb Ecol* 1984; **10**: 95–98.
37. Keya SO, Alexander M. Regulation of parasitism by host density: the *Bdellovibrio-Rhizobium* interrelationship. *Soil Biol Biochem* 1974; **7**: 231–237.
38. Shilo M, Bruff B. Lysis of Gram-negative bacteria by host-independent ectoparasitic *Bdellovibrio bacteriovorus* isolates. *J Gen Microbiol* 1965; **40**: 317–28.
39. Williams HN. The recovery of high numbers of *Bdellovibrios* from the surface water micro layer. *Can J Microbiol* 1987; **33**.
40. Williams HN, Schoeffield AJ, Guether D, J K, Shah D, Falkler JWA. Recovery of *Bdellovibrios* from submerged surfaces and other aquatic habitats. *Microb Ecol* 1995; **29**: 39–48.
41. Shemesh Y, Jurkevitch E. Plastic phenotypic resistance to predation by *Bdellovibrio* and like organisms in bacterial prey. *Environ Microbiol* 2004; **6**: 12–18.
42. Im H, Dwidar M, Mitchell RJ. *Bdellovibrio bacteriovorus* HD100, a predator of Gram-negative bacteria, benefits energetically from *Staphylococcus aureus* biofilms without predation. *ISME J* 2018; **12**: 2090–2095.
43. Lambert C, Fenton AK, Hobley L, Sockett RE. Predatory *Bdellovibrio* bacteria use gliding motility to scout for prey on surfaces. *J Bacteriol* 2011; **193**: 3139–41.

- 494 44. Hobley L, Lerner TR, Williams LE, Lambert C, Till R, Milner DS, et al. Genome
495 analysis of a simultaneously predatory and prey-independent, novel *Bdellovibrio*
496 bacteriovorus from the River Tiber, supports in silico predictions of both ancient and
497 recent lateral gene transfer from diverse bacteria. *BMC Genomics* 2012; **13**: 670.
- 498 45. Dias FF, Bhat J V. Microbial ecology of activated sludge. II. bacteriophages,
499 *Bdellovibrio*, coliforms, and other organisms. *Appl Microbiol* 1965; **13**: 257–261.
- 500 46. Seliavko A V., Lambina VA. Distribution of microorganisms lysing *Pseudomonas*
501 *aeruginosa* and *Staphylococcus aureus* in the system of sewage waters. *Nauchnye*
502 *Doki Vyss Shkoly Biol Nauk* 1985; **4**: 84–8.
- 503 47. Staples DG, Fry JC. Factors which influence the enumeration of *Bdellovibrio*
504 *bacteriovorus* in sewage and river water. *J Appl Bacteriol* 1973; **36**: 1–11.
- 505 48. Klein DA, Casida LEJ. Occurrence and enumeration of *Bdellovibrio bacteriovorus* in
506 soil capable of parasitizing *Escherichia coli* and indigenous soil bacteria. *Can J*
507 *Microbiol* 1967; **13**: 1235–41.
- 508 49. Jurkevitch E, Minz D, Ramati B, Barel G. Prey range characterization, ribotyping, and
509 diversity of soil and rhizosphere *Bdellovibrio spp.* isolated on phytopathogenic
510 bacteria. *Appl Environ Microbiol* 2000; **66**: 2365–71.
- 511 50. Chauhan A, Cherrier J, Williams HN. Impact of sideways and bottom-up control
512 factors on bacterial community succession over a tidal cycle. *Proc Natl Acad Sci U S*
513 *A* 2009; **106**: 4301–6.
- 514 51. Williams HN, Kelley JI, Baer ML, Turng B-F. The association of *Bdellovibrios* with
515 surfaces in the aquatic environment. *Can J Biochem* 1995; **41**: 1142–1147.
- 516 52. Fry JC, Staples DG. Distribution of *Bdellovibrio bacteriovorus* in sewage works, river
517 water, and sediments. *Appl Environ Microbiol* 1976; **31**: 469–474.
- 518 53. Kutter EM, Sulakvelidze A. Bacteriophages biology and applications. In: Kutter E,

- 519 Sulakvelidze A (eds). *Bacteriophages Biology and Applications*. 2004. CRC Press,
520 Boca Raton, pp 1–4.
- 521 54. Stolp H. The *Bdellovibrios*: Bacterial parasites of bacteria. *Annu Rev Phytopathol*
522 1973; **11**: 53–76.
- 523 55. Koonin E V., Wolf YI. Evolution of the CRISPR-Cas adaptive immunity systems in
524 prokaryotes: models and observations on virus–host coevolution. *Mol Biosyst* 2015;
525 **11**: 20–27.
- 526 56. Bayliss CD, Callaghan MJ, Moxon ER. High allelic diversity in the methyltransferase
527 gene of a phase variable type III restriction-modification system has implications for
528 the fitness of *Haemophilus influenzae*. *Nucleic Acids Res* 2006; **34**: 4046–59.
- 529 57. Luria SE, Delbrück M. Mutations of bacteria from virus sensitivity to virus resistance.
530 *Genetics* 1943; **28**: 491–511.
- 531 58. Feng S, Tan CH, Cohen Y, Rice SA. Isolation of *Bdellovibrio bacteriovorus* from a
532 tropical wastewater treatment plant and predation of mixed species biofilms
533 assembled by the native community members. *Environ Microbiol* 2016.
- 534 59. Dashiff A, Junka RA, Libera M, Kadouri DE. Predation of human pathogens by the
535 predatory bacteria *Micavibrio aeruginosavorus* and *Bdellovibrio bacteriovorus*. *J Appl*
536 *Microbiol* 2011; **110**: 431–444.
- 537 60. Cover WH, Martinez RJ, Rittenberg SC. Permeability of the boundary layers of
538 *Bdellovibrio bacteriovorus* 109J and its bdelloplasts to small hydrophilic molecules. *J*
539 *Bacteriol* 1984; **157**: 385–390.

540

Table 1 Baseline parameters for the principal model (Model 6) and range over which global sensitivity analysis was performed. All yields are expressed in the form $Y_{A/B}$, which is the yield of consumer A per resource B consumed. The affinity of consumer B for resource A is expressed as $K_{A,B}$.

Fig. 1 Principal model used to track predator and prey densities under chemostat conditions (Model 6, see Table S2). Substrate is consumed by prey to fuel growth. Prey and predators combine to form a bdelloplast. The bdelloplast matures to give new predators. Predators have mortality. This is the model used unless stated otherwise.

Fig. 2 Dynamic regimes of Model 6. **a** Analytically calculated regimes, as depending on the inflow substrate concentration S_0 and dilution rate D . **b-g** Simulations at $S_0 = 0.25 \text{ mg ml}^{-1}$ and increasing dilution rates, indicated by white crosses in **a**. **b** Damped oscillations that ended in steady state co-existence. **c** Damped oscillations of much shorter period than **b**. **d** Amplifying oscillations that ended in sustained, extreme oscillations. **e** The analytically predicted linearly unstable region gave sustained, extreme oscillations in the numerical simulations. **f** Stable co-existence of predator and prey. **g** Predator extinction.

Fig. 3 Minimal and optimal attack rate constant (μ_p). The average population densities and substrate concentrations, oscillatory periods and phase shifts strongly depend on the attack rate constant, shown at increasing inflow substrate concentrations (S_0) and two dilution rates (cf. phase diagram in Fig. 2a). Top row shows concentrations at steady state or averaged over one oscillatory cycle. Bottom row shows the oscillatory period (blue, left axis) and phase shifts (green, right axis) from substrate peak to peak of prey (solid line), free *Bdellovibrio* (dashed line) or bdelloplast (dotted line). Note that oscillations start above the optimal attack rate constant. In order to obtain accurate simulation results at all parameter values, the absolute tolerance of the ode45 solver had to be reduced from 1×10^{-9} to 1×10^{-12} .

Fig. 4 Increasing the maximal specific growth rate of the prey (μ_N) leads to a sharp drop of prey density and stabilizes the system at high inflow substrate concentrations (S_0). The system was more stable at low S_0 (cf. phase diagram in Fig. 2a). Top row shows concentrations at steady state or averaged over one oscillatory cycle. Bottom row shows the oscillatory period (blue, left axis) and phase shifts (green, right axis) from substrate peak to peak of prey (solid line), free *Bdellovibrio* (dashed line) or bdelloplast (dotted line). Note that oscillations occur at the higher inflow substrate concentration and below a critical μ_N . Prey density is much higher in the oscillatory regimes where it decreases with increasing prey growth rates. Since prey density will of course be 0 if μ_N is zero, there is an optimal prey growth rate.

Fig. 5 T4 phage caused oscillations and outcompeted *Bdellovibrio* at two very different inflow substrate concentrations. Dilution rate was 0.02 h^{-1} . Note that units had to be changed to particle densities from the biomass densities used in the other sections. Panel c shows a zoomed in version of a peak in panel b, the substrate concentration appears to be constant on the scale needed to show the other variables.

Fig. 6 A slow, but efficient (high $Y_{P/B}$) predator outcompetes a fast (high k_P) but wasteful predator at different inflow substrate concentrations (S_0). Dilution rate = 0.02 h^{-1} .

Fig. 7 Minimal and optimal prey size, relative to predator size (0.14 fl). Predator density showed a broad optimum at low inflow substrate concentrations (S_0) that became narrower at higher S_0 and dilution rate. Too large prey caused oscillations. Top row shows concentrations at steady state or averaged over one oscillatory cycle. Bottom row shows the oscillatory period (blue, left axis) and phase shifts (green, right axis) from substrate peak to peak of prey (solid line), free *Bdellovibrio* (dashed line) or bdelloplast (dotted line).

Fig. 1 Principal model used to track predator and prey densities under chemostat conditions (Model 6, see Table S2). Substrate is consumed by prey to fuel growth. Prey and predators combine to form a bdelloplast. The bdelloplast matures to give new predators. Predators have mortality. This is the model used unless stated otherwise.

Fig. 2 Dynamic regimes of Model 6. **a** Analytically calculated regimes, as depending on

the inflow substrate concentration S_0 and dilution rate D . **b-g** Simulations at $S_0 = 0.25$

mg ml⁻¹ and increasing dilution rates, indicated by white crosses in **a**. **b** Damped

oscillations that ended in steady state co-existence. **c** Damped oscillations of much

shorter period than **b**. **d** Amplifying oscillations that ended in sustained, extreme

oscillations. **e** The analytically predicted linearly unstable region gave sustained,

extreme oscillations in the numerical simulations. **f** Stable co-existence of predator and

prey. **g** Predator extinction.

Fig. 3 Minimal and optimal attack rate constant (μ_p). The average population densities and substrate concentrations, oscillatory periods and phase shifts strongly depend on the attack rate constant, shown at increasing inflow substrate concentrations (S_0) and two dilution rates (cf. phase diagram in Fig. 2a). Top row shows concentrations at steady state or averaged over one oscillatory cycle. Bottom row shows the oscillatory period (blue, left axis) and phase shifts (green, right axis) from substrate peak to peak of prey (solid line), free *Bdellovibrio* (dashed line) or bdelloplast (dotted line). Note that oscillations start above the optimal attack rate constant. In order to obtain accurate simulation results at all parameter values, the absolute tolerance of the ode45 solver had to be reduced from 1×10^{-9} to 1×10^{-12} .

Fig. 4 Increasing the maximal specific growth rate of the prey (μ_N) leads to a sharp drop of prey density and stabilizes the system at high inflow substrate concentrations (S_0). The system was more stable at low S_0 (cf. phase diagram in Fig. 2a). Top row shows concentrations at steady state or averaged over one oscillatory cycle. Bottom row shows the oscillatory period (blue, left axis) and phase shifts (green, right axis) from substrate peak to peak of prey (solid line), free *Bdellovibrio* (dashed line) or bdelloplast (dotted line). Note that oscillations occur at the higher inflow substrate concentration and below a critical μ_N . Prey density is much higher in the oscillatory regimes where it decreases with increasing prey growth rates. Since prey density will of course be 0 if μ_N is zero, there is an optimal prey growth rate.

Fig. 5 T4 phage caused oscillations and outcompeted *Bdellovibrio* at two very different inflow substrate concentrations. Dilution rate was 0.02 h^{-1} . Note that units had to be changed to particle densities from the biomass densities used in the other sections. Panel c shows a zoomed in version of a peak in panel b, the substrate concentration appears to be constant on the scale needed to show the other variables.

Fig. 6 A slow, but efficient (high $Y_{P/B}$) predator outcompetes a fast (high k_p) but wasteful predator at different inflow substrate concentrations (S_0). Dilution rate = 0.02 h^{-1} .

Fig. 7 Minimal and optimal prey size, relative to predator size (0.14 fl). Predator density showed a broad optimum at low inflow substrate concentrations (S_0) that became narrower at higher S_0 and dilution rate. Too large prey caused oscillations. Top row shows concentrations at steady state or averaged over one oscillatory cycle. Bottom row shows the oscillatory period (blue, left axis) and phase shifts (green, right axis) from substrate peak to peak of prey (solid line), free *Bdellovibrio* (dashed line) or bdelloplast (dotted line).

Table 1 – Baseline parameters for the principal model and range over which global sensitivity analysis was performed (Model 6). All yields are expressed in the form $Y_{A/B}$, which is the yield of consumer A per resource B consumed. The affinity of consumer B for resource A is expressed as $K_{A,B}$.

Parameter	Units	Default value	Minimum	Maximum	Source
Inflow substrate concentration (S_0)	$g\ l^{-1}$	5×10^{-3}	5×10^{-3}	2.5	Assumed to be $\sim 2 \times K_{S,N}$
Prey maximum specific growth rate (μ_N)	h^{-1}	1.23	$\frac{3}{4} \ln(2)$	$3 \ln(2)$	[1]
Predator attack rate constant (μ_P)	$g\ bdelloplast\ g\ predator^{-1}\ h^{-1}$	0.38	0.3	6	See cover letter
Predator affinity for prey ($K_{N,P}$)	$g\ prey\ l^{-1}$	8.6×10^{-4}	1.5×10^{-6}	6.5×10^{-3}	See cover letter
Yield of bdelloplasts from predators ($Y_{B/P}$)	$g\ bdelloplast\ g\ predator^{-1}$	8	5	20	Relative sizes of <i>E. coli</i> and <i>Bdellovibrio</i> cells [2, 3]
Bdelloplast maturation rate (k_P)	$g\ predator\ g\ bdelloplast^{-1}\ h^{-1}$	0.109	0.075	0.3	[4]
Predator mortality rate (m)	h^{-1}	0.06	0.03	0.09	[5]
Dilution rate (D)	h^{-1}	Always 0.125			

Prey affinity for substrate ($K_{S,N}$)	$g\ l^{-1}$	Always 2.34×10^{-3}	[1]
Yield of prey from substrate ($Y_{N/S}$)	$g\ dry\ mass\ prey\ g\ substrate^{-1}$	Always 0.4444	[1]
Yield of bdelloplasts from prey ($Y_{B/N}$)	$g\ bdelloplast^{-1}\ g\ prey^{-1}$	Always $Y_{B/P}/(Y_{B/P} - 1)$	Relative sizes of <i>E. coli</i> and <i>Bdellovibrio</i> cells [2, 3]
Yield of predators from bdelloplasts ($Y_{P/B}$)	$g\ predator\ g\ bdelloplast^{-1}$	Always 0.438	Relative sizes of <i>E. coli</i> and <i>Bdellovibrio</i> cells and burst size [2–4]

1. Kreft J-U, Booth G, Wimpenny JWT. BacSim, a simulator for individual-based modelling of bacterial colony growth. *Microbiology* 1998; **144**: 3275–3287.
2. Stolp H, Starr MP. *Bdellovibrio bacteriovorus* gen. et sp. n., a predatory, ectoparasitic, and bacteriolytic microorganism. *Antonie Van Leeuwenhoek* 1963; **29**: 217–248.
3. Kubitschek HE, Friske JA. Determination of bacterial cell volume with the Coulter Counter. *J Bacteriol* 1986; **168**: 1466–7.
4. Seidler RJ, Starr MP. Factors affecting the intracellular parasitic growth of *Bdellovibrio bacteriovorus* developing within *Escherichia coli*. *J Bacteriol* 1969; **97**: 912–23.
5. Hespell RB, Thomashow MF, Rittenberg SC. Changes in cell composition and viability of *Bdellovibrio bacteriovorus* during starvation. *Arch Microbiol* 1974; **97**: 313–327.