1 **RT:** Telomere length in reptiles

2

3 **Telomere length covaries with age across an elevational gradient in a**

4 **Mediterranean lizard**

5

6

7 **Pablo Burraco^{1,2*}, Mar Comas^{3,4}, Senda Reguera³, Francisco Javier Zamora-Camacho⁵,**

8 **Gregorio Moreno-Rueda³**

9 ¹ Animal Ecology, Department of Ecology and Genetics, Evolutionary Biology Centre,

10 Uppsala University, Norbyvägen 18D, SE-75236 Uppsala Sweden

11 ² Ecology, Evolution and Development Group, Doñana Biological Station (CSIC, Spain).

12 ³ Departamento de Zoología, Facultad de Ciencias, Universidad de Granada (Spain)

13 ⁴ Estación Biológica de Doñana (EBD-CSIC), Avda. Américo Vespucio 26, Seville E-41092,

14 (Spain).

15 ⁵ Departamento de Biogeografía y Cambio Global, Museo Nacional de Ciencias Naturales

16 (CSIC, Spain)

17

18 *Corresponding authors: pablo.burraco@ebc.uu.se

19

20 **Keywords:** altitude, body condition, ectotherms, growth, reptiles, senescence, telomere

21 shortening

22

23

24

25

26 **Abstract**

27 The timing of organisms' senescence is developmentally programmed but also shaped by the
 28 interaction between environmental inputs and life-history traits. In ectotherms, ageing
 29 dynamics are still poorly understood despite their particularities concerning thermoregulation,
 30 regeneration capacity, or growth trajectory. Here, we investigate the role of life-history traits
 31 such as age, sex, body size, body condition, and tail autotomy (i.e self-amputation) in shaping
 32 telomere length of six populations of the Algerian sand lizard (*Psammodromus algirus*)
 33 distributed across an elevational gradient from 300 to 2500 meters above the sea level.
 34 Additionally, we show in a review table the available information on reptiles' telomere
 35 length. We found that telomeres elongated with lizards' age. We also observed that body size
 36 and age class showed a positive relationship, suggesting that cell replication did not shorten
 37 lizards' telomeres by itself. Elevation affected telomere length in a non-linear way, a pattern
 38 that mirrored the variation in age structure across elevation. Telomere length was unaffected
 39 by tail autotomy, and was sex-independent, but positively correlated with body condition. Our
 40 results show that telomeres elongate throughout the first four years of lizards' lifetime, a
 41 process that stress the role of telomerase in maintaining ectothermic telomeres, and, likely, in
 42 extending lifespan in organisms with indeterminate growth. Regarding the non-linear impact
 43 that elevation had on telomere length of lizards, our results suggest that habitat (mainly
 44 temperature) and organisms' condition might play a key role in regulation ageing rate. Our
 45 findings emphasize the relevance of understanding species' life histories (e.g. age and body
 46 condition) and habitat characteristics for fully disentangling the causes and consequences of
 47 lifespan trajectory.

48

49

50

51 **Introduction**

52 The match between phenotypes and the environment not only defines species distribution
 53 (Sih, Ferrari, & Harris, 2011; Zamudio, Bell, & Mason, 2016) but also individual lifespan
 54 (Monaghan, 2007; Marasco et al., 2017; Ratikainen & Kokko, 2019). The study of the
 55 evolutionary underpinnings of ageing has been a long-standing topic both in ecological and
 56 medical research, pointing out mitochondrial activity and its underlying damages as the main
 57 drivers of individual variation in life expectancy across taxa (Selman, Blount, Nussey, &
 58 Speakman, 2012; Ziegler, Wiley, & Velarde, 2015; Vágási et al., 2019). Most studies on
 59 ageing in vertebrates have been conducted in endotherms whereas ectotherms have received
 60 scarce attention despite their singularities concerning thermoregulation, regeneration capacity,
 61 metabolism, or growth trajectory (Olsson, Wapstra, & Friesen, 2018; Monaghan, Eisenberg,
 62 Harrington, & Nussey, 2018). Understanding the role of environmental conditions and life-
 63 histories on shaping senescence in ectothermic vertebrates would increase the current
 64 knowledge about their evolutionary and ecological dynamics.

65 A reliable way for addressing individual ageing is through estimates of telomere
 66 length (Horn et al., 2011; Bateson, 2015; Angelier, Weimerskirch, Barbraud, & Chastel,
 67 2019). Telomeres are non-coding repeated sequences (TTAGGG_n in vertebrates) located at
 68 the termini of chromosomes, essential for maintaining genomic stability and for protecting
 69 cells from chromosome degradation and fusion (O'Sullivan & Karlseder, 2010). Telomeric
 70 sequences shorten after each cell replication due to the *end replication problem*, which occurs
 71 once the last piece of RNA primer is removed during replication and DNA cannot be
 72 extended (Allsopp et al., 1995). Hence, more cell replications may involve shorter length of
 73 telomeres. Consequently, telomere length should decrease with age, as often observed in
 74 mammals and birds (Haussmann et al., 2003; Heidinger et al., 2012). When telomeres become
 75 very short, apoptosis is induced (Aubert & Lansdrop, 2008). However, the expression of

76 telomerase, a reverse transcriptase enzyme that adds repeat sequence to the 3' end of
 77 telomeres, can either partially or fully restore telomere erosion. Telomerase expression is
 78 often detected in the germline and in embryonic tissues both in endotherms and ectotherms
 79 (Ingles & Deakin, 2016). Particularly in ectothermic vertebrates, telomerase is not only active
 80 early in life, but also in adult somatic tissues, like in the fish medaka (*Oryzias latipes*;
 81 Klapper, Heidorn, Kühne, Parwaresch, & Krupp, 1998) or in the African water frog (*Xenopus*
 82 *laevis*; Bousman, Schneider, & Shampay, 2003). In this line, telomere elongation has been
 83 found throughout larval development of the Atlantic salmon (*Salmo salar*, McLennan et al.,
 84 2016) and of the common water frog (*Rana temporaria*; Burraco et al., submitted), and also
 85 during the first years of life in some reptiles (e.g. Olsson, Pauliny, Wapstra, & Blomqvist,
 86 2010; Ujvari et al., 2017). In endotherms, telomere elongation after birth is not widespread
 87 and takes place under very particular conditions, as during the active season of the edible
 88 dormouse (*Glis glis*; Hoelzl, Cornils, Smith, Moodley, & Ruf, 2016), or in some long-lived
 89 birds (“elongation hypothesis”, see Haussmann & Mauck, 2007).

90 Such differences in telomere dynamics between ectothermic and endothermic
 91 vertebrates might be linked to organisms’ thermoregulation capacity and growth trajectories
 92 (typically, indeterminate growth in ectotherms versus determinate growth in endotherms), and
 93 explain lifespan across species (Jones et al., 2014). In reptiles, a paraphyletic group, 11
 94 studies have investigated the variation in telomere length across individuals’ lifetime (Table
 95 1). Four studies found that telomeres shorten with age, whereas in three cases telomere length
 96 increased with age in any of the two sexes (Table 1). A quadratic sex-dependent relationship
 97 between telomere length and age was observed in two studies, i.e. telomeres increase their
 98 length until a certain age, and then shorten (Table 1). Meanwhile, three studies found no
 99 effect of age on telomere length in reptiles (Table 1). The high inter-species variation

regarding the relation between telomere length across reptiles' lifetime highlights the need of further research to unravel it.

Telomere length, at a given ontogenetic point, is not only a function of cell replication but also of the organisms' ability to cope with stress across their life. In vertebrates, harmful conditions often enhance glucocorticoids secretion, which involve metabolic processes that provoke the overproduction of reactive oxygen species (ROS). This overproduction of ROS induces an oxidative state in cells that can damage essential biomolecules like lipids, proteins or DNA (Isaksson, 2015; Luceri et al., 2018), including telomeres (Haussman & Marchetto, 2010; Monaghan, 2014; Angelier, Costantini, Blevin, & Chastel, 2018). As a consequence, telomere length is a reliable indicator of the amount of stress accumulated by organisms across time (Young, 2018). Indeed, positive relationships between telomere length and organisms' life expectancy (Barret, Burke, Hammers, Komdeur, & Richardson, 2013; Wilbourn et al., 2018), reproductive outcome (Eastwood et al., 2019), or immunocompetence (Alder et al., 2018) are well established. In ectothermic vertebrates, telomere shortening is commonly associated with increased growth rate, bold personality, or predator exposure (reviewed in Olsson et al. 2018). Particularly in reptiles, telomere length positively correlates with social signalling, reproductive output and mode, or lifespan whereas it is unaffected by foraging behaviour or by ectoparasite load (Table 1). One might expect telomeres to shorten as body size increases across lifetime since it implies more cellular replications. However, only a few studies on reptiles have observed a significant effect of body size or growth rate on telomere length (Table 1), unlike in fish (McLennan et al. 2016) or amphibians (Burraco, Díaz-Paniagua, & Gomez-Mestre, 2017a).

Here, we aim to understand the role of life-history traits and environmental conditions on ageing of a lizard species. To this end, we investigated the effect of age, sex, body size, and body condition on telomeres across six populations of the Algerian sand lizard

(*Psammodromus algirus*) across a substantial mountain gradient (from 300 to 2500 metres above sea level, m.a.s.l thereafter). We predicted an effect of age class and body size on telomere length, either in a positive, negative, or quadratic way, which may be sex-linked, regarding the available literature on reptile telomeres (Table 1). The elevational gradient studied here allows to determine to which extent environment can influence telomere dynamics in lizard populations. As we ascend in altitude, temperature and activity time decrease while hibernation time increases (Zamora-Camacho, Reguera, Moreno-Rueda, & Pleguezuelos, 2013), which may induce lower telomere attrition (Hoelzl et al. 2016; Kirby, Johnson, Alldredge, & Pauli, 2019). Also, at higher elevations, lizards are exposed to conditions that might reduce telomere shortening such as low risk of overheating (Zamora-Camacho, Reguera, & Moreno-Rueda, 2016) or low ectoparasitism (Álvarez et al., 2018). In addition, oxidative damage –one of the main drivers of telomere attrition (Reichert & Stier, 2017)– decreases at higher elevation (Reguera, Zamora-Camacho, Trenzado, Sanz & Moreno-Rueda, 2014, Reguera et al., 2015). Therefore, we predict that, once corrected for age, populations at higher elevation would have longer telomeres.

Material and methods

General procedures

The lizard *P. algirus* is a medium-large lacertid (53-80 mm snout-vent length –SVL– in our study area) that inhabits shrubby habitats in the Mediterranean region from southwestern Europe and north-western Africa (Salvador, 2015). In the Sierra Nevada mountain system (SE Spain), we sampled individuals from six populations, which inhabit at 300, 700, 1200, 1700, 2200, and 2500 m.a.s.l. (Fig. 1). In total, we caught 106 individuals (50 males and 56 females): 7 in 2010, 28 in 2011, 65 in 2012 and 6 in 2013. Because lizards were part of a long-term study, we marked individuals by toe clipping, a marking method frequently

used in lizards, and that have limited impact on their fitness (Perry, Wallace, Perry, Curzer, & Muhlberger, 2011). We conserved toe samples in ethanol and used them for age class determination using phalanx skeletochronology (more details below). We collected a portion of the terminal region of lizards' tail (~ 1 cm) in the field and immersed it in an Eppendorf tube filled with 1.5 mL of absolute ethanol for genetic analyses. Lizards regenerate lost tails, so the small portion we sampled should have had no effects on lizard fitness or welfare. We took special care to disinfect the wounds caused by both toe clipping and tail sampling with chlorohexidine closing the wounds with a tissue adhesive glue (Dermabond®).

We measured lizard body mass with a digital balance (Model Radwag WTB200; to the nearest 0.01 g) and SVL with a metal ruler (to the nearest 1 mm). We estimated the body condition index (BCI) as the residuals of the regressing log mass on log SVL. This widely used index represents the relative energy reserves of an animal (Schulte-Hostedde, Zinner, Milar, & Hickling, 2005). We also recorded whether the tail was intact or regenerated. Males were distinguished from females mainly because they have more femoral pores in their hind limbs (Iraeta, Monasterio, Salvador, & Diaz, 2011) and an orange spot in the corners of their mouths (Carretero, 2002). Gravid females, recognized by palpation of developing eggs inside the trunk, were translated to a lab and placed in individual terrariums (100×20×40 cm) with a heat cable at one end of the cage, switched on three hours a day (11 h-14 h) to allow thermoregulation, indirect access to sun light, and water (in form of aqueous nutritious gel) and food (*Tenebrio molitor* larvae) *ad libitum*. Substrate was bare soil from the study area. We maintained eggs laid in terrariums until hatching. Then, we took a portion of tail of hatchlings for genetic analyses (see below). In order to avoid pseudoreplication, only one neonate per litter (N = 37) was used in the analyses. Females and their neonates were released at the point the female was caught. No lizard died or suffered permanent pain during the study.

175

176 *Telomere length measurement*

177 Once in the laboratory, we stored tail samples at -20 °C until assayed. We extracted
178 DNA from epidermis using a high-salt DNA extraction protocol. Since storage conditions,
179 extraction method, or tissue type can affect telomere length measures (Nussey et al., 2014) we
180 used the same conditions for all samples to avoid confounding factors.

181 We quantified relative telomere length through quantitative polymerase chain
182 reactions (qPCRs), which is one of the most widely used method for estimating telomere
183 length (Nussey et al., 2014). We compared the cycle threshold (C_t) of telomeric sequences
184 with the C_t of a control sequence that is autosomal and non-variable in copy number
185 (Cawthon, 2002, Nussey et al., 2014). As a reference sequence, we amplified GAPDH
186 sequences using 5'-AACCAGCCAAGTACGATGACAT-3' (GAPDH-F) and 5'-
187 CCATCAGCAGCAGCCTTCA-3' (GAPDH-R) as forward and reverse primers, respectively.
188 For telomere sequences, we used
189 5'CGGTTTGTGGTTTGGGTTTGGGTTTGGGTTTGGGTT-3' (Tel1b) and 5'-
190 GGCTTGCCTTACCCTTACCCTTACCCTTACCCTTACCCT-3' (Tel2b) as forward and
191 reverse primers, respectively. Conditions of qPCR for GAPDH fragment consisted of 10 min
192 at 95 °C and 40 cycles of 10 sec at 95 °C, 20 secs at 58 °C, and 1 min at 72 °C, and for
193 telomere fragment of 10 min at 95 °C, and 10 secs at 95 °C, 20 secs at 58 °C, and 1 min at 72
194 °C. We conducted qPCR assays for each gene in separate plates on a LightCycler 480
195 (Roche) and ran a melting curve from 65 to 95 °C, as a final step in each qPCR to check for
196 specific amplicons. For each sample, we added 20 ng of genomic DNA and used both set of
197 primers at a final concentration of 100nM in a 20 µL master mix containing 10 µL of Brilliant
198 SYBR Green (QPCR Master Mix, Roche). All samples were run in duplicate. Samples with
199 coefficient of variation higher than 5 % were measured again. We calculated qPCR-plates

efficiency by including five serial diluted standards in triplicate, obtained from a *golden standard sample* containing a pool of samples from all populations. We calculated the relative telomere length by applying the following formula (Pfaffl, 2001): $[(E_{\text{telomere}})^{\Delta\text{Ct telomere (control-sample)}}] / [(E_{\text{GAPDH}})^{\Delta\text{Ct GAPDH (control-sample)}}]$; where E_{telomere} and E_{GAPDH} are the qPCR efficiency of telomere and GAPDH fragment, respectively; $\Delta\text{Ct telomere (control-sample)}$ and $\Delta\text{Ct GAPDH (control-sample)}$ are the deviation of standard – telomere or GAPDH sequences for each sample, respectively. Efficiencies of qPCR were 1.99 ± 0.02 and 1.93 ± 0.02 for GAPDH and telomere fragments, respectively.

Estimation of age class with skeletochronology

We estimated individual age class by phalanx skeletochronology (Comas, Reguera, Zamora-Camacho, Salvadó, & Moreno-Rueda, 2016), one of the most accurate techniques to estimate age in many vertebrates, including reptiles (Zhao, Klaassen, Lisovski, & Klaassen, 2019). Vertebrate ectotherms show indeterminate growth, and consequently present a cyclic growth pattern in hard body structures such as bones, corresponding to alternate periods of growth and resting. This pattern is particularly marked in temperate climates, where age can be fairly estimated by counting annual growth rings in the phalanges (Comas et al. 2016). Growth rings are called lines of arrested growth (LAGs). Toes sampled were decalcified in 3% nitric acid for 3 h and 30 min. Cross-sections (10 μm) were prepared using a freezing microtome (CM1850 Leica), stained with Harris hematoxylin for 20 min and dehydrated through an alcohol chain (more details in Comas et al. 2016). Next, cross-sections were fixed with DPX (mounting medium for histology), mounted on slides, and examined for the presence of LAGs using a light microscope (Leitz Dialux20) at magnifications from 50 to 125x. We took several photographs (with a ProgresC3 camera, at the University of Barcelona UB) of various representative cross-sections, discarding those photographs in which cuts

were unsuitable for observing LAGs. The number of LAGs detected in the periosteal bone was independently and blindly counted three times by a single observer (MC) on three independent occasions.

Statistical analysis

We confirmed that the residuals of the models met parametric assumptions (Quinn & Keough 2002). In order not to violate those assumptions, we log-transformed relative telomere length, body mass, and body condition data. We examined the presence of outliers through a Cleveland plot (Zuur, Ieno, & Elphick, 2010), which revealed that an individual had an extremely abnormal low value of relative telomere length, so we decided to omit this data from all the analyses. We performed a linear model to check for differences in telomere length according to the year of capture in order to evaluate possible cohort effects. Since we sampled lizards with intact tail (n = 44) and regenerated tail (n = 58), and tail regeneration could affect the length of telomeres in tail tissue (Anchelin, Murcia, Alcaraz-Perez, Garcia-Navarro, & Cayuela, 2011; Tan et al. 2012; Alibardi, 2016), we tested whether there were differences in relative telomere length between lizards with intact or regenerated tail through linear models. We also performed linear models to examine the effect of sex, age class, and elevation on the relative telomere length. We examined the relationship between relative telomere length and body mass or length, and also between telomere length and age class Pearson correlations. We checked for the independent effect of elevation and age class on relative telomere length by conducting a linear model, in which the effect of sex nor autotomy were not significant and we discarded this variable from the full model. Additionally, we conducted a model selection attending to Akaike information criterion (AIC; Akaike, 1973), and following the recommendations of Burnham & Anderson (2002). The full model contained all measured variables, i.e. elevation, body condition, age class, SVL, sex, tail

autotomy, and year of capture. We generated 128 models and calculated the model average of the top 2-AIC models. All statistical analyses were conducted in Statistica software (version 8.0).

Results

Lizards did not show sexual dimorphism in body mass ($F_{1, 102} = 0.11$, $P = 0.74$) and sexes did not differ in the distribution of age classes ($F_{1, 103} = 1.70$, $P = 0.20$). The frequency of male and female lizards did not differ across lizard populations ($\chi^2 = 7.35$, $P = 0.69$). Relative telomere length did not differ between sexes ($F_{1, 103} = 0.30$, $P = 0.59$; Figure 2A) and neither with the year of capture ($F_{3, 101} = 0.45$, $P = 0.72$). The frequency of lizards with autotomized tails did not vary among lizard populations ($\chi^2_5 = 1.36$, $P = 0.93$), and tail autotomy did not affect lizard telomere length ($F_{1, 100} = 0.00$, $P = 0.99$; Figure 2B).

Relative telomere length increased with age class, at least until the fourth year ($F_{5, 136} = 3.21$, $P = 0.009$, Fig. 3). In individuals with five years, telomeres tended to shorten, but we only sampled two lizards with this age class. Spearman correlations between telomere length and age showed similar results ($r = 0.24$, $P = 0.003$, including neonates, $N = 147$; $r = 0.19$, $P = 0.047$, without neonates, $N = 105$). Age correlated positively with body mass ($r = 0.57$, $P < 0.001$), which is common in organisms with indeterminate growth like lizards. Likewise, larger individuals had longer telomeres ($r = 0.26$, $P = 0.007$; Fig. 4), which confirms that telomeres did not shorten by cell replication, but elongated, in those larger and older individuals. Relative telomere length tended to increase with body condition ($r = 0.18$, $P = 0.067$). This relationship became significant ($r = 0.20$, $P = 0.043$) when a possible outlier –an individual with very high body condition, indicated by the Cleveland plot– was removed (Fig. 5A). Body condition increased with elevation ($F_{5, 98} = 3.03$, $P = 0.014$; Fig. 5B).

Lizard telomere length among lizard populations inhabiting across an elevational gradient, but following a non-linear pattern ($F_{5,136} = 2.52$; $P = 0.03$ for all individuals, and $F_{5,99} = 2.07$; $P = 0.070$ when excluding neonates; Fig. 6). Individuals at 300 and at 2200 m.a.s.l. had the longest telomeres. However, average age varied with elevation in a similar way ($F_{5,131} = 5.44$; $P < 0.001$; Figure S1). When we tested the combined effect of age and elevation on telomere length, the effect of age remained significant ($F_{5,131} = 2.32$; $P = 0.047$), but the effect of elevation was no longer significant ($F_{5,131} = 1.67$; $P = 0.15$). Model selection showed that body length (SVL) had the highest explanatory power to understand variation in telomere length in our study system, which is presumably explainable by the positive relationship between telomere length and age.

Neonate telomere length, an indicator of the baseline telomere length at birth, varied among populations ($F_{5,31} = 2.91$; $P = 0.03$), but with no clear pattern; lizard neonates showed the longest telomeres at 2200 m.a.s.l., but the shortest at 1700 m.a.s.l. (Fig. S2).

Discussion

Life-history trade-offs and environmental conditions can shape ageing across taxa (Wilbourn et al., 2018; Eastwood et al., 2019). Here, we show that, in the Algerian sand lizard, telomeres elongated with age until their fourth year. Additionally, larger lizards had longer telomeres. Intriguingly, although lizard populations across this substantial elevational gradient differed in their telomere length, differences were not linear and the variation in telomere length mirrored the variation in the distribution of age classes across the elevational gradient. On the other hand, telomere elongation was sex-independent, unlike in adults of other sand lizard species (*Lacerta agilis*, Olsson et al., 2011). Sex differences in telomere length may result from sex differences in growth rate, body size, and/or age (Olsson et al., 2018). However, in our study system, lizards did not show sexual dimorphism in size or age

structure. Autotomy did not affect telomere length despite the fact that differences in the regulation of telomere length may be driven by evolutionary pressures such as predation (Olsson et al., 2010), and also by enhanced cell replication during tissue regeneration. Moreover, no cohort effect was detected, as telomere length did not differ with year of sampling, which validates our cross-sectional study.

Telomere elongation observed in lizards across their first four years of life agrees with previous studies in snakes and lizards (Ujvari & Madsen, 2009 and Ujvari et al., 2017, respectively). At the fifth year, telomeres tended to shorten, although this result should be interpreted carefully because we only collected two five-year-old individuals. Telomere length showed a positive relationship both with age and body size, suggesting that cell replication does not shorten telomeres by itself. This finding adds to previous studies showing that ectotherms, unlike endotherms, often show telomere elongation along their lifetime (Olsson et al., 2018). Such contrasting pattern of telomere dynamics in ectotherms may be related to a higher telomerase expression after birth in somatic cells in ectotherms than in endotherms (Gomes, Shay, & Wright, 2010). Hence, telomerase may be relevant for buffering downstream effects of ROS in organisms with indeterminate growth such as lizards (Jones et al. 2014). However, telomerase expression may not be enough to protect from telomere shortening in ectothermic vertebrates. For instance, telomerase is expressed in tissues of adult medaka fish (Klapper et al. 1998) but telomeres shorten with age (Hatakeyama et al., 2008). Furthermore, the maintenance of telomerase expression in species with indeterminate growth can imply a trade-off suggested by a higher cancer occurrence in ectotherms (Gomes et al., 2010; Olsson et al. 2018), however, the knowledge about cancer in wildlife is still meagre.

Mountains cover *circa* a quarter of the Earth's surface (Körner, 2007). Elevational gradients are characterised by deep changes both in biotic and physical conditions, such as competitor and/or predators' abundance, temperature, or ultraviolet radiation (Barry, 2008).

Previous research demonstrated a broad number of physiological adaptations to divergent habitats across altitude (Bozinovic, Calosi, & Spicer, 2011; Keller, Alexander, Holderegger, & Edwards, 2013; Boyle, Sandercock, & Martin, 2016). Such adjustments often imply elevational variation in energy expenditure devoted to reproduction and somatic maintenance, then affecting telomere dynamics (Stier et al., 2016). Also, variation in temperature can involve deep physiological shifts in ectotherms across elevations since their body temperature greatly depends on environmental heat (Angilleta, 2009; Gunderson & Stillman, 2015). In our study, we expected to find shorter telomeres in lizard populations at higher elevation, as we know that higher-altitude lizards undergo reduced activity time and oxidative damage (Zamora-Camacho et al., 2013; Reguera et al., 2014, 2015). However, we found a non-linear variation in telomere length with elevation. The most plausible explanation is that telomere length across elevation mirrored the altitudinal distribution of lizard age. Contrary to our results, Dupoué et al., (2017) found that populations of the common lizard (*Zootoca vivipara*) inhabiting at low elevations have shorter telomeres and higher extinction risk. In our study system, lowland populations also suffer poor habitat quality, such as low thermal quality (risk of overheating, Zamora-Camacho, Reguera, & Moreno-Rueda, 2016), high ectoparasitism (Álvarez et al., 2018), low food availability (Moreno-Rueda et al., 2018), high oxidative damage (Reguera et al., 2014, 2015), and even high risk of wildfire (Moreno-Rueda, Melero, Reguera, Zamora-Camacho, & Comas, 2019). Additionally, at low elevations, lizards increase their activity time while hibernation time decreases (Zamora-Camacho, Reguera, Moreno-Rueda, & Pleguezuelos, 2013). In spite of all this, lizard populations at lowland did not have shorter telomeres than populations at high elevations.

Lizard body condition, temperature, and telomerase expression might have shaped telomere length of lizards inhabiting at different elevations. In this study, body condition of lizards was higher in populations at higher elevation, and correlated positively with telomere

length. Telomere length is positively correlated to body condition in other reptiles (*Thamnophis sirtalis*; Rollings et al., 2017). In addition, it is likely a temperature-mediated regulation of telomerase expression, thus at low elevation telomerase might show a higher expression, then compensating for telomere erosion (Olsson et al. 2018). At the highest elevations (mainly at 2200 m.a.s.l.), the reduction in metabolic rate due to cold conditions may have favoured a reduction in the rate of telomere erosion due to a reduced production of ROS, involving an adaptive downregulation of telomerase. Indeed, increases in lifespan are often orchestrated by reductions in metabolic rate (Speakman, 2005), as for example suggested by the straightforward influence of latitude on lifespan of *Rana temporaria* frogs across the Swedish latitudinal gradient (Hjernquist et al., 2012). Furthermore, the variation in the pace-of-life as a consequence of facing particular environmental conditions is also known to alter telomeres, then resulting in complex or unexpected patterns (Giraudeau, Algelier, & Sepp, 2019). For example, shorter telomeres are associated with higher survival in migratory Atlantic salmon (McLennan et al. 2017), thus benefits of experiencing intense telomere erosion can be higher than costs of responding poorly to certain scenarios, such as during migration. Likewise, amphibian larvae surviving predators, which have larger bodies and larger fat reserves, experience telomere shortening as a consequence of growing faster due to relaxed intraspecific competition (Burraco et al. 2017a). In our system, other factors like diseases or intraspecific interactions might have also modulated ageing in lizards. A cross-fostering approach would help to fully clarify the evolutionary impact of both environment and life-history traits on telomeres of this lizard metapopulation.

Conclusions

Our results show that telomeres elongate throughout the first four years of lizards' lifetime, a process that stress the role of telomerase in maintaining ectothermic telomeres, and, likely, in

extending lifespan in organisms with indeterminate growth. Habitat features and repair mechanisms at different habitats may be relevant for understanding telomere dynamics in ectothermic vertebrates. This study also shows that telomere length can follow a complex trajectory across habitats occupied by an ectothermic vertebrate, as across a substantial altitudinal gradient. Our results emphasize the relevance of understanding species' life histories (e.g. age and body condition) and habitat characteristics for disentangling the causes and consequences of lifespan trajectory.

Authors' contribution

GMR, MC and PB conceived the idea; SR and FJZC performed the sampling; MC and SR carried out the histological labwork and data analysis of skeletochronology; PB carried out the telomere length analysis; GMR and PB performed the statistical analyses; PB wrote the manuscript with inputs from GMR, MC, SR and FJZC.

Acknowledgements

PB was supported by fellowship F.P.U.-AP2010-5373, and by the Carl Tryggers Foundation project CT 16:344. FJZC (F.P.U.-AP2009-3505) and SR (F.P.U.-AP2009-1325) also were supported by respective fellowships. MC was supported by a Severo Ochoa contract (SVP-2014-068620). This study was economically supported by the Ministerio de Ciencia e Innovación (project CGL2009-13185). The study complies with the current laws of Spain, and were performed in accordance with the Junta de Andalucía and Sierra Nevada National Park research permits (references GMN/GyB/JMIF and ENSN/JSG/JEGT/MCF). We are grateful to Concepción Hernández (Centre of Scientific Instrumentation of the University of Granada) for her help with the freezing microtome, Humbert Salvadó (Universitat of Barcelona) for let us to use his microscopy for this study, and Francisco Miranda

(Ecophysiology Laboratory of Doñana Biological Station) for assistance with the telomere assays. We also thank the personnel from the Espacio Natural de Sierra Nevada for their constant support.

Data accessibility

Data will be accessible at FigShare upon manuscript acceptance.

References

- Akaike, H. (1973). Maximum likelihood identification of Gaussian autoregressive moving average models. *Biometrika*, 60(2), 255-265.
- Alder, J. K., Hanumanthu, V. S., Strong, M. A., DeZern, A. E., Stanley, S. E., Takemoto, C. M., ... & Brodsky, R. A. (2018). Diagnostic utility of telomere length testing in a hospital-based setting. *Proceedings of the National Academy of Sciences*, 115(10), E2358-E2365. <https://doi.org/10.1073/pnas.1720427115>
- Alibardi, L. (2016) Immunocalization of telomerase in cells of lizard tail after amputation suggests cell activation for tail regeneration. *Tissue Cell*, 48, 63–71. <https://doi.org/10.1016/j.tice.2015.10.004>
- Allsopp, R. C., Chang, E., Kashefi-Aazam, M., Rogaev, E. I., Piatyszek, M. A., Shay, J. W., & Harley, C. B. (1995). Telomere shortening is associated with cell division in vitro and in vivo. *Experimental Cell Research*, 220(1), 194-200. <https://doi.org/10.1006/excr.1995.1306>
- Álvarez-Ruiz, L., Megía-Palma, R., Reguera, S., Ruiz, S., Zamora-Camacho, F. J., Figuerola, J., & Moreno-Rueda, G. (2018). Opposed elevational variation in prevalence and intensity of endoparasites and their vectors in a lizard. *Current Zoology*, 64(2), 197-204. <https://doi.org/10.1093/cz/zoy002>
- Anchelin, M., Murcia, L., Alcaraz-Perez, F., Garcia-Navarro, E. M., Cayuela, M. L. (2011) Behaviour of telomere and telomerase during aging and regeneration in zebrafish. *PLoS One*, 6, e16955. <https://doi.org/10.1371/journal.pone.0016955>
- Angelier, F., Costantini, D., Blevin, P., & Chastel, O. (2018). Do glucocorticoids mediate the link between environmental conditions and telomere dynamics in wild vertebrates? A

- review. *General and Comparative Endocrinology*, 256, 99-111.
- <https://doi.org/10.1016/j.ygcen.2017.07.007>
- Angelier, F., Weimerskirch, H., Barbraud, C., & Chastel, O. (2019). Is telomere length a molecular marker of individual quality? Insights from a long-lived bird. *Functional Ecology*. <https://doi.org/10.1111/1365-2435.13307>
- Angilletta Jr, M. J. (2009). *Thermal adaptation: a theoretical and empirical synthesis*. Oxford University Press.
- Aubert, G., & Lansdorp, P. M. (2008). Telomeres and aging. *Physiological Reviews*, 88(2), 557-579. <https://doi.org/10.1152/physrev.00026.2007>
- Barrett, E. L., Burke, T. A., Hammers, M., Komdeur, J., & Richardson, D. S. (2013). Telomere length and dynamics predict mortality in a wild longitudinal study. *Molecular Ecology*, 22(1), 249-259. <https://doi.org/10.1111/mec.12110>
- Barry, R. G. (2008). Mountain weather and climate, – Cambridge University Press. *New York*.
- Bateson, M. (2016). Cumulative stress in research animals: telomere attrition as a biomarker in a welfare context? *BioEssays*, 38(2), 201-212. <https://doi.org/10.1002/bies.201500127>
- Bousman, S., Schneider, G., & Shampay, J. (2003). Telomerase activity is widespread in adult somatic tissues of *Xenopus*. *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution*, 295(1), 82-86. <https://doi.org/10.1002/jez.b.7>
- Boyle, W. A.; Sandercock, B. K.; Martin, K. (2016). Patterns and drivers of intraspecific variation in avian life history along elevational gradients: a meta-analysis. *Biological Reviews* 91(2), 469–482. doi: 10.1111/brv.12180. <https://doi.org/10.1111/brv.12180>
- Bozinovic, F., Calosi, P., & Spicer, J.I. (2011). Physiological correlates of geographic range in animals. *Annual Review of Ecology, Evolution, and Systematics*, 42,155-179. <https://doi.org/10.1146/annurev-ecolsys-102710-145055>
- Burnham, K. P., & Anderson, D. R. (2004). Multimodel inference: understanding AIC and BIC in model selection. *Sociological methods & research*, 33(2), 261-304.
- Burraco, P., Díaz-Paniagua, C., & Gomez-Mestre, I. (2017a). Different effects of accelerated development and enhanced growth on oxidative stress and telomere shortening in amphibian larvae. *Scientific Reports*, 7(1), 7494. <https://doi.org/10.1038/s41598-017-07201-z>
- Burraco, P., Valdés, A. E., Johansson, F., & Gomez-Mestre, I. (2017b). Physiological mechanisms of adaptive developmental plasticity in *Rana temporaria* island

populations. *BMC Evolutionary Biology*, 17(1), 164. <https://doi.org/10.1186/s12862-017-1004-1>

Carretero, M. A. (2002). Sources of colour pattern variation in Mediterranean *Psammmodromus algirus*. *Netherlands Journal of Zoology*, 52(1), 43-60. <https://doi.org/10.1163/156854202760405177>

Cawthon, R. M. (2002). Telomere measurement by quantitative PCR. *Nucleic Acids Research*, 30(10), e47-e47. <https://doi.org/10.1093/nar/30.10.e47>

Comas, M., Reguera, S., Zamora-Camacho, F. J., Salvadó, H., & Moreno-Rueda, G. (2016). Comparison of the effectiveness of phalanges vs. humeri and femurs to estimate lizard age with skeletochronology. *Animal Biodiversity and Conservation*, 39(2), 237-240. <https://doi.org/10.32800/abc.2016.39.0237>

Dupoué, A., Rutschmann, A., Le Galliard, J. F., Clobert, J., Angelier, F., Marciau, C., ... & Meylan, S. (2017). Shorter telomeres precede population extinction in wild lizards. *Scientific Reports*, 7(1), 16976. <https://doi.org/10.1038/s41598-017-17323-z>

Eastwood, J. R., Hall, M. L., Teunissen, N., Kingma, S. A., Hidalgo Aranzamendi, N., Fan, M., ... & Peters, A. (2019). Early-life telomere length predicts lifespan and lifetime reproductive success in a wild bird. *Molecular Ecology*, 28, 1127-1137. <https://doi.org/10.1111/mec.15002>

Giraudeau, M., Angelier, F., & Sepp, T. (2019). Do telomeres influence pace-of-life-strategies in response to environmental conditions over a lifetime and between generations? *BioEssays*, 41, 1800162. <https://doi.org/10.1002/bies.201800162>

Gomes, N. M., Shay, J. W., & Wright, W. E. (2010). Telomere biology in Metazoa. *FEBS Letters*, 584(17), 3741-3751. <https://doi.org/10.1016/j.febslet.2010.07.031>

Gunderson, A. R., & Stillman, J. H. (2015). Plasticity in thermal tolerance has limited potential to buffer ectotherms from global warming. *Proceedings of the Royal Society B: Biological Sciences*, 282(1808), 20150401. <https://doi.org/10.1098/rspb.2015.0401>

Hatakeyama, H., Nakamura, K. I., Izumiyama-Shimomura, N., Ishii, A., Tsuchida, S., Takubo, K., & Ishikawa, N. (2008). The teleost *Oryzias latipes* shows telomere shortening with age despite considerable telomerase activity throughout life. *Mechanisms of Ageing and Development*, 129(9), 550-557. <https://doi.org/10.1016/j.mad.2008.05.006>

Hausmann, M. F., Winkler, D. W., O'Reilly, K. M., Huntington, C. E., Nisbet, I. C., & Vleck, C. M. (2003). Telomeres shorten more slowly in long-lived birds and mammals

- than in short-lived ones. *Proceedings of the Royal Society of London B: Biological Sciences*, 270(1522), 1387-1392. <https://doi.org/10.1098/rspb.2003.2385>
- Haussmann, M. F., & Mauck, R. A. (2007). Telomeres and longevity: testing an evolutionary hypothesis. *Molecular Biology and Evolution*, 25(1), 220-228. <https://doi.org/10.1093/molbev/msm244>
- Haussmann, M. F., & Marchetto, N. M. (2010). Telomeres: linking stress and survival, ecology and evolution. *Current Zoology*, 56(6), 714-727.
- Heidinger, B. J., Blount, J. D., Boner, W., Griffiths, K., Metcalfe, N. B., & Monaghan, P. (2012). Telomere length in early life predicts lifespan. *Proceedings of the National Academy of Sciences*, 109(5), 1743-1748. <https://doi.org/10.1073/pnas.1113306109>
- Hjernquist, M. B., Söderman, F., Jönsson, K. I., Herczeg, G., Laurila, A., & Merilä, J. (2012). Seasonality determines patterns of growth and age structure over a geographic gradient in an ectothermic vertebrate. *Oecologia*, 170(3), 641-649. <https://doi.org/10.1007/s00442-012-2338-4>
- Hoelzl, F., Cornils, J. S., Smith, S., Moodley, Y., & Ruf, T. (2016). Telomere dynamics in free-living edible dormice (*Glis glis*): the impact of hibernation and food supply. *Journal of Experimental Biology*, 219(16), 2469-2474. <https://doi.org/10.1242/jeb.14087>
- Horn, T., Robertson, B. C., Will, M., Eason, D. K., Elliott, G. P., & Gemmell, N. J. (2011). Inheritance of telomere length in a bird. *PLoS One*, 6(2), e17199. <https://doi.org/10.1371/journal.pone.0017199>
- Ingles, E. D., & Deakin, J. E. (2016). Telomeres, species differences, and unusual telomeres in vertebrates: presenting challenges and opportunities to understanding telomere dynamics. *AIMS Genetics*, 3, 1-24. <https://doi.org/10.3934/genet.2016.1.1>
- Iraeta, P., Monasterio, C., Salvador, A., & Diaz, J. A. (2011). Sexual dimorphism and interpopulation differences in lizard hind limb length: locomotor performance or chemical signalling? *Biological Journal of the Linnean Society*, 104(2), 318-329. <https://doi.org/10.1111/j.1095-8312.2011.01739.x>
- Isaksson, C. (2015). Urbanization, oxidative stress and inflammation: a question of evolving, acclimatizing or coping with urban environmental stress. *Functional Ecology*, 29(7), 913-923. <https://doi.org/10.1111/1365-2435.12477>

- 528 Jones, O. R., Scheuerlein, A., Salguero-Gómez, R., Camarda, C. G., Schaible, R., Casper, B.
529 B., ... & Quintana-Ascencio, P. F. (2014). Diversity of ageing across the tree of
530 life. *Nature*, 505(7482), 169. <https://doi.org/10.1038/nature12789>
- 531 Keller, I., Alexander, J. M., Holderegger, R., & Edwards, P. J. (2013). Widespread
532 phenotypic and genetic divergence along altitudinal gradients in animals. *Journal of*
533 *Evolutionary Biology* 26, 2527–2543. <https://doi.org/10.1111/jeb.12255>
534
- 535 Kirby, R., Johnson, H. E., Alldredge, M. W., & Pauli, J. N. (2019). The cascading effects of
536 human food on hibernation and cellular aging in free-ranging black bears. *Scientific*
537 *Reports*, 9(1), 2197. <https://doi.org/10.1038/s41598-019-38937>
- 538 Klapper, W., Heidorn, K., Kühne, K., Parwaresch, R., & Krupp, G. (1998). Telomerase
539 activity in ‘immortal’ fish. *FEBS Letters*, 434(3), 409-412.
540 [https://doi.org/10.1016/S0014-5793\(98\)01020-5](https://doi.org/10.1016/S0014-5793(98)01020-5)
- 541 Körner, C., Paulsen, J., & Spehn, E. M. (2011). A definition of mountains and their
542 bioclimatic belts for global comparisons of biodiversity data. *Alpine Botany*, 121(2),
543 73-78. <https://doi.org/10.1007/s00035-011-0094-4>
- 544 Luceri, C., Bigagli, E., Femia, A. P., Caderni, G., Giovannelli, L., & Lodovici, M. (2018).
545 Aging related changes in circulating reactive oxygen species (ROS) and protein
546 carbonyls are indicative of liver oxidative injury. *Toxicology Reports*, 5, 141-145.
547 <https://doi.org/10.1016/j.toxrep.2017.12.017>
- 548 Marasco, V., Stier, A., Boner, W., Griffiths, K., Heidinger, B., & Monaghan, P. (2017).
549 Environmental conditions can modulate the links among oxidative stress, age, and
550 longevity. *Mechanisms of Ageing and Development*, 164, 100-107.
551 <https://doi.org/10.1016/j.mad.2017.04.012>
- 552 McLennan, D., Armstrong, J. D., Stewart, D. C., Mckelvey, S., Boner, W., Monaghan, P., &
553 Metcalfe, N. B. (2016). Interactions between parental traits, environmental harshness
554 and growth rate in determining telomere length in wild juvenile salmon. *Molecular*
555 *Ecology*, 25(21), 5425-5438. <https://doi.org/10.1111/mec.13857>
- 556 McLennan, D., Armstrong, J. D., Stewart, D. C., Mckelvey, S., Boner, W., Monaghan, P., &
557 Metcalfe, N. B. (2017). Shorter juvenile telomere length is associated with higher
558 survival to spawning in migratory Atlantic salmon. *Functional ecology*, 31(11), 2070-
559 2079. <https://doi.org/10.1111/1365-2435.12939>

- 560 Monaghan, P. (2007). Early growth conditions, phenotypic development and environmental
561 change. *Philosophical Transactions of the Royal Society B: Biological*
562 *Sciences*, 363(1497), 1635-1645. <https://doi.org/10.1098/rstb.2007.0011>
- 563 Monaghan, P. (2014). Organismal stress, telomeres and life histories. *Journal of Experimental*
564 *Biology*, 217(1), 57-66. <https://doi.org/10.1242/jeb.090043>
- 565 Monaghan, P., Eisenberg, D. T., Harrington, L., & Nussey, D. (2018). Understanding
566 diversity in telomere dynamics. *Philosophical Transactions of the Royal Society B:*
567 *Biological Sciences*, 373, 20160435. <https://doi.org/10.1098/rstb.2016.0435>
- 568 Moreno-Rueda, G., Melero, E., Reguera, S., Zamora-Camacho, F. J., & Comas, M. (2019).
569 Short-term impact of a small wildfire on the lizard *Psammodromus algirus* (Linnaeus,
570 1758): a before-after-control-impact study. *Herpetozoa*, 31, 173-182.
- 571 Nussey, D. H., Baird, D., Barrett, E., Boner, W., Fairlie, J., Gemmell, N., ... & Turbill, C.
572 (2014). Measuring telomere length and telomere dynamics in evolutionary biology and
573 ecology. *Methods in Ecology and Evolution*, 5(4), 299-310.
574 <https://doi.org/10.1111/2041-210X.12161>
- 575 Olsson, M., Pauliny, A., Wapstra, E., & Blomqvist, D. (2010). Proximate determinants of
576 telomere length in sand lizards (*Lacerta agilis*). *Biology Letters*, 6(5), 651-653.
577 <https://doi.org/10.1098/rsbl.2010.0126>
- 578 Olsson, M., Pauliny, A., Wapstra, E., Uller, T., Schwartz, T., Miller, E., & Blomqvist, D.
579 (2011): Sexual differences in telomere selection in the wild. *Molecular Ecology* 20,
580 2085–2099. <https://doi.org/10.1111/j.1365-294X.2011.05085.x>
- 581 Olsson, M., Wapstra, E., & Friesen, C. (2018). Ectothermic telomeres: it's time they came in
582 from the cold. *Philosophical Transactions of the Royal Society B: Biological*
583 *Sciences*, 373(1741), 20160449. <https://doi.org/10.1098/rstb.2016.0449>
- 584 O'sullivan, R. J., & Karlseder, J. (2010). Telomeres: protecting chromosomes against genome
585 instability. *Nature Reviews Molecular Cell biology*, 11(3), 171-181.
586 <https://doi.org/10.1038/nrm2848>
- 587 Perry, G., Wallace, M.C., Perry, D., Curzer, H., Muhlberger, P. (2011): Toe clipping of
588 amphibians and reptiles: Science, ethics, and the law. *Journal of Herpetology*, 45(4),
589 547–555. <https://doi.org/10.2307/41415330>
- 590 Pfaffl, M. W. (2001). A new mathematical model for relative quantification in real-time RT–
591 PCR. *Nucleic Acids Research*, 29(9), e45-e45. <https://doi.org/10.1093/nar/29.9.e45>

- Quinn, G. P., & Keough, M. J. (2002). *Experimental design and data analysis for biologists*. Cambridge University Press.
- Ratikainen, I. I., & Kokko, H. (2019). The coevolution of lifespan and reversible plasticity. *Nature Communications*, 10(1), 538. <https://doi.org/10.5061/dryad.m7b43mm>
- Reichert, S., Stier, A. (2017). Does oxidative stress shorten telomeres in vivo? A review. *Biology Letters*, 13, 20170463. <https://doi.org/10.1098/rsbl.2017.0463>
- Reguera, S., Zamora-Camacho, F. J., Trenzado, C. E., Sanz, A., & Moreno-Rueda, G. (2014). Oxidative stress decreases with elevation in the lizard *Psammodromus algirus*. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 172, 52-56. <https://doi.org/10.1016/j.cbpa.2014.02.018>
- Reguera, S., Zamora-Camacho, F. J., Melero, E., García-Mesa, S., Trenzado, C. E., Cabrerizo, M. J., ... & Moreno-Rueda, G. (2015). Ultraviolet radiation does not increase oxidative stress in the lizard *Psammodromus algirus* along an elevational gradient. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 183, 20-26. <https://doi.org/10.1016/j.cbpa.2014.12.015>
- Rollings, N., Uhrig, E. J., Krohmer, R. W., Wayne, H. L., Mason, R. T., Olsson, M., ... & Friesen, C. R. (2017). Age-related sex differences in body condition and telomere dynamics of red-sided garter snakes. *Proceedings of the Royal Society B: Biological Sciences*, 284(1852), 20162146. <https://doi.org/10.1098/rspb.2016.2146>
- Salvador, A. (2015). Lagartija colilarga—*Psammodromus algirus*. *Enciclopedia Virtual de los Vertebrados Espanoles. Madrid: Museo Nacional de Ciencias Naturales, MNCN-CSIC*.
- Schulte-Hostedde, A. I., Zinner, B., Millar, J. S., & Hickling, G. J. (2005). Restitution of mass–size residuals: validating body condition indices. *Ecology*, 86(1), 155-163. <https://doi.org/10.1890/04-0232>
- Selman, C., Blount, J. D., Nussey, D. H., & Speakman, J. R. (2012). Oxidative damage, ageing, and life-history evolution: where now? *Trends in Ecology & Evolution*, 27(10), 570-577. <https://doi.org/10.1016/j.tree.2012.06.006>
- Sih, A., Ferrari, M. C., & Harris, D. J. (2011). Evolution and behavioural responses to human-induced rapid environmental change. *Evolutionary Applications*, 4(2), 367-387. <https://doi.org/10.1111/j.1752-4571.2010.00166.x>
- Speakman, J. R. (2005). Body size, energy metabolism and lifespan. *Journal of Experimental Biology*, 208(9), 1717-1730. <https://doi.org/10.1242/jeb.01556>

- 626 Stier, A., Delestrade, A., Bize, P., Zahn, S., Criscuolo, F., & Massemin, S. (2016).
627 Investigating how telomere dynamics, growth and life history covary along an
628 elevation gradient in two passerine species. *Journal of Avian Biology*, 47(1), 134-140.
629 <https://doi.org/10.1111/jav.00714>
- 630 Tan, T. C., Rahman, R., Jaber-Hijazi, F., Felix, D. A., Chen, C., Louis, E. J., & Aboobaker, A.
631 (2012) Telomere maintenance and telomerase activity are differentially regulated in
632 asexual and sexual worms. *Proceedings of the National Academy of Sciences*, 109,
633 4209–4214. <https://doi.org/10.1073/pnas.1118885109>
- 634 Ujvari, B., & Madsen, T. (2009). Short telomeres in hatchling snakes: erythrocyte telomere
635 dynamics and longevity in tropical pythons. *PloS One*, 4(10), e7493.
636 <https://doi.org/10.1371/journal.pone.0007493>
- 637 Ujvari, B., Biro, P. A., Charters, J. E., Brown, G., Heasman, K., Beckmann, C., & Madsen, T.
638 (2017). Curvilinear telomere length dynamics in a squamate reptile. *Functional*
639 *Ecology*, 31(3), 753-759. <https://doi.org/10.1111/1365-2435.12764>
- 640 Vágási, C. I., Vincze, O., Pátra, L., Osváth, G., Péntes, J., Haussmann, M. F., ... & Pap, P.
641 L. (2019). Longevity and life history coevolve with oxidative stress in
642 birds. *Functional Ecology*, 33:152–161. <https://doi.org/10.1111/1365-2435.13228>
- 643 Wilbourn, R. V., Moatt, J. P., Froy, H., Walling, C. A., Nussey, D. H., & Boonekamp, J. J.
644 (2018). The relationship between telomere length and mortality risk in non-model
645 vertebrate systems: a meta-analysis. *Philosophical Transactions of the Royal Society*
646 *B: Biological Sciences*, 373(1741), 20160447. <https://doi.org/10.1098/rstb.2016.0447>
- 647 Young, A. J. (2018). The role of telomeres in the mechanisms and evolution of life-history
648 trade-offs and ageing. *Philosophical Transactions of the Royal Society B: Biological*
649 *Sciences*, 37, 20160452. <https://doi.org/10.1098/rstb.2016.0452>
- 650 Zamora-Camacho, F. J., Reguera, S., Moreno-Rueda, G., & Pleguezuelos, J. M. (2013).
651 Patterns of seasonal activity in a Mediterranean lizard along a 2200 m altitudinal
652 gradient. *Journal of Thermal Biology*, 38(2), 64-69.
653 <https://doi.org/10.1016/j.jtherbio.2012.11.002>
- 654 Zamora-Camacho, F. J., Reguera, S., & Moreno-Rueda, G. (2016). Thermoregulation in the
655 lizard *Psammodromus algirus* along a 2200-m elevational gradient in Sierra Nevada
656 (Spain). *International Journal of Biometeorology*, 60(5), 687-697. <https://doi.org/10.1007/s00484-015-1063-1>
- 658 Zamudio, K. R., Bell, R. C., & Mason, N. A. (2016). Phenotypes in phylogeography: species'
659 traits, environmental variation, and vertebrate diversification. *Proceedings of the*

National Academy of Sciences, 113(29), 8041-8048.

<https://doi.org/10.1073/pnas.1602237113>

Zhao, M., Klaassen, C. A., Lisovski, S., & Klaassen, M. (2019). The adequacy of aging techniques in vertebrates for rapid estimation of population mortality rates from age distributions. *Ecology and Evolution*, 9(3), 1394-1402.

<https://doi.org/10.1002/ece3.4854>

Ziegler, D. V., Wiley, C. D., & Velarde, M. C. (2015). Mitochondrial effectors of cellular senescence: beyond the free radical theory of aging. *Aging Cell*, 14(1), 1-7.

<https://doi.org/10.1111/accel.12287>

Zuur, A. F., Ieno, E. N., & Elphick, C. S. (2010). A protocol for data exploration to avoid common statistical problems. *Methods in Ecology and Evolution*, 1(1), 3-14.

Figure legends

Figure 1. Sampling locations used in this study across an altitudinal mountain gradient (Sierra Nevada mountain system, SE Spain). Numbers from one to six correspond with each location, i.e. 300, 700, 1200, 1700, 2200, and 2500 m.a.s.l. respectively.

Figure 2. Variation in relative telomere length between sexes (A) or in response to autotomy (B) in individuals of the Algerian sand lizard (*Psammodromus algirus*). The red point shows the mean value at each age and the boxplot the interquartile range. The kernel density plot shows the probability density of data at different values.

Figure 3. Variation in relative telomere length across lifetime of in individuals of the Algerian sand lizard (*Psammodromus algirus*). The red point shows the mean value at each age and the boxplot the interquartile range. The kernel density plot shows the probability density of data at different values.

Figure 4. Regression between body mass and relative telomere length ($r = 0.26$, $P = 0.007$). Regression line shows the correlation between both parameters in all individuals of the Algerian sand lizard (*Psammodromus algirus*) sampled in this study, and indicates that telomeres did not shorten, but elongated with cell replication, as observed in larger (and older) individuals. The grey region indicates the 95% confidence intervals.

Figure 5. (A) Regression between body condition and relative telomere length ($r = 0.20$, $P = 0.043$). The asterisk indicates a possible outlier (B) Variation in lizards' body condition across altitude ($F_{5, 98} = 3.03$, $P = 0.014$) in individuals of the Algerian sand lizard (*Psammodromus algirus*). The red point shows the mean value at each elevation and the boxplot the

interquartile range. The kernel density plot shows the probability density of data at different values.

Figure 6. Variation in relative telomere length across altitude in individuals of the Algerian sand lizard (*Psammodromus algirus*). The red point shows the mean value at each age and the boxplot the interquartile range. The kernel density plot shows the probability density of data at different values.

723 **Table 1.** Summary of the studies describing the relationship between telomere length (TL) with age and/or other traits in reptiles. References are
724 included in the Supplementary Information S2.
725

reference	measured traits	relationship between TL and age	relationship between TL and trait(s)	type of study	technique	tissue	common name	species
Girondot & Garcia 1998	age	n.s. (embryos vs. adults)		field	TRF	blood	European freshwater turtle	<i>Emys orbicularis</i>
Scott et al. 2006	body size, sex		body length (-), sex (ns)	field	TRF	blood	American alligator	<i>Alligator mississippiensis</i>
Bronikowski 2008	age	(-) (1-13 y/o)		from the field, lab until hibernation	TRF	blood	western terrestrial garter snake	<i>Thamnophis elegans</i>
Hatase et al. 2008	age	n.s. (0-36 y/o)		from public aquarium	qPCR	blood, epidermis	loggerhead sea turtle	<i>Caretta caretta</i>
Ujvari & Madsen 2009	age, survival	(+) for both sexes if including hatchlings (0-20 y/o), the same in longitudinal study (N=8)	longer TL females than in males, hatchling sexes (ns), recaptured hatchlings (ns), recaptured old pythons (+)	field	TRF	blood	water python	<i>Liasis fuscus</i>
Xu et al. 2009	age, sex	(-) for both sexes (3-10 y/o)		field	TRF	blood	Chinese alligator	<i>Alligator sinensis</i>
Hatase et al. 2010	foraging behaviour		foraging behaviour (ns)	field	qPCR	epidermis	loggerhead turtle	<i>Caretta caretta</i>
Olsson et al. 2010	length, age, activity, ticks, tail	(+) for females, (-) trend for males (2-8 y/o)	females: length (ns), activity (ns), ticks (ns), tail regeneration (ns); males:	field	TRF	blood	sand lizard	<i>Lacerta agilis</i>

	regeneration		ticks (ns), badge size (ns), activity (+), length (-), tail regeneration (-)					
Olsson et al. 2011a	heritability, paternal age, offspring survival, malformations	(-) for sires, n.s. for sons (3-7 y/o)	capture probability of sires (+), offspring sex-ratio (+) TL of sons and paternal age at conception (-)	field	TRF	blood	sand lizard	<i>Lacerta agilis</i>
Olsson et al. 2011b	sex		longer TL in females than in males; females: lifespan (+), lifetime reproductive sucess (+), males: lifespan (ns), lifetime reproductive sucess (ns)	field	TRF	blood	sand lizard	<i>Lacerta agilis</i>
Ballen et al. 2012	maternal and offspring TL, body mass, superoxide		offspring TL with maternal TL (+), maternal reproductive investment (+), offspring mass (-), offspring superoxide (-)	from the field, hatching in lab	PNA Kit/FITC flow cytometry	blood	painted dragon	<i>Ctenophorus pictus</i>
Plot et al. 2012	sex, age, reproduction	n.s. (hatchlings vs. adults)	reproductive output (+), time to first breeding (+)	field	qPCR	blood	leatherback sea turtle	<i>Dermochelys coriacea</i>
Giraudeau et al. 2016	colour fading		colour fading (-)	field	qPCR	blood	painted dragon	<i>Ctenophorus pictus</i>
Rollings et al. 2017a	head color, bib presence		competition ability (-), bib presence (-)	field	qPCR	blood	painted dragon	<i>Ctenophorus pictus</i>
Dupoué et al. 2017	body size, sex, altitude, extinction risk, Tmin		body size (ns), sex (ns), extinction risk (-), altitude(+), Tmin (+)	field	TRF	blood	common lizard	<i>Zootoca vivipara</i>
Rollings et al. 2017b	age, sex	quadratic for males, n.s. for females (2-6 y/o)	shorter TL in males, body length (ns), growth (ns), body condition (+)	field	qPCR	blood	red-sided garter snake	<i>Thamnophis sirtalis parietalis</i>

Ujvari et al. 2017	age, survival	quadratic (1-8 y/o)	survival (ns)	field	qPCR	blood	frillneck lizard	<i>Chlamydosaurus kingii</i>
Pauliny et al. 2018	paternity probability		probability of siring offspring (+)	field	qPCR	blood	sand lizard	<i>Lacerta agilis</i>
Zhang et al. 2018	temperature		temperature (-)	laboratory	qPCR	heart	desert toad-headed agama	<i>Phrynocephalus przewalskii</i>
Rollings et al. 2019	sex, tissue		TL varies between sexes and among tissues	laboratory	qPCR	several	painted dragon	<i>Ctenophorus pictus</i>
Mclennan et al. 2019	reproductive mode		longer TL in viviparous mothers and offspring than in oviparous				common lizard	<i>Zootoca vivipara</i>
Burraco et al. 2019 (this study)	age, sex, altitude, autotomy, body mass, body condition	(+) for both sexes	longer TL at low and high altitudes, sex (ns), autotomy (ns), body mass (+), body condition (+)	field	qPCR	epidermis	Algerian sand lizard	<i>Psammodromus algerius</i>

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6