1 **Culture of salivary methanogens assisted by chemically produced hydrogen.**

2

3 Guindo CO^{1,2}, Terrer E^{2,3}, Chabrière E², Aboudharam G^{2,3}, Drancourt M^{2*}, Grine G^{1,2,3}.

4

5 1. IHU Méditerranée Infection, Marseille, France.

6 2. Aix-Marseille Univ., IRD, MEPHI, IHU Méditerranée Infection, Marseille, France.

7 3. Aix-Marseille-Univ., Faculty of Odontology, Marseille, France.

8

9

10 *Corresponding author: Prof. Michel Drancourt

11 IHU Méditerranée Infection, UMR MEPHI

12 19-21, Bd Jean Moulin 13005 Marseille, France. Tel: +33 (0)4 13 73 24 01, fax: + 33 (0) 13

13 73 24 02.

14 Email : michel.drancourt@univ-amu.fr.

15 **ABSTRACT**

16 Methanogen cultures require hydrogen produced by fermentative bacteria such as
17 *Bacteroides thetaiotaomicron* (biological method). We developed an alternative method for
18 hydrogen production using iron filings and acetic acid with the aim of cultivating
19 methanogens more efficiently and more quickly (chemical method). We developed this new
20 method with a reference strain of *Methanobrevibacter oralis*, compared the method to the
21 biological reference method with a reference strain of *Methanobrevibacter smithii* and finally
22 applied the method to 50 saliva samples. Methanogen colonies counted using ImageJ
23 software were identified using epifluorescence optical microscopy, real-time PCR and PCR
24 sequencing. For cultures containing the pure strains of *M. oralis* and *M. smithii*, colonies
25 appeared three days postinoculation with the chemical method versus nine days with the
26 biological method. The average number of *M. smithii* colonies was significantly higher with
27 the chemical method than with the biological method. There was no difference in the delay of
28 observation of the first colonies in the saliva samples between the two methods. However, the
29 average number of colonies was significantly higher when using the biological method than
30 when using the chemical method at six days and nine days postinoculation (Student's test, $p =$
31 0.005 and $p = 0.04$, respectively). The chemical method made it possible to isolate four
32 strains of *M. oralis* and three strains of *M. smithii* from the 50 saliva samples. Establishing
33 the chemical method will ease the routine isolation and culture of methanogens.

34

35

36 INTRODUCTION

37 Methanogenic archaea (referred to herein as methanogens) are acknowledged members of the
38 digestive tract microbiota, and they have been detected by PCR-based methods and cultured
39 from the oral cavity and the stools of apparently healthy individuals (1). More particularly,
40 *Methanobrevibacter oralis*, *Methanobrevibacter smithii* and *Methanobrevibacter massiliense*
41 have been isolated from the oral cavity, whereas *M. smithii*, *Methanosphaera stadtmanae*,
42 *Methanomassiliicoccus luminiensis*, *Methanobrevibacter arboriphilicus*, *M. oralis*, *Ca.*
43 *Methanomethylophilus alvus* and *Ca. Methanomassiliicoccus intestinalis* have been isolated
44 from stools (1). Accordingly, we previously showed that virtually all apparently healthy
45 individuals would carry methanogens in the digestive tract microbiota (2). This observation
46 indeed corroborated the pivotal role of methanogens, which detoxify hydrogen produced by
47 bacterial fermentations into methane (3, 4).
48 Moreover, methanogens are increasingly implicated in diseases; their presence or absence is
49 associated with dysbioses such as those observed in the gut microbiota in cases of chronic
50 constipation (5), obesity (6) and colonic diseases including ulcerative colitis, Crohn's disease
51 and colorectal cancer (7-9), in the vaginal microbiota in cases of vaginosis (10, 11), or their
52 presence in anaerobic pus abscesses such as brain abscesses (12, 13) and muscular abscesses
53 (14).
54 As is usual in clinical microbiology, isolation and culture of methanogens is the gold standard
55 to assess the detection of living methanogens in microbiota and in pathological clinical
56 specimens collected by puncture or biopsy (1). The routine application of methanogen
57 culturing is hampered by the fact that methanogens are strictly aero-intolerant microbes and
58 require hydrogen for culturing (15, 16). To facilitate the isolation and culture of methanogens
59 from a routine perspective, we previously designed a new process in which methanogens are
60 cultivated in the presence of hydrogen-producing *Bacteroides thetaiotaomicron* (17, 18).

61 Here, we tested the conditions to replace the biological production of hydrogen with a
62 chemical production of hydrogen and applied it to isolate methanogens from saliva as a
63 proof-of-concept.

64

65 MATERIALS AND METHODS

66 **Chemical production of hydrogen.** The immediate chemical production of hydrogen
67 resulting from the oxidation of iron by a weak acid has long been known (19). We used acetic
68 acid in our experiment because of its ability to oxidize iron, resulting in sufficient production
69 of hydrogen (20). We used two solutions for our experiments, namely, SAB culture broth
70 (21) and distilled water (Fresenius, Bad Homburg vor Hoehe, Germany) with pH values of
71 7.3 and pH values of 7, respectively. We used 5 mL of each solution in our experiments. The
72 experiments were performed in 10 Hungate tubes (Dominique Dutscher, Brumath, France)
73 for each solution. Iron filings (Amazon, Clichy, France) were used at increasing quantities:
74 0.5 g for tube n°1, 1 g for tube n°2, and a 0.25 g increase from tube n°3 to tube n°10. The
75 amount of acetic acid (VWR International, Pennsylvania, USA) used was 100 µL for tube n°1
76 to tube n°7 and 50 µL for tube n°8 to tube n°10. We first put the iron filings in the tubes,
77 followed by the solution (SAB broth or water) and then the acetic acid. We used three
78 negative control Hungate tubes, one tube with iron filings in SAB culture medium, another
79 tube with iron filings in distilled water and finally a third tube containing iron filings alone.
80 Gas chromatography was then performed using a Clarus 580 FID chromatograph
81 (PerkinElmer, Villebon-sur-Yvette, France) to measure hydrogen production in the different
82 tubes, and the pH was monitored (Fisher Scientific, Illkirch, France).

83

84 **Culture of methanogen strains using the chemical method for hydrogen production.** *M.*

85 *oralis* CSUR P9633, a human saliva isolate (22), was used for the development of the

86 chemical method. We used the mini-double-chamber flask technique, derived from the one

previously described (21). SAB agar plates (5-cm plates, Fisher Scientific) inoculated with 200 μ L of a *M. oralis* suspension at 10^7 colony-forming units (CFUs) was placed in the upper compartment of the mini-double-chamber flask, which was sealed with parafilm to ensure an anaerobic atmosphere. We then put 1 g of iron filings and 100 μ L of acetic acid in 10 mL of distilled water in the lower compartment of the mini-double-chamber flask in a microbiological safety station. We used an SAB agar plate inoculated with 200 μ L of sterile phosphate buffered saline (PBS) (Fisher Scientific) as a negative control. We then incubated the plates at 37°C for nine days with visual inspection on day 3, day 6 and day 9 postinoculation.

Then, the growth of *M. smithii* CSUR P9632, a human stool isolate, was compared using the chemical method with the reference biological method using a double-chamber system as previously described for the aerobic culture of methanogens (17). For the chemical method, we put 1.5 g of iron filings and 150 μ L of acetic acid in 200 mL of distilled water in the lower compartment instead of *Bacteroides thetaiotaomicron*, which was used in the biological method. SAB agar plates inoculated with 200 μ L of a *M. smithii* suspension at 10^6 CFU or 200 μ L of PBS for the negative controls were placed in the upper compartment.

The pH of the SAB agar plates was measured at day 3, day 6 and day 9 postinoculation using pH indicator strips (Macherey-Nagel SARL, Hœrdt, France). We placed the pH indicator strip directly on the SAB agar plate for two minutes. We then took the value corresponding to the color obtained on the strip from the values indicated by the supplier. The pH and the redox potential of the broth in the lower chamber were measured at day 3, day 6 and day 9 postinoculation using the Accumet™ AE150 apparatus (Fisher Scientific). The probe was rinsed thoroughly with distilled water and then immersed in the lower chamber until the device displayed the value on the reading screen. The rinsing step was performed after each use to avoid any error in reading the device. We used the pH and redox potential of distilled

water alone and the SAB culture broth alone as controls. We also measured the redox potential of the upper compartment. For the two methods, we used one double-chamber system, in which we put one Petri plate with 10 mL of distilled water and one Petri plate with 10 mL of the SAB culture broth. These double-chamber systems were then incubated at 37°C, and the redox potential was measured in each of the Petri dishes at day 3, day 6 and day 9 postinoculation. The effectiveness of the two methods for the growth of methanogens was compared by observing the appearance of the first colonies as well as the average number of colonies at day 3, day 6 and day 9 postinoculation. Colonies were confirmed by autofluorescence optical microscopy as follows. Briefly, the colony deposited on a microscopy slide was observed at 63X using an epifluorescence microscope (Leica DMI 3000, Wetzlar, Germany) and at 100X magnification using another epifluorescence microscope (Leica DMI 6000). Colonies were enumerated at day 3, day 6 and day 9 postinoculation by focusing the image of the entire 5-cm plate on a black background using a portable camera without flash (Lenovo L38011, Beijing, China). Numbered images were analyzed and counted using ImageJ version 8 (<https://imagej.nih.gov/ij/download.html>) (Wayne Rasband, Java, National Institutes of Health, USA) as follows: each image was imported into ImageJ software and analyzed in a first step using the blue, diamidino phenylindole filter to normalized the image. In a second step, the normalized image was analyzed using the green fluorescein isothiocyanate filter and saved in the Joint Photographic Experts Group format. We standardized the images using the elliptical selection tabulation of the software. We then proceeded to manually count the enlarged colonies using the magnifying glass and the multipoint tabulation of the software.

Comparison of the two methods on saliva samples. The study was previously approved by the Ethics Committee of the IHU Méditerranée infection under n°2016-011. After having

informed and collected the consent of the participants, we collected saliva samples from 50 people, including 25 tobacco smokers (15 men, 10 women, median age of 30) and 25 nonsmokers (12 men, 13 women, median age of 28). The collection and processing of samples were performed as previously described (22). We used SAB agar plates inoculated with 200 μ L of PBS as negative controls and then proceeded in the same manner as described above for the *M. smithii* strain.

Molecular analysis. PCR was performed to confirm the identity of the colonies. Colonies were suspended in 200 μ L of ultrapure water (Fisher Scientific), and a sonication step was performed for 30 minutes. DNA was then extracted with the EZ1 Advanced XL Extraction Kit (QIAGEN, Hilden, Germany) using 200 μ L as the sample volume and 200 μ L as the elution volume. Gene amplification and PCR sequencing were performed as previously described (22–24). Real-time PCR analyses were performed as previously described (25).

RESULTS AND DISCUSSION

We report on the use of chemically produced hydrogen for the isolation and culture of methanogens as an alternative to the currently used biological method (17). The method we report is based on the production of hydrogen resulting from the chemical reaction between iron filings and a weak acid such as acetic acid. All reported results were validated by negative controls that remained negative and identification of colonies by autofluorescence and PCR-based analyses.

We first determined an optimal balance between the concentration of iron filings and acetic acid to maintain the production of hydrogen over several days at a relative concentration compatible with the isolation and culture of methanogens. These preliminary

experiments enabled the development of a safe and controlled production of hydrogen that was shown to be efficient in the culture of *M. oralis* and *M. smithii* reference strains.

In the first step, we measured the average redox potential and pH values of the various media investigated here. The pH and redox potential of the controls consisting of distilled water and SAB broth culture medium were 7 and + 81.3 mV and 7.3 and -11.9 mV, respectively. After incubation at 37°C for 3, 6 and 9 days, these values were 6.98, 6.92 and 6.87 units and +64.6 mV, +67.3 mV and +79.9 mV for distilled water; and 6.83, 6.78 and 6.27 units and +11.1 mV, +12.9 mV and +14.8 mV for the SAB broth culture medium, respectively (Table 1). These results agree with previous observations that most types of water, including tap water and bottled water, are oxidizing agents, and the value of their redox potential is positive, ≥ 20 mV (26). The addition of antioxidants in the SAB medium makes this culture medium reductive. Then, the pH and redox potential of distilled water enriched with iron filings and acetic acid (chemical method) were measured at 5.23, 5.16 and 5.08 units and +109.77 mV, +108.6 mV and +112.7 mV, respectively (Table 2). These values were 5.8, 5.25, and 5.24 units and +113.8 mV, +113.8 mV and +114.5 mV at day 3, day 6 and day 9 for the SAB culture medium incubated with *B. thetaiotaomicron*, respectively (Table 2). These data indicated that the cultivation of *B. thetaiotaomicron* in the SAB medium releases fermenting compounds such as CO₂, hydrogen and acetate, which could oxidize the SAB medium (26). Then, we measured these values in the Petri dishes placed in the upper compartment of the double-chamber system, which hosted the methanogen culture (Fig. 1). The redox potential values were -28.1 mV, -81.2 mV and -108.9 mV at day 3, day 6 and day 9 for distilled water, respectively, and +13.5 mV, -116.5 mV and -77.5 mV at day 3, day 6 and day 9 for the SAB culture broth in the chemical method, respectively; and for the biological method, the redox potential values were -4.5 mV, -66 mV and -91.6 mV at day 3, day 6 and day 9 for distilled water, respectively, and +19.1 mV, -7.1 mV and -34.9 mV at day

3, day 6 and day 9 for the SAB culture broth in the biological method, respectively (Table 3). These data indicated that the hydrogen, being a reducer released into the bottom compartment, reaches the upper compartment and reduces the SAB medium or the water placed into Petri dishes in the upper compartment (27).

In a second step, by culturing the reference methanogen strains in the presence of the negative and positive controls, we obtained hydrogen production in all tubes and observed a correlation between hydrogen production and the amount of iron filings used and the acid acetic concentration. The hydrogen production using the chemical method was maintained until day 5, and the average pH of the mixture was 4.09 ± 0.15 and 4.35 ± 0.23 in the SAB culture broth and distilled water, respectively. *M. oralis* colonies that were confirmed by PCR sequencing were visible on day 3 postinoculation using the chemical method (Supplementary Fig. 1). The negative control consisting of PBS inoculation instead of *M. oralis* remained negative.

To compare the chemical method with the biological method, *M. smithii* was inoculated in SAB agar plates using both methods. Using the chemical method, inoculation of SAB agar plates with *M. smithii* yielded colonies as early as day 3 postinoculation, whereas they appeared on day 9 using the biological method (Supplementary Fig. 1, Fig. 2). All negative controls remained negative. The colonies were confirmed using PCR sequencing and real-time PCR. The average number of colonies of *M. smithii* was significantly higher using the chemical method than the biological method, regardless of the day of follow-up (Student's test, $p < 0.0001$, $p < 0.001$ and $p = 0.0001$ for day 3, day 6 and day 9 postinoculation, respectively) (Table 4). Moreover, cultures of these two strains were obtained in three days rather than in nine days using the biological method (17). In addition, the chemical method was easier to set up than the biological method because it incorporated distilled water instead of the SAB culture broth used in the chemical method, in contrast to

the biological method in which only the SAB culture broth can be used (Supplementary Table 1). Additionally, the biological method requires maintaining a *B. thetaiotaomicron* culture in the exponential growth phase to release efficient hydrogen production, which is not the case for the chemical method.

Finally, we compared the chemical method with the biological method for isolating methanogens from 50 saliva samples collected from 50 individuals after 9 days of incubation. The number of positive samples was 6/50 (12%) at day 3 postinoculation using both methods, and one additional saliva sample was positive at day 6, so that a total of 7/50 (14%) were positive using both methods. No additional sample was positive at day 9 postinoculation, regardless of the culture method. In these 7 culture-positive saliva samples, the average number of colonies at day 6 and day 9 postinoculation was significantly higher using the biological method than using the chemical method (Student's test, $p = 0.005$ and $p = 0.04$), whereas there was no significant difference in the average number of colonies on day 3 (Student's test, $p = 0.09$) (Table 4). Cultures from 6/25 (24%) of the samples from tobacco smokers yielded colonies as early as day 3 in both methods versus 1/25 (4%) of the samples from nonsmokers at day 6 in both methods. Colonies yielded autofluorescence in both methods (Fig. 2 and Fig. 3), whereas no colony or autofluorescence was observed in the negative controls. PCR sequencing and real-time PCR identified three *M. oralis*-positive and three *M. smithii*-positive samples in the tobacco smoker samples and one *M. oralis*-positive sample in one nonsmoker sample. Applying the chemical method for the isolation of *M. oralis* from saliva samples yielded significantly more colonies more rapidly than the biological method used in parallel. We hypothesized that the chemical method allowed mastering the kinetics of hydrogen production in such a way to optimize the atmosphere in the double-chamber system used in our experiments. These results show that the chemical method is more effective than the biological method for the rapid culture of methanogens.

Accordingly, the observation that almost a quarter of the saliva samples grew *M. oralis* and *M. smithii* is in agreement with a previously published report (22). Interestingly, we confirmed that methanogens are more prevalent in the saliva from tobacco smokers than in the saliva collected from nonsmokers. These results are consistent with those found previously (22), in which there was also a predominance of *M. oralis* over *M. smithii* in saliva samples. These observations also confirm that *M. oralis* is the most prevalent methanogen in the oral cavity. We measured the redox potential in the culture using the chemical method to monitor whether this chemical reaction created anaerobiosis. Our results indicate that this chemical reaction is reducing and confirm that acetic acid is an efficient oxidizer of iron filings (20). The pH of the agar plates remained neutral throughout the inoculation period both in the chemical method and the biological method; this observation suggests that the acidity produced in the lower compartment of the double-chamber device in the chemical method does not reach the upper compartment.

The present study highlighted some advantages of the chemical method over the biological method (Supplementary Table 1). The chemical method is simple to set up, the hydrogen production is controlled so that the time of appearance of the first colonies is fast due to the immediate hydrogen production, and this chemical method can be done with both distilled water and SAB culture broth. Given the simplicity, speed and efficiency of this chemical method of hydrogen production, it could replace the biological method incorporating *B. thetaiotaomicron* in the isolation and culture of methanogens. Using the chemical method will ease the routine isolation and culture of methanogens in microbiology laboratories to ease and speed the isolation and culture of these opportunistic pathogens (1).

ACKNOWLEDGEMENTS

COG and GG benefit from Ph.D. grants from the Fondation Méditerranée Infection, Marseille, France. This work was supported by the French Government under the «Investissements d’avenir» (Investments for the Future) program managed by the Agence Nationale de la Recherche (ANR, fr: National Agency for Research), (reference: Méditerranée Infection 10-IAHU-03). This work was supported by Région Sud (Provence Alpes Côte d’Azur) and European funding FEDER PRIMMI.

REFERENCES

1. Sogodogo E, Drancourt M, Grine G. 2019. Methanogens as emerging pathogens in anaerobic abscesses. *Eur J Clin Microbiol Infect Dis* 2:19-25.
2. Dridi B, Henry M, El Khéchine A, Raoult D, Drancourt M. 2009. High prevalence of *Methanobrevibacter smithii* and *Methanosphaera stadtmanae* detected in the human gut using an improved DNA detection protocol. *PLoS ONE* 4:e7063.
3. Deppenmeier U, Lienard T, Gottschalk G. 1999. Novel reactions involved in energy conservation by methanogenic archaea. *FEBS Lett* 457:291–297.
4. Deppenmeier U. 2002. Redox-driven proton translocation in methanogenic Archaea. *Cell Mol Life Sci* 59:1513–1533.
5. Chatterjee S, Park S, Low K, Kong Y, Pimentel M. 2007. The Degree of breath methane production in IBS correlates with the severity of constipation. *AJG* 102:837–841.
6. Zhang H, DiBaise JK, Zuccolo A, Kudrna D, Braidotti M, Yu Y, Parameswaran P, Crowell MD, Wing R, Rittmann BE, Krajmalnik-Brown R. 2009. Human gut microbiota in obesity and after gastric bypass. *Proc Natl Acad Sci* 106:2365–2370.
7. Piqué JM, Pallarés M, Cusó E, Vilar-Bonet J, Gassull MA. 1984. Methane production and colon cancer. *Gastroenterology* 87:601–605.
8. Holma R, Korpela R, Sairanen U, Blom M, Rautio M, Poussa T, Saxelin M, Osterlund P. 2013. Colonic Methane Production Modifies Gastrointestinal Toxicity Associated With Adjuvant 5-Fluorouracil Chemotherapy for Colorectal Cancer. *J Clin G* 47:45–51.

- 288 9. Ghavami SB, Rostami E, Sephay AA, Shahrokh S, Balaii H, Aghdaei HA, Zali MR.
289 2018. Alterations of the human gut *Methanobrevibacter smithii* as a biomarker for
290 inflammatory bowel diseases. Microb Patho 117:285–289.

- 291 10. Belay N, Mukhopadhyay B, Conway de Macario E, Galask R, Daniels L. 1990.
292 Methanogenic bacteria in human vaginal samples. J Clin Microbiol 28:1666–1668.

- 293 11. Grine G, Drouet H, Fenollar F, Bretelle F, Raoult D, Drancourt M. 2019. Detection of
294 *Methanobrevibacter smithii* in vaginal samples collected from women diagnosed with
295 bacterial vaginosis. Eur J Clin Microbiol Infect Dis 21:34-36.

- 296 12. Drancourt M, Nkanga VD, Lakhe NA, Régis J-M, Dufour H, Fournier P-E, Bechah Y,
297 Michael Scheld W, Raoult D. 2017. Evidence of Archaeal methanogens in brain
298 abscess. Clin Infect Dis 65:1–5.

- 299 13. Nkanga VD, Lotte R, Chirio D, Lonjon M, Roger P-M, Drancourt M, Ruimy R. 2018.
300 *Methanobrevibacter oralis* detected along with *Aggregatibacter actinomycetemcomitans*
301 in a series of community-acquired brain abscesses. Clin Microb Infect 24:207–208.

- 302 14. Nkanga VD, Lotte R, Roger P-M, Drancourt M, Ruimy R. 2016. *Methanobrevibacter*
303 *smithii* and *Bacteroides thetaiotaomicron* cultivated from a chronic paravertebral muscle
304 abscess. Clin Microb Infect 22:1008–1009.

- 305 15. Garcia JL, Patel BK, Ollivier B. 2000. Taxonomic, phylogenetic, and ecological
306 diversity of methanogenic Archaea. Anaerobe 6:205–226.

- 307 16. Ferrari A, Brusa T, Rutili A, Canzi E, Biavati B. 1994. Isolation and characterization of
308 *Methanobrevibacter oralis* sp. nov. Curr Microb 29:7–12.

- 309 17. Khelaifia S, Lagier J-C, Nkanga VD, Guilhot E, Drancourt M, Raoult D. 2016. Aerobic
310 culture of methanogenic archaea without an external source of hydrogen. Eur J Clin
311 Microbiol Infect Dis 35:985–991.
- 312 18. Traore SI, Khelaifia S, Armstrong N, Lagier JC, Raoult D. 2019. Isolation and culture of
313 *Methanobrevibacter smithii* by co-culture with hydrogen-producing bacteria on agar
314 plates. Clin Microb Infect 12:45-47.
- 315 19. Boddien A, Loges B, Gärtner F, Torborg C, Fumino K, Junge H, Ludwig R, Beller M.
316 2010. Iron-catalyzed hydrogen production from formic acid. J Am Chem Soc 132:8924–
317 8934.
- 318 20. Crouzet C, Brunet F, Recham N, Findling N, Lanson M, Guyot F, Ferrasse J-H, Goffé
319 B. 2017. Hydrogen production by hydrothermal oxidation of FeO under acidic
320 conditions. Inter J Hyd E 42:795–806.
- 321 21. Khelaifia S, Raoult D, Drancourt M. 2013. A Versatile Medium for Cultivating
322 Methanogenic Archaea. PLoS ONE 8:e61563.
- 323 22. Grine G, Terrer E, Boualam MA, Aboudharam G, Chaudet H, Ruimy R, Drancourt M.
324 2018. Tobacco-smoking-related prevalence of methanogens in the oral fluid microbiota.
325 Sci Rep 8:23-26.
- 326 23. Grine G, Boualam MA, Drancourt M. 2017. *Methanobrevibacter smithii*, a methanogen
327 consistently colonising the newborn stomach. Eur J Clin Microbiol Infect Dis 36:2449–
328 2455.

- 329 24. Nkamga VD, Huynh HTT, Aboudharam G, Ruimy R, Drancourt M. 2015. Diversity of
330 human-associated *Methanobrevibacter smithii* isolates revealed by multispacer sequence
331 typing. *Curr Microbiol* 70:810–815.
- 332 25. Bringuier A, Khelaifia S, Richet H, Aboudharam G, Drancourt M. 2013. Real-time PCR
333 quantification of *Methanobrevibacter oralis* in periodontitis. *J Clin Microbiol* 51:993–
334 994.
- 335 26. Adamberg S, Tomson K, Vija H, Puurand M, Kabanova N, Visnapuu T, Jaugi E, Alame
336 T, Adamberg K. 2014. Degradation of fructans and production of propionic acid by
337 *Bacteroides thetaiotaomicron* are enhanced by the shortage of amino acids. *Front Nut*
338 1:54-60.
- 339 27. Thauer RK, Kaster A-K, Seedorf H, Buckel W, Hedderich R. 2008. Methanogenic
340 archaea: ecologically relevant differences in energy conservation. *Nat Rev Microbiol*
341 6:579–591.

342

343 TABLES

344 **Table 1.** Controls of pH and redox potential.

Controls	Redox potential (mV)				pH			
	D0	D3	D6	D9	D0	D3	D6	D9
Distilled water	+81.3	+64.6	+67.3	+79.9	7	6.98	6.92	6.87
SAB medium	-8.7	+11.1	+12.9	+14.8	7.3	6.83	6.78	6.27

345

346

Table 2. pH and redox potential of the lower compartment.

Methods	Redox potential (mV)			pH		
	D3	D6	D9	D3	D6	D9
Chemical method	+109.77	+108.6	+112.7	5.23	5.16	5.08
Biological method	+113.8	+113.8	+114.5	5.8	5.25	5.24

Table 3. Redox potential of the upper compartment.

Redox potential (mv)	Chemical method			Biological method		
	D3	D6	D9	D3	D6	D9
Distilled water	-28.1	-81.2	-108.9	-4.5	-66	-91.6
SAB medium	+13.5	-116.5	-77.5	+19.1	-7.1	-34.9

Table 4. Comparison of the two methods according to the average of the colonies counted by ImageJ.

Methods	Chemical method			Biological method			P-value		
	D3	D6	D9	D3	D6	D9	D3	D6	D9
<i>M. smithii</i>	377±72	659±168	1116±193	0	0	436±74	<0.0001	<0.001	0.0001
Saliva samples	190±90	509±133	700 ± 232	305±139	722±96	936±136	0.09	0.005	0.04
Negative controls	0	0	0	0	0	0	NA	NA	NA

NA : Not adapted

FIGURES

Figure 1. Aerobic culture of methanogens in a double-chamber system on agar plates using either a microbiological method for hydrogen production (red arrow) or a chemical method for hydrogen production investigated here (blue arrow).

Figure 2. (A) Fluorescence emitted by *M. smithii* by applying the blue filter set. (B) Fluorescence emitted by *M. smithii* by applying the green filter set.

Figure 3. (A) Fluorescence emitted by *M. oralis* by applying the blue filter set. (B) Fluorescence emitted by *M. oralis* by applying the green filter set.

FIGURE 2

A

B

FIGURE 3