Post-Transcriptional Bone Morphogenetic Protein 2 (BMP2) Gene Regulation in Aorta

Tapan A. Shah¹, Ying Tang¹, Edward J. Yurkow² and Melissa B. Rogers^{1*}

¹Rutgers - New Jersey Medical School, Microbiology, Biochemistry, & Molecular Genetics,
Newark, NJ

²Rutgers University Molecular Imaging Center (RUMIC), Rutgers University, Piscataway, NJ

Running title: Bmp2 Gene Regulation in the Aorta

* To whom should correspondence and reprint request be addressed to: Melissa B. Rogers,
Ph.D., Microbiology, Biochemistry & Molecular Genetics, Rutgers - NJ Medical School, Bldg. UH
Cancer Center, Room F1216, 205 South Orange Ave., Newark, NJ 07103. Email:
rogersmb@njms.rutgers.edu, telephone: 973 972 2984

Keywords: BMP, Klotho, gene regulation, signaling, cardiovascular, calcification,
atherosclerosis, renal physiology, sex differences

Grant Support: Funding was provided by the National Heart, Lung, and Blood Institute
R01HL114751 and National Institutes of Aging R56AG050762 to MBR

Conflict of Interests Disclosure: The authors declare that they have no conflicts of interest
with the contents of this article.

Abstract

Deletion of an “ultra-conserved sequence” (UCS) within the *Bone Morphogenetic Protein*
(*Bmp*)2 mRNA previously revealed that the sequence represses *Bmp2* reporter gene
expression in vascular cells. The objective was to determine the impact of the endogenous
UCS on *Bmp2* mRNA levels, BMP signaling, and calcification in the healthy control aorta and in
the calcified aorta of mice with renal disease. We compared the phenotypes of mice bearing a
wild type *Bmp2* allele or the UCS deletion allele in mice with normal kidney function or in *Klotho*
mutant mice with reduced kidney function. BMP signaling and calcium levels were normally
higher in control females relative to males. UCS deletion induced aortic *Bmp2* mRNA and BMP
signaling in control males, but not in females. UCS deletion significantly increased BMP

signaling in both male and female *Klotho* homozygotes. Inheritance of the *Bmp2* UCS deletion and *Klotho* alleles was skewed from Mendelian expectations suggesting that these alleles influence interacting pathways. Analyses of body and heart weight supported these interactions. The *Bmp2* UCS represses BMP signaling in control males and in mice of both sexes with abnormal mineralization associated with kidney disease. Disease and sex-specific differences in *Bmp2* gene control may influence the onset and progression of cardiovascular diseases.

Introduction

The pro-osteogenic bone morphogenetic protein 2 (BMP2) is a potent pro-calcific signal (3-10). Moreover, BMP2 and its downstream effectors, e.g., phosphorylated SMAD1/5/9(8), are implicated strongly in pathological calcification (5,6,9-13). Various parallels exist between osteogenesis and cardiovascular calcification *via* the BMP2 link. BMP2 can induce osteogenic factors in human aortic valve interstitial cells (13) and aortic smooth muscle cells (14). BMP2 induces ossification in diseased aortic valves (9,11,15) and in atherosclerotic plaques (6) and induces calcification *in vitro* (10,13,16-18). Despite an explicit role of BMP2 in cardiovascular calcification, the mechanisms regulating the patterns of BMP2 and its downstream effectors in the healthy and diseased aorta are incompletely understood. Significant and unanswered questions are: What restrains *Bmp2* expression in healthy cardiovascular tissues? Why is *Bmp2* induced in physiologies such as aging and reduced renal function that promote pathological calcification? Elucidating the mechanisms that control BMP2 synthesis leading to pathological calcification may reveal new therapeutic strategies.

Several *cis* and *trans*-acting factors can regulate *Bmp2* gene expression either transcriptionally or post-transcriptionally [reviewed in (19) and (20)]. The *Bmp2* gene may be “active”; i.e., transcribed, but a post-transcriptional block may prevent BMP2 synthesis in specific cell types. Our studies showed that a unique ultra-conserved sequence (UCS) in the 3' untranslated region (UTR) of the transcript mediates this repression (21-23). The UCS repressed reporter genes in mesenchymal and other types of non-transformed cells *in vitro* (24-27) as well as in the

Bmp2 Gene Regulation in the Aorta

coronary vasculature, valves, and the aorta *in vivo* (25,27). The fact that these tissues are prone to calcification in patients with CAVD and atherosclerosis risk factors, suggests the hypothesis that UCS-mediated repression may protect against pathological levels of BMP2 synthesis leading to calcification.

Our previous findings in embryos demonstrated that the UCS limits *Bmp2* mRNA abundance and BMP signaling and that disturbing *Bmp2* 3'UTR-mediated events negatively impacted embryonic development. Here we describe the impact of an allele lacking the UCS (*Bmp2*^{ΔUCS}) on *Bmp2* expression and BMP signaling in the adult aorta of healthy control mice and in mice with a mutation that causes renal failure and premature aging.

BMP2 levels are normally low in the healthy vasculature but are induced in pathologically calcified tissues. Mutation of the *Klotho* gene provides an experimental model in which we could compare the mechanisms that repress or activate *Bmp2*. KLOTHO is a protein largely synthesized in the kidney that controls mineral metabolism. KLOTHO deficiency promotes hyperphosphatemia that leads to rapid and dramatic calcification of the aorta and aortic valve by 6-7 weeks of age (28,29). In contrast, calcification in other models, e.g., hyperlipidemia, is quite slow (30). Furthermore, BMP2 protein was observed in the calcified aortic valves of *Klotho* null mice and BMP signaling was shown to be required for aortic valve calcification (29). In this study, we describe how the UCS affects *Bmp2* RNA abundance, BMP signaling, and vascular calcification in the aorta and the effect on the overall fitness of control mice and *Klotho* mutant mice of both sexes.

Materials and Methods

Mouse strains

The background of all mice bearing the *Bmp2* and *Klotho* mutations was a mixture of strains 129, C57Bl/6J, and C3H/J. The UCS deletion allele (*Bmp2*^{ΔUCS}) was described in Shah *et al.* (31). Mice bearing the *Klotho* mutation (28) were a gracious gift from Dr. Makoto Kuro-o (Jichi Medical University) by way of Dr. Sylvia Christakos (Rutgers New Jersey Medical School).

C57Bl/6 female mice (47 days, 6 months, 12 months, 18 months and 22 or 23 months old) and male mice (47 days, 6 months, 12 months, 18 months and 21 months old) were obtained from the National Institute of Aging (NIA) aged rodent colonies (Bethesda, MD).

Mice Handling

Animals were handled in accordance with the Guidelines for Care and Use of Experimental Animals and approved by the NJ Medical School Institutional Animal Care and Use Committee (IACUC protocol #15069). Control and *Klotho* homozygote mice were fed regular chow and euthanized at 47 ± 6 days of age. After weaning, *Klotho* homozygotes received softened chow on the floor of the cage. Aortas were obtained from 10-week-old C57Bl/6 females that were ovariectomized for another study (Khariv and Elkabes in preparation, IACUC protocol #15038). Briefly, mice were anesthetized and both ovaries surgically removed. Sham mice underwent surgery without removal of ovaries. Three weeks post-surgery, mice were euthanized and necropsied as described below.

On the day of necropsy, mice were killed with an inhalation overdose of isoflurane. Immediately thereafter, the heart was perfused *via* the left ventricle with phosphate buffered saline (PBS, pH 7.3), to remove excess blood. The heart and aorta down to the aortic abdominal bifurcation into the left and right common iliac arteries were removed intact. After cutting the aorta at the surface of the heart, both tissues were rinsed in PBS, blot dried, and weighed. The heart was fixed in neutral buffered formalin. The aorta was snap-frozen in liquid nitrogen and stored at -80°C . For biochemical assays, frozen tissues were ground in liquid nitrogen using a mortar and pestle. The frozen powder was split to be used for different assays. To facilitate the handling of small tissues such as the diseased *Klotho* aortas, glass beads (Millipore-SIGMA, St. Louis, MO, # G1277) were added during the grinding. Glass beads did not affect the biochemical assays (Fig. 1A, B).

Genotyping

Bmp2 Gene Regulation in the Aorta

Genomic DNA was isolated and *Bmp2* genotypes were determined by semi-quantitative PCR as described in Shah *et al* (31). *Klotho* genotypes were determined by semi-quantitative PCR using LA-Tag DNA polymerase and TaKaRa buffer with Mg⁺² (TaKaRa Shuzo, Tokyo, Japan) as follows: initial denaturation at 97°C for 2 min; followed by 32 cycles of denaturation at 94°C for 30 secs; annealing at 55°C for 30 secs; and extension at 72°C for 1 min 30 secs; ending with a final extension at 72°C for 10 min. A common primer (TGGAGATTGGAAGTGGACG, 0.2 μM final concentration) and a wild type specific primer (TTAAGGACTCCTGCATCTGC, 0.05 μM final concentration) amplified a 458 bp fragment from the wild type *Klotho* allele. The common primer and a mutation specific primer (CAAGGACCAGTTCATCATCG, 0.2 μM final concentration) amplified a 920 bp fragment from the *Klotho* mutant allele (32). Some *Bmp2* and *Klotho* genotyping was performed by Transnetyx, Inc, Cordova, TN.

RNA Isolation and Reverse Transcription and Quantitative Real Time PCR (RT qPCR)

Total RNA was isolated using the miRNeasy kit (Qiagen Inc., Germantown, MD, # 217004). RNA quantity and quality (A260/280) were determined using a NanoDrop spectrophotometer (NanoDrop Technologies, Wilmington, DE). cDNA was synthesized using the QuantiTect® Reverse Transcription kit (Qiagen Inc., Germantown, MD, # 205313). Quantitative PCR was performed using the QuantiTect® SYBR® Green PCR kit (Qiagen Inc., Germantown, MD, #204145) and a CFX96 Touch™ Real-Time PCR Detection System (Bio-Rad Laboratories, Hercules, CA, #1855196). Relative *Bmp2* mRNA expression was calculated using CFX96 Manager software (Bio-Rad Laboratories, Hercules, CA, # 1845000) with actin as the reference gene. Intron-spanning primers were used to eliminate amplicons generated from any contaminating genomic DNA. The primer sequences used were *Bmp2* - Forward (TAGATCTGTACCGCAGGCA) and Reverse (GTTCTCTCCACGGCTTCTTC) and *Actin* - Forward (CGCCACCAGTTCGCCATGGA) and reverse (TACAGCCCGGGGAGCATCGT).

Western blots

Bmp2 Gene Regulation in the Aorta

Frozen ground tissue was solubilized in RIPA buffer, sonicated, and subjected to western blot analyses as described in Shah et al (31). BMP signaling was measured using a monoclonal phospho-SMAD 1/5/9(8) antibody (Cell Signaling Technology, Danvers, MA, #13820) at a dilution of 1:1000. The pSMAD antibody was authenticated as described in (31) and Fig. 1C, D. Polyclonal total SMAD 1/5/9(8) (Santa Cruz Biotechnology, Inc., Santa Cruz, CA, #sc-6031-R) and a polyclonal actin antibody (Santa Cruz Biotechnology, Inc., Santa Cruz, CA, #sc-1615-R) were subsequently used at a dilution of 1:1000. In all cases, the secondary antibody was Goat Anti-Rabbit HRP (Abcam, Cambridge, MA, # ab97080) at a dilution of 1: 20,000. Antibody-bound proteins were detected using SuperSignal™ West Femto Maximum Sensitivity Substrate (ThermoFisher Scientific, Waltham, MA, # 34096) and imaged using a FluoroChem M (Protein Simple, San Jose, California).

Calcium assays

Ground tissue was solubilized and lysed by sonication on ice in PBS, pH 7.3 containing 0.16 mg/mL heparin. The Cayman Chemical Calcium Assay kit was used to measure calcium levels (Ann-Arbor, MI, #701220). Calcium levels were normalized to protein levels measured using the Bradford assay (Bio-Rad Laboratories, Hercules, CA, # 5000006).

Spatial mapping of calcified structures

The patterns of mineralization in *Klotho* heterozygote vs. *Klotho* mutant mice were determined using microcomputerized tomography (microCT) at the Rutgers Molecular Imaging Center (<http://imaging.rutgers.edu/>). Mice were scanned using the Albira® PET/CT (Carestream, Rochester, NY) at standard voltage and current settings (45kV and 400µA) with a minimal voxel size of <35 µm. Voxel intensities in the reconstructed images were evaluated and segmented with VivoQuant image analysis software (version 1.23, inviCRO LLC, Boston).

Statistical Analysis

The statistical significance was determined using student's *t* test, Chi-square test, linear regression and two-way ANOVA analysis. A *p* value of less 0.05 was considered statistically significant.

Results

Adult mice lacking the *Bmp2* UCS

We previously demonstrated that the *Bmp2* UCS represses reporter gene expression *in vitro* in mesenchymal cells (20,24,25) and *in vivo* in the aorta and coronary vasculature (25). These tissues are prone to pathological calcification. Furthermore, we showed that the *Bmp2* UCS represses *Bmp2* RNA abundance and BMP signaling in mid-gestation embryos (31). Based on these findings, we hypothesized that the *Bmp2* UCS represses *Bmp2* RNA levels, BMP signaling and calcium levels in the aorta. Although mice completely lacking the UCS were underrepresented (31), some mice did survive to adulthood. Consequently, we tested the impact of the UCS in the aorta of these adults. In all cases, both sexes were assayed, because sex hormones can influence the expression of various members of the BMP signaling pathway (33). We also tested UCS function in pathologically calcified aorta from mice with renal failure and premature aging due to KLOTHO deficiency (28). To obtain aorta from control or diseased mice with three *Bmp2* genotypes (wild type *Bmp2*^{+/+}, heterozygous *Bmp2*^{+/ Δ UCS}, or homozygous *Bmp2* ^{Δ UCS/ Δ UCS} for the UCS deletion), we mated parents that were heterozygous for the *Bmp2* allele lacking the UCS (*Bmp2*^{+/ Δ UCS}) and for the *Klotho* mutant allele (*Kl*^{kl/+}). The expected and actual fractions of each genotype are shown in Fig. 2A. In mice with adequate KLOTHO levels (*Kl*^{+/+} or *Kl*^{kl/+}), no statistically significant skewing from the Mendelian inheritance of the *Bmp2* mutant allele was observed (Fig. 2B).

The *Klotho* mutant allele is recessive. The first visible phenotype of homozygous *Klotho* mutant mice (*Kl*^{kl/kl}) is runting relative to control littermates at the time of weaning. Although significant prenatal or perinatal lethality has not been reported, we observed underrepresentation of *Klotho* homozygotes, although the difference from Mendelian expectations was not quite significant

Bmp2 Gene Regulation in the Aorta

(Fig. 2C). However, segregation of the *Bmp2* genotypes was significantly skewed in the *Klotho* homozygotes (Fig. 2D) with a significant overrepresentation of heterozygotes (*Bmp2*^{+/ Δ UCS}) bearing one wild type and one UCS deletion allele (*Bmp2*^{+/ Δ UCS}, Chi-squared equals 7.15 with 2 degrees of freedom, two-tailed *p*-value < 0.03). These results suggest that changes in BMP2 levels associated with this regulatory mutation alter the fitness of the mice with KLOTHO deficiency. Indeed, as will be discussed below, *Bmp2* genotype was associated with significant differences in weight specifically in the diseased *Klotho* homozygotes.

To confirm that the *Klotho* mutant allele is fully recessive, we compared wild type (*Kl*^{+/+}) mice and heterozygotes that inherited one wild type and one mutated *Klotho* allele (*Kl*^{kl/+}). We found that the following parameters: *Bmp2* RNA levels, BMP signaling, calcium levels, and organ and body weights did not differ between *Klotho* heterozygotes and mice with two wild type *Klotho* alleles (Table 1). Therefore, the results from *Klotho* “control” mice presented below include data from both wild type and *Klotho* heterozygotes. The influence of *Bmp2* genotype will be discussed first in control mice and then in the diseased homozygotes.

The *Bmp2* UCS influences *Bmp2* RNA and BMP signaling levels in control aorta

We first tested the effect of UCS deletion in aorta from healthy control mice. Using RT qPCR, we measured relative *Bmp2* RNA levels in aorta from mice that were wild type, heterozygous, or homozygous for the *Bmp2* UCS deletion allele. Fig. 3A shows that aortic *Bmp2* RNA abundance increased with the number of UCS deletion alleles in males. Specifically, aortic *Bmp2* RNA abundance was 1.3-fold higher in heterozygotes (*Bmp2*^{+/ Δ UCS}) and 2-fold higher in homozygotes (*Bmp2* ^{Δ UCS/ Δ UCS}, *p* = 2.6 x 10⁻⁵) relative to wild type mice (*Bmp2*^{+/+}). Interestingly, although the UCS acted as a repressor in males, UCS deletion did not affect *Bmp2* RNA abundance in females (Fig. 3A).

We then measured the impact of UCS deletion on BMP signaling as assessed by the phosphorylation of SMAD1/5/9(8), the canonical BMP signaling intermediates, using an antibody authenticated as described previously ((31), Fig. 1C, D). The ratio of phosphorylated

Bmp2 Gene Regulation in the Aorta

protein pSMAD1/5/9(8) relative to total SMAD1/5/9(8) levels was 1.7-fold higher in aorta from male heterozygotes (*Bmp2*^{+/ Δ UCS}, $p = 0.008$) and 2.2-fold higher in male homozygotes (*Bmp2* ^{Δ UCS/ Δ UCS}, $p = 3.5 \times 10^{-4}$) relative to wild type male mice (Fig. 3B, C). These results are consistent with the hypothesis that the *Bmp2* UCS represses BMP2 synthesis in the aorta. In contrast to males, deleting the *Bmp2* UCS failed to induce either *Bmp2* RNA or BMP signaling in females (Fig. 3A - C). This result and those described below suggest that sex influences *Bmp2* gene expression.

Estrogen was previously shown to directly induce *Bmp2* transcription in cultured bone marrow mesenchymal stem cells (34). However, the impact of estrogen on *Bmp2* expression has not been tested *in vivo*. To test whether or not this sex steroid impacts *Bmp2* RNA levels in the aorta, we measured *Bmp2* mRNA levels in aged females who do not undergo the pre-ovulatory rise in circulating estrogen levels (35,36). A simple linear regression calculation determined that aortic *Bmp2* RNA levels declined with increasing age (Fig. 3F, $R^2 = 0.35$, $p = 0.003$). We also observed reduced *Bmp2* RNA levels in aorta from ovariectomized (OVX) mice relative to mice subjected to a sham operation (Fig. 3F, $p = 0.05$). Interestingly, in males, *Bmp2* mRNA abundance was significantly higher in aorta from 21 months old mice relative to younger male mice (Fig. 3E, $p = 8.5 \times 10^{-5}$) or similarly aged females ($p = 0.02$). These observations are consistent with both transcriptional and post-transcriptional differences in the mechanisms that regulate BMP2 synthesis in males and females.

We then measured calcium levels normalized to protein concentration in whole aorta lysate from control males and females. We observed that UCS deletion did not elevate calcium levels in aorta from either sex (Fig. 3G). This suggests that regulatory mechanisms in the osteogenic pathway between BMP and calcium deposition may counter the pro-calcific effect of increased BMP signaling. Intriguingly, BMP signaling (Fig. 3B, $p = 0.04$) and calcium levels (Fig. 3G, $p = 0.0004$) in control aorta from females were higher than that in control aorta from males. In summary, sex-dependent differences in the regulatory mechanisms that control aortic

calcification include increased BMP signaling and basal calcification levels in females. These increases are accompanied by an apparent lack of UCS-associated repression in females.

***Klotho* mutant mice: an inducible model of *Bmp2* gene expression**

Increased BMP2 mRNA levels and BMP signaling are required for calcification in the aortic valve of mice homozygous for the *Klotho* null allele (29). We tested the role of BMP2 and BMP signaling in the calcified aorta of mice homozygous for the original hypomorphic *Klotho* allele (28). First, we confirmed that *Klotho* mRNA levels in the kidneys of mice homozygous for the *Klotho* mutation were less than 1% of that observed in healthy control mice (Fig. 4A, $p = 0.0009$). *Bmp2* genotype did not significantly alter *Klotho* RNA abundance (Fig. 4A). BMP signaling and calcium levels in aorta from control mice that were either wild type ($Kl^{+/+}$) or heterozygous ($Kl^{kl/+}$) for the *Klotho* mutation were compared to the calcified aorta from *Klotho* mutant homozygotes ($Kl^{kl/kl}$). BMP signaling was induced nearly 2-fold in aorta from male and female *Klotho* homozygous mice relative to control mice (Fig. 4B, C; $p = 0.03$). Calcium levels in the aorta from *Klotho* mutant homozygotes were significantly elevated by over 2-fold in males ($p = 1.4 \times 10^{-6}$) and females ($p = 5.0 \times 10^{-4}$) relative to control aorta (Fig. 4D). PET-CT imaging revealed profound mineralization in the aortic sinus and ascending aorta of *Klotho* mutant ($Kl^{kl/kl}$) mice, but not in a control heterozygous ($Kl^{kl/+}$) littermate (Fig. 4E, F). Together, these results confirmed that homozygosity for the *Klotho* mutation amplifies BMP signaling and calcification in the aorta.

The *Bmp2* UCS represses BMP signaling levels in calcified aorta from *Klotho* mutant mice

We showed earlier that the *Bmp2* UCS repressed mRNA abundance and BMP signaling in non-calcified control aorta from males, but not females (Fig. 3A - C). Consistent with our findings in control aorta from males (Fig. 3B, C), UCS deletion significantly stimulated aortic BMP signaling in male *Klotho* homozygotes ($Kl^{kl/kl}$). BMP signaling was about 2-fold higher in heterozygotes ($Bmp2^{+/ΔUCS}$, $p = 0.01$) and homozygotes ($Bmp2^{ΔUCS/ΔUCS}$, $p = 0.02$) relative to wild type

(*Bmp2*^{+/+}) mice (Fig. 5A, B). In contrast to control aorta from females, UCS deletion also stimulated aortic BMP signaling by 1.6-fold in female *Klotho* homozygotes relative to wildtype females (Fig. 5A, B; $p = 0.02$). The observed elevation in BMP signaling did not lead to an obvious stimulation of calcium levels in mice lacking the UCS (Fig. 5C).

The *Bmp2* UCS influences body and heart weights

We observed that the inheritance of UCS deletion allele varied with *Klotho* genotype (Fig. 2D). Therefore, we tested whether deleting the *Bmp2* UCS impacts overall health as assessed by the body weights of the mice. *Bmp2* genotype did not impact the body weight for control animals of both sexes (Fig. 6A). As expected, female control mice weighed about 20% less than males for all 3 *Bmp2* genotypes ($p = 3.96 \times 10^{-18}$). The severe runting phenotype caused homozygous *Klotho* mutant mice with the wild type *Bmp2* genotype to weigh about a third that of the control mice. Curiously, the typical male to female weight ratio was reversed in *Klotho* mutant homozygotes. Specifically, female *Klotho* homozygotes with the wild type *Bmp2* genotype weighed about 22% more than male homozygotes (Fig. 6A, $p = 0.04$). Homozygous deletion of the UCS (*Bmp2*^{ΔUCS/ΔUCS}) increased the weight of male *Klotho* homozygotes by 40% (Fig. 6A, $p = 0.0003$). Because UCS deletion failed to impact females similarly, the mice lacking the *Bmp2* UCS exhibited the usual male to female weight ratio with males weighing 23% more than females (Fig. 6A, $p = 0.008$). These results suggest a sex-specific difference in the impact of *Bmp2* UCS on the overall physiology of *Klotho* homozygotes. To test if this *Bmp2* allele influenced overall cardiovascular health, we next compared the heart weights of male and female mice with all *Bmp2* and *Klotho* genotypes (Fig. 6B). A modest increase in the heart weights of control mice was associated with UCS deletion. Because heart weight is normally proportional to body weight, we calculated relative heart weights. The small, but significant, increase in the heart weight of female *Klotho* controls lacking the UCS was retained after normalizing to body weight (Fig. 6C, $p = 0.04$).

Bmp2 Gene Regulation in the Aorta

Deletion of the UCS affected absolute heart weight more dramatically in *Klotho* homozygotes of both sexes. In male and female *Klotho* homozygotes, homozygous UCS deletion (*Bmp2*^{ΔUCS/ΔUCS}) significantly increased absolute heart weight by 75% in males ($p = 0.0002$) and 40% in females ($p = 0.004$, Fig. 6B). Female heart weight normalized to body weight also was increased by nearly 50% ($p = 0.005$, Fig. 6C). In male *Klotho* homozygotes, a trend towards higher relative weight remained (Fig. 6C) despite the increased overall body weight associated with UCS deletion (Fig. 6A). These results suggest that an intact *Bmp2* UCS protects from an enlarged heart.

Discussion

Our objective was to test the role of an extraordinarily conserved *Bmp2* regulatory element, the UCS, in the aorta from control adult mice and in mice with genetically induced vascular calcification associated with severely reduced renal function. We used a new *Bmp2* allele without the UCS that increased *Bmp2* RNA and BMP signaling in embryos (31). As we observed in embryos, the UCS can repress BMP synthesis in the adult aorta. However, we also discovered that the UCS functions differently between male and female mice. We will first discuss our findings in males and then compare and contrast these findings to those in females. Our previous study using a *lacZ* reporter gene controlled by the distal *Bmp2* promoter and the *Bmp2* 3'UTR showed that Cre-mediated deletion of the *Bmp2* UCS induced robust gene expression in the aorta, coronary vasculature, and cardiac valves (25). However, this reporter gene lacked the proximal promoter, intronic regulatory elements, and long-range *cis*-regulatory elements. Thus, the reporter gene only partly recapitulated endogenous *Bmp2* gene expression patterns (19,27). Using our new *Bmp2* allele lacking the UCS (*Bmp2*^{ΔUCS}), we showed that the UCS represses *Bmp2* mRNA abundance and BMP signaling in embryos (31). In the present study, we demonstrated for the first time that the UCS also represses *Bmp2* RNA abundance and BMP signaling in control adult male aorta (Fig. 3A - C). This observation, which is consistent with all previous studies using reporter genes *in vitro* and *in vivo* (20,24,25,27),

Bmp2 Gene Regulation in the Aorta

indicates that the UCS can restrain this pro-calcific growth factor in healthy cardiovascular tissues.

In contrast to healthy vascular cells, increased levels of BMP2 are tightly associated with pathological calcification of the coronary vasculature (5,6,11,12). To understand why this pro-calcific protein is induced in physiologies such as aging and reduced renal function, we also assessed the capacity of the UCS to restrain BMP signaling in aortic tissue undergoing pathological calcification. In mice bearing the wild type *Bmp2* genotype, but homozygous for the hypomorphic *Klotho* mutation, BMP signaling, and calcium levels were increased in the aorta (Fig. 4B - F). This mirrors the increased levels observed in aortic valves from *Klotho* null mice (29). As in control males, the *Bmp2* UCS repressed BMP signaling in the aorta from male *Klotho* mutant mice (Fig. 5A, B). A trend towards increased aortic calcium levels with *Bmp2* UCS deletion was observed in male *Klotho* mutant mice (Fig. 5C). Thus, in both control and diseased male aorta, the UCS limits BMP signaling. Our observations in males correspond with the simple hypothesis that a functional UCS protects against excessive BMP signaling and maybe against calcification.

Interestingly, results in female mice suggest a more complex story. First, basal BMP signaling was 2-fold higher in aorta from control females relative to males (Fig. 3B). Second, unlike in males, the *Bmp2* UCS failed to repress *Bmp2* mRNA and BMP signaling in control non-calcified aorta from females (Fig. 3A - C). Finally, basal aortic calcium was 36% higher in females relative to males (Fig. 3G). However, as in both healthy control and *Klotho* mutant male mice, the UCS did repress aortic BMP signaling in the female *Klotho* homozygotes. This suggests that *Klotho*-associated renal disease stimulates the braking action of the UCS on aortic BMP signaling in females. Potential sex-related differences in *Bmp2* gene regulatory and BMP signaling mechanisms will be further investigated (33).

The female hormone estrogen may play a role in these differences. Indeed, previous findings in cell culture indicate that estrogen directly induces *Bmp2* transcription (34,37). We confirmed

Bmp2 Gene Regulation in the Aorta

that estrogen stimulates *Bmp2* RNA abundance in the intact aorta (Fig. 3F). The differential impact of deleting the UCS suggests additional dissimilarities in the post-transcriptional processes that regulate BMP2 synthesis in males and females. Sex-related differences in *Bmp2* gene regulation may be clinically relevant as significant disparities in incidence, prognosis, and response to treatments for arterial diseases occur between men and women (38-40).

BMPs were defined by their ability to induce bone from mesenchymal tissues (4). Indeed, all major forms of cardiovascular calcium deposition (aortic valve, medial artery, and atherosclerotic calcification) proceed by mechanisms resembling bone formation (5,9,12,42). However, the increased BMP signaling associated with UCS deletion did not lead clearly to a corresponding increase in aortic calcification. Thus, loss of the UCS-mediated repression of *Bmp2* may not be a sufficient trigger to induced aortic calcification. Indeed, this post-transcriptional repression is only one of the numerous mechanisms that control *Bmp2* gene expression and BMP signaling (19). BMP signaling can activate repressors such as the extracellular antagonist NOGGIN, the intracellular repressor SMAD6, and the miRNAs that repress BMP receptors (43-45). Furthermore, a multitude of positive and negative regulators control the differentiation program leading from bone progenitors to committed osteoprogenitors to differentiating osteoblasts and finally to mature osteoblasts that cause mineralization (46). The redundant feedback mechanisms that dampen BMP signaling along with downstream regulators of osteogenesis would buffer the impact of reduced UCS function on vascular calcification.

Our focus in this study was the regulatory impact of the UCS on *Bmp2* gene expression, BMP signaling, and calcification. However, we observed that the UCS deletion allele modified the *Klotho* phenotype more generally and in a manner also distinguished by sex. First, deletion of the UCS significantly increased the weight of male *Klotho* homozygotes, but not female homozygotes (Fig. 6A). A second phenotype was an increase in heart weight. Although most

notable in *Klotho* homozygotes of both sexes, the weights of control female hearts were significantly elevated by deletion of the UCS within only 6 -7 weeks of age (Fig. 6B, C). The UCS deletion allele dramatically altered embryonic morphogenesis and viability, but only in a subset of offspring (31). Because the penetrance of severe embryonic malformations correlated with the level of BMP signaling, we proposed that other BMP pathway regulators compensated for deletion of the repressive UCS. The surviving pups – the subjects of this study – are those with adequate regulation. In the survivors, subtler congenital defects, e.g., of the valves, are possible. Indeed, the edema observed in some embryos is consistent with reduced cardiac function (31). The impact of such developmental anomalies may appear sooner in mice subjected to additional cardiovascular stress, such as the renal failure associated with KLOTHO loss of function. The extraordinary conservation of the UCS is consistent with evolutionary drive to maintain BMP signaling within a narrow developmentally tolerated range. If the remaining redundant regulatory mechanisms are severely inadequate, then catastrophic embryonic malformations may occur. Less severe regulatory changes may reveal themselves only in the contexts of aging or pathological stresses.

Acknowledgments: We thank Drs. Stella Elkabes and Li Ni for the gift of the aorta from ovariectomized females. We also appreciate the part-time assistance of students Annica Tehim (Rutgers New Jersey Medical School), Amy Song (The College of NJ), Irina Kleiman and Risha Patel (Rutgers School of Graduate Studies).

References

1. Benjamin EJ, Virani SS, Callaway CW, et al. Heart Disease and Stroke Statistics—2018 Update: A Report From the American Heart Association. *Circulation*. 2018.
2. National Chronic Kidney Disease Fact Sheet. . Available at.
3. Vattikuti R, Towler DA. Osteogenic regulation of vascular calcification: an early perspective. *Am J Physiol Endocrinol Metab*. 2004;286(5):E686-696.
4. Urist MR. Bone: formation by autoinduction. *Science*. 1965;150(3698):893-899.
5. Towler DA. Commonalities Between Vasculature and Bone: An Osseocentric View of Arteriosclerosis. *Circulation*. 2017;135(4):320-322.
6. Bostrom K, Watson KE, Horn S, Wortham C, Herman IM, Demer LL. Bone morphogenetic protein expression in human atherosclerotic lesions. *The Journal of clinical investigation*. 1993;91(4):1800-1809.

Bmp2 Gene Regulation in the Aorta

7. Balachandran K, Sucosky P, Jo H, Yoganathan AP. Elevated cyclic stretch induces aortic valve calcification in a bone morphogenetic protein-dependent manner. *Am J Pathol*. 2010;177(1):49-57.
8. Kaden JJ, Bickelhaupt S, Grobholz R, et al. Expression of bone sialoprotein and bone morphogenetic protein-2 in calcific aortic stenosis. *J Heart Valve Dis*. 2004;13(4):560-566.
9. Miller JD, Weiss RM, Serrano KM, et al. Evidence for active regulation of pro-osteogenic signaling in advanced aortic valve disease. *Arterioscler Thromb Vasc Biol*. 2010;30(12):2482-2486.
10. Seya K, Yu Z, Kanemaru K, et al. Contribution of bone morphogenetic protein-2 to aortic valve calcification in aged rat. *Journal of pharmacological sciences*. 2011;115(1):8-14.
11. Mohler ER, 3rd, Gannon F, Reynolds C, Zimmerman R, Keane MG, Kaplan FS. Bone formation and inflammation in cardiac valves. *Circulation*. 2001;103(11):1522-1528.
12. Goumans MJ, Zwijsen A, Ten Dijke P, Bailly S. Bone Morphogenetic Proteins in Vascular Homeostasis and Disease. *Cold Spring Harb Perspect Biol*. 2017;10:2.
13. Yang X, Meng X, Su X, et al. Bone morphogenetic protein 2 induces Runx2 and osteopontin expression in human aortic valve interstitial cells: role of Smad1 and extracellular signal-regulated kinase 1/2. *The Journal of thoracic and cardiovascular surgery*. 2009;138(4):1008-1015.
14. Cheng SL, Shao JS, Charlton-Kachigian N, Loewy AP, Towler DA. MSX2 promotes osteogenesis and suppresses adipogenic differentiation of multipotent mesenchymal progenitors. *The Journal of biological chemistry*. 2003;278(46):45969-45977.
15. Wirrig EE, Hinton RB, Yutzey KE. Differential expression of cartilage and bone-related proteins in pediatric and adult diseased aortic valves. *Journal of molecular and cellular cardiology*. 2011;50(3):561-569.
16. Osman L, Yacoub MH, Latif N, Amrani M, Chester AH. Role of human valve interstitial cells in valve calcification and their response to atorvastatin. *Circulation*. 2006;114(1 Suppl):1547-552.
17. Nigam V, Srivastava D. Notch1 represses osteogenic pathways in aortic valve cells. *Journal of molecular and cellular cardiology*. 2009;47(6):828-834.
18. Yu Z, Seya K, Daitoku K, Motomura S, Fukuda I, Furukawa K. Tumor necrosis factor- α accelerates the calcification of human aortic valve interstitial cells obtained from patients with calcific aortic valve stenosis via the BMP2-Dlx5 pathway. *The Journal of pharmacology and experimental therapeutics*. 2011;337(1):16-23.
19. Rogers MB, Shah TA, Shaikh NN. Turning Bone Morphogenetic Protein 2 (BMP2) on and off in Mesenchymal Cells. *Journal of cellular biochemistry*. 2015;116(10):2127-2138.
20. Fotinos A, Fritz DT, Lisica S, Liu Y, Rogers MB. Competing Repressive Factors Control Bone Morphogenetic Protein 2 (BMP2) in Mesenchymal Cells. *Journal of cellular biochemistry*. 2016;117(2):439-447.
21. Abrams KL, Xu J, Nativelle-Serpentini C, Dabirshahsahebi S, Rogers MB. An evolutionary and molecular analysis of Bmp2 expression. *J Biol Chem*. 2004;279(16):15916-15928.
22. Fritz DT, Jiang S, Xu J, Rogers MB. A polymorphism in a conserved posttranscriptional regulatory motif alters bone morphogenetic protein 2 (BMP2) RNA:protein interactions. *Mol Endocrinol*. 2006;20(7):1574-1586.
23. Fritz DT, Liu D, Xu J, Jiang S, Rogers MB. Conservation of Bmp2 post-transcriptional regulatory mechanisms. *J Biol Chem*. 2004;279(47):48950-48958.
24. Devaney JM, Tosi LL, Fritz DT, et al. Differences in fat and muscle mass associated with a functional human polymorphism in a post-transcriptional BMP2 gene regulatory element. *Journal of cellular biochemistry*. 2009;107(6):1073-1082.
25. Kruithof BP, Fritz DT, Liu Y, et al. An autonomous BMP2 regulatory element in mesenchymal cells. *Journal of cellular biochemistry*. 2011;112(2):666-674.

Bmp2 Gene Regulation in the Aorta

26. Jiang S, Fritz DT, Rogers MB. A conserved post-transcriptional BMP2 switch in lung cells. *Journal of cellular biochemistry*. 2010;110(2):509-521.
27. Kruithof BP, Xu J, Fritz DT, Cabral CS, Gaussin V, Rogers MB. An in vivo map of bone morphogenetic protein 2 post-transcriptional repression in the heart. *Genesis*. 2011;49(11):841-850.
28. Kuro-o M, Matsumura Y, Aizawa H, et al. Mutation of the mouse klotho gene leads to a syndrome resembling ageing. *Nature*. 1997;390(6655):45-51.
29. Gomez-Stallons MV, Wirrig-Schwendeman EE, Hassel KR, Conway SJ, Yutzey KE. Bone Morphogenetic Protein Signaling Is Required for Aortic Valve Calcification. *Arterioscler Thromb Vasc Biol*. 2016;36(7):1398-1405.
30. Neven E, D'Haese PC. Vascular calcification in chronic renal failure: what have we learned from animal studies? *Circ Res*. 2011;108(2):249-264.
31. Shah TA, Zhu Y, Shaikh NN, Harris MA, Harris SE, Rogers MB. Characterization of new bone morphogenetic protein (Bmp)-2 regulatory alleles. *Genesis*. 2017;55(7):e23035.
32. Morishita K, Shirai A, Kubota M, et al. The progression of aging in klotho mutant mice can be modified by dietary phosphorus and zinc. *J Nutr*. 2001;131(12):3182-3188.
33. Shah TA, Rogers MB. Unanswered Questions Regarding Sex and BMP/TGF- β Signaling. *Journal of Developmental Biology*. 2018;6(2)(14).
34. Zhou S, Turgeman G, Harris SE, et al. Estrogens activate bone morphogenetic protein-2 gene transcription in mouse mesenchymal stem cells. *Molecular endocrinology (Baltimore, Md)*. 2003;17(1):56-66.
35. Frick KM. Estrogens and age-related memory decline in rodents: what have we learned and where do we go from here? *Horm Behav*. 2009;55(1):2-23.
36. Nelson JF, Felicio LS, Osterburg HH, Finch CE. Altered profiles of estradiol and progesterone associated with prolonged estrous cycles and persistent vaginal cornification in aging C57BL/6J mice. *Biol Reprod*. 1981;24(4):784-794.
37. Zhou S, Zilberman Y, Wassermann K, Bain SD, Sadovsky Y, Gazit D. Estrogen modulates estrogen receptor alpha and beta expression, osteogenic activity, and apoptosis in mesenchymal stem cells (MSCs) of osteoporotic mice. *J Cell Biochem Suppl*. 2001;Suppl 36:144-155.
38. Robinet P, Milewicz DM, Cassis LA, Leeper NJ, Lu HS, Smith JD. Consideration of Sex Differences in Design and Reporting of Experimental Arterial Pathology Studies-Statement From ATVB Council. *Arterioscler Thromb Vasc Biol*. 2018;38(4).
39. Chow JT, Khosla S, Melton LJ, 3rd, Atkinson EJ, Camp JJ, Kearns AE. Abdominal aortic calcification, BMD, and bone microstructure: a population-based study. *J Bone Miner Res*. 2008;23(10):1601-1612.
40. Jayalath RW, Mangan SH, Golledge J. Aortic calcification. *Eur J Vasc Endovasc Surg*. 2005;30(5):476-488.
41. Rigor and Reproducibility. NIH Website. . Available at.
42. Shao JS, Cai J, Towler DA. Molecular mechanisms of vascular calcification: lessons learned from the aorta. *Arterioscler Thromb Vasc Biol*. 2006;26(7):1423-1430.
43. Gazzero E, Minetti C. Potential drug targets within bone morphogenetic protein signaling pathways. *Curr Opin Pharmacol*. 2007;7(3):325-333.
44. Sun Q, Mao S, Li H, Zen K, Zhang CY, Li L. Role of miR-17 family in the negative feedback loop of bone morphogenetic protein signaling in neuron. *PLoS One*. 2013;8(12):e83067.
45. Mulloy B, Rider CC. The Bone Morphogenetic Proteins and Their Antagonists. *Vitamins and hormones*. 2015;99:63-90.
46. Salazar VS, Ohte S, Capelo LP, Gamer L, Rosen V. Specification of osteoblast cell fate by canonical Wnt signaling requires Bmp2. *Development*. 2016;143(23):4352-4367.

Bmp2 Gene Regulation in the Aorta

47. Carreira AC, Alves GG, Zambuzzi WF, Sogayar MC, Granjeiro JM. Bone Morphogenetic Proteins: structure, biological function and therapeutic applications. *Arch Biochem Biophys*. 2014;561:64-73.
48. Shao Y, Chen QZ, Zeng YH, et al. All-trans retinoic acid shifts rosiglitazone-induced adipogenic differentiation to osteogenic differentiation in mouse embryonic fibroblasts. *Int J Mol Med*. 2016;38(6):1693-1702.
49. Shen J, James AW, Zhang X, et al. Novel Wnt Regulator NEL-Like Molecule-1 Antagonizes Adipogenesis and Augments Osteogenesis Induced by Bone Morphogenetic Protein 2. *Am J Pathol*. 2016;186(2):419-434.
50. Vanhatupa S, Ojansivu M, Autio R, Juntunen M, Miettinen S. Bone Morphogenetic Protein-2 Induces Donor-Dependent Osteogenic and Adipogenic Differentiation in Human Adipose Stem Cells. *Stem Cells Transl Med*. 2015;4(12):1391-1402.
51. Kolodgie FD, Yahagi K, Mori H, et al. High-risk carotid plaque: lessons learned from histopathology. *Semin Vasc Surg*. 2017;30(1):31-43.
52. Simard L, Cote N, Dagenais F, et al. Sex-Related Discordance Between Aortic Valve Calcification and Hemodynamic Severity of Aortic Stenosis: Is Valvular Fibrosis the Explanation? *Circ Res*. 2017;120(4):681-691.

Bmp2 Gene Regulation in the Aorta

Figure 1. Experimental Controls. A, B. Grinding tissue with glass beads does not significantly affect calcium or protein yield. Whole aorta was ground in liquid nitrogen either with or without glass beads. Frozen ground powder tissue was split two ways for protein and calcium assays.

A. Effect of glass beads on calcium, protein and pSMAD 1/5/9(8) levels. Duplicate measurements are presented with range. Experiments were repeated twice with similar results.

B. Representative blots showing pSMAD 1/5/9(8), actin levels and the Ponceau S stained membrane after transfer. The positive control lane (+ Ctrl) was loaded with lysate from MC3T3-E1 cells transfected with a *Bmp2* expression plasmid. **C, D.** Validation of the phospho-SMAD 1/5/9(8) antibody. MC3T3-E1 (**C**) and C3H10T1/2 (**D**) cells were transfected with an expression plasmid encoding BMP2 (B2) or luciferase (Luc) (25). Cells were then lysed in RIPA buffer and subjected to western blotting as described in the experimental procedures section. These representative blots show that pSMAD1/5/9(8) levels were induced in cells transfected with the *Bmp2* expressing plasmid relative to the luciferase plasmid. Actin levels and a Ponceau S stained membrane are shown as loading controls.

Bmp2 Gene Regulation in the Aorta

526

527

528 **Figure 2. Apparent interaction between *Bmp2* and *Klotho* mutant alleles.** Mice

529 heterozygous for the *Bmp2* allele lacking the UCS (*Bmp2*^{+/ Δ UCS}) and for the *Klotho* mutant allele

530 (*Kl*^{kl/+}) were bred. The resulting pups were genotyped prior to weaning at approximately 4 weeks

531 of age. *Bmp2* genotypes: wild type (+/+), heterozygous (+/ Δ), or homozygous UCS deletion ($\Delta\Delta$)

532 and *Klotho* genotypes: wild type (+/+), heterozygous (kl/+), or homozygous *Klotho* mutation

533 (kl/kl). **A.** Expected and observed genotypes for the breeding scheme described above.

534 Occasionally, pups were found dead between birth and weaning, but the frequency did not differ

535 from that typically observed, nor was any particular genotype obviously over-represented. **B-D**

536 illustrate relative fractions for each genotype.

537

Bmp2 Gene Regulation in the Aorta

A.

Bmp2	+/+	+/ Δ	Δ/Δ	+/+	+/ Δ	Δ/Δ	+/+	+/ Δ	Δ/Δ
Klotho	+/+	+/+	+/+	kl/+	kl/+	kl/+	kl/kl	kl/kl	kl/kl
Expected %	6.25	12.5	6.25	12.5	25	12.5	6.25	12.5	6.25
Observed n	19	31	18	31	47	30	7	36	10
Observed %	8.3	13.5	7.9	13.5	20.5	13.1	3.1	15.7	4.4

B.

All healthy controls
(Kl $^{+/+}$ and Kl $^{kl/+}$)
Total = 176

C.

All Bmp2
genotypes
Total = 229

D.

All Klotho
Homozygotes (Kl $^{kl/kl}$)
Total = 53

Figure 3. *Bmp2* RNA, BMP signaling, and calcium levels in aorta from control mice with different *Bmp2* genotypes. Aortas were isolated from mice bearing the wild type (Wt), heterozygous (+/Δ), or homozygous UCS deletion (ΔΔ) *Bmp2* genotypes between 40 and 50 days of age. Mice were healthy with normal kidney function ($KI^{+/+}$ and $KI^{kl/+}$). Average parameter values are presented with the standard error of the mean (SEM). **A.** *Bmp2* RNA levels normalized to actin RNA levels (males, n = 12 – 21; females, n = 9 – 11). **B.** BMP signaling levels as assessed by phosphorylated SMAD1/5/9(8) (pSMAD) levels normalized to total SMAD 1/5/9(8) (tSMAD) levels (males, n = 6 – 8; females, n = 6 – 8). **C, D.** Representative western blot panels showing pSMAD1/5/9(8) and total SMAD1/5/9(8) levels, males (left) and females (right). The positive control lane (+ Ctrl) was loaded with lysate from MC3T3-E1 cells transfected with a *Bmp2* expression plasmid (Fig. 1). **E, F.** *Bmp2* RNA levels normalized to actin RNA levels in the aorta from male and female mice (ages below each bar, n = 4 - 6) and ovariectomized (OVX) and sham control mice (euthanized at 69 days, 3 weeks after operation, n = 3). **G.** Calcium levels normalized to protein concentration (males, n = 18 – 26; females, n = 6 – 18). All experiments were repeated at least twice with similar results. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.005$, **** $p < 0.001$.

Bmp2 Gene Regulation in the Aorta

Figure 4. *Klotho* RNA, BMP signaling levels and calcification in aorta from homozygous *Klotho* mutant mice. Aortas were isolated from control (C) mice with normal kidney function ($kl^{+/+}$ and $kl^{kl/+}$) or mice homozygous for the *Klotho* mutation with renal disease (KK) between 40 and 50 days of age. Average parameter values are presented with SEM. **A.** *Klotho* RNA levels normalized to actin RNA levels (males, n = 4 – 18; females, n = 3 – 8). *Bmp2* genotypes are indicated below each bar. **B.** BMP signaling levels as assessed by phosphorylated SMAD1/5/9(8) (pSMAD) levels normalized to total SMAD 1/5/9(8) (tSMAD) (males, n = 3 – 6; females, n = 5 – 7). All mice were wild type for *Bmp2* genotype. **C.** Representative western blot panels showing pSMAD1/5/9(8) and total SMAD1/5/9(8) levels, males (top) and females (bottom). **D.** Average calcium levels normalized to protein concentration (males, n = 9 – 18; females, n = 6 – 7). All experiments were repeated at least twice with similar results. * $p < 0.05$, **** $p < 0.001$. **E, F.** Male littermates were scanned with an Albira PET/CT Imaging System (Carestream, Rochester, NY) set at 45 kV, 400 μ A, and <35 μ m voxel size. Voxel intensities in the reconstructed images were evaluated and segmented with VivoQuant image analysis software (version 1.23, inviCRO LLC, Boston MA). The dashed white lines mark mineralized areas of the aortic sinus and ascending aorta present in the heart (**Ht**) of the *Klotho* mutant homozygote (**F**), but not in the heterozygous control $kl^{kl/+}$ (**E**).

Bmp2 Gene Regulation in the Aorta

579

580

581 **Figure 5. BMP signaling and calcium levels in aorta from *Klotho* mutant mice with *Bmp2***
 582 **mutations.** Aortas were isolated from mice bearing the wild type (Wt), heterozygous (+/ Δ), or
 583 homozygous UCS deletion ($\Delta\Delta$) *Bmp2* genotypes between 40 and 50 days of age. All mice were
 584 homozygous for the *Klotho* mutation with renal disease (*Kl^{kl/kl}*). Average parameter values are
 585 presented as averages with SEM. **A.** BMP signaling levels as assessed by phosphorylated
 586 SMAD1/5/9(8) (pSMAD) levels normalized to total SMAD 1/5/9(8) (tSMAD) (males, n = 3 – 6;
 587 females, n = 5 – 6). **B.** Representative western blot panels showing pSMAD1/5/9(8) and total
 588 SMAD1/5/9(8) levels, males (top) and females (bottom). **C.** Calcium levels normalized to protein
 589 concentration (males, n = 4 – 9; females, n = 5 – 8). All experiments were repeated at least
 590 twice with similar results. * $p < 0.05$.

591

Bmp2 Gene Regulation in the Aorta

593

594 **Figure 6. Body and heart weights from healthy control and *Klotho* mutant mice with**

595 ***Bmp2* mutations.** Body (A) and heart weights (B) from control mice with normal kidney function

596 (*Kl*^{+/+} and *Kl*^{kl/+}) or diseased mice homozygous for the *Klotho* mutation (*Kl*^{kl/kl}) bearing the wild

597 type (Wt), heterozygous (+/Δ), or homozygous UCS deletion (ΔΔ) *Bmp2* genotypes. Mice were

598 of age 47.1 ± 6 days. Heart weight was normalized to body weight in C. Parameter values are

599 presented as averages with SEM. The number of animals is indicated under the respective

600 bars. * *p* < 0.05, ** *p* < 0.01, ****p* < 0.005, **** *p* < 0.001.

601

Bmp2 Gene Regulation in the Aorta

Bmp2 Gene Regulation in the Aorta

Table 1. *Klotho* wild type and *Klotho* heterozygous mice are equivalent controls. Two-way ANOVA analysis was performed to confirm that there was no significant difference between *Klotho* wildtype ($K^{+/+}$) and *Klotho* mutant heterozygotes ($K^{kl/+}$) mice. Parameters compared included *Bmp2* RNA, pSMAD normalized to total SMAD, aortic calcium levels, body weight, heart weight, heart weight normalized to body weight and age at euthanasia. The average (Avg.), standard deviation (SD), and number of mice assayed (n) is shown.

Bmp2 Gene Regulation in the Aorta

	MALES							FEMALES					
Bmp2 genotype	Wt							Wt					
Klotho genotype	Wt			K+				Wt			K+		
Two-way Anova (p val)	0.29							0.21					
Parameters	Avg.	SD	n	Avg.	SD	n		Avg.	SD	n	Avg.	SD	n
Bmp2 RNA	0.72	0.24	7	0.78	0.46	16		0.57	0.09	5	0.89	0.31	6
pSMAD/Total SMAD	0.61	0.30	4	0.58	0.05	2		1.16	0.50	3	1.15	0.79	4
Calcium/Protein	42.1	21.8	7	51.2	21.2	11		64.8	23.4	5	53.2	14.5	3
Body Weight (g)	22.3	2.8	13	22.3	1.4	18		18.2	1.6	8	18.8	2.3	10
Heart Weight (g)	0.097	0.027	13	0.111	0.021	16		0.082	0.014	8	0.093	0.013	10
Heart / Body Weight	0.0044	0.0012	13	0.0049	0.0010	15		0.0045	0.0004	8	0.0049	0.0006	10
Age (d) at euthanasia	47.8	3.7	13	48.9	1.8	19		49.4	3.1	8	48.5	3.5	10
	MALES							FEMALES					
Bmp2 genotype	+/ Δ UCS							+/ Δ UCS					
Klotho genotype	Wt			K+				Wt			K+		
Two-way Anova (p val)	0.96							0.81					
Parameters	Avg.	SD	n	Avg.	SD	n		Avg.	SD	n	Avg.	SD	n
Bmp2 RNA	0.98	0.48	10	1.02	0.45	11		0.66	0.36	3	1.17	0.34	6
pSMAD/Total SMAD	1.00	0.30	5	1.00	0.06	3		1.00	0.61	5	1.00	0.33	3
Calcium/Protein	50.1	22.1	10	50.1	25.2	16		63.6	30.0	7	66.6	18.5	11
Body Weight (g)	22.4	3.0	16	21.9	3.3	26		19.5	3.2	15	19.3	2.2	22
Heart Weight (g)	0.11	0.02	15	0.11	0.02	23		0.09	0.02	11	0.10	0.02	13

Bmp2 Gene Regulation in the Aorta

Heart / Body Weight	0.0051	0.0007	15	0.0051	0.0012	22		0.0045	0.0010	11	0.0052	0.0008	12
Age (d) at euthanasia	46.9	5.8	16	46.9	6.2	27		49.1	2.6	15	48.5	5.9	23
	MALES							FEMALES					
Bmp2 genotype	ΔUCS/ΔUCS							ΔUCS/ΔUCS					
Klotho genotype	Wt			K+				Wt			K+		
Two-way Anova (p val)	0.26							0.52					
Parameters	Avg.	SD	n	Avg.	SD	n		Avg.	SD	n	Avg.	SD	n
Bmp2 RNA	1.71	0.16	4	1.48	0.58	8		2.12	0.75	2	0.65	0.26	8
pSMAD/Total SMAD	1.20	0.17	4	1.43	0.34	3		1.10	0.33	3	0.87	0.74	3
Calcium/Protein	36.1	11.0	8	45.6	18.3	11		67.9	27.7	4	60.5	30.7	9
Body Weight (g)	22.9	2.6	15	22.7	3.5	23		18.9	2.4	12	18.2	1.6	24
Heart Weight (g)	0.115	0.014	11	0.114	0.021	15		0.099	0.023	8	0.098	0.019	19
Heart / Body Weight	0.0051	0.0008	11	0.0052	0.0006	15		0.0052	0.0010	8	0.0054	0.0011	18
Age (d) at euthanasia	47.7	4.4	15	47.3	5.5	23		47.5	5.6	12	48.7	5.4	25

610