SCIENTIFIC REPORT: Controlling the Mitochondrial Protonmotive Force with Light to Impact Cellular Stress Resistance

Running title: Energizing Mitochondria with Light

Brandon J. Berry¹, Adam J. Trewin², Alexander S. Milliken¹, Aksana Baldzizhar², Andrea M. Amitrano^{3,4}, Minsoo Kim^{3,4}, Andrew P. Wojtovich^{1,2*}

¹University of Rochester Medical Center, Department of Pharmacology and Physiology, 575 Elmwood Ave., Rochester NY, 14642. Box 711/604. United States of America. Phone: 585-275-4613.

²University of Rochester Medical Center, Department of Anesthesiology and Perioperative Medicine, 575 Elmwood Ave., Rochester NY, 14642 Box 711/604. United States of America. Phone: 585-275-4613.

³Department of Pathology, University of Rochester Medical Center, Box 609, 601 Elmwood Ave., Rochester, NY 14642, USA

⁴Department of Microbiology and Immunology, University of Rochester Medical Center, Box 609, 601 Elmwood Ave., Rochester, NY 14642, USA

brandon_berry@urmc.rochester.edu

a.trewin@deakin.edu.au

alexander_milliken@urmc.rochester.edu

aksana_baldzizhar@urmc.rochester.edu

Andrea_Amitrano@urmc.rochester.edu

Minsoo_Kim@urmc.rochester.edu

*Corresponding author

andrew_wojtovich@urmc.rochester.edu

Keywords: metabolism, anoxia, hypoxia, uncoupling, ischemia reperfusion

FINAL CHARACTER COUNT: 25,817

ABSTRACT

Mitochondrial respiration generates an electrochemical proton gradient across the mitochondrial inner membrane called the protonmotive force (PMF) to drive diverse functions and make ATP. Current techniques to manipulate the PMF are limited to its dissipation; there is no precise, reversible method to increase the PMF. To address this issue, we used an optogenetic approach and engineered a mitochondria-targeted light-activated proton pumping protein we called mitochondria-ON (mtON) to selectively increase the PMF. Here, mtON increased the PMF light dose-dependently, supported ATP synthesis, increased resistance to mitochondrial toxins, and modulated energy-sensing behavior in *Caenorhabditis elegans*. Moreover, transient mtON activation during hypoxia prevented the well-characterized adaptive response of hypoxic preconditioning. Our novel optogenetic approach demonstrated that a decreased PMF is both necessary and sufficient for hypoxia-stimulated stress resistance. Our results show that optogenetic manipulation of the PMF is a powerful tool to modulate metabolic and cell signaling outcomes.

INTRODUCTION

Mitochondria generate an electrochemical proton gradient known as the protonmotive force (PMF). This consists of an electrical charge gradient, or membrane potential ($\Delta\psi_m$), and a pH gradient (ΔpH) that drives energy availability and controls diverse physiologic outputs (Garrido, Galluzzi et al., 2006, Rizzuto, De Stefani et al., 2012, Shadel & Horvath, 2015). The PMF is generated by proton pumping respiratory complexes of the electron transport chain (ETC) located in the mitochondrial inner membrane (IM). ETC dysfunction can lead to loss of PMF and a diverse range of pathologies (Berry, Trewin et al., 2018, Dingley, Polyak et al., 2010). For example, because the ETC consumes O_2 to establish the PMF, the PMF is

decreased under pathologic hypoxic conditions. The mechanistic link between acute changes in the PMF and downstream physiologic changes are poorly understood (Chalmers, Saunter et al., 2015, Santo-Domingo, Giacomello et al., 2013), and research is focused on developing new techniques to elucidate these pathways (Chalmers, Caldwell et al., 2012, Glancy, Hartnell et al., 2018).

Stroke is a common pathology in which cells undergo hypoxia and rapid reoxygenation that causes changes in the PMF and compromises mitochondrial functions. Changes in the PMF during hypoxia and reoxygenation influence cell-survival outcomes via mechanisms that are not fully understood (Murphy & Hartley, 2018, Solaini, Baracca et al., 2010, Yang, Mukda et al., 2018, Zhang, Trushin et al., 2016). Selectively increasing the PMF to distinguish cause and effect in hypoxic models is necessary to open new avenues of investigation that may reveal important metabolic changes that occur in stroke.

There are several techniques that allow experimental modulation of the PMF, but most are pharmacologic and therefore irreversible and not cell- or tissue-specific. Herein, we take a novel approach to overcome these barriers to precisely control the PMF by using optogenetics, adapting an approach used recently to dissipate the PMF (Ernst, Xu et al., 2019, Tkatch, Greotti et al., 2017). One family of widely-used optogenetic proteins are bacteriorhodopsin-related light-activated proton pumps. These proteins pump protons across membranes in response to specific wavelengths of light, and are often used to study physiology by modulating electrochemical gradients at the plasma membrane (Chow, Han et al., 2010, Husson, Liewald et al., 2012, Kandori, 2015). Only recently have precise optogenetic techniques been applied to compartmentalized cellular events using light-activated proteins targeted to organelles (Ernst, Xu et al., 2018, Rost, Schneider et al., 2015, Tkatch et al., 2017, Trewin, Bahr et al., 2019, Trewin, Berry et al., 2018a). Rather than using a non-specific cation channel to permeabilize the IM and dissipate the PMF, here we target the light-activated proton pump from the fungal organism *Leptosphaeria maculans* (Chow et al., 2010, Waschuk, Bezerra et al., 2005) to

mitochondria and selectively increase the PMF. We call this optogenetic tool mitochondria-ON (mtON) due to its ability to mimic the proton pumping activity of the ETC in response to light, independent of oxygen or substrate availability.

We validated mtON using the well-characterized genetic model organism, *C. elegans* (Butler, Ventura et al., 2010, Dingley et al., 2010, Tsang & Lemire, 2003). Using hypoxia and reoxygenation, we tested the hypothesis that hypoxia adaptation through preconditioning requires a decreased PMF. Our data demonstrate that transient loss of PMF during preconditioning is necessary and sufficient for resistance to hypoxia. By probing the evolutionarily conserved hypoxia adaptation response (Pena, Sherman et al., 2016, Wang, Lim et al., 2019, Wojtovich, Nadtochiy et al., 2012b, Wojtovich, Nadtochiy et al., 2013), we show that tools like mtON allow precise determination of cause and effect in physiologic models.

RESULTS & DISCUSSION

Light-activated proton pump mitochondria-ON (mtON) is expressed in mitochondria.

Using a ubiquitously-expressed gene promoter (*P_{eft-3}*), we directed expression of a light-activated proton pump to the mitochondrial IM in *C. elegans*. Mitochondrial localization was achieved by fusion of the proton pump to an N-terminal mitochondrial targeting sequence of the IMMT1 protein (Fischer, Igoudjil et al., 2011, John, Shang et al., 2005) in an orientation that allows proton pumping from the mitochondrial matrix towards the intermembrane space to increase the PMF in response to light (Fig. 1A). Using a C-terminal GFP fusion for subcellular visualization, and MitoTrackerTM CMXRos, we observed overlap of green and red fluorescence in *C. elegans* tissues, indicating the intended mitochondrial targeting (Fig. 1B & Fig. EV 1). We confirmed the expression of mtON in isolated mitochondrial preparations by immunoblot against GFP and observed a band at the predicted molecular weight of 82 kDa (Fig. 1C).

mtON activation increases the PMF.

Proton pumping activity of mtON requires the cofactor all trans-retinal (ATR) (Okazaki & Takagi, 2013, Sumii, Furutani et al., 2005, Waschuk et al., 2005). Because *C. elegans* do not produce ATR endogenously, exogenous supplementation is required for the light-activated proton pump to function (Kandori, 2015, Takahashi & Takagi, 2017). Thus, we were able to control for the expression of mtON (and the included GFP) in functional and non-functional forms, depending on whether ATR was supplemented, in addition to the use of light controls (Fig. EV 2A-D).

To test if mtON was capable of generating a PMF we measured the $\Delta\psi_m$ component using the indicator tetramethylrhodamine ethyl ester (TMRE) in isolated *C. elegans* mitochondria. TMRE is a fluorescent lipophilic cation that will accumulate in mitochondria proportionally with the $\Delta\psi_m$. At the quenching concentration of TMRE used, fluorescence is low at the $\Delta\psi_m$ of intact, energized isolated mitochondria. Upon loss of the $\Delta\psi_m$, TMRE redistributes to the extramitochondrial space resulting in increased fluorescence due to dequenching of TMRE (Fig. 2A). Under non-phosphorylating conditions in the absence of added substrate to fuel ETC activity, mtON activation was able to polarize the $\Delta\psi_m$ as much as ETC-driven respiration (Fig. 2B) light dose dependently (Fig. EV 3A). These data show that more photons result in increased polarization of the PMF, which is in line with the biophysical studies of the proton pump (Chow et al., 2010). While using the non-functional pump (no ATR), light had no effect on $\Delta\psi_m$ (Fig. 2B).

When protons are pumped out of mitochondria during respiration the matrix pH increases (Porcelli, Ghelli et al., 2005). Thus, we measured pH changes in the mitochondrial matrix in response to light to test this parameter as another readout of mtON activity. Using a ratiometric pH indicator, BCECF-AM, we found that matrix pH increased in response to mtON activation, mimicking the effect of succinate driven respiration (Fig. 2C&D). mtON-stimulated

changes in matrix pH and $\Delta\psi_m$ were sensitive to the protonophore FCCP, demonstrating that mtON polarizes the PMF, as measured by both the $\Delta\psi_m$ and the ΔpH .

Figure 1 - Light-activated proton pump mitochondria-ON (mtON) is expressed in mitochondria.

A Schematic depicting the targeting strategy to localize mtON to the mitochondrial inner membrane (IM). The electron transport chain (ETC) complexes generate the endogenous mitochondrial PMF by proton pumping, represented by the + and - across the IM. Mitochondrial ATP synthase utilizes the PMF to convert ADP to ATP. The N-terminal mitochondria target sequence from the IMMT1 protein shown in red; GFP shown in green. In response to light, mtON pumps protons from the mitochondrial matrix to the intermembrane space (IMS).

B Confocal images demonstrate overlap of GFP tagged mtON with MitoTrackerTM Red CMXRos stained *C. elegans* hypodermal mitochondria. Scale bar 10 μm .

C Immunoblot comparing the cytosolic supernatant and the mitochondria-enriched pellet of isolation fractions. GFP-tagged mtON migrates at the predicted molecular weight of 82 kDa accounting for the mitochondria-target sequence, the proton pump, and GFP. mtON is observed only in the mitochondrial fraction compared to marker proteins HSP60 (mitochondria) and actin (cytosol). All blots are from the same lanes on one membrane.

Figure 2 - mtON activation increases the PMF.

A Representative TMRE fluorescence traces in arbitrary units (a.u.) before and after $\Delta\psi_m$ dissipation with FCCP. Dashed lines indicate where FCCP was added. +/- ATR traces were performed in the absence of succinate. mtON activation was continuous throughout the traces. Light green trace is from mitochondria with ATR, gray traces are without.

B Quantification of change in fluorescence (ΔF) normalized to the maximum change in fluorescence given by succinate respiration ($\Delta F_{\max}$) for each mitochondrial preparation. Data are from the maximum light dose in Fig. EV 3A. Two-way ANOVA with Sidak's test for multiple comparisons, * $p = 0.0469$, +ATR succinate vs. +ATR +light $p = 0.9978$, $n = 6$ mitochondrial isolations. Bars are means $\pm$ SEM.

C Representative BCECF-AM 490/440 nm ratio trace. Ratio of 545 nm fluorescence intensity at either 440 or 490 nm excitation. Dashed lines are where light or FCCP treatment occurred. Light green trace is from mitochondria with ATR, gray traces are without.

D Quantification of change in BCECF-AM fluorescence ratio normalized to maximum change given by succinate matrix alkalization. Two-way ANOVA with Sidak's test for multiple comparisons, -ATR succinate vs. -ATR, +light * $p = 0.0212$, +ATR succinate vs. +ATR light $p = 0.999$, +ATR succinate vs. +ATR, -light $p = 0.0237$, +ATR, +light vs. +ATR, -light $p = 0.0474$, $n = 3$ mitochondrial isolations.

E ATP levels normalized to total ATP synthesis given by succinate respiration. Data from Fig. EV 3B. Succinate data shown for comparison after normalization. Two-way ANOVA with Sidak's test for multiple comparisons, * $p = 0.0011$, +ATR succinate vs +ATR light $p = 0.5680$, $n = 3 - 7$ assays across two mitochondrial isolations.

F O_2 required to consume 50 nmoles of ADP after mtON activation. Data are the maximum illumination from Fig. EV 4D using time-matched dark conditions, One-way ANOVA * $p = 0.013$ (-ATR, -light vs. +ATR, +light $p = 0.010$. -ATR, +light vs. +ATR, +light $p = 0.013$.) -ATR, -light $n = 4$, rest $n = 5$ mitochondrial preparations.

mtON increases ATP synthesis without respiration.

To further test if mtON could increase energetics, we measured ATP levels from isolated mitochondria that were supplied with ADP to phosphorylate, since ATP levels are highly regulated *in vivo* (Balaban, Kantor et al., 1986, Glancy, Hartnell et al., 2015, Viola & Hool, 2017, Wang, Zhang et al., 2017). As expected, mtON activation increased ATP levels (Fig. 2E) light-dose dependently (Fig. EV 3B), similar to the $\Delta\psi_m$ results. In mitochondria, the amount of ADP converted to ATP is reliant on O_2 consumption by the ETC (Brand & Nicholls, 2011). We tested if we could drive ADP conversion to ATP by generating a PMF with mtON, bypassing the requirement for O_2 consumption. Respiration was consistent across all control conditions (Fig. EV 4A&B), indicating no baseline differences in mitochondrial quality. Activation of mtON decreased the amount of O_2 required to phosphorylate a given amount of ADP (Fig. 2F & Fig. EV 4C) light dose dependently (Fig. EV 4D), indicating mtON-driven ATP production does not rely on O_2 consumption. Until now, interventions to increase the PMF have involved fueling the ETC, where here we show mtON generates a PMF independent of O_2 consumption or metabolic

substrates to provide electrons. mtON's ability to augment the PMF via optic control opens a new avenue for discovery in metabolic research where complex interactions between physiology and bioenergetics often obscure molecular mechanisms.

mtON increases survival during acute ETC dysfunction.

Given that mtON activity decreased reliance on the ETC *in vitro*, we tested if mtON could compensate for acute ETC dysfunction *in vivo*. We exposed *C. elegans* to toxic inhibitors of specific sites within ETC complexes and scored their survival in response to mtON activation. Light alone had no effect on survival after inhibitor exposure in all cases (Fig. 3A-C). For transgenic animals, when exposed to the complex I inhibitor rotenone, mtON was able to improve survival (Fig. 3A). Rotenone toxicity is mediated chiefly through oxidative damage (Ishiguro, Yasuda et al., 2001, Schmeisser, Priebe et al., 2013), and ATR alone exhibited a protective effect under these conditions, possibly due to its antioxidant properties (Lee, Casadesus et al., 2009, Palace, Khaper et al., 1999, Siddikuzzaman & Grace, 2013). This effect was also present in wild type control experiments (Fig. EV 5A). However, the effect of mtON was significantly greater than the ATR effect (Fig. EV 5A, one-way ANOVA, $p = 0.0002$), with no effect in wild type animals (Fig. EV 5A&B). Activation of mtON also protected animals exposed to antimycin A, a complex III inhibitor (Ishiguro et al., 2001) (Fig. 3B), and azide, a complex IV inhibitor (Fig. 3C). Unlike with rotenone treatment, ATR alone did not mitigate antimycin A or azide toxicity. Overall, these data indicate that mtON can partially overcome inhibited ETC activity in whole organisms.

Figure 3 - mtON improves survival during acute ETC dysfunction.

A Day 1 adult animals were exposed to 50 μ M rotenone (ETC complex I inhibitor) for 5 hours and survival was scored. Illumination was continuous throughout toxin exposure (see methods). ATR alone was protective (-ATR, -light vs. +ATR, -light $p = 0.016$.) The effect of mtON activation was greater than the ATR alone effect * $p = 0.01$. (-ATR, -light vs +ATR, +light $p = 0.0002$. -ATR, +light vs. +ATR, +light $p = 0.0009$. $n = 3$ plates each condition).

B Animals were exposed to 50 μ M antimycin A (complex III inhibitor) and survival was scored 18 hours later. * $p = 0.02$ (-ATR, -light vs +ATR, +light $p = 0.03$, -ATR, +light vs. +ATR, +light $p = 0.01$. $n = 5$ plates each condition).

C Animals were exposed to 0.25 M azide (complex IV inhibitor) for 1 hour and scored for survival 1 hour after recovering. * $p = 0.0002$. (-ATR, -light vs +ATR, +light $p < 0.0001$, -ATR, +light vs. +ATR, +light $p < 0.0001$. $n = 3$ plates each condition). Statistics are one-way ANOVA with Tukey's post hoc test. Bars are means $\pm$ SEM.

mtON affects whole-animal energy sensing.

We next asked whether mtON could affect metabolic signaling. One way organisms sense energy availability and preserve energy homeostasis is through the AMP-activated protein kinase (AMPK) (Hardie, Ross et al., 2012). In *C. elegans* the *aak-2* gene encodes the catalytic subunit ortholog of mammalian AMPK α 2. Mutation of *aak-2* has well-characterized phenotypic outputs linked to energy availability, and serves as a regulator of whole-organism energy sensing (Apfeld, O'Connor et al., 2004, Cunningham, Bouagnon et al., 2014). We hypothesized that mtON activity would signal energy availability and decrease phosphorylation of AMPK, its activated state. As such, we exposed animals expressing mtON to light in the absence of food, where AMPK should be phosphorylated, and immunoblotted against phosphorylated AMPK. We found that removal from food increases AMPK phosphorylation as expected, and mtON activation prevented this phosphorylation (Fig. 4A). In *C. elegans*, the

AMPK homologue AAK-2 regulates a behavioral response to food availability (Lee, Cho et al., 2008), where in the absence of food animals will increase locomotion to search for food. Based on our phosphorylation results, we hypothesized that mtON activation would attenuate the energy deficit signal for animals off food, suppressing their movement, and thus mimicking the *aak-2* loss-of-function phenotype. We first confirmed that *aak-2* loss-of-function mutant animals had decreased locomotion in the absence of food compared to wild type (Fig. 4B). In response to acute mtON activation in wild type background, we observed decreased locomotion in the absence of food (Fig. 4C). This effect was reversed using the AMPK activator 5-aminoimidazole-4-carboxamide ribonucleotide (AICAR), indicating the mtON suppression of locomotion was rescued by AMPK activity (Fig. 4C). These data suggest that mtON activity can modify downstream metabolic signaling. As changing the PMF affects many aspects of metabolism that are able to activate AMPK, the exact mechanism of AMPK activation in our experiments is unclear. For example, AMPK is activated by increased energy demand, but also redox signaling (Garcia & Shaw, 2017), and calcium signaling (Hardie, 2011). Upon further characterization in these contexts mtON could be used to modulate the many cellular processes that AMPK regulates (Hinchy, Gruszczuk et al., 2018, Jeon, 2016, Mihaylova & Shaw, 2011, Trewin, Berry et al., 2018b).

Figure 4 - mtON affects whole-animal energy sensing.

A Top: Immunoblot assessing the effect of mtON activation on AMPK phosphorylation status. Top bands (~62 kDa) are phosphorylated AMPK signal, and bottom bands (~43 kDa) are actin signal. Image is from the same membrane cut to separately probe for phosphorylated AMPK (pAMPK) and actin. Bottom: Densitometry analysis showing decreased pAMPK to actin ratio, as there is no known antibody directed against total AMPK in *C. elegans*. Phosphorylation increases in the absence of food, but low phosphorylation is preserved when mtON is activated, two-way ANOVA with Sidak's test for multiple comparisons, * $p = 0.0203$, $n = 3$ -4 biological replicates.

B Locomotion was assessed by counting body bends per minute. Wild type animals were compared to *aak-2(ok524)* mutant animals. 2-sample, 2-tailed unpaired t-test * $p < 0.0001$, wild type $n = 35$, *aak-2* $n = 39$ animals across at least 3 days. Bars are means $\pm$ standard deviation.

C Locomotion in response to mtON activation. Illumination was continuous throughout body bends measurement (see methods). For AAK-2 activation, animals were exposed to 1 mM AICAR for 4 hours before body bends measurement. One-way ANOVA with Tukey's post hoc test, -ATR, +light vs +ATR, +light * $p < 0.0001$, +ATR, +light vs AICAR * $p < 0.0001$. $n =$ (in order, left to right) 36, 39, 37, 46, 36 animals across at least 3 days. Bars are means $\pm$ standard deviation.

mtON inhibits hypoxia-adaptation.

We next tested if mtON could impact stress resistance. Hypoxia and reoxygenation (HR) is a pathologic insult that involves changes in the PMF that can contribute to injury and survival (Chouchani, Pell et al., 2014, Michiels, 2004, Murphy & Hartley, 2018, Sanderson, Reynolds et al., 2013, Solaini et al., 2010, Xu, Wang et al., 2001, Yang et al., 2018) depending on the context and degree of (de)polarization. In addition, the phenomenon of preconditioning (PC) is effectively modeled with hypoxia in *C. elegans* (Dasgupta, Patel et al., 2007, Hayakawa, Kato et

al., 2011, Jia & Crowder, 2008, Pena et al., 2016, Wojtovich, DiStefano et al., 2012a), where a short period of hypoxia protects against a later pathologic exposure (Fig. 5B). A decreased PMF before HR also mediates a protective effect (Brennan, Berry et al., 2006, Brennan, Southworth et al., 2006, Ozcan, Palmeri et al., 2013), but mechanisms are debated (Shabalina & Nedergaard, 2011), as understanding is complicated by lack of temporal control when using pharmacologic approaches. In support of this, the protonophore FCCP, which can dissipate the PMF, protected wild type animals against HR (Fig. 5C), suggesting hypoxia resistance may be mediated by a decreased PMF. We combined these lines of evidence and hypothesized that increasing the PMF during PC would reverse the protection against hypoxia afforded by PC. To test this hypothesis we activated mtON selectively during PC (Fig. 5A). To quantify protection, we subtracted the percent survival after pathologic hypoxia from the percent survival after an intervention, giving the percent alive above baseline (Fig. 5C). Further highlighting the importance of our control conditions, we found that light on its own was as protective as PC in mtON expressing animals, and the combination of light and PC was additive (Fig. 5D). However, activation of mtON during PC decreased the protection (Fig. 5D), supporting our hypothesis that some of the protective effect of PC is mediated by decreasing the PMF. In effect, mtON counteracts a PC-induced decrease in the PMF by restoring it to normal levels. These data suggest that loss of PMF during PC is necessary for hypoxic adaptation. The FCCP data corroborate our mtON findings, and demonstrate the sufficiency of PMF loss to protect against hypoxia. Using mtON only during PC to boost the PMF provided temporal control that had not been tested before, allowing us to determine when PMF changes have physiologic effects in HR.

Figure 5 - mtON inhibits hypoxia-adaptation.

A Schematic of HR experiments. Top shows control hypoxia and reoxygenation. Below is the hypoxic preconditioning (PC) protocol, survival for these two timelines shown in panel B. Third from the top shows FCCP treatment protocol, calculated protection shown in panel C. Bottom two show light treatment protocol + and - PC, calculated protection data from these shown in panel D.

B Survival after HR in day 1 adult animals is shown with hypoxic preconditioning (PC, represented by diagonal stripes) 2-sample 2-tailed unpaired t-test, ctrl vs. PC * $p = 0.0035$, $n = 4$, each averaged from 3 technical replicates. Bars are means $\pm$ SEM.

C Protection (%) is percent survival minus percent survival of control condition. PC data calculated from panel B. FCCP final concentrations 0.001, 0.01, 0.1, 1 nM. One-way ANOVA with Tukey's post hoc test, PC vs. 0.001 nM FCCP, $p = 0.0012$, 0.001 nM FCCP vs 0.01 FCCP, $p = 0.0142$, $n = 3 - 4$ independent experiments, each averaged from 3 technical replicates. Bars are means $\pm$ SEM.

D Illumination was continuous throughout PC alone, control illumination was for the same duration under normoxic conditions (see methods & panel A). Two-way ANOVA comparing +ATR vs -ATR in each group. * $p = 0.016$, $n =$ (in order, left to right) 11,11,4,4,6,6 independent experiments, each averaged from 3 technical replicates. Bars are means $\pm$ SEM.

Our findings that protection afforded by PC relies on the transient loss of PMF suggest that a decreased PMF is sufficient to elicit stress resistance at a later period. This implies that interventions at the level of the PMF alone can impact cellular responses to stress. Combining the mtON approach with tissue-specific gene promoters, precise spatiotemporal control could be achieved in discerning the effects of mitochondrial function across tissues in whole organisms. Taken together, mtON appears to be a useful tool to enable discrimination of cause and effect in complex (patho)physiologic contexts that involve modest changes in mitochondrial function and metabolism. Defining when changes in the PMF are adaptive and when they are detrimental may advance our understanding of many pathologies and inform novel therapeutic strategies that target mitochondrial function. Our findings from this approach suggest the PMF is a keystone of metabolism that senses cellular stress and elicits appropriate adaptive responses to maintain homeostasis.

MATERIALS & METHODS

Molecular biology

The light-activated proton pump from *Leptosphaeria maculans* (Mac) fused to eGFP was amplified from plasmid DNA pFCK-Mac-GFP, a gift from Edward Boyden (Addgene plasmid #22223 (Chow et al., 2010)) (forward amplification primer: ACACCTGCAGGCTTGATCGTGGACCAGTTCGA, reverse amplification primer: CACAGCGGCCGCTTACTTGTACAGCTCGTCCA). The N-terminal 187 amino acids of the *Immt1* gene was amplified by PCR from mouse cDNA (forward amplification primer: ACAACCGGTAAAAATGCTGCGGGCCTGTCAGTT, reverse amplification primer: CACCCTGCAGGTTCTCTGTGGTTTCAGACG). The ubiquitously expressed gene promoter *Peft-3* (also known as *Peef-1A.1*) was amplified by PCR from pDD162 (forward amplification

primer: AACAAAGCTTGCACCTTTGGTCTTTTA, reverse amplification primer: ACATCTAGAGAGCAAAGTGTTTCCCA). The body wall muscle promoter *Pmyo-3* was PCR amplified from pDJ16 (forward amplification primer: ACAGCTAGCTGTGTGTGATTGCT, reverse amplification primer: ACAACCGGTGCGGCAATTCTAGATGG). PCR fragments were ligated into pFH6.II (pPD95.81 with a modified multi-cloning site) for *C. elegans* expression using restriction digest cloning. Resulting plasmids were pBJB20 (*Peft-3::IMMT1*(N-terminal 187 amino acids)::Mac::GFP), pBJB16 (*Pmyo-3::IMMT1*(N-terminal 187 amino acids)::Mac::GFP), Sanger sequencing was used to confirm plasmid sequences (Eurofins Genomics). Animals were transformed by plasmid DNA microinjection with *pha-1*(+) selection in a *pha-1(e2123ts)* temperature-sensitive mutant strain, where transgenic animals were selected for growth at 20° C.

C. elegans strains growth and maintenance

All animals were maintained at 20° C on nematode growth medium (NGM) seeded with OP50 *E. Coli*. Young adult hermaphrodite animals were used for all experiments. Transgenic strains were generated by plasmid DNA microinjection as described (Mello, Kramer et al., 1991). For a complete strain list see Table 1. Where indicated, OP50 was supplemented with all-trans-Retinal to a final concentration of 100 µM on seeded NGM plates. Animals were cultured on ATR-containing plates for at least one generation.

Fluorescence microscopy

Images were taken on a FV1000 Olympus laser scanning confocal microscope using a 60x oil objective (Olympus, N.A. 1.42). Diode laser illumination was at 561 nm for red fluorescence and 488 nm for green fluorescence. Where indicated, animals were stained with 10 µM MitoTracker™ Red CMXRos for 4 hours. MitoTracker™ stain was dissolved in DMSO, diluted in

M9 media (22 mM KH_2PO_4 , 42 mM Na_2HPO_4 , 86 mM NaCl, 1 mM MgSO_4 , pH 7) and added to OP50 seeded NGM plates (DMSO < 0.02% final) and allowed to dry. Line scan pixel intensity was performed using ImageJ software.

Mitochondria isolation

C. elegans mitochondria were isolated from day 1 adults as previously described (Wojtovich et al., 2012a) using differential centrifugation in mannitol and sucrose-based media. Animals from three 15 cm culture plates were transferred into 50 mL of M9 media in a conical tube and allowed to settle by gravity on ice. Pelleted animals were rinsed with ice-cold M9 twice, and once with ice-cold mitochondrial isolation media (220 mM mannitol, 70 mM sucrose, 5 mM MOPS, 2 mM EGTA, pH 7.4) with 0.04% BSA. After settling by gravity, supernatant was removed and worms were transferred to an ice-cold mortar containing ~2 g of pure sea sand per 1 mL of animals. Animals were ground with an ice-cold pestle for 1 minute and extracted from the sand using mitochondrial isolation media and transferred to a 10 mL conical tube. The suspension was then transferred to an ice-cold glass Dounce homogenizer and homogenized with 40 strokes. The homogenate was centrifuged at 600 g for 5 minutes. Supernatant was transferred to a new tube and centrifuged at 700 g for 10 minutes. The pellet was resuspended in 1 mL of mitochondrial isolation media without BSA, which was centrifuged at 7000 g for 5 minutes. The pellet was finally resuspended in 50 μL of mitochondrial isolation media without BSA. Protein concentration was quantified using the Folin-phenol method.

Light sources

Illumination sources included a 580 nm Quantum SpectraLife LED Hybrid lamp by Quantum Devices, Barneveld WI, USA (abbreviated Quantum LED), a 540-600 nm GYX module, X-Cite LED1 by Excelitas, Waltham MA, USA (abbreviated XCite LED), and a 540-580 nm excitation filter MVX10 Fluorescence MacroZoom dissecting microscope by Olympus (abbreviated MVX).

Light intensities are indicated for each experimental condition and were determined with a calibrated thermopile detector (818P-010-12, Newport Corporation, Irvine, CA) and optical power meter (1916-R, Newport Corporation).

Immunoblotting

Synchronized young adult animals were exposed to 1 Hz light (Quantum LED, 0.02 mW/mm²) for 4 hours, and immediately harvested with ice-cold M9 media and centrifuged at 1,000 x g for 1 minute. Animals were ground by plastic pestle disruption in lysis buffer (20mM Tris-HCl, 100 mM NaCl, 1mM EDTA, 1mM DTT, 10% glycerol, 0.1% SDS, pH 7.6, 1X Halt™ protease inhibitor cocktail, Thermo78429) and diluted 1:1 in sample loading buffer (100 mM Tris HCl, 10% v/v glycerol, 10% SDS, 0.2% w/v bromophenol blue, 2% v/v β-mercaptoethanol). These samples were heated at 95 °C for 5 minutes. Isolated mitochondrial samples were prepared as described above and diluted 1:1 in sample loading buffer with 1% SDS. 30 µg samples were loaded to 7.5% polyacrylamide gels and separated by SDS-PAGE. Proteins were transferred to nitrocellulose membranes and blocked in 5% nonfat milk/TBST (50 mM Tris, 150 mM NaCl, 0.05% Tween 20, pH 8.0) for 1 h at room temperature. Membranes were incubated at 4°C in primary antibodies diluted 1:1000 in 5% bovine serum albumin: anti-GFP (ClonTech Living Colours #ab632375), anti-HSP60 (Department of Biology, Iowa City, IA 52242. Developmental Studies Hybridoma Bank, University of Iowa Department of Biology, Iowa City, IA 52242), (Cell Signaling, #4188), anti-Actin (Abcam #ab14128), and 1:10,000 anti-phospho-AMPKα). Membranes were washed in TBST and incubated in horseradish peroxidase-conjugated secondary antibodies: anti-rabbit IgG (Cell Signaling #7074S) or anti-mouse IgG (Thermo Scientific #32430, lot #RF234708) for 1 hour at room temperature. Proteins were visualized using ECL (Clarity Western ECL Substrate, Bio Rad) by chemiluminescence (ChemiDoc, Bio Rad). Densitometry was performed using Image Lab software (version 5.2.1).

Mitochondrial membrane potential measurement

Isolated mitochondria at 0.5 mg/mL were stirred in mitochondrial respiration buffer (MRB: 120 mM KCl, 25 mM sucrose, 5 mM MgCl₂, 5 mM KH₂PO₄, 1 mM EGTA, 10 mM HEPES, 1 mg/mL FF-BSA, pH 7.35) at 25° C in the presence of 2 μM rotenone, and 5 mM succinate where indicated. 300 nM tetramethylrhodamine, ethyl ester (TMRE, ThermoFisher, T669) was added to observe mitochondrial membrane potential in quench mode. Under quenching conditions TMRE fluorescence is low in the presence of a $\Delta\psi_m$. Upon addition of a protonophore (e.g. FCCP), TMRE will exit mitochondria and dequench and increase total fluorescence (Chouchani et al., 2014, Perry, Norman et al., 2011). TMRE signal was measured by Cary Eclipse Fluorescence Spectrophotometer (Agilent Technologies) using a 335-620 nm excitation filter and a 550-1100 nm emission. Illumination was performed continuously throughout all measurements (555 nm, 0.0016 mW/mm²). Increasing illumination time exposed mitochondria to more photons (calculated as fluence, J/cm²). After stable baseline measurements with or without succinate, 2 μM FCCP was added to completely depolarize mitochondria. The average fluorescence intensity after addition of 2 μM FCCP (maximum fluorescence in quench mode) was subtracted from the starting test condition to give a change in fluorescence corresponding to changes in $\Delta\psi_m$ (ΔF for conditions without succinate, and $\Delta F_{\max}$ for conditions with succinate). To represent polarization of the $\Delta\psi_m$, we used the ratio of change in fluorescence (ΔF , FCCP fluorescence minus experimental fluorescence) to the maximum change in fluorescence generated by succinate-driven $\Delta\psi_m$ ($\Delta F_{\max}$).

Mitochondrial matrix pH measurement

The ratiometric pH indicator BCECF-AM (ThermoFisher, B1170) was used to measure pH changes in the mitochondrial matrix (Aldakkak, Stowe et al., 2010) in response to succinate

respiration or mtON activation. Isolated mitochondria (~200 μ L per isolation) were incubated at room temperature with 50 μ M BCECF-AM for 10 minutes with periodic mixing. Mitochondria were then pelleted at 7000 g for 5 minutes at 4° C, isolation media replaced and pelleted again to remove extramitochondrial BCECF-AM. Isolated mitochondrial suspensions were then assayed under the same conditions as in the mitochondrial membrane potential measurements described above. Ratiometric fluorescent signal was measured by Cary Eclipse Fluorescence Spectrophotometer (Agilent Technologies) using 440 and 490 nm excitation wavelengths and 545 nm emission. The fluorescence intensity ratio at 545 nm of 490/440 nm excitation wavelengths was used to represent pH changes in the mitochondrial matrix. Light treatment was 0.16 J/cm² (XCite LED, 0.02 mW/mm²), and 2 μ M FCCP was used at the end of each trace to establish baseline signal.

ATP measurement

Relative ATP levels were determined in isolated mitochondria using a luciferase bioluminescence kit according to manufacturer's instructions (Invitrogen™ Molecular Probes™, A22066). Mitochondria were stirred in MRB at 0.5 mg/mL with 1 mg/mL fat free BSA, 600 μ M ADP, 2 μ M rotenone. 5 mM succinate for a control for maximum ATP level, and 0.001 mg/mL oligomycin A was used as a zero ATP synthesis control. Mitochondrial suspensions were immediately frozen with liquid nitrogen after 1, 5, or 10 minutes light exposure (XCite LED, 0.02 mW/mm²). Samples were then thawed on ice, centrifuged at 14,800 g, and supernatant was collected and run at 1:100 dilution in MRB in the luminescence assay. Oligomycin A control values were subtracted from experimental reads, and data were then normalized to luminescent signal from succinate control samples (complete ADP conversion confirmed by monitoring O₂ consumption rate transitions).

Mitochondrial O₂ consumption

O₂ consumption was measured using a Clark-type O₂ electrode (S1 electrode disc, DW2/2 electrode chamber and Oxy-Lab control unit, Hansatech Instruments, Norfolk UK) at 25° C. Isolated mitochondria were stirred in MRB at 1 mg/mL with 1 mg/mL fat free BSA. Substrates and inhibitors were added by syringe port (100 μM ADP, 2 μM rotenone, 5 mM succinate). Given excess succinate as substrate for ETC respiration, we measured the amount of O₂ required to convert 50 nmol of ADP to ATP and established a baseline for comparison. To test mtON activity, we illuminated mitochondria as in the ATP measurement, for 1, 5, or 10 minutes in the presence of ADP without succinate to allow for mtON conversion of ADP to ATP. Any remaining ADP was then converted using O₂-dependent ETC respiration upon succinate addition (Fig. EV4C). Slopes were calculated from plots of O₂ concentration versus time to give rates of O₂ consumption during ADP respiration, ADP+succinate respiration, and respiration after ADP had been entirely consumed. The intersections of these three rates were used to calculate total amount of O₂ consumed during ADP+succinate respiration. Light activation of mitochondria (XCite LED, 0.02 mW/mm²) during ADP respiration alone was carried out for differing lengths of time to test the ability of mtON to drive ADP consumption before succinate was added. Dark control was 10 minutes of no illumination before addition of succinate.

ETC inhibitor assays

Experiments were performed on at least three separate days, using 15-100 young adult animals per plate. Seeded plates were supplemented with rotenone (50 μM final concentration), or antimycin A (50 μM final concentration) 24 hours before animals were transferred onto them. For azide toxicity animals were placed in M9 buffer with 250 mM azide. Control plates were kept in the dark and experimental plates were exposed to 1 Hz light (Quantum LED, 0.02 mW/mm²) for the duration of the experiment. For rotenone, surviving animals were scored after 5 hours. For antimycin A, animals were scored 16 hours after exposure. For azide, animals were exposed for one hour in M9 and allowed to recover for 1 hour on a seeded culture plate, then

survival was scored. Azide experimental plates were exposed to light (XCite LED, 0.19 mW/mm²) for the duration of azide treatment and recovery. For all toxins, animals that were moving or those that moved in response to a light touch to the head were scored as alive.

Locomotion assay

Locomotion was scored by counting the number of body bends in 15 seconds immediately after being transferred off OP50 food (Sawin, Ranganathan et al., 2000) (n = 36-46 animals scored on at least 2 separate days). 1 body bend was scored as a deflection of direction of motion of the posterior pharyngeal bulb (Tsalik & Hobert, 2003). For AMPK activation, animals were placed on plates containing 1 mM AICAR 4 hours before counting body bends. AICAR was dissolved in M9 buffer, added directly onto OP50 seeded plates and allowed to dry. Illumination was continuous through measurements (MVX, 0.265 mW/mm²).

Hypoxia and reoxygenation

Experiments were carried out using a hypoxic chamber (Coy Lab Products, 5%/95% H₂/N₂ gas, palladium catalyst) at 26° C with 50-100 animals per plate. O₂ was less than 0.01%. Hypoxic preconditioning (PC) duration was 4 hours, and control animals were incubated at 26° C in room air for the same time. 1 Hz illumination (Quantum LED, 0.02 mW/mm²) was carried out during PC period alone. Hypoxic exposure was for 18.5 hours, 21 hours after PC. 24 hours after hypoxia exposure, animals that were moving or those that moved in response to a light touch to the head were scored as alive. Data from days where PC was at least 15% effective for both +ATR and -ATR were used. Animals supplemented with ATR were allowed to lay eggs onto plates with no ATR that were subsequently used for HR experiments to minimize confounding effects of ATR. ATR from the parent animal was sufficient to provide active mtON in progeny as tested by the azide toxin assay described above. Only progeny from parents supplemented with ATR were significantly protected against azide upon illumination (tested by One-way ANOVA, -

ATR, -light [17.3% survival] vs. +ATR, +light [66.8% survival] $p = 0.0054$, -ATR, +light [32.5% survival] vs. +ATR, +light $p = 0.0395$, +ATR, -light [27.5% survival] vs. +ATR, +light $p = 0.0199$, $DF = 11$, $F = 8.94$). PC experiments were represented as protection (%), where baseline survival was subtracted to correct for protection from ATR (-ATR survival: $34.4 \pm 14.4\%$, +ATR survival: $49.7 \pm 20.3\%$, 2 sample 2-tailed paired t-test, $p = 0.033$, $n = 12$ independent experiments). FCCP dissolved in ethanol was deposited onto seeded NGM plates to the final concentrations indicated in the figure legend (Fig. 5C).

Statistics

One-way ANOVA was used when comparing only our four experimental conditions (Fig. EV2). Two-way ANOVA was used when comparing our conditions with other variables, such as +/- succinate. Shapiro-Wilkes normality tests were used to determine whether parametric or non-parametric tests should be used. See figure legends for detailed statistical information and post-hoc tests.

DATA AVAILABILITY

All data are presented in the manuscript, and raw files will be made available upon request.

ACKNOWLEDGEMENTS

Work in the laboratory of A.P.W. is supported by a grant from National Institutes of Health (R01 NS092558) and institutional funds from the University of Rochester; B.J.B. is supported by an American Heart Association Predoctoral Fellowship (18PRE33990054) and an Institutional Ruth L. Kirschstein National Research Service Award (NIH T32 GM068411). A.J.T current address: Institute for Physical Activity and Nutrition (IPAN), Deakin University, Burwood, Australia. We thank the members of the mitochondrial research groups at University of Rochester Medical

Center for helpful discussions, suggestions, and guidance. Some strains were provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440).

AUTHOR CONTRIBUTIONS

B.J.B. and A.P.W. designed the research. B.J.B., A.J.T., A.S.M., A.B. and A.M.A. performed experiments. M.K. provided critical review of the results and manuscript. B.J.B. analyzed the data and wrote the manuscript. All authors approved the final version of the manuscript.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

References

- Aldakkak M, Stowe DF, Cheng Q, Kwok WM, Camara AK (2010) Mitochondrial matrix K⁺ flux independent of large-conductance Ca²⁺-activated K⁺ channel opening. *Am J Physiol Cell Physiol* 298: C530-41
- Apfeld J, O'Connor G, McDonagh T, DiStefano PS, Curtis R (2004) The AMP-activated protein kinase AAK-2 links energy levels and insulin-like signals to lifespan in *C. elegans*. *Genes Dev* 18: 3004-9
- Balaban RS, Kantor HL, Katz LA, Briggs RW (1986) Relation between work and phosphate metabolite in the in vivo paced mammalian heart. *Science* 232: 1121-3
- Berry BJ, Trewin AJ, Amitrano AM, Kim M, Wojtovich AP (2018) Use the Protonmotive Force: Mitochondrial Uncoupling and Reactive Oxygen Species. *J Mol Biol*
- Brand MD, Nicholls DG (2011) Assessing mitochondrial dysfunction in cells. *Biochem J* 435: 297-312
- Brennan JP, Berry RG, Baghai M, Duchon MR, Shattock MJ (2006) FCCP is cardioprotective at concentrations that cause mitochondrial oxidation without detectable depolarisation. *Cardiovasc Res* 72: 322-30
- Brennan JP, Southworth R, Medina RA, Davidson SM, Duchon MR, Shattock MJ (2006) Mitochondrial uncoupling, with low concentration FCCP, induces ROS-dependent cardioprotection independent of KATP channel activation. *Cardiovasc Res* 72: 313-21
- Butler JA, Ventura N, Johnson TE, Rea SL (2010) Long-lived mitochondrial (Mit) mutants of *Caenorhabditis elegans* utilize a novel metabolism. *FASEB J* 24: 4977-88

Chalmers S, Caldwell ST, Quin C, Prime TA, James AM, Cairns AG, Murphy MP, McCarron JG, Hartley RC (2012) Selective uncoupling of individual mitochondria within a cell using a mitochondria-targeted photoactivated protonophore. *J Am Chem Soc* 134: 758-61

Chalmers S, Saunter CD, Girkin JM, McCarron JG (2015) Flicker-assisted localization microscopy reveals altered mitochondrial architecture in hypertension. *Sci Rep* 5: 16875

Chouchani ET, Pell VR, Gaude E, Aksentijevic D, Sundier SY, Robb EL, Logan A, Nadtochiy SM, Ord ENJ, Smith AC, Eyassu F, Shirley R, Hu CH, Dare AJ, James AM, Rogatti S, Hartley RC, Eaton S, Costa ASH, Brookes PS et al. (2014) Ischaemic accumulation of succinate controls reperfusion injury through mitochondrial ROS. *Nature* 515: 431-435

Chow BY, Han X, Dobry AS, Qian X, Chuong AS, Li M, Henninger MA, Belfort GM, Lin Y, Monahan PE, Boyden ES (2010) High-performance genetically targetable optical neural silencing by light-driven proton pumps. *Nature* 463: 98-102

Cunningham KA, Bouagnon AD, Barros AG, Lin L, Malard L, Romano-Silva MA, Ashrafi K (2014) Loss of a neural AMP-activated kinase mimics the effects of elevated serotonin on fat, movement, and hormonal secretions. *PLoS Genet* 10: e1004394

Dasgupta N, Patel AM, Scott BA, Crowder CM (2007) Hypoxic preconditioning requires the apoptosis protein CED-4 in *C. elegans*. *Curr Biol* 17: 1954-9

Dingley S, Polyak E, Lightfoot R, Ostrovsky J, Rao M, Greco T, Ischiropoulos H, Falk MJ (2010) Mitochondrial respiratory chain dysfunction variably increases oxidant stress in *Caenorhabditis elegans*. *Mitochondrion* 10: 125-36

Ernst P, Xu N, Qu J, Chen H, Goldberg MS, Darley-Usmar V, Zhang JJ, O'Rourke B, Liu X, Zhou L (2019) Precisely Control Mitochondria with Light to Manipulate Cell Fate Decision. *Biophys J*

Ernst P, Xu N, Song J, Qu J, Chen H, Goldberg MS, Zhang J, O'Rourke B, Liu X, Zhou L (2018) Precisely control mitochondrial membrane potential with light to manipulate cell fate decisions. *bioRxiv*: 469668

Fischer LR, Igoudjil A, Magrane J, Li Y, Hansen JM, Manfredi G, Glass JD (2011) SOD1 targeted to the mitochondrial intermembrane space prevents motor neuropathy in the Sod1 knockout mouse. *Brain* 134: 196-209

Garcia D, Shaw RJ (2017) AMPK: Mechanisms of Cellular Energy Sensing and Restoration of Metabolic Balance. *Mol Cell* 66: 789-800

Garrido C, Galluzzi L, Brunet M, Puig PE, Didelot C, Kroemer G (2006) Mechanisms of cytochrome c release from mitochondria. *Cell Death Differ* 13: 1423-33

Glancy B, Hartnell LM, Combs CA, Femnou A, Sun J, Murphy E, Subramaniam S, Balaban RS (2018) Power Grid Protection of the Muscle Mitochondrial Reticulum. *Cell Rep* 23: 2832

Glancy B, Hartnell LM, Malide D, Yu ZX, Combs CA, Connelly PS, Subramaniam S, Balaban RS (2015) Mitochondrial reticulum for cellular energy distribution in muscle. *Nature* 523: 617-20

Hardie DG (2011) AMP-activated protein kinase: an energy sensor that regulates all aspects of cell function. *Genes Dev* 25: 1895-908

Hardie DG, Ross FA, Hawley SA (2012) AMPK: a nutrient and energy sensor that maintains energy homeostasis. *Nat Rev Mol Cell Biol* 13: 251-62

Hayakawa T, Kato K, Hayakawa R, Hisamoto N, Matsumoto K, Takeda K, Ichijo H (2011) Regulation of anoxic death in *Caenorhabditis elegans* by mammalian apoptosis signal-regulating kinase (ASK) family proteins. *Genetics* 187: 785-92

Hinchey EC, Gruszczyk AV, Willows R, Navaratnam N, Hall AR, Bates G, Bright TP, Krieg T, Carling D, Murphy MP (2018) Mitochondria-derived ROS activate AMP-activated protein kinase (AMPK) indirectly. *J Biol Chem* 293: 17208-17217

Husson SJ, Liewald JF, Schultheis C, Stirman JN, Lu H, Gottschalk A (2012) Microbial light-activatable proton pumps as neuronal inhibitors to functionally dissect neuronal networks in *C. elegans*. *PLoS One* 7: e40937

Ishiguro H, Yasuda K, Ishii N, Ihara K, Ohkubo T, Hiyoshi M, Ono K, Senoo-Matsuda N, Shinohara O, Yosshii F, Murakami M, Hartman PS, Tsuda M (2001) Enhancement of oxidative damage to cultured cells and *Caenorhabditis elegans* by mitochondrial electron transport inhibitors. *IUBMB Life* 51: 263-8

Jeon SM (2016) Regulation and function of AMPK in physiology and diseases. *Exp Mol Med* 48: e245

Jia B, Crowder CM (2008) Volatile anesthetic preconditioning present in the invertebrate *Caenorhabditis elegans*. *Anesthesiology* 108: 426-33

John GB, Shang Y, Li L, Renken C, Mannella CA, Selker JM, Rangell L, Bennett MJ, Zha J (2005) The mitochondrial inner membrane protein mitofilin controls cristae morphology. *Mol Biol Cell* 16: 1543-54

Kandori H (2015) Ion-pumping microbial rhodopsins. *Front Mol Biosci* 2: 52

Lee H, Cho JS, Lambacher N, Lee J, Lee SJ, Lee TH, Gartner A, Koo HS (2008) The *Caenorhabditis elegans* AMP-activated protein kinase AAK-2 is phosphorylated by LKB1 and is required for resistance to oxidative stress and for normal motility and foraging behavior. *J Biol Chem* 283: 14988-93

Lee HP, Casadesus G, Zhu X, Lee HG, Perry G, Smith MA, Gustaw-Rothenberg K, Lerner A (2009) All-trans retinoic acid as a novel therapeutic strategy for Alzheimer's disease. *Expert Rev Neurother* 9: 1615-21

Mello CC, Kramer JM, Stinchcomb D, Ambros V (1991) Efficient gene transfer in *C.elegans*: extrachromosomal maintenance and integration of transforming sequences. *EMBO J* 10: 3959-70

Michiels C (2004) Physiological and pathological responses to hypoxia. *Am J Pathol* 164: 1875-82

Mihaylova MM, Shaw RJ (2011) The AMPK signalling pathway coordinates cell growth, autophagy and metabolism. *Nat Cell Biol* 13: 1016-23

Murphy MP, Hartley RC (2018) Mitochondria as a therapeutic target for common pathologies. *Nat Rev Drug Discov*

Okazaki A, Takagi S (2013) An optogenetic application of proton pump ArchT to *C. elegans* cells. *Neurosci Res* 75: 29-34

Ozcan C, Palmeri M, Horvath TL, Russell KS, Russell RR, 3rd (2013) Role of uncoupling protein 3 in ischemia-reperfusion injury, arrhythmias, and preconditioning. *Am J Physiol Heart Circ Physiol* 304: H1192-200

Palace VP, Khaper N, Qin Q, Singal PK (1999) Antioxidant potentials of vitamin A and carotenoids and their relevance to heart disease. *Free Radic Biol Med* 26: 746-61

Pena S, Sherman T, Brookes PS, Nehrke K (2016) The Mitochondrial Unfolded Protein Response Protects against Anoxia in *Caenorhabditis elegans*. *PLoS One* 11: e0159989

Perry SW, Norman JP, Barbieri J, Brown EB, Gelbard HA (2011) Mitochondrial membrane potential probes and the proton gradient: a practical usage guide. *Biotechniques* 50: 98-115

Porcelli AM, Ghelli A, Zanna C, Pinton P, Rizzuto R, Rugolo M (2005) pH difference across the outer mitochondrial membrane measured with a green fluorescent protein mutant. *Biochem Biophys Res Commun* 326: 799-804

Rizzuto R, De Stefani D, Raffaello A, Mammucari C (2012) Mitochondria as sensors and regulators of calcium signalling. *Nat Rev Mol Cell Biol* 13: 566-78

Rost BR, Schneider F, Grauel MK, Wozny C, Bentz C, Blessing A, Rosenmund T, Jentsch TJ, Schmitz D, Hegemann P, Rosenmund C (2015) Optogenetic acidification of synaptic vesicles and lysosomes. *Nat Neurosci* 18: 1845-1852

Sanderson TH, Reynolds CA, Kumar R, Przyklenk K, Huttemann M (2013) Molecular mechanisms of ischemia-reperfusion injury in brain: pivotal role of the mitochondrial membrane potential in reactive oxygen species generation. *Mol Neurobiol* 47: 9-23

Santo-Domingo J, Giacomello M, Poburko D, Scorrano L, Demarex N (2013) OPA1 promotes pH flashes that spread between contiguous mitochondria without matrix protein exchange. *EMBO J* 32: 1927-40

Sawin ER, Ranganathan R, Horvitz HR (2000) *C. elegans* locomotory rate is modulated by the environment through a dopaminergic pathway and by experience through a serotonergic pathway. *Neuron* 26: 619-31

Schmeisser S, Priebe S, Groth M, Monajembashi S, Hemmerich P, Guthke R, Platzer M, Ristow M (2013) Neuronal ROS signaling rather than AMPK/sirtuin-mediated energy sensing links dietary restriction to lifespan extension. *Mol Metab* 2: 92-102

Shabalina IG, Nedergaard J (2011) Mitochondrial ('mild') uncoupling and ROS production: physiologically relevant or not? *Biochem Soc Trans* 39: 1305-9

Shadel GS, Horvath TL (2015) Mitochondrial ROS signaling in organismal homeostasis. *Cell* 163: 560-9

Siddikuzzaman, Grace VM (2013) Antioxidant potential of all-trans retinoic acid (ATRA) and enhanced activity of liposome encapsulated ATRA against inflammation and tumor-directed angiogenesis. *Immunopharmacol Immunotoxicol* 35: 164-73

Solaini G, Baracca A, Lenaz G, Sgarbi G (2010) Hypoxia and mitochondrial oxidative metabolism. *Biochim Biophys Acta* 1797: 1171-7

Sumii M, Furutani Y, Waschuk SA, Brown LS, Kandori H (2005) Strongly hydrogen-bonded water molecule present near the retinal chromophore of Leptosphaeria rhodopsin, the bacteriorhodopsin-like proton pump from a eukaryote. *Biochemistry* 44: 15159-66

Takahashi M, Takagi S (2017) Optical silencing of body wall muscles induces pumping inhibition in *Caenorhabditis elegans*. *PLoS Genet* 13: e1007134

Tkatch T, Greotti E, Baranauskas G, Pendin D, Roy S, Nita LI, Wettmarshausen J, Prigge M, Yizhar O, Shirihaï OS, Fishman D, Hershfinkel M, Fleidervish IA, Perocchi F, Pozzan T, Sekler I (2017) Optogenetic control of mitochondrial metabolism and Ca(2+) signaling by mitochondria-targeted opsins. *Proc Natl Acad Sci U S A* 114: E5167-E5176

Trewin AJ, Bahr LL, Almast A, Berry BJ, Wei AY, Foster TH, Wojtovich AP (2019) Mitochondrial Reactive Oxygen Species Generated at the Complex-II Matrix or Intermembrane Space Microdomain Have Distinct Effects on Redox Signaling and Stress Sensitivity in *Caenorhabditis elegans*. *Antioxid Redox Signal* 31: 594-607

Trewin AJ, Berry BJ, Wei AY, Bahr LL, Foster TH, Wojtovich AP (2018a) Light-induced oxidant production by fluorescent proteins. *Free Radic Biol Med* 128: 157-164

Trewin AJ, Berry BJ, Wojtovich AP (2018b) Exercise and Mitochondrial Dynamics: Keeping in Shape with ROS and AMPK. *Antioxidants (Basel)* 7

Tsalik EL, Hobert O (2003) Functional mapping of neurons that control locomotory behavior in *Caenorhabditis elegans*. *J Neurobiol* 56: 178-97

Tsang WY, Lemire BD (2003) The role of mitochondria in the life of the nematode, *Caenorhabditis elegans*. *Biochim Biophys Acta* 1638: 91-105

Viola HM, Hool LC (2017) Auto-regulation in the powerhouse. *Elife* 6

Wang X, Zhang X, Wu D, Huang Z, Hou T, Jian C, Yu P, Lu F, Zhang R, Sun T, Li J, Qi W, Wang Y, Gao F, Cheng H (2017) Mitochondrial flashes regulate ATP homeostasis in the heart. *Elife* 6

Wang YT, Lim Y, McCall MN, Huang KT, Haynes CM, Nehrke K, Brookes PS (2019) Cardioprotection by the mitochondrial unfolded protein response (UPR(mt)) requires ATF5. *Am J Physiol Heart Circ Physiol*

Waschuk SA, Bezerra AG, Jr., Shi L, Brown LS (2005) Leptosphaeria rhodopsin: bacteriorhodopsin-like proton pump from a eukaryote. *Proc Natl Acad Sci U S A* 102: 6879-83

Wojtovich AP, DiStefano P, Sherman T, Brookes PS, Nehrke K (2012a) Mitochondrial ATP-sensitive potassium channel activity and hypoxic preconditioning are independent of an inwardly rectifying potassium channel subunit in *Caenorhabditis elegans*. *FEBS Lett* 586: 428-34

Wojtovich AP, Nadtochiy SM, Brookes PS, Nehrke K (2012b) Ischemic preconditioning: the role of mitochondria and aging. *Exp Gerontol* 47: 1-7

Wojtovich AP, Nadtochiy SM, Urciuoli WR, Smith CO, Grunnet M, Nehrke K, Brookes PS (2013) A non-cardiomyocyte autonomous mechanism of cardioprotection involving the SLO1 BK channel. *PeerJ* 1: e48

Xu M, Wang Y, Ayub A, Ashraf M (2001) Mitochondrial K(ATP) channel activation reduces anoxic injury by restoring mitochondrial membrane potential. *Am J Physiol Heart Circ Physiol* 281: H1295-303

728 Yang JL, Mukda S, Chen SD (2018) Diverse roles of mitochondria in ischemic stroke. Redox Biol 16: 263-
729 275
730 Zhang L, Trushin S, Christensen TA, Bachmeier BV, Gateno B, Schroeder A, Yao J, Itoh K, Sesaki H, Poon
731 WW, Gyls KH, Patterson ER, Parisi JE, Diaz Brinton R, Salisbury JL, Trushina E (2016) Altered brain
732 energetics induces mitochondrial fission arrest in Alzheimer's Disease. Sci Rep 6: 18725
733
734

Expanded view figures

Figure EV1 - mtON is targeted to mitochondria.

A mtON tagged with GFP expressed in body wall muscle overlaps with MitoTrackerTM Red CMXRos stained mitochondria.

B Profile plot of a selected region demonstrating the overlapping signal intensity of GFP and MitoTrackerTM Red CMXRos determined using ImageJ software.

Figure EV2 - Schematic of experimental conditions.

A Expression control: baseline condition where animals express mtON, but have not been supplemented with ATR or exposed to light. This condition is represented by the color black throughout.

B Light control: condition exposed to light where mtON is illuminated but not supplemented with ATR, resulting in an inactive proton pump. This condition is represented by the color gray throughout.

C ATR control: condition supplemented with ATR but not exposed to light, where proton pumping is possible but no light activation has occurred. This condition is represented by dark green throughout.

D Active optogenetic condition: supplemented with ATR and exposed to light, where proton pumping activity is expected. This condition is represented by bright green throughout.

Figure EV3 - mtON increases the PMF and ATP synthesis light dose-dependently

A Quantification of change in TMRE fluorescence with increasing light dose (increasing fluence from left to right: 0, 0.016, 0.08, 0.096, 0.16, 0.24 J/cm²). Two-way ANOVA with Sidak's multiple comparisons test, +ATR: 0 vs. 0.096 J/cm² *p = 0.0232, 0 vs. 0.24 J/cm² *p = 0.0149, 0.016 vs. 0.096 J/cm² *p = 0.0162, 0.016 vs. 0.16 J/cm² *p = 0.0376, 0.016 vs. 0.24 J/cm² *p = 0.0101. n = 3 - 6 mitochondrial isolations.

B ATP levels in response to increasing illumination normalized to maximum ATP synthesis given by succinate respiration (dotted line). 10 minute illumination time from Fig. 2E. Two-way ANOVA with Sidak's multiple comparison test, -ATR: succinate vs. 1, 5 and 10 minutes light *p < 0.0001. +ATR: succinate vs. 1 minute light, *p < 0.0001, 1 minute light vs. 5 minutes light *p = 0.0098, 1 minute light vs. 10 minutes light *p = 0.0001. n = 3 - 7 light treatments across 2 mitochondrial isolations.

Figure EV4 - mtON effects on respiration

A Respiratory control ratios (indicative of ability of isolated mitochondria to respond to energy demand) were calculated by dividing the rate of O_2 consumption with ADP and succinate by the rate after ADP was depleted. One-way ANOVA with Tukey's post hoc test, $p = 0.72$, $n = 6$ mitochondrial preparations. Bars are means $\pm$ SEM.

B O_2 consumption rates under different states of respiration comparing the control conditions. One-way ANOVA with Tukey's post hoc tests performed for each respiration state. ADP rate $p = 0.66$, Succinate rate $p = 0.61$. ADP exhausted rate $p = 0.83$. Bars are means $\pm$ SEM. **C** Representative traces depicting O_2 consumption rate. +/- light for mitochondria with ATR present. Light exposure was 10 minutes before the addition of succinate. Traces show an initial rapid depletion of ADP before transitioning to a lower rate of O_2 consumption.

C Activation of mtON decreases the amount of O_2 required to consume 50 nmoles ADP light dose-dependently. Isolated mitochondria were exposed to light for the indicated time and then succinate was added (example traces in panel c). Dark and 10 minute illumination data was used for analysis in Fig. 2F. Linear regression showed a negative relationship between O_2 required to consume ADP and illumination time in mitochondria from animals with mtON that were supplemented with ATR, $R^2 = 0.98$ $p = 0.007$, $n = 6$ mitochondrial preparations. Linear regression shows no relationship between O_2 required to consume ADP and illumination in mitochondria from animals with mtON not supplemented with ATR, $R^2 = 0.756$ $p = 0.130$, dark $n = 5$, 1 min light $n = 5$, 5 min light $n = 6$, 10 min light $n = 6$. Data are means $\pm$ SEM.

Figure EV5 – ATR and light have no effect in wild type survival

A Same experimental conditions from Fig. 3A, here testing wild type animals. No difference in survival within ATR conditions, suggesting the interaction between ATR and light has no effect on its own. One-way ANOVA with Tukey's multiple comparison test. Again, ATR alone was protective (-ATR, -light vs. +ATR, -light $p = 0.0111$.)

B Same experimental conditions from Fig. 3B, here testing wild type animals, showing no difference in survival. One-way ANOVA with Tukey's multiple comparison test.

Table 1. *C. elegans* strains. Some strains were provided by the *C. elegans* Genetics Center (CGC).

Strain	Genotype	Abbreviation	Source	Notes
N2(Bristol)		wildtype	CGC	Wild type
APW32	<i>pha-1(e2123ts) III; jbmEx11 [pBJB20(P_{eff}-₃::Mitofilin(N' 187 aa)::Mac::GFP), pC1 (pha-1(+))]</i>	mtON	this study	Ubiquitously expressed mitochondria-targeted light-activated proton pump
APW16	<i>pha-1(e2123ts) III; him-5 (e1490)V, jbmEx1 [pBJB16(P_{myo-3}::Mitofilin(187N'aa)::Mac::GFP), pC1 (pha-1(+))]</i>		this study	Mitochondria-targeted light-activated proton pump expressed only in body wall muscle cells strain
RB754	<i>aak-2(ok524) X.</i>	<i>aak-2</i>	CGC	AAK-2 loss of function mutant from the <i>C. elegans</i> Gene Knockout Project at the Oklahoma Medical Research Foundation, which was part of the International <i>C. elegans</i> Gene Knockout Consortium.