Title:

Multilevel modelling, prevalence and predictors of hypertension in Ghana: Evidence from Wave 2 of the World Health Organization's Study on Global AGEing and adult health.

Authors:

Justice Moses K. Aheto^{1,2*} and Getachew A. Dagne²

Authors' Affiliations:

¹Department of Biostatistics, School of Public Health, College of Health Sciences, University of Ghana, Ghana.

²College of Public Health, University of South Florida, USA.

***Corresponding author:**

Justice Moses K. Aheto
Department of Biostatistics
School of Public Health
College of Health Sciences
University of Ghana
P. O. Box LG13, Legon-Accra.

E-mail: justiceaheto@yahoo.com; jmkaheto@ug.edu.gh

Abstract

Background

Hypertension is a major public health issue, a critical risk factor for cardiovascular diseases and stroke, especially in developing countries where the rates remain unacceptably high. In Africa, hypertension is the leading driver of cardiovascular disease and stroke deaths. Identification of critical risk factors of hypertension can help formulate targeted public health programmes and policies aimed at reducing the prevalence and its associated morbidity, disability and mortality. This study attempts to develop multilevel regression, an in-depth statistical model to identify critical risk factors of hypertension.

Methods

This study used data on 4381 individuals aged ≥ 18 years from the nationally representative World Health Organization Study on global AGEing and adult health (SAGE) Ghana Wave 2. Multilevel regression modelling was employed to identify critical risk factors for hypertension based on systolic blood pressure (SBP) (i.e. $SBP > 140$ mmHg).

Results

The data on 4381 individuals were analysed out of which 27.3% were hypertensive. Critical risk factors for hypertension identified were age, obesity, marital status, health state and difficulty with self-care. Strong unobserved household-level residual variations were found.

Conclusion

Hypertension remains high in Ghana. Addressing the problem of obesity, targeting specific interventions to those aged over 50 years, and improvement in the general health of Ghanaians are paramount to reducing the prevalence and its associated morbidity, disability and mortality. Lifestyle modification in the form of dietary intake, knowledge provision supported with strong public health message and political will could be beneficial to the management and prevention of hypertension.

Keywords: Multilevel modelling, binary logistic regression, Hypertension, Risk factors, Hypertension determinants, Developing countries, Sub-Saharan Africa, Ghana.

Background

Hypertension remains one of the biggest threats to public health globally, especially in the low- and middle-income countries where the prevalence is the highest as a result of more people residing in these countries, and with greater priority and interest in infectious diseases [1, 2]. It contributes significantly to the global burden of cardiovascular and its related illnesses like stroke, kidney and heart failures, and their resultant premature morbidity, disability and mortality. Hypertension is responsible for 51% of deaths from stroke and at least 45% from heart diseases and overall, it is responsible for about 50% of deaths from stroke and heart disease [1, 3]. The Global Burden of Disease Study Group reported that in 2017, cardiovascular diseases were the leading cause of death in Africa and were responsible for 1.42 million deaths or 16.4% of the total deaths in all ages [4]. The number of deaths represent 61.0% increase over the estimated number of cardiovascular deaths in 1990. High blood pressure, one of the Non-Communicable Diseases (NCDs), is the leading risk factor for deaths in Africa responsible for nearly two-thirds of the cardiovascular deaths in the region. Globally, Africa has the highest prevalence of high blood pressure (27%) [5] and a common cause of medical hospitalization in the region [6] and responsible for over 50% of first time acute stroke [7, 8]. Previous studies reported that with the ageing population and the rising urbanization and its associated stress, sedentary lifestyle and 'Western' diet, high blood pressure will continue to rise [2, 8-10].

Previous studies observe that older age, race, cigarette smoking, high salt intake, high body mass index (BMI), alcohol use, female sex, urban residence, physical inactivity and genetics are among the main factors associated with hypertension [8, 11-16]. Though not a significant problem previously in groups like young and rural populations, hypertension is now a critical public health problem in these groups [8, 17-19].

There is a renewed political will to address NCDs as a result of the third high-level meeting of the United Nations General Assembly in October 2018 during which Heads of State and Governments committed to reorienting health systems to respond to the needs of the rapidly ageing population in relation to NCDs [8, 20, 21]. However, sound statistical methods are required to analyse the barriers and facilitators of hypertension in order to achieve this ambitious goal. It is against this background that this study attempts to estimate the prevalence and to employ a multilevel regression model to identify critical risk factors for hypertension to inform sound and targeted policies aimed at improving cardiovascular health outcomes among this population.

Methods

Data source and study population

This work was based on the nationally representative World Health Organization Study on global AGEing and adult health (SAGE) Ghana Wave 2 conducted in the period 2014/2015. SAGE is a multi-country study that collects data to complement existing ageing data sources to inform policy and programmes. Information on subjective wellbeing, risk factors and preventive health behaviours, quality of life and health care utilization, perceived health status, household characteristics, socio-demographic and social cohesion were collected from individuals aged 50 years. A smaller sample of those aged 18-49 years were also included in the study. Detailed description of the methods is published elsewhere [22].

Outcome variable

The outcome variable of interest in the study is hypertension status based on systolic blood pressure measurement (SBP>140mmHg).

Covariates

This study considered several covariates based on literature on factors influencing hypertension, including other potential health variables yet to be established in the literature as risk factors. These include age, obesity, ethnicity, sex, marital status, type of toilet facility in household, health state and difficulty with self-care at the time of the interview, alcohol consumption, type of cooking fuel, household wall and floor types [2, 11-14, 23-25].

Statistical analysis

Descriptive statistics were used to summarize the distribution of selected background characteristics of respondents. Further analyses were conducted to examine individual and household-level factors that might be significantly associated with hypertension and explored unobserved household level effects on the outcome. Both single-level and multilevel (mixed effects) logistic regression models were applied on 4381 individuals residing in 3790 households with complete measurements on hypertension as well as complete measurements on potential explanatory variables considered in the final models. The extension of the single level logistic regression model to the multilevel logistic regression model is warranted because of the hierarchical structure of the SAGE dataset where we have individuals nested within households. Specifically, we applied multilevel logistic regression models to examine possible differences in hypertension among individuals across households while simultaneously identifying potential risk factors. Thus, the multilevel modelling approach [26] placed particular emphasis on

household level differences in the risk of hypertension among individuals and the extent of nesting of hypertension within a household which cannot be achieved through a single level logistic regression model.

For quantifying the proportion of total variation attributable to within-households differences, the household-level Variance Partition Coefficient (VPC) [27] were employed, which is defined as $VPC = (\text{household-level variance} / (\text{household-level variance} + \text{individual-level variance}))$. The individual-level residual is assumed to follow a standard logistic distribution with mean zero and variance $\pi^2 / 3$, where $\pi = 3.14$ [28].

Model parameters were obtained using maximum likelihood. Identity covariance structure provided a good fit to the data in the multilevel logistic model. The goodness of fit for the fitted models was examined using a likelihood ratio test (LRT), Akaike Information Criterion (AIC) and Bayesian Information Criterion (BIC). Variance inflation factor (VIF) was used to check multicollinearity, and a VIF value below 10 was considered acceptable [29]. All the analyses were performed using STATA Version 14 [30]. Backward elimination was employed to select candidate set of risk factors for multivariable logistic regression analysis. P-value < 0.05 was used to declare statistical significance.

Ethical approval and consent to participate

SAGE was approved by the World Health Organization's Ethical Review Board (reference number RPC149) and the Ethical and Protocol Review Committee, College of Health Sciences, University of Ghana, Accra, Ghana. Written informed consent was obtained from all study participants. All methods were performed in accordance with the relevant guidelines and regulations.

Results

Participant characteristics

Out of the 4381 individual respondents, 1273 (27.3%) had hypertension. Approximately 76% of the respondents were at least 50 years old and 58.9% were females. Over 48% of the respondents were Akan, with the Guan ethnicity in the minority (4.1%) and over 55% reported being currently married and about 74% had fathers with no formal education. Over 12% were obese and over 29% reported ever using a tobacco while over 32% reported that they did not know high salt diet can cause health problems. Household characteristics included durable material for walls (62.8%), non-flush toilets (84.6%), shared toilets (76.9%) and hard floor (85.1%) (Table 1).

Table 1: Summary of selected background characteristics of respondents

Variables	n (%)	Variables	n (%)
Hypertensive		Ever use alcohol	
No	3394 (72.7)	No	3308 (70.82)
Yes	1273 (27.3)	Yes	1363 (29.18)
Age		Health status	
18-49	1160 (24.5)	Good/very good	3192 (67.97)
50 or more	3575 (75.5)	Moderate	1114 (23.72)
Sex		Bad/very bad	390 (8.3)
Male	1948 (41.14)	High salt diet can cause health problems	
Female	2787 (58.86)	Yes	2912 (67.42)
Ethnicity		No	1407 (32.58)
Akan	2296 (48.62)	Self-care	
Ewe	276 (5.84)	None	3756 (79.93)
Ga-Adangbe	587 (12.43)	Mild	680 (14.47)
Guan	193 (4.09)	Moderate	207 (4.41)
Northern dialect	1370 (29.01)	Severe/extreme	56 (1.19)
Marital status		Wall	
Currently not married	2107 (44.5)	Durable materials	3348 (61.75)
Currently married	2628 (55.5)	Non-durable materials	2074 (38.25)
Father's education		Drinking water	
No formal education	3480 (73.84)	Piped source	2702 (49.84)
Primary or less	584 (12.39)	Non-piped source	2719 (50.16)
Secondary or higher	649 (13.77)	Toilets	
BMI		Flush toilets	836 (15.44)
No	3926 (87.44)	Non-flush toilets	4577 (84.56)
Yes	564 (12.56)	Cooking fuel	
Vigorous activity		Gas/electric	642 (11.91)
None	2046 (43.61)	Coal/charcoal/kerosene	1861 (34.53)
Mild	720 (15.35)	Wood	2887 (53.56)
Moderate	897 (19.12)	Shared toilets	
Severe	465 (9.91)	No	1046 (23.14)
Extreme	564 (12.02)	Yes	3474 (76.86)
Use tobacco		Floor	
Yes	283 (6.06)	Hard floor	4607 (85.13)
No	4388 (93.94)	Earth floor	805 (14.87)

n: Frequency. %: Percentage

Predictors of hypertension

The univariable analyses identified age, obesity, sex, marital status, toilet facility, health state, difficulty with self-care, wall type, floor type and ethnicity as significant predictors of hypertension. Significant predictors of hypertension in the multivariable model include age, obesity, marital status, toilet facility, health state and difficulty with self-care.

Comparing the single-level multivariable logistic regression (Table 2) to the multilevel logistic regression model (Table 3), the multilevel model provided a good fit to the data. Thus, the multilevel logistic regression is preferred to the single-level multivariable model.

Table 2: Risk factors for hypertension from single-level logistic regression models

Variables	Single-level univariable logistic		Single-level multivariable logistic	
	UOR (95% CI)	P-value	AOR (95% CI)	P-value
Age		<0.001***		
<50 years	ref		ref	
50 or more	5.12 (4.13-6.35)	<0.001***	4.85 (3.85-6.11)	<0.001***
Obesity		<0.001***		
No	ref		ref	
Yes	1.51 (1.25-1.83)	<0.001***	1.47 (1.19-1.83)	<0.001***
Sex		<0.001***		
Male	ref		ref	
Female	1.28 (1.12-1.46)	<0.001***	1.11 (0.94-1.31)	0.209
Marital status		<0.001***		
Not currently married	ref		ref	
Currently married	0.72 (0.63-0.82)	<0.001***	0.76 (0.66-0.89)	<0.001***
Toilet facility		<0.001***		
Flush toilets	ref		ref	
Non-flush toilets	0.74 (0.62-0.88)	0.001**	0.79 (0.63-0.99)	0.039*
Health state		<0.001***		
Good/ very good	ref		ref	
Moderate	1.91 (1.65-2.22)	<0.001***	1.34 (1.14-1.58)	<0.001***
Bad/very bad	2.08 (1.66-2.6)	<0.001***	1.32 (1-1.73)	0.047*
Difficulty with self-care		<0.001***		
Good/ very good	ref		ref	
Mild	1.18 (0.98-1.41)	0.076	1.05 (0.86-1.28)	0.646
Moderate	2.13 (1.59-2.85)	<0.001***	1.57 (1.09-2.25)	0.015*
Severe/extreme	3.32 (1.96-5.65)	<0.001***	1.78 (0.77-4.13)	0.177
Alcohol		0.605		
No	ref		ref	
Yes	1.04 (0.9-1.2)	0.605	1.09 (0.92-1.28)	0.317
Cooking fuel		0.368		
Gas/ electric	ref		ref	
Coal/charcoal/kerosene	0.96 (0.77-1.18)	0.681	0.97 (0.75-1.26)	0.816
Wood	0.88 (0.72-1.09)	0.24	0.87 (0.66-1.13)	0.296
Walls		0.005**	-	
Durable materials	ref			
Non-durable material	0.82 (0.72-0.94)	0.005**		
Floor		0.026*	-	
Hard floor	ref			
Earth floor	0.8 (0.66-0.97)	0.026*		
Ethnicity		<0.001***	-	
Akan	ref			
Ewe	1.33 (1.01-1.74)	0.04*		
Ga-Adangbe	1.31 (1.08-1.6)	0.007**		
Guan	0.8 (0.56-1.13)	0.203		
Northern dialect	0.87 (0.75-1.02)	0.08		

UOR: unadjusted odds ratio. AOR: adjusted odds ratio. CI: confidence interval. ref: reference category. *: P-value <0.05. **: P-value<0.01. ***: P-value <0.001

Significant predictors of hypertension in the multilevel model include age, obesity, marital status, health state and difficulty with self-care.

Significant unobserved household-level variations in hypertension were found. The results from the variation analyses showed that over 12% of variance in hypertension could be attributable to residual household-level variations after adjusting for individual and household level factors considered in the multilevel model.

Individuals aged ≥ 50 years were at increased risk of hypertension compared to those aged 18-49 years (OR=5.4, 95% CI: 4.11-7.09). There is a 51% increase in the odds of hypertension among individuals who are obese compared to their counterparts who are not obese (OR=1.51, 95% CI: 1.19-1.91). Individuals who are currently married had 25% less odds of having hypertension compared to those who are not currently married (OR=0.75, 95% CI: 0.64-0.89). Individuals who rated their health state as moderate or bad/very bad had 38% and 35% higher odds of having hypertension, respectively compared to those who rated themselves as good/very good (OR=1.38, 95% CI: 1.15-1.65 and OR=1.35, 95% CI: 1.0-1.83). Those who rated themselves as having moderate difficulty with self-care had 64% higher odds of having hypertension compared to those who rated themselves as having no difficulty OR=1.64, 95% CI: 1.1-2.44) (Table 3).

Table 3: Risk factors for hypertension from multilevel (mixed effect) logistic regression models

Variables	Mixed effect logistic		Parameter (95% CI)
	AOR (95% CI)	P-value	
Age			
<50 years	ref		
50 or more	5.4 (4.11-7.09)	<0.001***	
Obesity			
No	ref		
Yes	1.51 (1.19-1.91)	0.001**	
Sex			
Male	ref		
Female	1.11 (0.94-1.33)	0.225	
Marital status			
Not currently married	ref		
Currently married	0.75 (0.64-0.89)	0.001**	
Toilet facility			
Flush toilets	ref		
Non-flush toilets	0.79 (0.62-1.01)	0.06	
Health state			
Good/ very good	ref		
Moderate	1.38 (1.15-1.65)	<0.001***	
Bad/very bad	1.35 (1-1.83)	0.047*	
Difficulty with self-care			
None	ref		
Mild	1.06 (0.85-1.32)	0.62	
Moderate	1.64 (1.1-2.44)	0.016*	
Severe/extreme	1.88 (0.74-4.73)	0.182	
Alcohol			
No	ref		

Yes	1.1 (0.92-1.31)	0.282	
Cooking fuel			
Gas/ electric	ref		
Coal/charcoal/kerosene	0.96 (0.73-1.28)	0.799	
Wood	0.84 (0.63-1.14)	0.266	
Individual-level variance			3.29
Household-level variance		0.022	0.46 (0.15 – 1.48)

AOR: adjusted odds ratio. CI: confidence interval. ref: reference category. *: P-value <0.05. **: P-value<0.01. ***: P-value <0.001

Discussion

The study sets out to estimate hypertension prevalence and to develop a novel multilevel logistic regression model to identify critical risk factors of hypertension to help in formulating targeted policies that could improve cardiovascular health among the Ghanaian adults. In this study, a hypertension prevalence of 27.3% was observed, suggesting that hypertension among Ghanaian adults is still a serious public health issue. This prevalence exceeded the 18% prevalence observed in her neighbouring country Burkina Faso [25], 24.5% in Kenya [31], and 8% in Tanzania [32] but lower than the 31% observed in Nigeria [33]. Critical risk factors independently associated with hypertension while adjusting for the unobserved household-level effects were age, obesity, marital status, health state and difficulty with self-care.

Of critical importance to this study is the quantification of residual household-level effects on hypertension among adults, which represent differences in household-level hypertension outcomes that cannot be explained by the available household-level covariates. Generally, the health and the general wellbeing of individuals is heavily reliant on the households they belong to. Thus, the households determine the resources, opportunities and risks available to the individual over their life course [34-36]. We observed strong residual household-level variations in hypertension and that over 12% of variation in hypertension in adults could be attributable to unobserved household-level variations after adjusting for the risk factors in the model. This could be as a result of household-level, social and environmental factors not considered in our model.

The study broadly supports earlier studies that examined determinants of hypertension in developing countries. For instance, older age group, obesity, rating health state as moderate or bad/very bad, and rating level of difficulty for self-care as moderate were associated with increased odds of hypertension in adults, and those who were currently married had reduced odds of hypertension [3, 11-15, 25, 33, 37, 38]. The association between hypertension and the older age group observed in this study could be attributable to variations in the arterial structure and function, notably arterial stiffening with adverse consequences on cardiac structure and function [33, 39, 40]. Individuals who were currently married had reduced odds

of hypertension compared to those who were not currently married. Protective effects for marriage on health has been established in previous studies [41, 42]. However, this finding is not consistent with a previous study that observed that married women had increased odds of hypertension but no such association was found in men in the same study [3]. It also contradicts a previous study [43] that did not find an association between marital status and hypertension but this could be as a result of the setting or how the variable marital status was categorised. Marital status is a critical social characteristic which is well known to predict a range of health outcomes such as cardiovascular illnesses [44, 45] and mortality in general [42, 46].

Obesity, which is one of the modifiable risk factors considered in this study showed an association with hypertension. Individuals who were obese had higher odds of hypertension compared to their counterparts who were not obese, a finding which is consistent with previous studies [25, 31-33, 38, 47]. The association between hypertension and obesity had been established and well known, and reducing BMI is part of the advice provided in the treatment of hypertension [25, 48]. This could be due to metabolic and endocrine disorders as a result of increasing BMI [49].

Individuals who rated their health state as moderate or bad/very bad on the day of the interview had increased odds of hypertension compared to those who rated their health state as good/very good. Also, those who rated their difficulty with self-care as moderate had increased odds of hypertension compared to their counterparts who rated themselves as having no difficulty. These findings are plausible because in a previous study, a hypertensive group had significantly lower age-adjusted health status scores compared to non-hypertensive group [50]. To the best of our knowledge, this is the first study to have established an association between hypertension and health variables like self-reported health state and difficulty with self-care. The significant association between hypertension and these health variables suggest the need to improve the overall health of Ghanaian adults to reduce and prevent the high prevalence of hypertension among this group.

One of the major public health issues in Ghana presently is the rise in the prevalence of NCDs [51, 52]. The findings in the present study provided vital and current information on prevalence and critical risk factors for hypertension that can be used by policymakers and health practitioners for better understanding of hypertension and its prevention and management, which could lead to more effective prevention approaches, patient management and improved cardiovascular outcomes. The study highlights the need to address the problem of obesity, targeting specific interventions to those aged over 50 years, and improvement in the general health of the Ghanaian population as a primary intervention is warranted as part of an overall

strategy to reduce the hypertension prevalence and its resultant premature morbidity, disability and mortality.

Strength and limitation of the study

The strengths of this study include the fact that it utilised data from a nationally representative population-based survey which is globally respected for its sound survey methods and sound data quality on individuals, their households and communities in which they reside. The large samples drawn nationwide permits generalization of findings to the population of adults in Ghana and that of adults from other similar populations globally. The study also used a novel multilevel modelling approach, permitting the study of unobserved household-level effects on hypertension. Thus, providing much more information about why individuals from certain households are more likely to be hypertensive while others are not and at the same time investigating underlying associations between hypertension and the risk factors which could not have been possible using single-level logistic regression approach. Despite these strengths, the study has limitations and so the findings should be interpreted with caution. For example, the analytical techniques employed in the analysis of the data could not establish cause and effect relationship between hypertension and the risk factors considered. Also, some of the risk factors such as health state and difficulty with self-care were based on self-reports and so could introduce reporting bias.

Conclusion

Findings from the study show that prevalence of hypertension remains high among Ghanaian adults. This study developed a novel multilevel binary logistic regression model which captures unobserved household-level effects and identified critical risk factors of hypertension which can aid formulation of health policies and intervention strategies that will improve cardiovascular health outcomes of the Ghanaian adults. Lifestyle modification in the form of dietary intake, knowledge provision supported with strong public health messages and political will could be beneficial to the management and prevention of hypertension. Active screening for hypertension should be encouraged to identify undiagnosed cases to minimise the danger of stroke and cardiovascular diseases, but there is the need to improve the health systems and services in the country to reap the full benefits of such interventions. There is also the need to target younger populations to minimize their risk of developing hypertension during adulthood/old age. Further study to identify as-yet unidentified risk factors that might account for the substantial unexplained household-level variations in adult hypertension is warranted.

Abbreviations

AIC: Akaike Information Criterion
 BIC: Bayesian Information Criterion
 BMI: Body Mass Index
 LRT: Likelihood Ratio Test
 NDCs: Non-Communicable Diseases
 SBP: Systolic Blood Pressure
 VIF: Variance Inflation Factor
 VPC: Variance Partition Coefficient

Author contributions

JMKA developed the concept and analysed the data and wrote the first draft manuscript. JMKA and GAD contributed to the writing and reviewing of the various sections of the manuscript. All the authors reviewed the final version of the manuscript before submission. All authors read and approved the final manuscript.

Declarations

Competing interest

The authors declare that they have no competing interests.

Acknowledgements

This Fellowship was supported by the University of Ghana Building a New Generation of Academics in Africa (BANGA-Africa) Project with funding from the Carnegie Corporation of New York. The statements made and views are solely the responsibility of the authors. We are also grateful to Prof. Alfred Yawson and his Ghana Team and all respondents and interviewers who made the SAGE survey in Ghana possible. Financial support was provided by the US National Institute on Aging through Interagency Agreements (OGHA 04034785; YA1323-08-CN-0020; Y1-AG-1005-01) with the World Health Organization and a Research Project Grant (R01 AG034479- 64401A1). WHO contributed financial and human resources to SAGE. The Ministry of Health, Ghana, is supportive of SAGE. The University of Ghana's Department of Community Health contributed training facilities, data entry support, and storage of materials. The Ghana Statistical Office provided the sampling information for the sampling frame and updates.

Consent for publication

Not applicable

Funding

Not applicable

Data availability

Data is freely available upon making official request to WHO-SAGE Team through the WHO website at <http://www.who.int/healthinfo/sage/cohorts/en/>.

References

1. World Health Organization (WHO): **A Global Brief on Hypertension: Silent Killer, Global Public Health Crisis** (Accessed on 30/03/2019 and available at https://www.who.int/cardiovascular_diseases/publications/global_brief_hypertension/en/). In. Geneva: WHO; 2013.
2. van de Vijver S, Akinyi H, Oti S, Olajide A, Agyemang C, Aboderin I, Kyobutungi C: **Status report on hypertension in Africa--consultative review for the 6th Session of the African Union Conference of Ministers of Health on NCD's**. *Pan Afr Med J* 2013, **16**:38.
3. Tuoyire DA, Ayetey H: **GENDER DIFFERENCES IN THE ASSOCIATION BETWEEN MARITAL STATUS AND HYPERTENSION IN GHANA**. *J Biosoc Sci* 2018:1-22.
4. **Global Burden of Disease: GBD Compare Data Visualization** [<https://vizhub.healthdata.org/gbd-compare/>]
5. Zhou B, Bentham J, Di Cesare M, Bixby H, Danaei G, Cowan MJ, Paciorek CJ, Singh G, Hajifathalian K, Bennett JE: **Worldwide trends in blood pressure from 1975 to 2015: a pooled analysis of 1479 population-based measurement studies with 19· 1 million participants**. *Lancet* 2017, **389**(10064):37-55.
6. Etyang AO, Gerard Scott JAJGha: **Medical causes of admissions to hospital among adults in Africa: a systematic review**. *Glob Health Action* 2013, **6**(19090).
7. O'Donnell MJ, Chin SL, Rangarajan S, Xavier D, Liu L, Zhang H, Rao-Melacini P, Zhang X, Pais P, Agapay S: **Global and regional effects of potentially modifiable risk factors associated with acute stroke in 32 countries (INTERSTROKE): a case-control study**. *Lancet* 2016, **388**(10046):761-775.
8. Bosu WK, Reilly ST, Aheto JMK, Zucchelli E: **Hypertension in older adults in Africa: A systematic review and meta-analysis**. *PLoS One* 2019, **14**(4):e0214934.
9. Adeloye D, Basquill C: **Estimating the prevalence and awareness rates of hypertension in Africa: a systematic analysis**. *PLoS One* 2014, **9**(8):e104300.
10. Zhou B, Bentham J, Di Cesare M, Bixby H, Danaei G, Cowan MJ, Paciorek CJ, Singh G, Hajifathalian K, Bennett JE *et al*: **Worldwide trends in blood pressure from 1975 to 2015: a pooled analysis of 1479 population-based measurement studies with 19.1 million participants**. *Lancet* 2017, **389**(10064):37-55.
11. Dramé ML, Houehanou C, Sogbohossou P, Paré R, Ekambi A, Mizéhoun-Adissoda C, Houinato D, Gyselinck K, Marx M, Martins MRO *et al*: **Determinants of High Blood Pressure and Quality of Management in Three Regions of Benin**. *Open Journal of Epidemiology* 2018, **8**(01):14.
12. Bosu WK: **Epidemic of hypertension in Ghana: a systematic review**. *BMC Public Health* 2010, **10**:418.
13. Joseph-Shehu EM, Ncama BP: **Evaluation of health status and its predictor among university staff in Nigeria**. *BMC Cardiovasc Disord* 2018, **18**(1):183.
14. Mohamed SF, Mutua MK, Wamai R, Wekesah F, Haregu T, Juma P, Nyanjau L, Kyobutungi C, Ogola E: **Prevalence, awareness, treatment and control of hypertension and their determinants: results from a national survey in Kenya**. *BMC Public Health* 2018, **18**(3):1219.
15. Laxmaiah A, Meshram IL, Arlappa N, Balakrishna N, Rao KM, Reddy CG, Ravindranath M, Kumar S, Kumar H, Brahman GN: **Socio-economic & demographic determinants of hypertension & knowledge, practices & risk behaviour of tribals in India**. *Indian J Med Res* 2015, **141**(5):697-708.
16. Nyarko SH: **Prevalence and Sociodemographic Determinants of Hypertension History among Women in Reproductive Age in Ghana**. *Int J Hypertens* 2016, **2016**:3292938.
17. Addo J, Amoah AGB, Koram KA: **The changing patterns of hypertension in Ghana: a study of four rural communities in the Ga district**. *Ethn Dis* 2006, **16**:894-899.

18. Okeahialam BN, Ogbonna C, Otokwula AE, Joseph DE, Chuhwak EK, Isiguzoro IO: **Cardiovascular epidemiological transition in a rural habitat of Nigeria: the case of Mangu Local Government Area.** *West Afr J Med* 2012, **31**(1):14-18.
19. Awuah RB, Anarfi JK, Agyemang C, Ogedegbe G, Aikins Ad-G: **Prevalence, awareness, treatment and control of hypertension in urban poor communities in Accra, Ghana.** *J Hypertens* 2014, **32**(6):1203-1210.
20. United Nations General Assembly: **Political declaration of the third high-level meeting of the General Assembly on the prevention and control of non-communicable diseases.** In: *Resolution adopted by the General Assembly on 10 October 2018: 73rd Session, Agenda Item 119.* New York: UN; 2018.
21. Islam SM, Purnat TD, Phuong NT, Mwingira U, Schacht K, Fröschl G: **Non-communicable diseases (NCDs) in developing countries: a symposium report.** *Global Health* 2014, **10**:81.
22. Charlton K, Ware LJ, Menyana E, Biritwum RB, Naidoo N, Pieterse C, Madurai SL, Baumgartner J, Asare GA, Thiele E *et al*: **Leveraging ongoing research to evaluate the health impacts of South Africa's salt reduction strategy: a prospective nested cohort within the WHO-SAGE multicountry, longitudinal study.** *BMJ Open* 2016, **6**(11):e013316.
23. Echouffo-Tcheugui JB, Batty GD, Kivimäki M, Kengne AP: **Risk models to predict hypertension: a systematic review.** *PLoS One* 2013, **8**(7):e67370.
24. Fava C, Sjögren M, Montagnana M, Danese E, Almgren P, Engström G, Nilsson P, Hedblad B, Guidi GC, Minuz P *et al*: **Prediction of blood pressure changes over time and incidence of hypertension by a genetic risk score in Swedes.** *Hypertension* 2013, **61**(2):319-326.
25. Soubeiga JK, Millogo T, Bicaba BW, Doulougou B, Kouanda S: **Prevalence and factors associated with hypertension in Burkina Faso: a countrywide cross-sectional study.** *BMC Public Health* 2017, **17**(1):64.
26. Goldstein H: **Multilevel Statistical Models**, 3rd edn. London: Arnold; 2003.
27. Steele F, Goldstein H, Rao C, Sinharay S: **Multilevel Models in Psychometrics.** *Psychometrics* 2007, **26**:401-420.
28. Hedeker D, Gibbons RD: **MIXOR: a computer program for mixed-effects ordinal regression analysis.** *Comput Methods Programs Biomed* 1996, **49**(2):157-176.
29. Hair JFJ, Anderson RE, Tatham RL, Black WC: **Multivariate Data Analysis** 3edn. New York: Macmillan; 1995.
30. StataCorp: **Stata Statistical Software: Release 14.** In. College Station, TX: StataCorp LP. ; 2015.
31. Mohamed SF, Mutua MK, Wamai R, Wekesah F, Haregu T, Juma P, Nyanjau L, Kyobutungi C, Ogola E: **Prevalence, awareness, treatment and control of hypertension and their determinants: results from a national survey in Kenya.** *BMC Public Health* 2018, **18**(Suppl 3):1219.
32. Mosha NR, Mahande M, Juma A, Mboya I, Peck R, Urassa M, Michael D, Todd J: **Prevalence, awareness and factors associated with hypertension in North West Tanzania.** *Glob Health Action* 2017, **10**(1):1321279.
33. Ogah OS, Madukwe OO, Chukwuonye II, Onyeonoro UU, Ukegbu AU, Akhimien MO, Onwubere BJ, Okpechi IG: **Prevalence and determinants of hypertension in Abia State Nigeria: results from the Abia State Non-Communicable Diseases and Cardiovascular Risk Factors Survey.** *Ethn Dis* 2013, **23**(2):161-167.
34. Aheto JMK, Keegan TJ, Taylor BM, Diggle PJ: **Childhood Malnutrition and Its Determinants among Under-Five Children in Ghana.** In., vol. 29: Paediatric and Perinatal Epidemiology; 2015: 552-561.
35. Aheto JMK: **Predictive model and determinants of under-five child mortality: evidence from the 2014 Ghana demographic and health survey.** *BMC Public Health* 2019, **19**(1):64.

36. Adekanmbi VT, Kayode GA, Uthman OA: **Individual and contextual factors associated with childhood stunting in Nigeria: a multilevel analysis.** *Maternal and Child Nutrition* 2013, **9**(2):244-259.
37. Ware LJ, Chidumwa G, Charlton K, Schutte AE, Kowal P: **Predictors of hypertension awareness, treatment and control in South Africa: results from the WHO-SAGE population survey (Wave 2).** *J Hum Hypertens* 2019, **33**(2):157-166.
38. Bosu WK: **Determinants of Mean Blood Pressure and Hypertension among Workers in West Africa.** *Int J Hypertens* 2016, **2016**:3192149.
39. Aronow WS, Fleg JL, Pepine CJ, Artinian NT, Bakris G, Brown AS, Ferdinand KC, Ann Forciea M, Frishman WH, Jaigobin C *et al*: **ACCF/AHA 2011 expert consensus document on hypertension in the elderly: a report of the American College of Cardiology Foundation Task Force on Clinical Expert Consensus documents developed in collaboration with the American Academy of Neurology, American Geriatrics Society, American Society for Preventive Cardiology, American Society of Hypertension, American Society of Nephrology, Association of Black Cardiologists, and European Society of Hypertension.** *J Am Coll Cardiol* 2011, **57**(20):2037-2114.
40. Williams B: **Vascular ageing and interventions: lessons and learnings.** *Ther Adv Cardiovasc Dis* 2016, **10**(3):126-132.
41. Manzoli L, Villari P, M Pirone G, Boccia A: **Marital status and mortality in the elderly: a systematic review and meta-analysis.** *Soc Sci Med* 2007, **64**(1):77-94.
42. Kaplan RM, Kronick RG: **Marital status and longevity in the United States population.** *J Epidemiol Community Health* 2006, **60**(9):760-765.
43. Schwandt HM, Coresh J, Hindin MJ: **Marital Status, Hypertension, Coronary Heart Disease, Diabetes, and Death Among African American Women and Men: Incidence and Prevalence in the Atherosclerosis Risk in Communities (ARIC) Study Participants.** *J Fam Issues* 2010, **31**(9):1211-1229.
44. Brummett BH, Barefoot JC, Siegler IC, Clapp-Channing NE, Lytle BL, Bosworth HB, Williams RB, Mark DB: **Characteristics of socially isolated patients with coronary artery disease who are at elevated risk for mortality.** *Psychosom Med* 2001, **63**(2):267-272.
45. Lett HS, Blumenthal JA, Babyak MA, Strauman TJ, Robins C, Sherwood A: **Social support and coronary heart disease: epidemiologic evidence and implications for treatment.** *Psychosom Med* 2005, **67**(6):869-878.
46. House JS: **Social isolation kills, but how and why?** *Psychosom Med* 2001, **63**(2):273-274.
47. Ibrahim MM, Damasceno A: **Hypertension in developing countries.** *Lancet* 2012, **380**(9841):611-619.
48. Chalmers J: **The 1999 WHO-ISH Guidelines for the Management of Hypertension.** *Med J Aust* 1999, **171**(9):458-459.
49. Akintunde AA, Akinwusi PO, Adebayo RA, Ogunyemi S, Opadijo OG: **Burden of obesity in essential hypertension: pattern and prevalence.** *Niger J Clin Pract* 2010, **13**(4):399-402.
50. Lawrence WF, Fryback DG, Martin PA, Klein R, Klein BEK: **Health status and hypertension: A population-based study.** *Journal of Clinical Epidemiology* 1996, **49**(11):1239-1245.
51. de-Graft Aikins A, Addo J, Ofei F, Bosu W, Agyemang C: **Ghana's burden of chronic non-communicable diseases: future directions in research, practice and policy.** *Ghana Med J* 2012, **46**(2 Suppl):1-3.
52. Ministry of Health (MOH): **National Policy for the Prevention and Control of Chronic Non-Communicable Diseases in Ghana.** In. Accra: MOH; 2012.