Integrated annotations and analyses of small RNA-producing loci from 47 diverse plants

Alice Lunardon¹, Nathan R. Johnson^{1,2}, Emily Hagerott³, Tamia Phifer³, Seth Polydore^{1,2,4}, Ceyda Coruh^{1,2,5}, and Michael J. Axtell^{1,2}

Corresponding author: Michael J. Axtell (mja18@psu.edu)

¹ Department of Biology, The Pennsylvania State University, University Park, PA 16802 USA

² Huck Institutes of the Life Sciences, The Pennsylvania State University, University Park, PA 16802 USA

³ Department of Biology, Knox College, Galesburg, IL 61401 USA

⁴ Present address: Donald Danforth Plant Science Center, St. Louis, MO 63132 USA

⁵ Present address: Salk Institute for Biological Studies, La Jolla, CA 92037 USA

Running title

Annotation and analysis of sRNA loci from 47 plant species

Keywords: small RNAs, siRNAs, microRNAs, protein-coding genes, RdDM, transposons

Abstract

Plant endogenous small RNAs (sRNAs) are important regulators of gene expression. There are two broad categories of plant sRNAs: microRNAs (miRNAs) and endogenous short interfering RNAs (siRNAs). MicroRNA loci are relatively well-annotated but comprise only a small minority of the total sRNA pool; siRNA locus annotations have lagged far behind. Here, we used a large dataset of published and newly generated sRNA sequencing data (1,333 sRNA-seq libraries containing over 20 billion reads) and a uniform bioinformatic pipeline to produce comprehensive sRNA locus annotations of 47 diverse plants, yielding over 2.7 million sRNA loci. The two most numerous classes of siRNA loci produced mainly 24 nucleotide and 21 nucleotide siRNAs, respectively. 24 nucleotide-dominated siRNA loci usually occurred in intergenic regions, especially at the 5'-flanking regions of protein-coding genes. In contrast, 21 nucleotide-dominated siRNA loci were most often derived from double-stranded RNA precursors copied from spliced mRNAs. Genic 21 nucleotide-dominated loci were especially common from disease resistance genes, including from a large number of monocots. Individual siRNA sequences of all types showed very little conservation across species, while mature miRNAs were more likely to be conserved. We developed a web server where our data and several search and analysis tools are freely accessible at <http://plantsmallrnagenes.science.psu.edu>.

Introduction

Plant regulatory small RNAs (sRNAs) play important roles in almost all biological processes. Endogenous sRNAs are 20-24 nucleotides in length and derive from longer RNA precursors that are processed by DICER-LIKE (DCL) ribonucleases. Once processed, they are loaded into Argonaute (AGO) proteins to form the RNA-induced silencing complex (RISC). Then, sRNAs guide the RISC complex to complementary sites on target RNAs, inducing either post-transcriptional or transcriptional gene silencing.

Endogenous sRNAs can be grouped in two broad classes based on their biogenesis and typical functions: microRNAs (miRNAs) and small interfering RNAs (siRNAs) (Axtell 2013a). MiRNAs are typically 21-22 nucleotides long, processed from single-stranded RNA (ssRNA) stem-loop precursors by DCL1, and regulate gene expression post-transcriptionally, directing mRNA degradation and translational repression (Rogers and Chen 2013). SiRNAs are processed from double-stranded RNA (dsRNA) precursors and are categorized in multiple sub-classes. The most abundant sub-class of siRNAs participates in the RNA-directed DNA methylation (RdDM) pathway, involving 24 or 21-22 nucleotide siRNAs. 24 nucleotide siRNAs are derived from Polymerase IV (Pol IV) transcripts that are converted to dsRNAs by RNA-dependent RNA polymerase 2 (RDR2) which are then processed by DCL3. They act in "canonical" RdDM, primarily targeting transposable elements (TEs) and other repeats to induce DNA methylation and reinforce transcriptional silencing. 21-22 nucleotide siRNAs are derived from Pol II transcripts and are copied by RDR6 into dsRNAs and processed by DCL2/DCL4. They act in the non-canonical RdDM pathway to establish the silencing of young TEs, both transcriptionally and post-transcriptionally (Nuthikattu et al. 2013). Another major siRNA sub-class is secondary siRNAs. Their biogenesis is triggered by a miRNA-directed cleavage of a coding or non-coding transcript. The transcript is then converted to dsRNA by RDR6 and processed by DCL proteins

into secondary siRNAs in a phased pattern relative to the miRNA cut site. Phased secondary siRNAs (phasiRNAs) are typically 21 or 22 nucleotides long, however a specific population of 24 nucleotide phasiRNAs has been detected in anthers of many angiosperms (Xia et al. 2019). *TAS* genes are an example of loci generating non-coding RNA precursors that produce secondary siRNAs, which act *in trans* (trans-acting siRNAs, tasiRNAs) on other targets and direct their cleavage (Allen et al. 2005). Pentatricopeptide repeat (PPR) genes are the first reported protein-coding genes generating secondary siRNAs in *Arabidopsis thaliana* (Howell et al. 2007).

At the chromosomal level, sRNA distribution correlates with gene density, typically lower in the centromeric and pericentromeric regions and enriched in the distal euchromatic regions. This trend has been observed in maize (He et al. 2013), rice (Wei et al. 2014), tomato (The Tomato Genome Consortium 2012), hot pepper (Kim et al. 2014), upland cotton (Song et al. 2015) and sugar beet (Dohm et al. 2014). However, in a smaller number of species sRNAs mostly arise from centromeric and pericentromeric regions away from genes, as shown in *A. thaliana* (Kasschau et al. 2007; Ha et al. 2009), soybean (Schmitz et al. 2013), cucumber (Lai et al. 2017) and *Brachypodium distachyon* (The International Brachypodium Initiative 2010).

Despite differences in chromosomal distributions, the sRNA profiles near protein-coding genes are conserved amongst plant species, with 24 nucleotide siRNAs preferentially found in gene-proximal regions but depleted in gene bodies themselves. This pattern has been described in maize (Gent et al. 2013), rice (Wei et al. 2014), rapeseed (Shen et al. 2017), Chinese cabbage (Woodhouse et al. 2014), soybean (Song et al. 2013), upland cotton (Song et al. 2015) and *A. thaliana* (Kasschau et al. 2007; Ha et al. 2009). Depending on the species, sRNAs have opposite effects on the regulation of the proximal downstream genes. In maize, 24 nucleotide siRNAs are found with higher probability near expressed genes than non-expressed genes (Gent et al. 2013; Lunardon et al. 2016). Here, siRNAs participate in RdDM to reinforce the silencing of TEs that are inserted upstream of genes, where the chromatin is accessible, therefore repressing the potentially deleterious Pol II transcription of TEs (Gent et al. 2014). In contrast, the siRNA-mediated silencing of TEs near genes is linked to lower expression of the genes in *A. thaliana* and Chinese cabbage (Hollister et al. 2011; Woodhouse et al. 2014). In addition to target TEs near genes, 24 nucleotide siRNAs can also target TEs inserted inside genes, affecting their expression (Wei et al. 2014; Lunardon et al. 2016).

Genome-wide analyses in barley, soybean, *Medicago truncatula* and *Physcomitrella patens* showed that 21 nucleotide siRNAs are not enriched in gene body regions (Hackenberg et al. 2016; Schmitz et al. 2013; Lelandais-Brière et al. 2009; Coruh et al. 2015). Nevertheless, there are many cases of well characterized genes generating 21 nucleotide phasiRNAs in dicots: nucleotide binding/leucine-rich repeat (NB-LRR) and receptor like kinase (RLK) resistance genes, PPR genes, auxin-responsive factor (ARF) genes, MYB and NAC transcription factors and F-BOX genes (Arikiti et al. 2014; Hu et al. 2015a; Xia et al. 2015b). NB-LRR genes evolve rapidly by tandem duplication and they are controlled by sRNA-mediated silencing to avoid their over-expression and prevent autoimmune responses (Yang and Huang 2014). This mechanism is conserved in a large number of dicots: soybean, *M. truncatula*, common bean, chickpea, *Populus trichocarpa*, cassava, pima cotton, potato and Norway spruce (Zhai et al. 2011; Formey et al. 2015; Srivastava et al. 2015; Klevebring et al. 2009; Xia et al. 2014; Hu et al. 2015b; Xia et al. 2015a). Amongst monocots, 21 nucleotide phasiRNAs from NB-LRR genes have been only found in barley and wheat so far (Liu et al. 2014; Zhang et al. 2019). This is consistent with the

fact that monocots, in contrast to dicots, produce phasiRNAs mainly from non-coding RNAs (Komiya 2017; Zheng et al. 2015).

The conservation of sRNAs across plants has been widely investigated for miRNAs. There are deeply conserved miRNA families together with their targets, suggesting common functional regulatory networks (Axtell and Bowman 2008). However, the majority of miRNA sequences are species-specific, indicating the presence of numerous young or still evolving miRNAs (Cuperus et al. 2011; Chávez Montes et al. 2014). Much less is known about conservation of siRNAs but a study comparing *Arabidopsis thaliana* and *Arabidopsis lyrata* suggested that individual siRNA sequences are not conserved even between closely related species (Ma et al. 2010). Moreover, while in most species analyzed so far the 24 nucleotide siRNAs are the most abundant expressed group of sRNAs, mosses, lycophytes and conifers lack a strong peak of 24 nucleotide siRNAs (Axtell and Bartel 2005; Banks et al. 2011; Dolgosheina et al. 2008).

There are several existing web-based resources that serve sRNA sequencing (sRNA-seq) data for multiple plants. The Cereal small RNA Database contains maize and rice genome browsers with accessible sRNA-seq data (Johnson et al. 2007). The Pln24NT website stores annotations and sequences of 24 nucleotide siRNA reads and loci for 10 species (Liu et al. 2017). The Next-Gen Sequence Databases produced by the Meyers lab contain sRNA-seq and other high-throughput data with custom-built genome browsers and search functions for 27 species (Nakano et al. 2006). The miRBase database (Kozomara and Griffiths-Jones 2014) provides curated, comprehensive annotations of *MIRNA* loci in a very large number of species. An equivalent database for the storage and distribution of reference annotations of siRNA-producing loci in a vast number of plant genomes does not exist (Coruh et al. 2014).

In this study, we used a large dataset of published and newly generated sRNA-seq data, that we processed with a consistent pipeline, to create reference sRNA loci annotations for 47 plant species, including model plants and crops. We propose and use a systematic nomenclature and ontology for sRNA-producing loci that is consistent with their biology and easily traceable and updatable. We examined the genome-wide distribution of sRNA loci relative to protein-coding genes and compared it across species, providing insights into conserved sRNA functions. We organized the sRNA-seq alignment data and sRNA loci annotations in a freely available web-based database that represents an important public resource for future studies aimed to understand the biological function of sRNAs.

Results

Identification and classification of sRNA loci in 47 plants

We obtained and analyzed 48 plant genome assemblies, representing 47 different species (Table 1; two independent assemblies of *Cuscuta campestris* were analyzed). To facilitate succinct communication in figures and our database, a short code was designated for each assembly. The code begins with a three-letter prefix representing the genus and species, following the abbreviations established by miRBase (Kozomara et al. 2019). The second part of the code indicates the genome build ('-b') version in use. These genome assemblies varied widely in size, contiguity, protein-coding gene number, and repeat content (Supplemental Fig.

S1,S2). Most genome assemblies were from crops; others included the model plants *Arabidopsis thaliana* and *Medicago truncatula*, the parasitic plant *Cuscuta campestris*, and representatives of diverse lineages (*Amborella trichopoda* [basal angiosperm], *Picea abies* [gymnosperm], *Physcomitrella patens* [bryophyte], and *Marchantia polymorpha* [bryophyte]).

Table 1. Plants included in this study

Common Name	Binomial Name	Code	Group	Order	Family
Thale Cress	<i>Arabidopsis thaliana</i>	ath-b10	Core Eudicots - Rosids	Brassicales	Brassicaceae
Rapeseed	<i>Brassica napus</i>	bn-a-b1	Core Eudicots - Rosids	Brassicales	Brassicaceae
Cabbage	<i>Brassica oleracea var. capitata</i>	bol-b1.0	Core Eudicots - Rosids	Brassicales	Brassicaceae
Chinese Cabbage	<i>Brassica rapa var. pekinensis</i>	bra-b1	Core Eudicots - Rosids	Brassicales	Brassicaceae
Papaya	<i>Carica papaya</i>	cpa-b0.4	Core Eudicots - Rosids	Brassicales	Caricaceae
Watermelon	<i>Citrullus lanatus</i>	clt-b1	Core Eudicots - Rosids	Cucurbitales	Cucurbitaceae
Cucumber	<i>Cucumis sativus</i>	csa-b2	Core Eudicots - Rosids	Cucurbitales	Cucurbitaceae
Chickpea	<i>Cicer arietinum</i>	car-b2.0	Core Eudicots - Rosids	Fabales	Fabaceae
Soybean	<i>Glycine max</i>	gma-b1.0	Core Eudicots - Rosids	Fabales	Fabaceae
Barrel Medic	<i>Medicago truncatula</i>	mtr-b4.0	Core Eudicots - Rosids	Fabales	Fabaceae
Common Bean	<i>Phaseolus vulgaris</i>	pvu-b1.0	Core Eudicots - Rosids	Fabales	Fabaceae
Rubber Tree	<i>Hevea brasiliensis</i>	hbr-b0	Core Eudicots - Rosids	Malpighiales	Euphorbiaceae
Cassava	<i>Manihot esculenta</i>	mes-b6	Core Eudicots - Rosids	Malpighiales	Euphorbiaceae
Black Cottonwood	<i>Populus trichocarpa</i>	ptc-b3.0	Core Eudicots - Rosids	Malpighiales	Salicaceae

Pima Cotton	<i>Gossypium barbadense</i>	gba-b1.0	Core Eudicots - Rosids	Malvales	Malvaceae
Upland Cotton	<i>Gossypium hirsutum</i>	ghr-b1.1	Core Eudicots - Rosids	Malvales	Malvaceae
Cacao	<i>Theobroma cacao</i>	tcc-b1.1	Core Eudicots - Rosids	Malvales	Malvaceae
Strawberry	<i>Fragaria x ananassa</i>	fan-b1.0	Core Eudicots - Rosids	Rosales	Rosaceae
Woodland Strawberry	<i>Fragaria vesca</i>	fve-b2.0	Core Eudicots - Rosids	Rosales	Rosaceae
Apple	<i>Malus x domestica</i>	mdm- b3.0	Core Eudicots - Rosids	Rosales	Rosaceae
Peach	<i>Prunus persica</i>	ppe-b2.0	Core Eudicots - Rosids	Rosales	Rosaceae
Clementine	<i>Citrus clementina</i>	ccl-b1	Core Eudicots - Rosids	Sapindales	Rutaceae
Sweet Orange	<i>Citrus sinensis</i>	csi-b2	Core Eudicots - Rosids	Sapindales	Rutaceae
Carrot	<i>Daucus carota</i>	dca-b2.0	Core Eudicots - Asterids	Apiales	Apiaceae
Lettuce	<i>Lactuca sativa</i>	lsa-b8	Core Eudicots - Asterids	Asterales	Asteraceae
Olive Tree	<i>Olea europaea</i>	oeu-b6	Core Eudicots - Asterids	Lamiales	Oleaceae
Field Dodder	<i>Cuscuta campestris</i>	ccm- b0.32	Core Eudicots - Asterids	Solanales	Convolvulaceae
Field Dodder	<i>Cuscuta campestris</i>	ccm-b0.1	Core Eudicots - Asterids	Solanales	Convolvulaceae
Pepper	<i>Capsicum annuum</i>	can-b1.6	Core Eudicots - Asterids	Solanales	Solanaceae
Tobacco	<i>Nicotiana tabacum</i>	nta-b0	Core Eudicots - Asterids	Solanales	Solanaceae
Tomato	<i>Solanum lycopersicum</i>	sly-b2.5	Core Eudicots - Asterids	Solanales	Solanaceae

Potato	<i>Solanum tuberosum</i>	stu-b4.04	Core Eudicots - Asterids	Solanales	Solanaceae
Beet	<i>Beta vulgaris</i>	bvu-b1.2.2	Core Eudicots	Caryophyllales	Amaranthaceae
Quinoa	<i>Chenopodium quinoa</i>	cqi-b1.0	Core Eudicots	Caryophyllales	Amaranthaceae
Spinach	<i>Spinacia oleracea</i>	sol-b1	Core Eudicots	Caryophyllales	Amaranthaceae
African Oil Palm	<i>Elaeis guineensis</i>	egu-b5.1	Monocots	Arecales	Arecaceae
Stiff brome	<i>Brachypodium distachyon</i>	bdi-b1.0	Monocots	Poales	Poaceae
Barley	<i>Hordeum vulgare</i>	hvu-b1	Monocots	Poales	Poaceae
Rice	<i>Oryza sativa</i>	osa-b1.0	Monocots	Poales	Poaceae
Sorghum	<i>Sorghum bicolor</i>	sbi-b3.0	Monocots	Poales	Poaceae
Foxtail Millet	<i>Setaria italica</i>	sit-b2	Monocots	Poales	Poaceae
Wheat	<i>Triticum aestivum</i>	tae-b1	Monocots	Poales	Poaceae
Maize	<i>Zea mays</i>	zma-b4	Monocots	Poales	Poaceae
Banana	<i>Musa acuminata</i>	mac-b2	Monocots	Zingiberales	Musaceae
Amborella	<i>Amborella trichopoda</i>	atr-b1	Basal Angiosperms	Amborellales	Amborellaceae
Norway Spruce	<i>Picea abies</i>	pab-b1.0c	Gymnosperms	Pinales	Pinaceae
Spreading Earthmoss	<i>Physcomitrella patens</i>	ppt-b3.0	Bryophytes	Funariales	Funariaceae
Common Liverwort	<i>Marchantia polymorpha</i>	mpo-b3.0	Bryophytes	Marchantiales	Marchantiaceae

We gathered sRNA-seq libraries from each genome (Figure 1A). In most cases, these data were from public sequencing archives (Supplemental Table S1). In a few cases, we also generated novel sRNA-seq libraries (*Zea mays*, *Spinacia oleracea*, *Daucus carota*, *Theobroma cacao*; Supplemental Table S1). We sought to annotate the full diversity of sRNA loci and thus selected libraries with the goal of including as many different tissues and conditions as possible. However, we excluded low-depth sRNA-seq datasets (less than two million reads aligned to the genome) and also excluded sRNA-seq datasets from mutants known to affect sRNA biogenesis or stability. For each given genome assembly, all cognate sRNA-seq libraries were aligned and then merged into a single master sRNA alignment which we call the "reference set" (Figure 1A). Reference sets had considerable variation in both total number of sRNA reads (minimum: 2.1E6, median: 1.6E8, maximum: 4.1E9) and in number of contributing sRNA-seq libraries (minimum: 1, median: 11, maximum: 161)(Supplemental Fig. S3).

For annotation, we first identified genomic regions producing sRNAs, independently in all sRNA-seq libraries with ShortStack (Axtell 2013b; Johnson et al. 2016). Then we compared the sRNA expression from different samples of the same species and identified the regions that were robustly expressing sRNAs in at least three separate samples. Millions of discrete sRNA clusters were annotated in this way and defined as sRNA-producing loci, which were then

analyzed in the genome-aligned reference sets. Canonical plant miRNAs and siRNAs are between 20 and 24 nucleotides in length, while other types of sRNA loci produce a broader range of RNA sizes. For each locus, we computed the fraction of aligned sRNA-seq reads that were 20-24 nucleotides long. We found that these fractions had consistent bimodal distributions in the various genomes (Figure 1B). Based on these distributions, we used a cutoff of 80% to discriminate canonical siRNA/*MIRNA* loci from 'OtherRNA' loci (Figure 1C). We then developed a simplified ontology to describe the siRNA and *MIRNA* loci: 'MIRNA' loci were those that met all *MIRNA* annotation criteria, while 'nearMIRNA' loci met most criteria except for that the exact predicted miRNA*, the complementary strand to the mature miRNA in the miRNA-miRNA* duplex, was not sequenced. The remaining loci were classified as siRNA loci based on the predominant length of aligned sRNAs within each locus (Figure 1C). This ontology has the advantage of being applicable to any genome regardless of any other annotations or information. We also devised a simple nomenclature to systematically name the sRNA loci (Figure 1D). In total, we annotated approximately 2.7E6 sRNA-producing loci from the 48 genome assemblies (Supplemental Table S2; also see <http://plantsmallrnagenes.science.psu.edu> for easier access and more analysis options).

The 'OtherRNA' category of loci, defined by having less than 80% of aligned reads with sizes between 20-24 nucleotides in length, typically comprised less than half of all loci in the flowering plants (Supplemental Fig. S4A,B). In contrast, the majority of loci identified in one gymnosperm and two bryophyte genomes were annotated as OtherRNA (Supplemental Fig. S4A,B). Across all taxa, OtherRNA loci typically contributed large fractions of total read abundance (Supplemental Fig. S4C,D). This is because many of the OtherRNA loci represented clusters of short fragments derived from highly abundant, longer RNAs, such as rRNAs, tRNAs, and plastid-derived mRNAs. There is evidence that some plant RNAs longer than 24 nucleotides, or shorter than 20 nucleotides, may function as gene-regulatory factors (Martinez et al. 2017); such loci will have been annotated in the OtherRNA category by our procedure. Nonetheless, we focused our subsequent analyses on the MIRNA, nearMIRNA, and siRNA loci dominated by 20-24 nucleotide RNAs because these sizes are most clearly associated with production by DCL endonucleases and usage by AGO proteins. By default, ShortStack assigns a phasing score to the sRNA loci based on the algorithm described in Guo et al. 2015. However, an accurate annotation of the phasing would require a more complex study to avoid false positives that may be produced by the commonly used phasing-detecting algorithms (Polydore et al. 2018). Therefore, we did not further analyze the phasing of the sRNA loci in this analysis.

After excluding OtherRNA loci, the remaining loci were mostly designated siRNA24 in angiosperms (Figure 1E-F). In contrast, and consistent with prior reports (Dolgosheina et al. 2008; Axtell and Bartel 2005), gymnosperm and bryophyte loci were less dominated by the siRNA24 type and instead had more siRNA21 loci. When tallied by sRNA abundance, MIRNA and siRNA21 loci made substantial contributions in all taxa (Figure 1G-H). This indicates that a relatively small number of MIRNA and siRNA21 loci produce high levels of their respective sRNAs. In a number of species, the proportion of 22 nucleotide siRNAs was also substantial and this trend was particularly consistent amongst the asterids (Figure 1H). In most cases, angiosperms had more annotated sRNA loci compared to non-angiosperms (Supplemental Fig. S4A, Figure 1E). However, that comparison is potentially complicated by the different amounts of input sRNA reads used for each species (Supplemental Fig. S3).

221

222

223

Figure 1. Overview of sRNA locus annotation pipeline and summary of annotated sRNA loci. **(A)** Schematic illustrating how multiple sRNA libraries from diverse plant tissues are merged to create a 'reference set' of sRNAs for a given species. Accession numbers shown are fictional. **(B)** Distributions of the fractions of sRNAs between 20-24 nucleotides in length (inclusive) within all loci in each genome. Gray line at 80% represents the cutoff used to discriminate silencing-related RNA loci from other types of sRNA-producing loci. **(C)** Flowchart illustrating the ontology used to classify sRNA-producing loci. Colors designating different locus types are used throughout this work. **(D)** Schematic illustrating the nomenclature used to annotate sRNA-producing loci. **(E-H)** Summary of annotated sRNA loci, by species and locus type, excluding the category 'OtherRNA' **(E)** Counts of annotated loci. **(F)** Proportions of annotated loci. **(G)** Total counts of aligned small RNAs in reference sets. **(H)** Proportions of small RNA total read counts in reference sets. See Table 1 for species codes. Eu: eudicots, BA: basal angiosperm, G: gymnosperm, Br: bryophyte.

The plantsmallrnagenes.science.psu.edu server

All data and analyses from this study have been systematically organized and are freely available at <https://plantsmallrnagenes.science.psu.edu>. Users can search for loci of interest by sRNA sequence, *MIRNA* family name, locus name, or by BLAST-based homology searches. A JBrowse-based genome browser is available for each of the 48 genomes. Genome browsers are customized to display sRNA-seq data based on sRNA size, strand, and multi-mapping (Figure 2A). Genome browsers also allow users to highlight a region of interest and perform on the fly analyses, including ShortStack (Axtell 2013b; Johnson et al. 2016) and visualization of possible *MIRNA* hairpins (Figure 2B). Bulk data are also available in standard, widely used formats: sRNA-seq alignments are in the BAM format, while annotations of sRNA loci are in the GFF3 format. It is our intention to maintain and expand this resource for the benefit of anyone interested in the analysis of plant sRNA-producing loci.

Figure 2. Example screenshots from <https://plantsmallrnagenes.science.psu.edu> **(A)** Screenshot of genome browser for a region of *Phaseolus vulgaris* chromosome 9. Coverage track shows sRNA-seq alignment depths from the reference set, separated by sRNA lengths (indicated by colors). ShortStack sRNA loci track shows sRNA locus annotations. Alignments track shows individual sRNA reads from the reference set, with lengths indicated by colors. Hollow bars indicate multi-mapped reads; solid bars are uniquely mapped reads. A user-highlighted region is indicated in yellow. **(B)** Analysis of predicted RNA secondary structure with sRNA-alignment depths indicated by colors. This analysis is one of several that can be triggered by user selection of a region of interest (yellow region in panel A).

Chromosomal distribution of sRNA loci and association with protein-coding genes

Where feasible based on genome assembly quality, we compared the distribution of sRNA loci and genes across entire chromosomes and confirmed that the most common trend is a positive correlation between gene density and sRNA density (Supplemental Fig. S5), as has previously been shown in several prior species-specific studies (He et al. 2013; Wei et al. 2014; The Tomato Genome Consortium 2012; Kim et al. 2014; Song et al. 2015; Dohm et al. 2014). *A. thaliana* is unique in that it has a clear trend from telomeres to centromeres of decreasing gene density and increasing sRNA loci density (Kasschau et al. 2007; Ha et al. 2009). Surprisingly, rice showed a similar trend to *A. thaliana* (Supplemental Fig. S5). Chinese cabbage and sweet orange also showed a slight inverse correlation between the gene and the sRNA loci distributions. Finally, soybean had a general positive correlation between genes and sRNA loci but in the most distal segments of the chromosome arms it showed a local negative correlation (Supplemental Fig. S5).

We examined siRNA21 loci and siRNA24 loci locations relative to protein-coding genes. Other types of sRNA loci were excluded due to their lower frequencies. Coverage of protein-

coding genes and flanking 5kb regions by siRNA21 or siRNA24 loci was calculated and normalized. siRNA21 loci had a striking tendency in nearly all taxa to overlap with protein-coding genes (Figure 3A). In contrast, siRNA24 loci were strongly depleted in protein-coding genes in most angiosperms (Figure 3B). siRNA24 loci were often strongly enriched in the 5'-proximal regions upstream of protein-coding genes. There were, however, some notable exceptions to this pattern. There was no upstream peak of siRNA24 loci in bryophytes and the gymnosperm (Figure 3B), which is consistent with the generally low levels of siRNA24 loci in these taxa (Figure 1). The basal angiosperm *Amborella trichopoda* was unusual in that siRNA24 loci were not depleted in gene bodies at all (Figure 3B). Finally, the model plant *A. thaliana* also lacked a conspicuous upstream gene-proximal enrichment of siRNA24 loci. This observation, together with the unique chromosomal distribution of sRNA loci in *A. thaliana*, suggests that *A. thaliana* may not be representative of most angiosperms in its genome-wide patterns of sRNA loci.

Figure 3. Associations of siRNA21 loci and siRNA24 loci with protein-coding genes. **(A)** Heatmap showing normalized coverage of protein-coding genes +/- 5kb by siRNA21 loci. Each row is a given species (See Table 1 for species codes), grouped taxonomically. Eu: eudicots, BA: basal angiosperm, G: gymnosperm, Br: bryophyte. Negative and positive numbers are upstream and downstream regions, respectively (in nucleotides). The region from -1 to +1 represents the gene bodies, scaled to a uniform size of 1,000 nominal units of 0.1% each. **(B)** As in A, except for siRNA24 loci.

Distribution of sRNAs in exons and introns of protein-coding genes

We then analyzed the distribution of sRNAs mapped to protein-coding genes, relative to the mRNA exons/introns and relative to the coding/non-coding strand of the mRNA (Figure 4). Although siRNA24 loci were generally depleted in mRNAs (Figure 3), their very large numbers still resulted in many overlaps, and therefore they were included in this analysis (Supplemental

Fig. S6). For each species, we calculated the proportion of mRNAs that have 0% to 100% sRNAs mapped to the exons and the proportion of mRNAs that have 0% to 100% sRNAs mapped to the same strand of the mRNA. The proportions were plotted separately for mRNAs containing siRNA21 and siRNA24 loci. mRNAs containing siRNA21 loci showed a strong association with sRNAs arising from exons in the vast majority of the species (Figure 4A). These exonic 21 nucleotide siRNAs are most likely secondary siRNAs derived from the processing of the mRNAs. In contrast, in the mRNAs containing siRNA24 loci, sRNAs were primarily generated from introns in nearly all species (Figure 4B). Because 24 nucleotide siRNAs are known to be enriched in TEs, these intronic 24 nucleotide siRNAs could often be generated from intronic TE insertions. Some species showed a lesser association of siRNA24 loci with introns: this may be caused by differences in the annotation of TEs, which can sometimes be erroneously annotated as mRNAs. The siRNAs at both siRNA21 and siRNA24 loci typically originated from both strands of their associated genes (Figure 4C-D). This trend is consistent with processing from dsRNA precursors, as opposed to breakdown products from the mRNAs themselves.

Figure 4. Distribution of sRNAs in the body region of protein-coding mRNAs. **(A)** Heatmap showing the proportion of mRNAs containing siRNA21 loci that have 0% to 100% of their aligned sRNAs mapped to their exons. 0% means all sRNAs map to introns, 100% means all sRNAs map to exons. **(B)** as in A, except for siRNA24 loci. **(C)** Heatmap showing the proportion of mRNAs containing siRNA21 loci with 0% to 100% of their aligned sRNAs mapped to the coding strand of the mRNA: 0% means all sRNAs map to the non-coding strand, 100% blue means all sRNAs map to the coding strand of the mRNA. **(D)** as in C, except for siRNA24 loci. Each row is a given species (See Table 1 for species codes), grouped taxonomically. Eu: eudicots, BA: basal angiosperm, G: gymnosperm, Br: bryophyte.

Identity of genes associated with sRNA loci

To begin to understand the function of genes associated with siRNA loci, we performed GO enrichment analysis on the protein-coding genes that contained siRNA21 or siRNA24 loci, or siRNA24 loci in their 1 kb upstream region (Figure 5). For 38 of the 48 plant genomes, we were able to easily retrieve adequate GO annotations. These were used to perform Fisher's Exact Test in Blast2GO in each species (FDR < 0.05). We plotted the frequency at which the GO terms were found enriched amongst the species to find conserved terms (Figure 5A). Enriched GO terms commonly found in at least ten species were considered to be well conserved, because at this number the frequency distribution inverted after gradually decreasing to zero. Genes containing siRNA21 and siRNA24 loci had respectively two and eight well conserved GO terms. In contrast, genes with siRNA24 loci within their 1 kb upstream region had no enriched GO term shared by ten or more species. The species distribution of the well conserved GO terms (Figure 5B) revealed that the "ADP binding" term was enriched in genes containing siRNA21 loci in rosids, asterids, in *A. trichopoda* and only in one monocot (wheat). Genes associated with the ADP binding function corresponded in all species with NB-LRR type disease resistance genes, which are known to produce secondary siRNAs in many species and only in barley and wheat amongst the monocots (Liu et al. 2014; Zhang et al. 2019). The "protein binding" term was also enriched in genes containing siRNA21 loci, but the genes associated with this term had heterogeneous and variable annotations between species, therefore no single common pathway was identified. Nevertheless, a few gene families in the "protein binding" group were commonly found amongst species, for example F-box genes, PPR-containing genes, kinases and SET domain containing genes. Genes containing siRNA24 loci had well conserved enriched GO terms mostly found in all clades and with different molecular functions (Figure 5B): "terpene synthase" and "heme binding" (mostly cytochromes P450 and other peroxidases) were the most conserved, followed by five others, including the "ADP binding" function. We hypothesize that the genes with these specific functions might be particularly frequent targets of intronic TE insertions silenced by 24 nucleotide siRNAs.

Figure 5. GO enrichment analysis of protein-coding genes associated with siRNA21 or siRNA24 loci. **(A)** Frequency of enriched GO terms in the 38 species analyzed (hbr-b0, tcc-b1.1, fan-b1.0, mdm-b3.0, csi-b2, ccm-b0.32, ccm-b0.1, can-b1.6, nta-b0 and hvu-b1 were excluded because no gene annotation or no GO annotation was available). **(B)** Species distribution of well conserved enriched GO terms, common to ten or more plant species (car-b2.0, egu-b5.1, mac-b2, pab-b1.0c, ppt-b3.0 and mpo-b3.0 were not displayed because they were not enriched in any of these terms). Each row is a given species (See Table 1 for species codes), grouped taxonomically. Eu: eudicots, BA: basal angiosperm.

Disease resistance genes and other genes producing siRNAs in monocots

We further examined the nature of the genes containing exonic sRNA loci in monocots. This was of interest because the regulation of disease resistance genes by sRNAs in monocots has been described only in barley and wheat so far (Liu et al. 2014; Zhang et al. 2019). Genes containing exonic siRNA21 and siRNA22 loci were both studied, because within the monocots, maize produced high quantities of 22 nucleotide siRNAs (Figure 1), whose function is not well-understood. The genes were manually screened to discard those with stacks of sRNA reads mapped at only one or two unique positions, that could be alignment artifacts or miRNA-like sRNAs. Known miRNA matches, lowly expressed sRNA loci (< 1 RPM, reads per million), transposons and inverted repeats were also discarded. In total, 524 genes in the nine monocots were selected as containing robust siRNA21 and siRNA22 loci (Figure 6A, Supplemental Table S3). Maize was the only species where the majority of genic siRNA loci were siRNA22 loci; wheat also had some genic siRNA22 loci. This suggests that in maize and maybe wheat, the 22 nucleotide siRNAs could be a functionally active class of sRNAs in the regulation of genes, in addition to the 21 nucleotide siRNAs.

Evidence of sRNA expression from genes annotated as or having sequence homology with disease resistance genes, was found in seven species (Supplemental Table S3). Confirming

previous reports, 13 disease resistance genes in barley and 48 in wheat contained siRNA21 loci. In oil palm and banana, 33 and 34 disease resistance genes, produced 21 nucleotide siRNAs, respectively. Disease resistance genes evolve rapidly by tandem duplications (Yang and Huang 2014), whose expression may be controlled by siRNAs. In the banana genome we found an example of this where two clusters of disease resistance genes, both on chromosome 3, contained 23 and 15 genes in tandem in a range of ~137 and ~130kb, respectively, that were sources of 21 nucleotide siRNAs. In *B. distachyon* and sorghum, we found seven and six resistance genes producing 21 nucleotide siRNAs, respectively, while in maize only one resistance gene produced 22 nucleotide siRNAs. In rice and foxtail millet there were no disease resistance genes associated with exonic 21 or 22 nucleotide siRNAs. This result suggests that the siRNA-mediated regulation of resistance genes could be conserved in a larger number of monocots than just barley and wheat but be selectively absent in some other monocots like rice.

Genes with different functions than resistance genes also contained siRNA21 and siRNA22 loci in monocots and a few were conserved in multiple species (Supplemental Table S3). Example of these genes include: *TAS3* genes, auxin responsive genes, kinase genes, genes encoding transport inhibitor response 1-like (TIR1-like) proteins, predicted E3 ubiquitin ligase genes, genes encoding or similar to DNA-directed RNA polymerases, two-component response regulators and methyl-CpG-binding domain-containing proteins. Genes participating in sRNA pathways were also found to be sources of siRNAs: *HEN1 SUPPRESSOR1* (*HESO1*, Figure 6B) and *AGO108* in maize, *DOMAINS REARRANGED METHYLTRANSFERASE 2* (*DRM2*) in rice, a predicted *AGO1B* in sorghum and three predicted copies of *AGO2* in wheat. As it is visible by the sRNA alignment coverage in *HESO1* (Figure 6B), sRNAs were expressed from multiple adjacent exons. This pattern of sRNA expression that reflects the mature mRNA structure was observed in many genes and strongly suggests that these exonic 21 and 22 nucleotides sRNAs are secondary siRNAs, originated from the processing of the mRNA by a DCL protein.

Figure 6. Genes containing siRNA21 and siRNA22 loci in nine monocot species. **(A)** Counts of genes containing siRNA21 and siRNA22 loci in monocots. **(B)** Screenshot of genome browser for maize *HES01* (Zm00001d042885). Top row: mRNA structure: blue blocks for UTRs, yellow blocks for CDS and black lines for introns. Middle row: sRNA-seq coverage from the reference set across the gene. Bottom row: ShortStack sRNA loci annotation.

Analysis of sRNA conservation across plant species

Annotated sRNA loci were grouped into putative families based on the sequences of the most abundant single sRNA (the 'major RNA') produced by each locus (Supplemental Table S4). Loci were considered to be members of the same family if the sequences of their major RNAs

had up to two mismatches with each other; these criteria are similar to those commonly used to group *MIRNA* loci into families. Most of the resulting families (1,556,834; 85.3%) had only a single locus (Figure 7A) and relatively few families (38,794; 2.1%) were present in more than a single species (Figure 7B). Even fewer families (1,968; 0.1%) were present in more than one major taxonomic group (Figure 7C). In general, the proportions of MIRNA, nearMIRNA, and siRNA21 loci were higher for more extensively conserved families (Figure 7D-F); at the most extreme levels of conservation, MIRNA loci and siRNA21 loci predominated.

Figure 7. Conservation of sRNA loci in plants. **(A)** Frequency distribution of number of sRNA loci per putative sRNA family. **(B)** Frequency distribution of number of distinct plant species per putative sRNA family. **(C)** Frequency distribution of number of plant 'supergroups' per putative sRNA family. Supergroups defined in this study are: rosids, asterids, other eudicots, monocots, basal angiosperms, gymnosperms, and bryophytes. **(D)** Proportions of types by number of loci per putative sRNA family. **(E)** Proportions of types by number of distinct plant species per

putative sRNA family. **(F)** Proportions of types by number of plant 'supergroups' per putative sRNA family.

Discussion

A public resource on sRNAs for the scientific community

We created an extensive resource for a large number of plant genomes that allows users to freely and easily retrieve, visualize and analyze sRNA loci, including not only miRNA annotations but also siRNA annotations. Our research extended into non-model systems, including many species of horticultural importance. For three economically important plants, spinach, carrot and cacao, we annotated for the first time miRNA and siRNA loci. Recently, a study published the first miRNA annotation in carrot using high throughput sequencing, but the siRNAs were not examined (Bhan et al. 2019). In many published works, sRNA-seq is used to annotate and profile miRNAs but not individual siRNA loci. We used the vast amount of available sRNA-seq datasets to exploit all this unrevealed information and annotate the entire population of sRNAs in 48 plant genomes.

Our database and analyses are limited by the quality and quantity of the available genomic annotations and sRNA-seq data. For example, sRNAs expressed in specific tissues/cell types or growth conditions that were not represented in our sRNA-seq dataset are by consequence absent from our reference annotations. This might be the case for reproductive phasiRNAs (Zhai et al. 2015; Fei et al. 2016; Xia et al. 2019): for most of the analyzed species we did not have any or enough sRNA-seq libraries from male reproductive tissues to allow the specific annotation of this sRNA population. For this reason, we did not investigate the reproductive sRNAs in our work. Our database has the potential to be expanded in the future to include new plant genomes, new annotations and new sRNA-seq data that are of interest for the plant biology community.

Overall, we created a resource that will be useful for future sRNA studies. Thanks to the standard annotation and classification methods followed for all genomes, our sRNA annotations and alignments can be directly visualized or downloaded from our web-server and compared between species. Our web-server is a practical way to quickly interrogate existing plant sRNA data in a usable format and will enable scientists to rapidly search for evidence of sRNA expression in specific regions in a species or investigate the conservation of single sRNA sequences across species.

Multiple protein-coding gene families are sources of 21 nucleotide siRNAs in dicots and monocots

The best characterized case of protein-coding genes generating secondary siRNAs are the disease resistance genes, whose expression is kept under control by secondary siRNA production to avoid fitness loss (Yang and Huang 2014). We confirmed expression of 21 nucleotide siRNAs from exons of resistance genes in the rosoid and asterid clades and expanded the number of monocot species that also showed this evidence, suggesting that this pathway might be more broadly conserved than what is known. In none of the three studied

caryophyllales species the protein-coding genes containing siRNA21 loci were enriched in the GO:ADP binding term, characteristic of resistance genes. This could result from incomplete gene/GO annotations in spinach, sugar beet and quinoa, missing real resistance genes. Alternatively, these secondary siRNAs might be reduced in caryophyllales because the number of disease resistance genes in this clade is lower compared to the typical expansion of this gene family in rosids and asterids or because in caryophyllales, specific subfamilies of resistance genes have expanded that might be differentially regulated (Dohm et al. 2014; Xu et al. 2017; Funk et al. 2018).

In the literature, the number of known protein-coding genes producing secondary siRNAs in monocots is smaller than in dicots. Accordingly, from our analyses, the enrichment of siRNA21 loci in protein-coding genes was less evident in monocots compared to dicots and also the tendency of 21 nucleotide siRNAs to map to exons was smaller in monocots. For these reasons, we decided to manually screen the monocot species for evidence of 21 nucleotide siRNA production from protein-coding genes. We described a number of gene families, more or less conserved in the nine monocots, that produced 21 nucleotide siRNAs, and also 22 nucleotide siRNAs in maize and wheat. In many cases, the siRNAs were expressed specifically from multiple adjacent exons, supporting the hypothesis that they are secondary siRNAs processed from mature mRNAs. Some of the genes found were previously described as sources of secondary siRNAs in other species, for example kinase genes (Zheng et al. 2015; Reyes-Chin-Wo et al. 2017), TIR1-like genes (Si-Ammour et al. 2011; Seo et al. 2018; Xia et al. 2015a) and *AGO2* (Arikrit et al. 2014). In addition to *AGO2*, there are more genes participating in siRNA biogenesis and function that are themselves known targets of siRNA regulation: *DCL1* (Xie et al. 2003; Hu et al. 2015b; Xia et al. 2014), *DCL2* (Zhai et al. 2011; Arikrit et al. 2014), *AGO1* (Vaucheret et al. 2006) and *SUPPRESSOR OF GENE SILENCING 3* (Arikrit et al. 2014). We found evidence of siRNA expression from four additional genes involved in siRNA pathways: in maize, from *AGO108* (also named *AGO5d*), highly expressed in ears but not well functionally characterized (Zhai et al. 2014), and *HESO1*, a nucleotidyl transferase that uridylylates unmethylated sRNAs to trigger their degradation (Zhao et al. 2012); in sorghum, from a predicted *AGO1B* and in rice from *DRM2*. *DRM2* is a known target of miR820 in rice (Nosaka et al. 2012), which could be the trigger miRNA for the production of the observed 21 nucleotides siRNAs. We reported many more genes in monocots that spawned 21 or 22 nucleotide long siRNAs, belonging to different families. These genes represent an interesting set to research in the future to better characterize the nature of genic siRNAs. The next obvious step will be searching for possible miRNA triggers and examining the phasing pattern of siRNA expression in each specific gene, to confirm that these siRNAs are secondary siRNAs.

Different hypotheses on 22 nucleotide siRNA functions

We found that asterids consistently had considerable proportions of siRNA22 loci, while in the other clades, only certain species (soybean, cassava and maize) had this same trend. There are several hypotheses that could explain the presence of 22 nucleotide siRNAs in a genome: they could originate from *MIRNA* or *MIRNA*-like loci that were missed by our annotation method, from endogenous direct or inverted repeats (Kasschau et al. 2007), or from protein-coding genes, as we observed in maize. Alternatively, these siRNA22 loci could express

siRNAs involved in the non-canonical RdDM pathway to silence active TEs (Matzke and Mosher 2014), as it was proposed for maize (Nobuta et al. 2008). Active retrotransposons have been described in asterids, for example the Tto1 element or the Tnt1 element, which has many copies that are still transcriptionally active in tobacco (Casacuberta et al. 1997) and lettuce (Mazier et al. 2007). In this hypothesis, what still remains unclear is why we observed expression of 22 nucleotide siRNA most often in the asterids and not in the grasses, where retrotransposon transcription is very prevalent (Vicient et al. 2001). If the 22 nucleotide siRNAs come from active retrotransposons, then the ability to detect their expression could depend on the specific samples analyzed, because retrotransposons are only active during certain stages of plant development or stress conditions (Flavell et al. 1992). Lastly, 22 nucleotide siRNAs could target Endogenous Viral Elements, virus segments that are integrated in the host genome, that form inverted repeats (Pooggin 2018). To understand the role of the siRNA22 loci, the next step in future research will be the genome-wide profiling of the genomic regions where these loci map, discriminating between genes, intergenic regions and different classes of TEs.

Roles of 24 nucleotide siRNAs in regulating protein-coding gene expression

We assumed that the distribution of the total sRNA loci across the chromosome length reflected the distribution of the siRNA24 loci, because these accounted for the vast majority of loci in angiosperms. *A. thaliana* and Chinese cabbage are two of the few species where siRNA24 loci and gene densities were negatively correlated. In both species, siRNA regulation of TEs near genes was previously linked to lower expression of the genes (Hollister et al. 2011; Woodhouse et al. 2014). It would be informative to test if the same link occurs in the other species with inverse correlation between siRNA24 locus and gene densities, like sweet orange. Differences in siRNA24 locus distribution and influence on gene expression might be directly explained by differences in TE composition between genomes. Accordingly, it was previously suggested that the transcription of gene networks can be balanced by the genome distribution of TEs (Freeling et al. 2015), which can be highly variable among species (Vicient and Casacuberta 2017). In many cases, a few TE families have increased their copy number in one lineage (Baidouri and Panaud 2013). For example, a single type of LTR retrotransposon is responsible for most of the hot pepper genome expansion (Park et al. 2012).

The angiosperms analyzed were strongly enriched in siRNA24 loci in the 5'-proximal regions upstream of protein-coding genes. In *A. thaliana*, this distribution was much less strong but the enrichment of siRNA24 loci in the 5' upstream region compared to the gene body region was still evident. The function of siRNA24 loci at these sites has been widely studied in maize: near genes, 24 nucleotide siRNAs engage RdDM, blocking the spread of open, active chromatin into adjacent transposons (Li et al. 2015). In addition to silencing TEs, the RdDM activity near genes in *A. thaliana* can also affect the expression levels of the genes (Zheng et al. 2013; Zhong et al. 2012), likely by changing the chromatin landscape at gene promoters and influencing the ability of transcription factors to bind to the promoters and stimulate transcription. In maize on the contrary, no obvious direct effects on gene expression were detected as a consequence of the loss of gene proximal 24 nucleotide siRNAs (Lunardon et al. 2016). Finally, in *A. thaliana*, it has been speculated that the RdDM activity near genes can influence their expression by inhibiting interactions between the promoters and their potential

distant regulatory elements (Rowley et al. 2017). Similarly, most angiosperms were also enriched in siRNA24 loci at the 3'-proximal regions downstream of genes, where the RdDM activity seems to reduce the readthrough transcription by Pol II into neighboring genes or TEs (Erhard et al. 2015).

When siRNA24 loci were found inside protein-coding genes, they were mostly in introns. A few gene families were most commonly targeted by 24 nucleotide siRNAs in both dicots and monocots. Two possible reasons might explain why these specific genes were a common target of 24 nucleotide siRNAs. On one side, families like disease resistance genes evolve rapidly, creating high numbers of partial genes and pseudogenes (Luo et al. 2012) that might be suppressed by the activity of 24 nucleotide siRNAs (Kasschau et al. 2007). This could also be the case of polygalacturonases that are encoded by a large gene family. An accurate study of the protein-coding gene annotations, precisely separating genes from pseudogenes, would be necessary to verify this hypothesis. On the other side, gene families like disease resistance genes control adaptive responses to the environment, making them frequent targets of TE transposition events (Quadrana et al. 2016). Although the majority of TE insertions in genes are deleterious, they can be advantageous and therefore be retained as source of variability, which is essential in environmental response genes to adapt to the ever-changing environment. As a consequence, new TE insertions are overrepresented in genes that respond to environmental stresses (Grover et al. 2003; Miyao et al. 2003). Also in cytochrome P450s, a family known to participate in stress responses, frequent TE insertions were described as a strategy for variability (Chen and Li 2007) and this could explain why these genes were frequent targets of 24 nucleotide siRNAs. Likewise, serine-type carboxypeptidases, which participate in protein degradation, and xenobiotic transmembrane transporters, which work in xenobiotic detoxification pathways together with cytochromes P450, both play pivotal roles in plant defense responses and therefore could be frequent targets of TE insertions controlled by 24 nucleotide siRNAs. To verify if the intronic 24 nucleotide siRNAs influence the regulation of the genes that they target, it will be informative in the future to examine mutants lacking the production of 24 nucleotide siRNAs and observe if these gene families tend to be altered in their expression.

Conservation of siRNAs

The sequence comparison of the most abundant sRNA expressed from each locus revealed a very low level of conservation of siRNAs across species, not just between distant species but also between close relatives. Studying the conservation of siRNAs is complicated by the fact that the siRNA population can vary substantially between different organs of the same plant species (Ha et al. 2009). Nonetheless, our result is in line with previous observations (Ma et al. 2010). If we consider plants that all have a strong peak of 24 nucleotide siRNAs and have a functional RdDM pathway, the genomic TE composition and organization can significantly differ between different species and even between different varieties of the same species (Brunner et al. 2005; Quadrana et al. 2016). This might explain why the individual siRNA sequences that target the TEs are also poorly conserved. Much of our knowledge regarding sRNAs comes from model plants like *A. thaliana*, which has a low amount of TEs that are not active in wild-type plants. Crop genomes, instead, have high TE loads and some TEs are active in wild-type genetic

backgrounds in maize and rice (Jiang et al. 2003; Nakazaki et al. 2003; Lisch 2012). Due to these differences it is important to study sRNAs in non-model systems, because lineage- or species-specific sRNAs might be associated to traits that other plants lack or have not evolved (Chen et al. 2018).

Methods

Plant material and sRNA sequencing

Leaves of *Theobroma cacao* (line Scavina 6) were kindly provided by Dr. M. Guiltinan of The Pennsylvania State University, from plants grown in greenhouse conditions. The tips of leaves at the immature green leaf stage were collected. *Daucus carota* (cultivar 'Burpee') was grown in a growth room at 22°C, 16h light 8h dark regime and leaves and roots from 5- and 6-week-old plants, respectively, were sampled. *Spinacia oleracea* Sp75 inbred line seeds were kindly provided by Dr. Z. Fei of the Boyce Thompson Institute, Cornell University, and grown in a growth room at 22°C, 16h light 8h dark regime. Leaves from 3- and 5-week-old plants were collected. *Zea mays* B73 inbred line seeds were germinated on ProMix B, then transferred to soil in pots and grown in greenhouse conditions with occasional Osmocote fertilization. The fifth and the sixth leaves from V5 plants, mature pollen and 21-27 DAP (days after pollination) embryo tissue were collected from a pool of plants. All samples were flash frozen in liquid nitrogen, stored at -80°C and then ground with liquid nitrogen cooled mortar and pestle. For carrot, spinach and maize, the RNA was extracted with Tri-reagent (Sigma) as per manufacturer instructions, adding a second sodium-acetate-ethanol precipitation and ethanol wash step. For cacao, the RNA was extracted with PureLink Plant RNA Reagent (Life technologies) following manufacturer's suggestions. Sequencing libraries were prepared using the NEB Next sRNA-seq library preparation kit for Illumina (NEB, E7300S) following manufacturer's suggestions. Reactions were purified and size selected for sRNAs 15-40nt in length by PAGE. Extracted bands were quantified by qPCR and quality-controlled by high-sensitivity DNA chip (Agilent). Sequencing was performed on a HiSeq2500 (Illumina) in rapid run mode (50 nucleotides, single-end, single barcode) by the Penn State genomics core.

sRNA-seq data processing

sRNA-seq raw fastq files were downloaded from SRA and GEO databases (Supplemental Table S1). The libraries were processed to remove the 3' adapter with cutadapt (Martin 2011) (cutadapt -a 3'_adapter_sequence --discard-untrimmed -m 15 -o output_file.fastq input_file.fastq). Reads containing the 5' adapter were removed with cutadapt (cutadapt -g 5'_adapter_sequence --discard-trimmed -m 15 -o output_file.fastq input_file.fastq). Low quality reads were discarded with FASTX-Toolkit (Gordon and Hannon 2010) (fastq_quality_filter -q 20 -p 85 -Q 33 -v -i input_file.fastq -o output_file.fastq). Finally, reads quality was checked with FastQC (Andrews 2010): if additional sequencing adapters were overrepresented amongst reads, they were eliminated from the fastq files with a custom Perl script.

Pipeline to create reference sRNA loci annotations

For each species, the reference annotation of sRNA loci was created with the following steps. Each individual library was aligned to the genome (see <https://plantsmallrnagenes.science.psu.edu> for list of genome assemblies used) using ShortStack v3.8.1 (Axtell 2013b; Johnson et al. 2016) with default parameters. Libraries with less than 2 million mapped reads were discarded. Clusters of sRNAs were *de novo* identified in each library independently with ShortStack (ShortStack --bamfile *alignment_file.bam* --mincov 2rpm --genomefile *genome_file.fa*). The sRNA clusters files from all libraries of the same species were intersected with the bedtools function 'multiIntersectBed' (Quinlan and Hall 2010) with default parameters. Only genomic intervals with annotated sRNA clusters common to at least three libraries were kept and merged with bedtools, with 25 nucleotides as maximum distance allowed between the intervals to be merged into sRNA loci (mergeBed -d 25 -i *input_intervals_file.bed* > *output_merged_intervals_file.bed*). sRNA loci with length < 15 nucleotides were removed with a custom Perl script. Finally, sRNA loci whose expression was < 0.5 RPM in all libraries were also removed. The sRNA loci that were selected after applying these filters represented the reference annotation for each species.

Analysis of sRNA loci occupancy relative to protein-coding genes

Locations of protein-coding genes were determined from public GFF3 files from each genome. Intergenic regions were calculated using bedtools 'complement', computationally cut in half, and associated with their nearest protein-coding genes using bedtools 'closest'. The regions were marked as upstream or downstream based on the orientation of their nearest flanking gene. Per-nucleotide overlap between upstream, downstream, and gene-body regions vs. small RNA loci were calculated using bedtools 'overlap'. The lengths of gene-bodies were scaled to 1,000 arbitrary units (each such unit is 0.1% of the gene length). Coverage was summarized in 25 nucleotides / unit bins, and normalized to a scale of 0 to 1, where 1 represented the maximum fraction occupancy observed in that genome.

Analysis of sRNA distribution in exons and introns of protein-coding mRNAs

Only protein-coding mRNAs having at least one intron and overlapping with siRNA21 and siRNA24 loci were studied. Each mRNA was either classified as containing siRNA21 or siRNA24 loci: in case of overlap with both siRNA21 and siRNA24 loci, the longest sRNA locus was considered. The number of sRNAs mapped to exons and to the same strand of protein-coding mRNAs containing one or more introns were calculated with the bedtools function 'coverageBed -counts' (parameters added for exons: '-F 1'; for the same strand: '-F 1 -s'). The number of sRNAs mapped to introns and to the opposite strand of the mRNAs were also calculated for the final ratios (parameters added for the opposite strand: '-F 1 -S'). The percentage of sRNAs mapped to exons was calculated based on the ratio 'number of reads mapped to exons / (number of reads mapped to exons + number of reads mapped to introns)'. The percentage of sRNAs mapped to the same strand of the mRNA was calculated based on the ratio 'number of reads mapped to the same strand / (number of reads mapped to the same

strand + number of reads mapped to the opposite strand)'. Here and in the other analyses of siRNAs, sRNA loci classified as MIRNA and nearMIRNA or whose most abundant sequence had a perfect match with a high-confidence plant miRNA hairpin annotated in miRBase v22 (Kozomara and Griffiths-Jones 2014) were not included.

GO enrichment analysis

Protein-coding genes were classified as containing siRNA21 or siRNA24 loci and as flanked in their 1kb upstream region by siRNA21 or siRNA24 loci: when the same gene/upstream region overlapped with both siRNA21 and siRNA24 loci, the longest sRNA locus determined the classification. The GO enrichment analysis was performed with Blast2GO (Götz et al. 2008), using the Fisher's Exact Test with default parameters (FDR < 0.05). Only the species for which we were able to retrieve a GO annotation were analyzed, this excluded: hbr-b0, tcc-b1.1, fan-b1.0, mdm-b3.0, csi-b2, ccm-b0.32, ccm-b0.1, can-b1.6, nta-b0 and hvu-b1.

Analysis of genes containing siRNA21 and siRNA22 loci in monocots

To find all genes containing siRNA21 and siRNA22 loci in exons we used bedtools (intersectBed -wao -F 0.75 -a *exons_file.gff3* -b *sRNA_loci_file.gff3* > *output_intersection_file.txt*). When the same gene contained both siRNA21 and siRNA22 loci, if it contained a greater number of siRNA21 loci than siRNA22 loci it was classified as containing siRNA21 loci. In case there were the same number of siRNA21 and siRNA22 loci, the gene was classified based on the longest locus. The description of the genes (Supplemental Table S3) was copied from the gene annotation files retrieved from the same online resources used for the genome sequences (see <https://plantsmallrnagenes.science.psu.edu> for sources of genomes and gene annotations files). For species without available gene annotations, the function of the genes was predicted using BLAST (Camacho et al. 2009) on the gene sequence and considering the best result.

Data access

All sRNA-seq libraries used, published and newly generated, are available in GEO and SRA; see Supplemental Table S1 for accession numbers. All data and analyses are hosted at <https://plantsmallrnagenes.science.psu.edu>.

Acknowledgements

We thank the Penn State Genomics Core Facility for small RNA sequencing services, and the Eberly College of Science IT office for providing server hosting services for this project. We thank Matthew Jones-Rhoades for supervision and mentoring of undergraduate researchers and for insightful comments on the manuscript. This work was supported by an award from the US National Science Foundation (Award 1339207) to MJA.

Author Contributions

AL and MJA generated most primary annotations and most analyses, with some primary annotations also contributed by SP. The project website was developed by MJA. NRJ, EH, TP, and CC contributed small RNA sequencing results. The manuscript was written by AL and MJA with input from NRJ and CC.

References Cited

- Allen E, Xie Z, Gustafson AM, Carrington JC. 2005. microRNA-directed phasing during trans-acting siRNA biogenesis in plants. *Cell* **121**: 207–21.
- Andrews S. 2010. Babraham Bioinformatics - FastQC A Quality Control tool for High Throughput Sequence Data. *FastQC a Qual Control tool high throughput Seq data*. <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>.
- Arikiti S, Xia R, Kakrana A, Huang K, Zhai J, Yan Z, Valdés-López O, Prince S, Musket TA, Nguyen HT, et al. 2014. An Atlas of Soybean Small RNAs Identifies Phased siRNAs from Hundreds of Coding Genes. *Plant Cell* **26**: 4584–601.
- Axtell MJ. 2013a. Classification and Comparison of Small RNAs from Plants. *Annu Rev Plant Biol* **64**: 137–59.
- Axtell MJ. 2013b. ShortStack: Comprehensive annotation and quantification of small RNA genes. *RNA* **19**: 740–51.
- Axtell MJ, Bartel DP. 2005. Antiquity of MicroRNAs and Their Targets in Land Plants. *Plant Cell* **17**: 1658–73.
- Axtell MJ, Bowman JL. 2008. Evolution of plant microRNAs and their targets. *Trends Plant Sci* **13**: 343–9.
- Baidouri M El, Panaud O. 2013. Comparative genomic paleontology across plant kingdom reveals the dynamics of TE-driven genome evolution. *Genome Biol Evol* **5**: 954–65.
- Banks JA, Nishiyama T, Hasebe M, Bowman JL, Gribskov M, DePamphilis C, Albert VA, Aono N, Aoyama T, Ambrose BA, et al. 2011. The selaginella genome identifies genetic changes associated with the evolution of vascular plants. *Science (80-)* **332**: 960–3.
- Bhan B, Koul A, Sharma D, Manzoor MM, Kaul S, Gupta S, Dhar MK. 2019. Identification and expression profiling of miRNAs in two color variants of carrot (*Daucus carota* L.) using deep sequencing. *PLoS One* **14**: e0212746.
- Brunner S, Fengler K, Morgante M, Tingey S, Rafalski A. 2005. Evolution of DNA Sequence Nonhomologies among Maize Inbreds. *Plant Cell* **17**: 343–60.
- Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: Architecture and applications. *BMC Bioinformatics* **10**: 421.
- Casacuberta JM, Vernhettes S, Audeon C, Grandbastien M-A. 1997. Quasispecies in retrotransposons: a role for sequence variability in Tnt1 evolution. *Genetica* **100**: 109–17.
- Chávez Montes RA, De Fátima Rosas-Cárdenas F, De Paoli E, Accerbi M, Rymarquis LA, Mahalingam G, Marsch-Martínez N, Meyers BC, Green PJ, De Folter S. 2014. Sample sequencing of vascular plants demonstrates widespread conservation and divergence of microRNAs. *Nat Commun* **5**: 3722.
- Chen C, Zeng Z, Liu Z, Xia R. 2018. Small RNAs, emerging regulators critical for the development of horticultural traits. *Hortic Res* **5**: 63.
- Chen S, Li X. 2007. Transposable elements are enriched within or in close proximity to

803 xenobiotic-metabolizing cytochrome P450 genes. *BMC Evol Biol* **7**: 46.

804 Coruh C, Cho SH, Shahid S, Liu Q, Wierzbicki A, Axtell MJ. 2015. Comprehensive Annotation of
805 Physcomitrella patens Small RNA Loci Reveals That the Heterochromatic Short Interfering
806 RNA Pathway Is Largely Conserved in Land Plants. *Plant Cell* **27**: 2148–62.

807 Coruh C, Shahid S, Axtell MJ. 2014. Seeing the forest for the trees: Annotating small RNA
808 producing genes in plants. *Curr Opin Plant Biol* **18**: 87–95.

809 Cuperus JT, Fahlgren N, Carrington JC. 2011. Evolution and Functional Diversification of MIRNA
810 Genes. *Plant Cell* **23**: 431–42.

811 Dohm JC, Minoche AE, Holtgräwe D, Capella-Gutiérrez S, Zakrzewski F, Tafer H, Rupp O,
812 Sörensen TR, Stracke R, Reinhardt R, et al. 2014. The genome of the recently domesticated
813 crop plant sugar beet (*Beta vulgaris*). *Nature* **505**: 546–9.

814 Dolgosheina E V., Morin RD, Aksay G, Sahinalp SC, Magrini V, Mardis ER, Mattsson J, Unrau PJ.
815 2008. Conifers have a unique small RNA silencing signature. *RNA* **14**: 1508–15.

816 Erhard KF, Talbot JERB, Deans NC, McClish AE, Hollick JB. 2015. Nascent transcription affected
817 by RNA polymerase IV in *Zea mays*. *Genetics* **199**: 1107–25.

818 Fei Q, Yang L, Liang W, Zhang D, Meyers BC. 2016. Dynamic changes of small RNAs in rice
819 spikelet development reveal specialized reproductive phasiRNA pathways. *J Exp Bot* **67**:
820 6037–6049.

821 Flavell AJ, Dunbar E, Anderson R, Pearce SR, Hartley R, Kumar A. 1992. Ty1-copia group
822 retrotransposons are ubiquitous and heterogeneous in higher plants. *Nucleic Acids Res* **20**:
823 3639–44.

824 Formey D, Iñiguez LP, Peláez P, Li YF, Sunkar R, Sánchez F, Reyes JL, Hernández G. 2015.
825 Genome-wide identification of the *Phaseolus vulgaris* sRNAome using small RNA and
826 degradome sequencing. *BMC Genomics* **16**: 423.

827 Freeling M, Xu J, Woodhouse M, Lisch D. 2015. A solution to the c-value paradox and the
828 function of junk DNA: The genome balance hypothesis. *Mol Plant* **8**: 899–910.

829 Funk A, Galewski P, McGrath JM. 2018. Nucleotide-binding resistance gene signatures in sugar
830 beet, insights from a new reference genome. *Plant J* **95**: 659–71.

831 Gent JI, Ellis NA, Guo L, Harkess AE, Yao Y, Zhang X, Dawe RK. 2013. CHH islands: De novo DNA
832 methylation in near-gene chromatin regulation in maize. *Genome Res* **23**: 628–37.

833 Gent JI, Madzima TF, Bader R, Kent MR, Zhang X, Stam M, McGinnis KM, Dawe RK. 2014.
834 Accessible DNA and Relative Depletion of H3K9me2 at Maize Loci Undergoing RNA-
835 Directed DNA Methylation. *Plant Cell* **26**: 4903–17.

836 Gordon A, Hannon GJ. 2010. Fastx-toolkit. FASTQ/A short-reads pre-processing tools. *Unpubl*
837 http://hannonlab.cshl.edu/fastx_toolkit.

838 Götz S, García-Gómez JM, Terol J, Williams TD, Nagaraj SH, Nueda MJ, Robles M, Talón M,
839 Dopazo J, Conesa A. 2008. High-throughput functional annotation and data mining with
840 the Blast2GO suite. *Nucleic Acids Res* **36**: 3420–35.

841 Grover D, Majumder PP, Rao CB, Brahmachari SK, Mukerji M. 2003. Nonrandom distribution of
842 Alu elements in genes of various functional categories: Insight from analysis of human
843 chromosomes 21 and 22. *Mol Biol Evol* **20**: 1420–4.

844 Guo Q, Qu X, Jin W. 2015. PhaseTank: Genome-wide computational identification of phasiRNAs
845 and their regulatory cascades. *Bioinformatics* **31**: 284–6.

846 Ha M, Lu J, Tian L, Ramachandran V, Kasschau KD, Chapman EJ, Carrington JC, Chen X, Wang X-J,

Chen ZJ. 2009. Small RNAs serve as a genetic buffer against genomic shock in Arabidopsis interspecific hybrids and allopolyploids. *Proc Natl Acad Sci* **106**: 17835–40.

Hackenberg M, Rueda A, Gustafson P, Langridge P, Shi BJ. 2016. Generation of different sizes and classes of small RNAs in barley is locus, chromosome and/or cultivar-dependent. *BMC Genomics* **17**: 735.

He G, Chen B, Wang X, Li X, Li J, He H, Yang M, Lu L, Qi Y, Wang X, et al. 2013. Conservation and divergence of transcriptomic and epigenomic variation in maize hybrids. *Genome Biol* **14**: R57.

Hollister JD, Smith LM, Guo Y-L, Ott F, Weigel D, Gaut BS. 2011. Transposable elements and small RNAs contribute to gene expression divergence between Arabidopsis thaliana and Arabidopsis lyrata. *Proc Natl Acad Sci* **108**: 2322–7.

Howell MD, Fahlgren N, Chapman EJ, Cumbie JS, Sullivan CM, Givan SA, Kasschau KD, Carrington JC. 2007. Genome-Wide Analysis of the RNA-DEPENDENT RNA POLYMERASE6/DICER-LIKE4 Pathway in Arabidopsis Reveals Dependency on miRNA- and tasiRNA-Directed Targeting. *Plant Cell* **19**: 926–42.

Hu H, Rashotte AM, Singh NK, Weaver DB, Goertzen LR, Singh SR, Locy RD. 2015a. The complexity of posttranscriptional small RNA regulatory networks revealed by In Silico analysis of Gossypium arboreum L. leaf, flower and boll small regulatory RNAs. *PLoS One* **10**: e0127468.

Hu H, Yu D, Liu H. 2015b. Bioinformatics analysis of small rnas in pima (Gossypium barbadense L.). *PLoS One* **10**: e0116826.

Jiang N, Bao Z, Zhang X, Hirochika H, Eddy SR, McCouch SR, Wessler SR. 2003. An active DNA transposon family in rice. *Nature* **421**: 163–7.

Johnson C, Bowman L, Adai AT, Vance V, Sundaresan V. 2007. CSRDB: A small RNA integrated database and browser resource for cereals. *Nucleic Acids Res* **35**: D829–33.

Johnson NR, Yeoh JM, Coruh C, Axtell MJ. 2016. Improved placement of multi-mapping small RNAs. *G3 Genes, Genomes, Genet* **6**: 2103–11.

Kasschau KD, Fahlgren N, Chapman EJ, Sullivan CM, Cumbie JS, Givan SA, Carrington JC. 2007. Genome-wide profiling and analysis of Arabidopsis siRNAs. *PLoS Biol* **5**: e57.

Kim S, Park M, Yeom SI, Kim YM, Lee JM, Lee HA, Seo E, Choi J, Cheong K, Kim KT, et al. 2014. Genome sequence of the hot pepper provides insights into the evolution of pungency in Capsicum species. *Nat Genet* **46**: 270–8.

Klevebring D, Street NR, Fahlgren N, Kasschau KD, Carrington JC, Lundeberg J, Jansson S. 2009. Genome-wide profiling of Populus small RNAs. *BMC Genomics* **10**: 620.

Komiya R. 2017. Biogenesis of diverse plant phasiRNAs involves an miRNA-trigger and Dicer-processing. *J Plant Res* **130**: 17–23.

Kozomara A, Birgaoanu M, Griffiths-Jones S. 2019. MiRBase: From microRNA sequences to function. *Nucleic Acids Res* **47**: D155–D162.

Kozomara A, Griffiths-Jones S. 2014. MiRBase: Annotating high confidence microRNAs using deep sequencing data. *Nucleic Acids Res* **42**: D68–73.

Lai YS, Zhang X, Zhang W, Shen D, Wang H, Xia Y, Qiu Y, Song J, Wang C, Li X. 2017. The association of changes in DNA methylation with temperature-dependent sex determination in cucumber. *J Exp Bot* **68**: 2899–2912.

Lelandais-Brière C, Naya L, Sallet E, Calenge F, Frugier F, Hartmann C, Gouzy J, Crespi M. 2009.

Genome-Wide Medicago truncatula Small RNA Analysis Revealed Novel MicroRNAs and Isoforms Differentially Regulated in Roots and Nodules. *Plant Cell* **21**: 2780–96.

Li Q, Gent JI, Zynda G, Song J, Makarevitch I, Hirsch CD, Hirsch CN, Dawe RK, Madzima TF, McGinnis KM, et al. 2015. RNA-directed DNA methylation enforces boundaries between heterochromatin and euchromatin in the maize genome. *Proc Natl Acad Sci* **112**: 14728–33.

Lisch D. 2012. Regulation of transposable elements in maize. *Curr Opin Plant Biol* **15**: 511–6.

Liu J, Cheng X, Liu D, Xu W, Wise R, Shen QH. 2014. The miR9863 Family Regulates Distinct Mla Alleles in Barley to Attenuate NLR Receptor-Triggered Disease Resistance and Cell-Death Signaling. *PLoS Genet* **10**: e1004755.

Liu Q, Ding C, Chu Y, Zhang W, Guo G, Chen J, Su X. 2017. Pln24NT: A web resource for plant 24-nt siRNA producing loci. *Bioinformatics* **33**: 2065–2067.

Lunardon A, Forestan C, Farinati S, Axtell MJ, Varotto S. 2016. Genome-Wide Characterization of Maize Small RNA Loci and Their Regulation in the required to maintain repression6-1 (*rmr6-1*) Mutant and Long-Term Abiotic Stresses. *Plant Physiol* **170**: 1535–48.

Luo S, Zhang Y, Hu Q, Chen J, Li K, Lu C, Liu H, Wang W, Kuang H. 2012. Dynamic Nucleotide-Binding Site and Leucine-Rich Repeat-Encoding Genes in the Grass Family. *Plant Physiol* **159**: 197–210.

Ma Z, Coruh C, Axtell MJ. 2010. Arabidopsis lyrata Small RNAs: Transient MIRNA and Small Interfering RNA Loci within the Arabidopsis Genus. *Plant Cell* **22**: 1090–103.

Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* **17**: 10–12.

Martinez G, Choudury SG, Slotkin RK. 2017. TRNA-derived small RNAs target transposable element transcripts. *Nucleic Acids Res* **45**: 5142–5152.

Matzke MA, Mosher RA. 2014. RNA-directed DNA methylation: An epigenetic pathway of increasing complexity. *Nat Rev Genet* **15**: 394–408.

Mazier M, Botton E, Flamain F, Bouchet J-P, Courtial B, Chupeau M-C, Chupeau Y, Maisonneuve B, Lucas H. 2007. Successful Gene Tagging in Lettuce Using the Tnt1 Retrotransposon from Tobacco. *Plant Physiol* **144**: 18–31.

Miyao A, Tanaka K, Murata K, Sawaki H, Takeda S, Abe K, Shinozuka Y, Onosato K, Hirochika H. 2003. Target Site Specificity of the Tos17 Retrotransposon Shows a Preference for Insertion within Genes and against Insertion in Retrotransposon-Rich Regions of the Genome. *Plant Cell* **15**: 1771–80.

Nakano M, Nobuta K, Vemaraju K, Tej SS, Skogen JW, Meyers BC. 2006. Plant MPSS databases: signature-based transcriptional resources for analyses of mRNA and small RNA. *Nucleic Acids Res* **34**: D731-5.

Nakazaki T, Okumoto Y, Horibata A, Yamahira S, Teraishi M, Nishida H, Inoue H, Tanisaka T. 2003. Mobilization of a transposon in the rice genome. *Nature* **421**: 170–2.

Nobuta K, Lu C, Shrivastava R, Pillay M, De Paoli E, Accerbi M, Arteaga-Vazquez M, Sidorenko L, Jeong D-H, Yen Y, et al. 2008. Distinct size distribution of endogenous siRNAs in maize: Evidence from deep sequencing in the mop1-1 mutant. *Proc Natl Acad Sci* **105**: 14958–63.

Nosaka M, Itoh JI, Nagato Y, Ono A, Ishiwata A, Sato Y. 2012. Role of Transposon-Derived Small RNAs in the Interplay between Genomes and Parasitic DNA in Rice. *PLoS Genet* **8**: e1002953.

Nuthikattu S, McCue AD, Panda K, Fultz D, DeFraia C, Thomas EN, Slotkin RK. 2013. The Initiation of Epigenetic Silencing of Active Transposable Elements Is Triggered by RDR6 and 21-22 Nucleotide Small Interfering RNAs. *Plant Physiol* **162**: 116–31.

Park M, Park J, Kim S, Kwon JK, Park HM, Bae IH, Yang TJ, Lee YH, Kang BC, Choi D. 2012. Evolution of the large genome in *Capsicum annuum* occurred through accumulation of single-type long terminal repeat retrotransposons and their derivatives. *Plant J* **69**: 1018–29.

Polydore S, Lunardon A, Axtell MJ. 2018. Several phased siRNA annotation methods can frequently misidentify 24 nucleotide siRNA-dominated PHAS loci. *Plant Direct* **2**: e00101.

Pooggin MM. 2018. Small RNA-omics for plant virus identification, virome reconstruction, and antiviral defense characterization. *Front Microbiol* **9**: 2779.

Quadrana L, Silveira AB, Mayhew GF, LeBlanc C, Martienssen RA, Jeddeloh JA, Colot V. 2016. The *Arabidopsis thaliana* mobilome and its impact at the species level. *Elife* **5**: e15716.

Quinlan AR, Hall IM. 2010. BEDTools: A flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**: 841–2.

Reyes-Chin-Wo S, Wang Z, Yang X, Kozik A, Arikat S, Song C, Xia L, Froenicke L, Lavelle DO, Truco MJ, et al. 2017. Genome assembly with in vitro proximity ligation data and whole-genome triplication in lettuce. *Nat Commun* **8**: 14953.

Rogers K, Chen X. 2013. Biogenesis, turnover, and mode of action of plant microRNAs. *Plant Cell* **25**: 2383–2399.

Rowley MJ, Rothi MH, Böhmendorfer G, Kuciński J, Wierzbicki AT. 2017. Long-range control of gene expression via RNA-directed DNA methylation. *PLoS Genet* **13**: e1006749.

Schmitz RJ, He Y, Valdés-López O, Khan SM, Joshi T, Urich MA, Nery JR, Diers B, Xu D, Stacey G, et al. 2013. Epigenome-wide inheritance of cytosine methylation variants in a recombinant inbred population. *Genome Res* **23**: 1663–74.

Seo E, Kim T, Park JH, Yeom SI, Kim S, Seo MK, Shin C, Choi D, Tabata S. 2018. Genome-wide comparative analysis in Solanaceous species reveals evolution of microRNAs targeting defense genes in *Capsicum* spp. *DNA Res* **25**: 561–575.

Shen Y, Sun S, Hua S, Shen E, Ye CY, Cai D, Timko MP, Zhu QH, Fan L. 2017. Analysis of transcriptional and epigenetic changes in hybrid vigor of allopolyploid *Brassica napus* uncovers key roles for small RNAs. *Plant J* **91**: 874–893.

Si-Ammour A, Windels D, Arn-Bouldoires E, Kutter C, Ailhas J, Meins F, Vazquez F. 2011. miR393 and Secondary siRNAs Regulate Expression of the TIR1 / AFB2 Auxin Receptor Clade and Auxin-Related Development of *Arabidopsis* Leaves. *Plant Physiol* **157**: 683–91.

Song Q, Guan X, Chen ZJ. 2015. Dynamic Roles for Small RNAs and DNA Methylation during Ovule and Fiber Development in Allotetraploid Cotton. *PLoS Genet* **11**: e1005724.

Song QX, Xiang L, Li QT, Chen H, Hu XY, Ma B, Zhang WK, Chen SY, Zhang JS. 2013. Genome-Wide analysis of DNA methylation in soybean. *Mol Plant* **6**: 1961–74.

Srivastava S, Zheng Y, Kudapa H, Jagadeeswaran G, Hivrale V, Varshney RK, Sunkara R. 2015. High throughput sequencing of small RNA component of leaves and inflorescence revealed conserved and novel miRNAs as well as phasiRNA loci in chickpea. *Plant Sci* **235**: 46–57.

The International Brachypodium Initiative. 2010. Genome sequencing and analysis of the model grass *Brachypodium distachyon*. *Nature* **463**: 763–8.

The Tomato Genome Consortium. 2012. The tomato genome sequence provides insights into

979 fleshy fruit evolution. *Nature* **485**: 635–41.

980 Vaucheret H, Mallory AC, Bartel DP. 2006. AGO1 Homeostasis Entails Coexpression of MIR168
981 and AGO1 and Preferential Stabilization of miR168 by AGO1. *Mol Cell* **22**: 129–36.

982 Vicient CM, Casacuberta JM. 2017. Impact of transposable elements on polyploid plant
983 genomes. *Ann Bot* **120**: 195–207.

984 Vicient CM, Jääskeläinen MJ, Kalendar R, Schulman AH. 2001. Active Retrotransposons Are a
985 Common Feature of Grass Genomes. *Plant Physiol* **125**: 1283–92.

986 Wei L, Gu L, Song X, Cui X, Lu Z, Zhou M, Wang L, Hu F, Zhai J, Meyers BC, et al. 2014. Dicer-like
987 3 produces transposable element-associated 24-nt siRNAs that control agricultural traits in
988 rice. *Proc Natl Acad Sci* **111**: 3877–82.

989 Woodhouse MR, Cheng F, Pires JC, Lisch D, Freeling M, Wang X. 2014. Origin, inheritance, and
990 gene regulatory consequences of genome dominance in polyploids. *Proc Natl Acad Sci* **111**:
991 5283–8.

992 Xia J, Zeng C, Chen Z, Zhang K, Chen X, Zhou Y, Song S, Lu C, Yang R, Yang Z, et al. 2014.
993 Endogenous small-noncoding RNAs and their roles in chilling response and stress
994 acclimation in Cassava. *BMC Genomics* **15**: 634.

995 Xia R, Chen C, Pokhrel S, Ma W, Huang K, Patel P, Wang F, Xu J, Liu Z, Li J, et al. 2019. 24-nt
996 reproductive phasiRNAs are broadly present in angiosperms. *Nat Commun* **10**: 627.

997 Xia R, Xu J, Arikait S, Meyers BC. 2015a. Extensive families of miRNAs and PHAS loci in Norway
998 spruce demonstrate the origins of complex phasiRNA networks in seed plants. *Mol Biol*
999 *Evol* **32**: 2905–18.

1000 Xia R, Ye S, Liu Z, Meyers BC, Liu Z. 2015b. Novel and Recently Evolved MicroRNA Clusters
1001 Regulate Expansive F-BOX Gene Networks through Phased Small Interfering RNAs in Wild
1002 Diploid Strawberry. *Plant Physiol* **169**: 594–610.

1003 Xie Z, Kasschau KD, Carrington JC. 2003. Negative feedback regulation of Dicer-Like1 in
1004 Arabidopsis by microRNA-guided mRNA degradation. *Curr Biol* **13**: 784–9.

1005 Xu C, Jiao C, Sun H, Cai X, Wang X, Ge C, Zheng Y, Liu W, Sun X, Xu Y, et al. 2017. Draft genome
1006 of spinach and transcriptome diversity of 120 Spinacia accessions. *Nat Commun* **8**: 15275.

1007 Yang L, Huang H. 2014. Roles of small RNAs in plant disease resistance. *J Integr Plant Biol* **56**:
1008 962–70.

1009 Zhai J, Jeong DH, de Paoli E, Park S, Rosen BD, Li Y, González AJ, Yan Z, Kitto SL, Grusak MA, et
1010 al. 2011. MicroRNAs as master regulators of the plant NB-LRR defense gene family via the
1011 production of phased, trans-acting siRNAs. *Genes Dev* **25**: 2540–53.

1012 Zhai J, Zhang H, Arikait S, Huang K, Nan GL, Walbot V, Meyers BC. 2015. Spatiotemporally
1013 dynamic, cell-type-dependent premeiotic and meiotic phasiRNAs in maize anthers. *Proc*
1014 *Natl Acad Sci U S A* **112**: 3146–51.

1015 Zhai L, Sun W, Zhang K, Jia H, Liu L, Liu Z, Teng F, Zhang Z. 2014. Identification and
1016 characterization of Argonaute gene family and meiosis-enriched Argonaute during
1017 sporogenesis in maize. *J Integr Plant Biol* **56**: 1042–52.

1018 Zhang, Zhang, Hao, Song, Li, Li, Gao, Zheng, Li. 2019. Lineage-Specific Evolved MicroRNAs
1019 Regulating NB-LRR Defense Genes in Triticeae. *Int J Mol Sci* **20**: E3128.

1020 Zhao Y, Yu Y, Zhai J, Ramachandran V, Dinh TT, Meyers BC, Mo B, Chen X. 2012. The arabidopsis
1021 nucleotidyl transferase HESO1 uridylates unmethylated small RNAs to trigger their
1022 degradation. *Curr Biol* **22**: 689–94.

- Zheng Q, Rowley MJ, Böhmendorfer G, Sandhu D, Gregory BD, Wierzbicki AT. 2013. RNA polymerase v targets transcriptional silencing components to promoters of protein-coding genes. *Plant J* **73**: 179–89.
- Zheng Y, Wang Y, Wu J, Ding B, Fei Z. 2015. A dynamic evolutionary and functional landscape of plant phased small interfering RNAs. *BMC Biol* **13**: 32.
- Zhong X, Hale CJ, Law JA, Johnson LM, Feng S, Tu A, Jacobsen SE. 2012. DDR complex facilitates global association of RNA polymerase v to promoters and evolutionarily young transposons. *Nat Struct Mol Biol* **19**: 870–5.

Supplemental Tables

Supplemental Table S1. sRNA-seq libraries that were used as components in reference sets. Format: comma-separated values (csv).

Supplemental Table S2. Small RNA-producing loci from 48 plant genomes. Gzip-compressed, tab-separated text file. Note that the project website (<http://plantsmallrnagenes.science.psu.edu>) has the same data with search functions, more details, and alternative formats.

Supplemental Table S3. List of genes containing siRNA21 and siRNA22 loci in the nine monocot species studied.

Supplemental Table S4. Grouping of small RNA loci into putative families based on sequences of most abundant sRNA sequence. Gzip-compressed, tab-delimited text file.

Supplemental Figures

Supplemental Fig. S1. Properties of genome assemblies used in this study. See Table 1 for species codes. Eu: eudicots, BA: basal angiosperm, G: gymnosperm, Br: bryophyte. **(A)** Genome assembly sizes (log₁₀ scale, nucleotides). **(B)** Scaffold N50 lengths (log₁₀ scale, nucleotides). **(C)** Scaffold L50 ranks (log₁₀ scale).

Supplemental Fig. S2. Properties of gene annotations and genome assemblies used in this study. See Table 1 for species codes. Eu: eudicots, BA: basal angiosperm, G: gymnosperm, Br: bryophyte. **(A)** Counts of annotated genes that contain one or more CDS features. Note: data were unavailable for three assemblies (tcc-b1, mdm-b3.0, can-b1.6). **(B)** Repetitiveness, gaps, and single-copy regions, defined by k-mer analysis of genome assemblies with k=24. K-mers containing one or more ambiguous character were tallied as gaps. Non-ambiguous k-mers present more than once in the assembly were tallied as repetitive.

Supplemental Fig. S3. Summary of sRNA-seq libraries and reference sets. See Table 1 for species codes. Eu: eudicots, BA: basal angiosperm, G: gymnosperm, Br: bryophyte. **(A)** Total number of sRNA-seq reads in reference sets. **(B)** Number of individual sRNA-libraries in reference sets.

Supplemental Fig. S4. Summary of annotated sRNA loci, by species and locus type, including the category 'OtherRNA'. See Table 1 for species codes. Eu: eudicots, BA: basal angiosperm, G: gymnosperm, Br: bryophyte. **(A)** Counts of annotated loci. **(B)** Proportions of annotated loci. **(C)** Total counts of aligned small RNAs in reference sets. **(D)** Proportions of small RNA total read counts in reference sets.

Supplemental Fig. S5. Chromosomal distribution of genes and sRNA loci.

Species with a number of annotated sRNA loci high enough to have a continuous chromosomal distribution are shown, instead, species with a low number of annotated sRNA loci that are scattered across the chromosomes are not shown. For each species one representative chromosome is shown. Purple: distribution of genes. Green: distribution of sRNA loci, including the category 'OtherRNA', which represents a minor contribution on the total loci for the species reported here. See Table 1 for species codes. Eu: eudicots.

Supplemental Fig. S6. Number of protein-coding mRNAs that overlap with sRNA loci. Counts of mRNAs, containing at least one intron, that overlap with a sRNA locus for at least 25% of the length of the sRNA locus. In case of overlapping with multiple sRNA loci of different categories, the mRNA intersection was classified based on the longest overlapping sRNA locus. See Table 1 for species codes. Eu: eudicots, BA: basal angiosperm, G: gymnosperm, Br: bryophyte.