1 **Parallel clines in native and introduced ragweed populations are** 2 **likely due to adaptation**

3

4 Brechann V. McGoey¹, Kathryn A. Hodgins², and John R. Stinchcombe^{1, 3, *}

5

6 1. Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, ON

7 Canada M5S 3B2

8 2. School of Biological Sciences, Monash University, Clayton, VIC, Australia

9

10 3. Koffler Scientific Reserve, University of Toronto, Toronto, ON Canada M5S3B2

11 Emails: Brechann.mcgoey@utoronto.ca ; Kathryn.hodgins@monash.edu ;

12 john.stinchcombe@utoronto.ca

13 * Author for correspondence.

14 **Abstract**

15 As introduced species expand their ranges, they often encounter differences in climate which are
 16 often correlated with geography. For introduced species, encountering a geographically variable
 17 climate sometimes leads to the re-establishment of clines seen in the native range. However,
 18 clines can also be caused by neutral processes, and so it is important to gather additional
 19 evidence that population differentiation is the result of selection as opposed to non-adaptive
 20 processes. Here, we examine phenotypic and genetic differences in ragweed from the native
 21 (North America) and introduced (European) ranges. We used a common garden to assess
 22 phenotypic differentiation in size and flowering time in ragweed populations. We found
 23 significant parallel clines in flowering time in both North America and Europe. Height and
 24 branch number had significant clines in North America and, while not statistically significant,
 25 the patterns in Europe were the same. We used SNP data to assess population structure in both
 26 ranges and to compare phenotypic differentiation to neutral genetic variation. We failed to detect
 27 significant patterns of isolation by distance, geographic patterns in population structure, or
 28 correlations between the major axes of SNP variation and phenotypes or latitude of origin. We
 29 conclude that the clines seen for flowering time and size are most likely the result of adaptation.

30 **Introduction**

31 Invasive species are both biological disasters and curiosities. In addition to the dual economic
 32 and ecological damage introduced plants can cause when they proliferate, they also offer
 33 opportunities to study evolutionary ecology during colonization (Callaway & Maron 2006). For
 34 example, invasions provide ideal systems to study the effects of reproductive isolation, how
 35 dispersal affects species distributions and the effect of a new individual species on an ecosystem
 36 (Sax *et al.* 2007). Evolutionary biologists have garnered insights from introduced species for
 37 decades by studying patterns of variation, interactions with native species, and comparing native
 38 and introduced populations (Huey *et al.* 2005). Clines in introduced populations offer an
 39 opportunity to study parallel evolution, the rate and predictability of local adaptation, and
 40 whether phenotypic divergence is due to selection or stochastic, non-adaptive forces (Huey *et al.*
 41 2005; Samis *et al.* 2012; Colautti & Lau 2015). Here, we use a field common garden experiment
 42 and genotyping-by-sequencing (GBS) to investigate clines in introduced and native ragweed

43 populations with the goal of distinguishing between adaptive and non-adaptive mechanisms
 44 underlying clinal variation.

45 Incorporating an evolutionary perspective into invasion biology is critical to understanding the
 46 course an invasion has taken and how it might continue to unfold. For example, the capacity of
 47 an introduced population to adapt to its new climate is important for its capacity to persist and
 48 spread (Colautti & Barrett 2013). Adaptation to climate variables often lead to geographic
 49 differentiation, as climate and geography are strongly correlated (Endler 1977). The common
 50 pattern of a gradient in traits or alleles over a geographic range (Huxley 1938) is often
 51 interpreted as evidence of adaptive differentiation. Clines can be found in both Mendelian and
 52 quantitative traits and there are hundreds of examples across a wide variety of taxa (Campitelli
 53 2013). Clines in introduced species, especially those that mirror geographic variation in the
 54 native range, are often perceived as evidence that introduced populations have adapted to their
 55 new environments (Samis et al. 2012; Colautti and Barrett 2013). However, processes other than
 56 adaptation can also be responsible for both phenotypic and genetic clines, and need to be
 57 controlled for (Keller and Taylor 2008). For example, phenotypic clines observed *in situ* could
 58 be caused by plastic responses to environmental variables, especially in plant species (Huxley
 59 1938), and neutral processes present through colonization could also produce clines (Vasemägi
 60 2006; Keller *et al.* 2009; Santangelo *et al.* 2018).

61 Distinguishing between the possible forces underlying clines can be achieved in several ways.
 62 Plasticity can be excluded by using a common garden to ensure that any differences between
 63 populations have a genetic basis (Lucek *et al.* 2014). Linking clines with natural selection and
 64 demonstrating a correspondence between the direction of selection and variation in phenotypes
 65 can also strengthen the case that a cline is the result of adaptation (Etterson *et al.* 2008). Parallel
 66 clines can also be interpreted as evidence of adaptation (Samis et al. 2012). Neutral markers can
 67 be used to rule out drift and support hypotheses that differences are due to adaptation (Keller &
 68 Taylor 2008; Kooyers & Olsen 2012; Lima *et al.* 2012; Campitelli and Stinchcombe 2013; Le
 69 Gros *et al.* 2016). Likewise Q_{ST} - F_{ST} comparisons can be used to compare molecular and
 70 quantitative genetic variation. Whereas F_{ST} examines differentiation at neutral markers, Q_{ST} is
 71 analogous but quantifies population divergence for quantitative traits (Spitze 1993). Meta-
 72 analyses have found that genetic differentiation among introduced populations are common and
 73 on average do not differ much in magnitude from divergence between native populations

74 (Colautti and Lau 2015). By using these methods, researchers have bolstered the argument that
 75 rapid adaptation occurs in non-native species from plants to fruit flies (Huey *et al.* 2000;
 76 Montague *et al.* 2008; Samis *et al.* 2012; Colautti & Lau 2015).

77 *Ambrosia artemisiifolia* (ragweed) is a globally invasive species with a wide range in its native
 78 continent of North America and a presence in Europe, Asia and Australia (Friedman & Barrett
 79 2008). Past work on *A. artemisiifolia* has demonstrated parallel clinal patterns in flowering time
 80 in the native and introduced ranges (Hodgins & Rieseberg 2011). In this experiment, we examine
 81 variation in several quantitative traits across geography. To determine how quantitative variation
 82 may have been impacted by neutral processes, we use single neutral polymorphism (SNP) data to
 83 assess neutral genetic variation. We ask the questions: **Are there clinal patterns in quantitative**
 84 **traits, in the introduced and native ranges? Are the patterns consistent with a history of**
 85 **selection in the introduced range, or non-adaptive processes?** Our results corroborate, using
 86 independent biological samples, experiments, and analytical approaches, past results (Hodgins
 87 and Rieseberg 2011; van Boheemen *et al.* 2018) of clinal variation in ragweed being likely due to
 88 adaptive differentiation rather than stochastic processes.

89 **Methods**

90 *Study species*

91 *Ambrosia artemisiifolia* (common ragweed) is an annual outcrosser in the Asteraceae family
 92 (Bassett & Crompton 1975; Friedman & Barrett 2008). Ragweed is thought to have originated in
 93 the plains of North America, and then spread eastward (Bassett and Crompton 1975). In the
 94 modern era, ragweed has been accidentally introduced to Europe, Asia and Australia (MacKay &
 95 Kotanen 2008). In Europe, ragweed has been present since at least the mid-1800s, but propagule
 96 pressure (composite measure of the individuals or seeds released in an introduction and the
 97 number of introductions (Lockwood *et al.* 2005)) increased dramatically in the mid-20th century
 98 when ragweed seeds contaminated grains shipped from the Americas to Europe (Chauvel *et al.*
 99 2006). The geopolitical situation during that period meant that imports were coming from
 100 different areas into Western versus Eastern Europe. This resulted in two invasion centres, with
 101 distinct genetic origins (Gladieux *et al.* 2011). In France, the epicenter of the invasion is the
 102 Rhône valley, where ragweed grows in very large populations along riverbanks (Chauvel *et al.*

2006; Thibaudon *et al.* 2013). The most impacted nation is Hungary, and in Eastern Europe ragweed populations now extend north up into Poland and south into the Baltic states (Prank *et al.* 2013). Ragweed is considered one of the most problematic invaders in Europe: it causes allergies, and is a significant agricultural weed. In Hungary, it is the most widespread weed in surveys (Kiss & Béres 2006) and over 80% of arable land is affected (Buttenschön *et al.* 2010).

Seed collection and preparation

In the autumns of 2012 and 2013, we collected seeds from a total of 20 native and 18 introduced populations (Figure 1; population coordinates in Appendix Tables A1 and A2). In both ranges the populations spanned ~7.5 degrees in latitude. These populations ranged from small (5 individuals) up to tens of thousands of individuals. When populations had fewer than twenty individuals, we collected seeds from all the plants. For larger populations, we collected from a random subset. Using methods adapted from Willemsen (1975) and Jannice Friedman (Queen's University, pers. comm.), we stratified seeds at 4°C for six months in plastic bags filled with silica and distilled water.

Common garden

For this experiment we randomly chose 10 maternal families per population, except in cases where there were fewer than 10 families, in which case we used all available. We chose three germinants from each family (one for each of three blocks) for a total of 1110 plants. Each individual was randomly assigned a position within a block. We planted germinants in seedling trays and kept in a greenhouse where we sprayed and bottom watered them for three weeks. At the end of June 2014, we planted seedlings into a plowed field at the Koffler Scientific Reserve (www.ksr.utoronto.ca; 44.803°N, 79.829°W). Blocks were subdivided into plots, each containing 64 plants in an 8x8 configuration, except for the final plot. We continued to remove interspecific vegetation and provide the seedlings with water for four weeks after transplantation to promote establishment.

Phenotypic traits

We measured final height, final number of branches, and date of first flower. Since the vast majority of ragweed plants are monoecious (Bassett and Crompton 1975), we measured proxies

132 of both male and female fitness. Male reproductive effort was estimated as the total inflorescence
 133 length, which is correlated with pollen production (Fumanal *et al.* 2007). We estimated female
 134 reproductive output, using seed mass, which is highly correlated with seed number ($r^2=0.96$,
 135 $p<0.001$) (MacDonald and Kotanen 2010). We ran initial models including latitude and block to
 136 assess if block had a significant effect on traits; block was not significant for any trait, and was
 137 thus dropped in subsequent models. To test for clines in phenotypic traits, we used linear
 138 regressions. For each continent, we conducted regressions for three phenotypic traits (height,
 139 flowering time and branch number) and the two fitness traits on latitude. Similar results were
 140 obtained using population means for traits and latitude. Unless otherwise specified below,
 141 statistical analyses were conducted in R (R Development Core Team 2016).

142 *Neutral markers*

143 We collected leaf material from 180 ragweed plants from 26 populations (9 introduced, 17
 144 native) grown in seedling trays in growth chambers at the University of Toronto. These plants
 145 were from a subset of the populations used in the common garden, but were separate plants. GBS
 146 library prep was conducted by the Hodgins lab at Monash University Australia. In brief, DNA
 147 was digested from the dried leaves and adapters were ligated to the strands. A double enzyme
 148 digest with Pst1 and Msp1 was implementing using the same protocol as in van Boheemen et al
 149 (2018). DNA libraries were sent to Genome Quebec for sequencing on an Illumina HiSeq 2500,
 150 using PE125 sequencing.

151 We used Stacks (Catchen *et al.* 2013) and Bowtie 2 (Langmead & Salzberg 2012) to
 152 demultiplex, align to a reference genome provided by the Hodgins lab and to calculate
 153 population genetic metrics. We checked the sequence quality using FastQC (Andrews 2010) and
 154 samtools (Li *et al.* 2009). We converted between multiple formats using PGD Spider (Lischer &
 155 Excoffier 2011), samtools (Li et al. 2009), Bowtie 2 (Langmead and Salzberg 2012), admixr
 156 (Petr n.d.) and custom python and bash scripts (see Github). To prepare for STRUCTURE and
 157 isolation by distance (IBD) analysis we filtered snps using vcf tools (Danecek *et al.* 2011). We
 158 excluded snps with a minor allele count lower than 4 (equivalent to 2.2%) or with data missing in
 159 greater than 20% of samples. These thresholds were average to conservative based on those used
 160 in previous studies (McGrath 2014; Huang *et al.* 2014; Taylor *et al.* 2014; Sawler *et al.* 2015;
 161 Beck & Semple 2015; Ilut *et al.* 2015; Martin *et al.* 2016; Mondon *et al.* 2017). We set the

significance threshold for Hardy-Weinberg equilibrium at $1e-5$, which was the midpoint used in a review of past studies (Anderson *et al.* 2010). Since very rare alleles could still be important for assessing population structure (Linck & Battey 2017), we evaluated whether inclusion of them altered the results, and found that they did not. We present results with the exclusion of SNPs with rare minor allele counts $<2.2\%$.

Population Structure and Geography.

To conduct a STRUCTURE analysis (Pritchard *et al.* 2000) while taking advantage of parallelization, we used StrAuto (Chhatre & Emerson 2017). We conducted separate analyses for the two continents, with five replicates at $K=1-6$ for each. In addition, we ran a STRUCTURE analysis using data from all the populations together. To visualize the STRUCTURE output we used the default settings of the web-based program Cluster Markov Packager Across K (CLUMPAK) (Kopelman *et al.* 2015) and the R package *pophelper* (Francis 2017).

To examine isolation by distance (IBD), we used the R package *adeigenet* to test for IBD in each continent separately (Jombart 2008). *Adegenet* uses a Mantel test between matrices of genetic and geographic distances to assess if more spatially disparate populations are also more genetically divergent.

In addition to STRUCTURE and IBD, patterns in genetic data can also be understood with principal component analysis (Cavalli-Sforza *et al.* 1994; Patterson *et al.* 2006; McVean 2009; Josephs *et al.* 2018). We performed PCA on SNP data from the native and introduced ranges to test for correlations between major axes of SNP variation and the phenotypic traits of interest, and latitude. To extract principal components from SNPs we used the R package LEA (Frichot & François 2015). The vcf files were converted to geno files using the *vcf2geno* function. We then ran PCAs of all available SNPs separately for the two continents. To determine which principal components should be retained in subsequent analyses, we used Tracy-Widom tests (Patterson *et al.* 2006). For each population, we calculated a PC score along each significant axis of SNP variation, and then used these PC scores to test for associations with traits or geography. Specifically, to determine whether axes of neutral SNP variation were related to either geography or phenotype with linear regressions. For each continent, we regressed each significant principal component on latitude and the three phenotypic traits.

191 *Descriptive Population Genetic Statistics*

192 To explore population genetics of the native and introduced ragweed populations, we used the
 193 programs Genodive and Splitstree (Hudson 1998; Meirmans & Van Tienderen 2004). We
 194 converted vcf files for each continent, and for the entire dataset, to the genetix format and then
 195 imported them into Genodive. We then used Genodive to estimate F_{ST} , population genetic
 196 summary statistics (including observed and expected heterozygosity and fixation indices), and to
 197 run an AMOVA to partition genetic variation between the range, population and individual
 198 levels. We used Splitstree to construct a neighbornet for all the individuals (Hudson 1998).

199 *Data availability.* Upon acceptance, accession numbers for sequences deposited in the short read
 200 archive will be placed here (<https://www.ncbi.nlm.nih.gov/sra>). A summary of our
 201 bioinformatics pipeline, including code and the original fastq files is available here
 202 (<https://github.com/brechann-mcgoey/ragweedGBS>)

203 **Results**

204 *Phenotypic traits*

205 Plants from more southern latitudes flowered later and grew larger both in total height and
 206 branch number (see Figure 2). These clines were all significant for the North American
 207 populations, while only flowering time had a statistically significant association with latitude in
 208 Europe (see Table 1). There was only one significant cline for fitness traits, with more southern
 209 North American plants producing more fruits than those in more northern populations (Figure 3).
 210 Since more southern plants were also larger, this correlation may be driven by a relationship
 211 between size and latitude.

212 *Neutral markers*

213 The Illumina analysis resulted in 250 033 952 total sequences which all passed a FastQC quality
 214 check. We started with 258 540 SNPs across all the populations. After filtering, we had 20 843
 215 sites for Europe and 28 643 for North America. Our STRUCTURE analysis indicated that the
 216 European samples clustered in three groups, while the North American lines clustered in five
 217 groups. In Europe, there was no obvious geographic pattern to ancestry (see Figure 4A). In North
 218 America, two populations seem distinct, but otherwise there is not much geographic structure

(see Figure 4B). There was no significant isolation by distance in either continent (p-value of 0.84 for Europe and 0.342 for North America). Our Splitstree analysis was consistent with the above results, showing populations were not clustered into distinct subgroups (Appendix Figure A1). The global STRUCTURE analysis for all populations found the most support for $k=5$. As with the by-continent analyses, little of the ancestry was geographically structured (see Appendix Figure A2).

Tracy-Widom tests indicated that there was one significant SNP principal component in Europe and seven for the North American dataset (North America: TW Statistic ≥ 1.029 , $P \leq 0.04671$ for PCs 1-7; Europe: TW Statistic = 6.255, $P = 1.013E-06$ for PC1). The significant European principal component explained 4.5% of genetic variation. Altogether, the significant North America principal components explained 14.6% of genetic variation. There were no significant relationships between significant SNP principal component scores and either latitude or phenotype (all p values >0.05). These results suggest that axes of (neutral) SNP variation are not related to either latitude, or the phenotypic traits, suggesting that latitudinal clines are unlikely to be generated by neutral or stochastic processes.

In keeping with the lack of geographic structure revealed by the Splitstree and Structure results, F_{ST} values were universally low (mean of 0.0639 (range of 0.003 to 0.213) for North America and 0.0566 (range of 0.018 to 0.12) for Europe, see Appendix tables A13& A4 for population specific results). Other population level statistics were consistent with the conclusion that there is low population differentiation in both ranges (see Appendix tables A5 & A6). An AMOVA of the global dataset indicated that very little variance was at the between continent or between population levels (see Appendix table A7). Total heterozygosity was lower in native (0.150) as compared to the invasive range (0.202). The expected heterozygosity within populations was also higher in the invasive range (0.192 vs 0.141 for the native range).

Discussion

Introduced species offer a unique opportunity to address important questions in evolutionary biology (Sax et al. 2007; Yoshida et al. 2007). Adaptation is an important and sometimes overlooked aspect of invasions (Huey et al. 2005; Barrett 2015). New species introductions offer an excellent avenue to explore the rate and predictability of adaptation, a topic of great interest to

evolutionary biologists (Huey et al. 2005). The reemergence of genetic clines in introduced ranges represents evidence that adaptation may occur quickly and result in predictable phenotypic differentiation (Leger & Rice 2007). Adaptation to local habitats and population differentiation can be critical to the ability of an invasive species to expand its range (Colautti and Barrett 2013). Here we detected phenotypic differentiation among native and introduced populations of common ragweed, and evidence that these clines are the results of local adaptation as opposed to neutral processes. We also examined the population structure and genetic diversity of both native and invasive populations.

Clines as evidence of local adaptation

Past studies of ragweed have found geographic patterns in phenotype, including parallel clines in the native and introduced ranges (Hodgins and Rieseberg 2011, van Boheemen et al. 2019). With populations from different parts of both Europe and North America than those used in previous studies, we corroborated their conclusions that flowering time patterns have been reproduced in the introduced range. We present evidence that these patterns are almost certainly the results of selection as opposed to neutral processes. Unlike flowering time, patterns of SNP variation were not correlated with latitude, and principal components of genetic variation were not correlated with the phenotypic traits we examined, including flowering time. Overall there was little population structure and low F_{ST} values were identified in both ranges. Van Boheemen et al. (2019) detected repeated latitudinal divergence in a host of life history and size traits in Australian, North American, and European samples of ragweed, as well. In their case, they detected these clines when controlling for structure coefficients (so-called q-values) as a measure of neutral population genetic structure. Here we used independent biological samples, phenotyped plants in a field common garden in the native range (as opposed to growth chamber), and used alternative means of assessing neutral population genetic structure (PCA of the SNP matrix). That we found qualitatively similar results—clinal differentiation above and beyond what could be explained by neutral processes—all make it unlikely that the cline in flowering time is the results of drift.

One caveat of this study is that all plants were grown from seed and therefore subject to maternal effects. However, we think it unlikely that maternal effects could explain the phenotypic patterns presented here. The population level maternal effects would all need to be in a consistent

direction across latitude. Maternal effects are more pervasive in early-life history stages (Rossiter 1996; Montague *et al.* 2008), while here we focused on traits at the end of the annual life cycle. It is also unlikely that maternal effects would be responsible for differentiation at such large geographic scales (Montague *et al.* 2008). In addition, the clines found here were consistent with those of our previous work (McGoey and Stinchcombe 2018), which included fewer populations but did remove the impact of maternal effects.

Ragweed has been present in Europe for ~250 years (Chauvel *et al.* 2006). In as many generations, it has spread and proliferated, especially in France, Italy and the Pannonian Plain (Thibaudon *et al.* 2013). Our previous research demonstrated that invasive ragweed populations possess ample additive genetic variation, in spite of any bottlenecks and founder effects during the colonization process. Given the results shown here, it must also be concluded that this quantitative genetic variation has been preserved in the face of selection and adaptation as well.

Past studies of population differentiation, including one examining ragweed in its introduced range, have used Q_{ST} - F_{ST} comparisons to assess geographic variation (Leinonen *et al.* 2008; Chun *et al.* 2009). As with our study, Chun and colleagues found the invasive ragweed populations had low F_{ST} values and geographic structure (Chun *et al.* 2009). Their Q_{ST} - F_{ST} analysis indicated there was significant diversifying selection acting on ragweed in its introduced range. The conclusions of Chun and colleagues align with our own, that population differentiation in the invasive range is the result of adaptive evolution as opposed to neutral processes. Q_{ST} - F_{ST} comparisons are very interesting but fraught with technical challenges and statistical difficulties (Whitlock 2008). For example, true Q_{ST} comparisons must obtain estimates of additive genetic variance, which means controlled crosses of dozens of pairs from each population. Doing so for six populations for our previous work was challenging, doing so for more than thirty populations would be prohibitive. Given the caveats that would be necessary with this dataset (i.e. having to use P_{ST} instead of Q_{ST}), we decided not to pursue this analysis here.

Population genetics

Our population genetic results indicate that there is very little geographic population structure in both continents. Population differentiation in neutral markers as measured by F_{ST} was low for

both the native (0.0639) and invasive (0.0566) ranges. Past work on ragweed has found similarly low F_{ST} values (Genton *et al.* 2005; Chun *et al.* 2010; 2011; Martin 2011; Martin *et al.* 2016; van Boheemen *et al.* 2017). Martin and colleagues used SNPs to assess population differentiation of North American ragweed, and found that a solitary genetic cluster was the most likely population structure (Martin *et al.* 2016). Our STRUCTURE analyses showed the highest likelihoods for multiple clusters, but there was not a geographic pattern to the ancestral groupings, especially for the European populations. The consistent findings of low isolation by distance and population structure may not be surprising given that ragweed is a wind-pollinated outcrosser (Friedman and Barrett 2008).

The assumption that introduced species will always face significant deficits in genetic variation has been challenged by numerous counter-examples in the literature (Colautti & Lau 2015; Estoup *et al.* 2016). In some cases, due to multiple introductions and subsequent admixture, molecular diversity is actually higher in invasive populations when compared to their native counterparts (Novak & Mack 1993; Dlugosch & Parker 2008; Keller & Taylor 2008; 2010). Here, we estimated slightly higher metrics of genetic diversity (i.e., expected heterozygosity) in the invasive range for the represented populations. These findings corroborate a study in the French and North American ranges using microsatellite genetic variability where within population diversity was higher in the invasive range and overall genetic diversity was comparable between the two ranges (Genton *et al.* 2005). The high genetic diversity in the invasive range is likely the result of high propagule introduction from multiple sources in the native range (Genton *et al.* 2005a). The persistence of large, genetically diverse populations in the worst affected areas of Europe are dangerous sources for the spread of ragweed into currently unaffected areas. Roads and railway tracks are ideal corridors and habitats for ragweed and could facilitate multiple introductions and gene flow (Lavoie *et al.* 2007).

Conclusions

Adaptation in introduced environments is not just theoretically interesting, but also has extremely important ecological and economic implications. Gradients in abiotic variables can lead to divergent selection across an introduced range and, if populations have sufficient genetic variation, create clines in traits (Maron *et al.* 2004). This adaptation can exacerbate the negative impacts of introduced species (Maron *et al.* 2004; Huey *et al.* 2005)

337 Ragweed has already caused economic and health impacts in Europe (Buttenschön et al. 2010).
 338 Climate change is expected to extend its growing season, and it continues to spread its
 339 geographic range (Ziska et al. 2011). Our research corroborates past findings that ragweed has
 340 been able to adapt to its invasive range (Chun et al. 2011; Hodgins and Rieseberg 2011). The
 341 genetic, phenotypic and ecological traits of introduced ragweed make it very likely that the
 342 invasion will worsen.

343 **Acknowledgments**

344 Bruce Hall, Bill Cole and Andrew Petrie provided excellent greenhouse support. Stephan
 345 Schneider provided much helpful assistance and encouragement at KSR. Antonio DiTomasso,
 346 Kathleen Howard, Rakesh Chandran, Quentin Martinez and Barbara Tokarska-Guzik helped to
 347 locate populations, assisted with collections and generously offered their very considerable
 348 expertise. Bruno Chauvel, Jannice Friedman, Peter Kotanen and all provided invaluable ragweed
 349 advice. Ali Parker, Amanda Gorton, Elisabeth McGregor, Rafal Janik and Valérie Kaakeh helped
 350 with collections. We are very grateful to all the authors of the tools that we used for this study,
 351 and help from many dedicated undergraduates. We thank Laura Galloway, Emily Josephs, Peter
 352 Kotanen, Art Weis, Stephen Wright for comments that improved this manuscript.

Literature Cited.

- Anderson, C.A., Pettersson, F.H., Clarke, G.M., Cardon, L.R., Morris, A.P. & Zondervan, K.T. (2010). Data quality control in genetic case-control association studies. *Nature Protocols*, **5**, 1564–1573.
- Andrews, S. (2010). *FastQC: a quality control tool for high throughput sequence data*.
- Bassett, I.J. & Crompton, C.W. (1975). The biology of Canadian weeds: 11. *Ambrosia artemisiifolia* L. and *A. psilostachya* DC. *Canadian Journal of Plant Science*, **55**, 463–476.
- Beck, J.B. & Semple, J.C. (2015). Next-generation sampling: Pairing genomics with herbarium specimens provides species-level signal in *Solidago* (asteraceae). *Applications in Plant Sciences*, **3**, 1500014–11.
- Buttenschön, R.M., Waldispühl, S. & Bohren, C. (2010). *Guidelines for management of common ragweed*.
- Callaway, R.M. & Maron, J.L. (2006). What have exotic plant invasions taught us over the past 20 years? *Trends in Ecology & Evolution*, **21**, 369–374.
- Campitelli, B. (2013). *Ecological, physiological and molecular population genetics of a single-locus leaf shape cline in ivyleaf morning glory, Ipomea hederacea*.
- Campitelli, B.E. and J.R. Stinchcombe. (2013). Natural selection maintains a single-locus leaf shape cline in the Ivyleaf morning glory, *Ipomoea hederacea*. *Molecular Ecology* **22**: 552–564.
- Catchen, J., Hohenlohe, P.A., Bassham, S., Amores, A. & Cresko, W.A. (2013). Stacks: an analysis tool set for population genomics. *Molecular Ecology*, **22**, 3124–3140.
- Cavalli-Sforza, L.L., Menozzi, P. & Piazza, A. (1994). *The History and Geography of Human Genes*. Princeton, New Jersey.
- Chauvel, B., Dessaint, F., Cardinal-Legrand, C. & Bretagnolle, F. (2006). The historical spread of *Ambrosia artemisiifolia* L. in France from herbarium records. *Journal of Biogeography*, **33**, 665–673.
- Chhatre, V. & Emerson, K. (2017). StrAuto: Automation and Parallelization of STRUCTURE Analysis. 1–5.
- Chun, Y.J., Fumanal, B., Laitung, B. & Bretagnolle, F. (2010). Gene flow and population admixture as the primary post-invasion processes in common ragweed (*Ambrosia artemisiifolia*) populations in France. *New Phytologist*, **185**, 1100–1107.
- Chun, Y.J., Le Corre, V.V. & Bretagnolle, F. (2011). Adaptive divergence for a fitness-related trait among invasive *Ambrosia artemisiifolia* populations in France. *Molecular Ecology*, **20**, 1378–1388.
- Chun, Y.J., Nason, J.D. & Moloney, K.A. (2009). Comparison of quantitative and molecular

389 genetic variation of native vs. invasive populations of purple loosestrife (*Lythrum salicaria*
390 L., Lythraceae). *Molecular Ecology*, **18**, 3020–3035.

391 Colautti, R.I. & Barrett, S.C.H. (2013). Rapid adaptation to climate facilitates range expansion of
392 an invasive plant. *Science*, **342**, 364–366.

393 Colautti, R.I. & Lau, J.A. (2015). Contemporary evolution during invasion: evidence for
394 differentiation, natural selection, and local adaptation. *Molecular Ecology*, **24**, 1999–2017.

395 Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., Handsaker, R.E.,
396 Lunter, G., Marth, G.T., Sherry, S.T., McVean, G., Durbin, R. 1000 Genomes Project
397 Analysis Group. (2011). The variant call format and VCFtools. *Bioinformatics*, **27**, 2156–
398 2158.

399 Dlugosch, K.M. & Parker, I.M. (2008). Founding events in species invasions: Genetic variation,
400 adaptive evolution, and the role of multiple introductions. *Molecular Ecology*, **17**, 431–449.

401 Endler, J.A. (1977). *Geographic variation, speciation, and clines*. Princeton Univ. Press,
402 Princeton, NJ.

403 Estoup, A., Ravigné, V., Hufbauer, R., Vitalis, R., Gautier, M. & Facon, B. (2016). Is there a
404 genetic paradox of biological invasion? *Annu Rev Ecol Evol Syst*, **47**, 51–72.

405 Etterson, J.R., Delf, D.E., Craig, T.P., Ando, Y. & Ohgushi, T. (2008). Parallel patterns of clinal
406 variation in *Solidago altissima* in its native range in central USA and its invasive range in
407 Japan. *Botany*, **86**, 91–97.

408 Frichot, E. & François, O. (2015). LEA: An Rpackage for landscape and ecological association
409 studies. *Methods in Ecology and Evolution*, **6**, 925–929.

410 Friedman, J. & Barrett, S.C.H. (2008). High outcrossing in the annual colonizing species
411 *Ambrosia artemisiifolia* (Asteraceae). *Annals of Botany*, **101**, 1303–1309.

412 Fumanal, B., Chauvel, B. & Bretagnolle, F. (2007). Estimation of pollen and seed production of
413 common ragweed in France. *Annals of Agricultural and Environmental Medicine*, **14**, 233–
414 236.

415 Genton, B.J., Shykoff, J.A. & Giraud, T. (2005). High genetic diversity in French invasive
416 populations of common ragweed, *Ambrosia artemisiifolia*, as a result of multiple sources of
417 introduction. *Molecular Ecology*, **14**, 4275–4285.

418 Gladieux, P., Giraud, T., Kiss, L., Genton, B.J., Jonot, O. & Shykoff, J.A. (2011). Distinct
419 invasion sources of common ragweed (*Ambrosia artemisiifolia*) in Eastern and Western
420 Europe. *Biological Invasions*, **13**, 933–944.

421 Hodgins, K.A. & Rieseberg, L.H. (2011). Genetic differentiation in life-history traits of
422 introduced and native common ragweed (*Ambrosia artemisiifolia*) populations. *Journal of*
423 *Evolutionary Biology*, **24**, 2731–2749.

424 Huang, Y.-F., Poland, J.A., Wight, C.P., Jackson, E.W. & Tinker, N.A. (2014). Using

- genotyping-by-sequencing (GBS) for genomic discovery in cultivated oat. *PLoS ONE*, **9**, e102448–16.
- Hudson, D.H. (1998). SplitsTree: a program for analyzing and visualizing evolutionary data, Version 2.4. *Bioinformatics*, **14**, 68–73.
- Huey, R.B., Gilchrist, G.W. & Hendry, A.P. (2005). Using Invasive Species to Study Evolution. (eds D.F. Sax, J.J. Stachowicz & S.D. Gaines), pp. 139–164. Sinauer, Sunderland, MA.
- Huey, R.B., Gilchrist, G.W., Carlson, M.L., Berrigan, D. & Serra, L. (2000). Rapid evolution of a geographic cline in size in an introduced fly. *Science*, **287**, 308–309.
- Huxley, J. (1938). Clines: An Auxiliary Taxonomic Principle. *Nature*, **142**, 491–520.
- Ilut, D.C., Sanchez, P.L., Costich, D.E., Friebe, B., Coffelt, T.A., Dyer, J.M., Jenks, M.A. & Gore, M.A. (2015). Genomic diversity and phylogenetic relationships in the genus *Parthenium* (Asteraceae). *Industrial Crops and Products*, **76**, 920–929.
- Jombart, T. (2008). adegenet: a R package for the multivariate analysis of genetic markers. *Bioinformatics*, **24**, 1403–1405.
- Josephs, E.B., Berg, J.J., Ross-Ibarra, J. & Coop, G. (2018). Detecting adaptive differentiation in structured populations with genomic data and common gardens. *Genetics* 211: 989-1004.
- Keller, S.R. & Taylor, D.R. (2010). Genomic admixture increases fitness during a biological invasion. *Journal of Evolutionary Biology*, **23**, 1720–1731.
- Keller, S.R. & Taylor, D.R. (2008). History, chance and adaptation during biological invasion: separating stochastic phenotypic evolution from response to selection. *Ecology Letters*, **11**, 852–866.
- Keller, S., Sowell, D., Neiman, M., Wolfe, L.M. & Taylor, D.R. (2009). Adaptation and colonization history affect the evolution of clines in two introduced species. *New Phytologist*, **183**, 678–690.
- Kiss, L. & Béres, I. (2006). Anthropogenic factors behind the recent population expansion of common ragweed (*Ambrosia artemisiifolia* L.) in Eastern Europe: Is there a correlation with political transitions? *Journal of Biogeography*, **33**, 2156–2157.
- Kooyers, N.J. & Olsen, K.M. (2012). Rapid evolution of an adaptive cyanogenesis cline in introduced North American white clover (*Trifolium repens* L.). *Molecular Ecology*, **21**, 2455–2468.
- Kopelman, N.M., Mayzel, J., Jakobsson, M., Rosenberg, N.A. & Mayrose, I. (2015). Clumpak: a program for identifying clustering modes and packaging population structure inferences across K. *Molecular Ecology Resources*, **15**, 1179–1191.
- Langmead, B. & Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature Methods*, **9**, 357–359.

- 461 Lavoie, C., Jodoin, Y. & Merlis, D. (2007). How did common ragweed (*Ambrosia artemisiifolia*
462 L.) spread in Québec? A historical analysis using herbarium records. *Journal of*
463 *Biogeography*, **34**, 1751–1761.
- 464 Le Gros, A., Clergeau, P., Zuccon, D., Cornette, R., Mathys, B. & Samadi, S. (2016). Invasion
465 history and demographic processes associated with rapid morphological changes in the Red-
466 whiskered bulbul established on tropical islands. *Molecular Ecology*, **25**, 5359–5376.
- 467 Leger, E.A. & Rice, K.J. (2007). Assessing the speed and predictability of local adaptation in
468 invasive California poppies (*Eschscholzia californica*). *Journal of Evolutionary Biology*, **20**,
469 1090–1103.
- 470 Leinonen, T., O'Hara, R.B. & Cano, J.M. (2008). Comparative studies of quantitative trait and
471 neutral marker divergence: a meta-analysis. *Journal of evolutionary ...*
- 472 Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
473 Durbin, R. 1000 Genome Project Data Processing Subgroup. (2009). The Sequence
474 Alignment/Map format and SAMtools. *Bioinformatics*, **25**, 2078–2079.
- 475 Lima, M.R., Macedo, R.H.F., Martins, T.L.F., Schrey, A.W., Martin, L.B. & Bensch, S. (2012).
476 Genetic and morphometric divergence of an invasive bird: The introduced house sparrow
477 (*Passer domesticus*) in Brazil. *PLoS ONE*, **7**, e53332–13.
- 478 Linck, E.B. & Battey, C.J. (2017). Minor allele frequency thresholds strongly affect population
479 structure inference with genomic datasets. *bioRxiv*, 1–26.
- 480 Lischer, H.E.L. & Excoffier, L. (2011). PGDSpider: an automated data conversion tool for
481 connecting population genetics and genomics programs. *Bioinformatics*, **28**, 298–299.
- 482 Lockwood, J.L., Cassey, P. & Blackburn, T. (2005). The role of propagule pressure in explaining
483 species invasions. *Trends in Ecology & Evolution*, **20**, 223–228.
- 484 Lucek, K., Sivasundar, A. & Seehausen, O. (2014). Disentangling the role of phenotypic
485 plasticity and genetic divergence in contemporary ecotype formation during a biological
486 invasion. *Evolution*, **68**, 2619–2632.
- 487 MacKay, J. & Kotanen, P.M. (2008). Local escape of an invasive plant, common ragweed
488 (*Ambrosia artemisiifolia* L.), from above-ground and below-ground enemies in its native
489 area. *Journal of Ecology*, **96**, 1152–1161.
- 490 Maron, J.L., Vilà, M., Bommarco, R., Elmendorf, S. & Beardsley, P. (2004). Rapid evolution of
491 an invasive plant. *Ecological Monographs*, **74**, 261–280.
- 492 Martin, M.D. (2011). *Evolutionary mechanisms of Ambrosia artemisiifolia invasion*. Baltimore.
- 493 Martin, M.D., Olsen, M.T., Samaniego, J.A., Zimmer, E.A. & Gilbert, M.T.P. (2016). The
494 population genomic basis of geographic differentiation in North American common ragweed
495 (*Ambrosia artemisiifolia* L.). *Ecology and Evolution*.
- 496 McGrath, K. (2014). Investigating the North American *Crepis* (Asteraceae) agamic complex for

- 497 evidence of homoploid hybridization. MSc. thesis, University of British Columbia
- 498 McGoey, B.V. and J.R. Stinchcombe. (2018). Introduced populations of ragweed show as much
499 evolutionary potential as native populations. *bioRxiv* 305540.
- 500 McVean, G. (2009). A Genealogical Interpretation of Principal Components Analysis (M.
501 Przeworski, Ed.). *PLoS genetics*, **5**, e1000686–10.
- 502 Meirmans, P.G. & Van Tienderen, P.H. (2004). Genotype and Genodive: two programs for the
503 analysis of genetic diversity of asexual organisms. *Molecular Ecology Notes*, **4**, 792–794.
- 504 Mondon, A., Owens, G.L., Poverene, M., Cantamutto, M. & Rieseberg, L.H. (2017). Gene flow
505 in Argentinian sunflowers as revealed by genotyping-by-sequencing data. *Evolutionary*
506 *Applications*, **11**, 193–204.
- 507 Montague, J., Barrett, S.C.H. & Eckert, C. (2008). Re-establishment of clinal variation in
508 flowering time among introduced populations of purple loosestrife (*Lythrum salicaria*,
509 *Lythraceae*). *Journal of Evolutionary Biology*, **21**, 234–245.
- 510 Novak, S.J. & Mack, R.N. (1993). Genetic variation in *Bromus tectorum* (Poaceae): comparison
511 between native and introduced populations. *Heredity*, **71**, 167–176.
- 512 Patterson, N., Price, A.L. & Reich, D. (2006). Population Structure and Eigenanalysis. *PLoS*
513 *genetics*, **2**, e190–20.
- 514 Prank, M., Chapman, D.S., Bullock, J.M., Belmonte, J., Berger, U., Dahl, A., Jäger, S.,
515 Kovtunen, I., Magyar, D., Niemelä, S., Rantio-Lehtimäki, A., Rodinkova, V., Sauliene, I.,
516 Severova, E., Sikoparija, B. & Sofiev, M. (2013). An operational model for forecasting
517 ragweed pollen release and dispersion in Europe. *Agricultural and Forest Meteorology*, **182-**
518 **183**, 43–53.
- 519 Pritchard, J.K., Stephens, M. & Donnelly, P. (2000). Inference of population structure using
520 multilocus genotype data. *Genetics*, **155**, 945–959.
- 521 Rossiter, M. (1996). Incidence and consequences of inherited environmental effects. *Annual*
522 *Review of Ecology and Systematics*, **27**, 451–476.
- 523 Samis, K.E., Murren, C.J., Bosssdorf, O., Donohue, K., Fenster, C.B., Malmberg, R.L.,
524 Purugganan, M.D. & Stinchcombe, J.R. (2012). Longitudinal trends in climate drive
525 flowering time clines in North American *Arabidopsis thaliana*. *Ecology and Evolution*, **2**,
526 1162–1180.
- 527 Santangelo, J.S., Johnson, M.T.J. & Ness, R.W. (2018). Modern spandrels: the roles of genetic
528 drift, gene flow and natural selection in the evolution of parallel clines. *Proceedings of the*
529 *Royal Society of London. Series B. Biological Sciences*, **285**, 20180230–9.
- 530 Sawler, J., Stout, J.M., Gardner, K.M., Hudson, D., Vidmar, J., Butler, L., Page, J.E. & Myles, S.
531 (2015). The Genetic Structure of Marijuana and Hemp. *PLoS ONE*, **10**, e0133292.
- 532 Sax, D.F., Stachowicz, J.J., Brown, J.H., Bruno, J.F., Dawson, M.N., Gaines, S.D., Grosberg,

533 R.K., Hastings, A., Holt, R.D., Mayfield, M.M., O'Connor, M.I. & Rice, W.R. (2007).
534 Ecological and evolutionary insights from species invasions. *Trends in Ecology & Evolution*,
535 **22**, 465–471.

536 Spitze, K. (1993). Population structure in *Daphnia obtusa*: Quantitative genetic and allozymic
537 variation. *Genetics*, **135**, 367–374.

538 Taylor, S.A., Curry, R.L., White, T.A., Ferretti, V. & Lovette, I. (2014). Spatiotemporally
539 consistent genomic signatures of reproductive isolation in a moving hybrid zone. *Evolution*,
540 **68**, 3066–3081.

541 Thibaudon, M., Šikoparija, B., Oliver, G. & Smith, M. (2013). Ragweed pollen source inventory
542 for France - the second largest centre of *Ambrosia* in Europe. *Atmospheric Environment*.

543 van Boheemen, L.A., Lombaert, E., Nurkowski, K.A., Gauffre, B., Rieseberg, L.H. & Hodgins,
544 K.A. (2017). Multiple introductions, admixture and bridgehead invasion characterize the
545 introduction history of *Ambrosia artemisiifolia* in Europe and Australia. *Molecular Ecology*,
546 **26**, 5421–5434.

547 van Boheemen, L.A., D.Z. Atwater, and K.A. Hodgins. 2019. Rapid and repeated local
548 adaptation to climate in an invasive plant. *New Phytologist* **222**: 614-627.

549 Vasemägi, A. (2006). The adaptive hypothesis of clinal variation revisited: single-locus clines as
550 a result of spatially restricted gene flow. *Genetics*, **173**, 2411–2414.

551 Whitlock, M.C.C. (2008). Evolutionary inference from Q_{ST} . *Molecular Ecology*, **17**, 1885–1896.

552

553 **Table 1. Result of regressions of *Ambrosia artemisiifolia* traits on latitude** in the invasive
554 (European) and native (North American) ranges. *** Indicates P<0.01.

555

	Flowering time	Height	Total branches	Male fitness	Female fitness
EUROPE					
Estimate	-1.545***	-0.672	-0.120	-9.523	-0.578
Std. error	(0.319)	(0.578)	(0.180)	(7.236)	(0.387)
Degrees of freedom	231	228	227	228	194
NORTH AMERICA					
Estimate	-4.694***	-5.570***	-0.333***	-10.557	-2.243***
Std. error	(0.262)	(0.495)	(0.129)	(6.776)	(0.339)
Degrees of freedom	287	285	284	285	262

556

Figures.

Figure 1. Map of population *Ambrosia artemisiifolia* collection sites from (A) Native North American range (B) Invasive European range. Collection sites used only for the common garden are shown in blue. Sites used for both the common garden and SNP study are shown in red.

Figure 2. Phenotypic results for ragweed plants grown in common garden. (A) Flowering time (B) Height (C) Total branches on the y axes, the latitudes of origin on the x-axis.

Figure 3. Female and male reproductive output results for ragweed plants grown in common garden. (A) Estimated male fitness (B) Estimated female fitness.

590 **Figure 4. STRUCTURE results for ragweed populations (A) Europe (B) North America.**

591 Populations are ordered from lowest to highest latitudes (south to north).

592 Appendices

593 **Table A1.** Population coordinates for 18 European (invasive) ragweed populations.

594

Population Abbreviation	Continent	Latitude	Longitude
PG	Europe	43.9475	4.536631
RM	Europe	44.74194	4.940783
LH	Europe	45.66117	4.966539
HK	Europe	46.89164819	19.65085249
HR	Europe	47.13845	19.47464
HM	Europe	47.62666192	19.44151603
HJ	Europe	47.70273145	20.07493272
HB	Europe	47.8226962	19.089972
HC	Europe	47.962356	20.894481
HS	Europe	48.18515189	20.92252985
SH	Europe	48.21292953	18.93131109
SZ	Europe	48.74195476	19.13689272
SB	Europe	49.13566482	21.55096851
SL	Europe	49.222985	19.302752
PV	Europe	49.8798343	19.89245121
PB	Europe	50.12194	18.81494
PH	Europe	50.22392429	19.17026453
PE	Europe	51.35332297	19.60609006

595

596 **Table A2.** Population coordinates for 20 North American (native) ragweed populations.

597

Population Abbreviation	Continent	Latitude	Longitude
OM	North American	37.77932728	-75.61065442
VN	North American	37.92026101	-75.47660198
MF	North American	38.40708263	-75.56824697
DS	North American	38.64183334	-75.445
SC	North American	38.68109706	-75.07450333
HL	North American	38.92013797	-75.44832488
DH	North American	39.32390625	-75.62242479
BP	North American	39.64569554	-75.71813984
WV	North American	39.6584	-79.90525
PA	North American	39.930872	-75.583156
AV	North American	40.20018743	-76.76244026
PR	North American	40.65946579	-76.91820217
SU	North American	41.45751046	-77.13565654
IH	North American	42.44996	-76.46113
CB	North American	43.63519	-79.34576
WC	North American	44.003197	-79.393311
MI	North American	44.37683	-79.74113
FL	North American	44.750014	-79.704694
SD	North American	45.225364	-79.681836
PS	North American	45.3347	-79.956231

598 **Table A3.** Pairwise F_{st} values for North American ragweed populations

	VN	MF	HL	BP	WV	PA	AV	PR	SU	CB	MI	FL	SD	PS	DH	IH	WC
VN	--	0.061	0.101	0.18	0.078	0.082	0.18	0.089	0.093	0.075	0.073	0.061	0.13	0.085	0.086	0.066	0.06
MF	0.061	--	0.033	0.118	0.02	0.018	0.116	0.031	0.029	0.017	0.018	0.022	0.049	0.027	0.02	0.007	0.01
HL	0.101	0.033	--	0.154	0.051	0.052	0.149	0.059	0.061	0.049	0.05	0.054	0.092	0.057	0.053	0.04	0.0
BP	0.18	0.118	0.154	--	0.131	0.131	0.213	0.137	0.145	0.129	0.139	0.13	0.188	0.145	0.139	0.117	0.1
WV	0.078	0.02	0.051	0.131	--	0.034	0.125	0.044	0.038	0.024	0.019	0.026	0.049	0.03	0.043	0.02	0
PA	0.082	0.018	0.052	0.131	0.034	--	0.125	0.043	0.043	0.033	0.032	0.035	0.068	0.044	0.042	0.021	0.0
AV	0.18	0.116	0.149	0.213	0.125	0.125	--	0.133	0.135	0.124	0.129	0.121	0.171	0.137	0.136	0.112	0.1
PR	0.089	0.031	0.059	0.137	0.044	0.043	0.133	--	0.051	0.04	0.039	0.042	0.072	0.049	0.053	0.031	0.03
SU	0.093	0.029	0.061	0.145	0.038	0.043	0.135	0.051	--	0.035	0.036	0.036	0.079	0.044	0.05	0.025	0.03
CB	0.075	0.017	0.049	0.129	0.024	0.033	0.124	0.04	0.035	--	0.009	0.009	0.046	0.02	0.043	0.015	0.00
MI	0.073	0.018	0.05	0.139	0.019	0.032	0.129	0.039	0.036	0.009	--	0.01	0.041	0.011	0.042	0.016	0.00
FL	0.061	0.022	0.054	0.13	0.026	0.035	0.121	0.042	0.036	0.009	0.01	--	0.04	0.017	0.041	0.017	0.00

SD	0.13	0.049	0.092	0.188	0.049	0.068	0.171	0.072	0.079	0.046	0.041	0.04	--	0.054	0.071	0.047	0.03
PS	0.085	0.027	0.057	0.145	0.03	0.044	0.137	0.049	0.044	0.02	0.011	0.017	0.054	--	0.048	0.025	0.01
DH	0.086	0.02	0.053	0.139	0.043	0.042	0.136	0.053	0.05	0.043	0.042	0.041	0.071	0.048	--	0.031	0.03
IH	0.066	0.007	0.04	0.117	0.02	0.021	0.112	0.031	0.025	0.015	0.016	0.017	0.047	0.025	0.031	--	0.00
WC	0.063	0.015	0.047	0.122	0.02	0.029	0.114	0.036	0.031	0.007	0.003	0.006	0.036	0.012	0.037	0.009	--

599

600 **Table A4.** Pairwise F_{st} values for European ragweed populations

	PG	LH	HK	HR	HJ	HB	HC	HS	SZ
PG	--	0.021	0.050	0.026	0.091	0.051	0.043	0.038	0.054
LH	0.021	--	0.035	0.018	0.083	0.034	0.035	0.030	0.047
HK	0.050	0.035	--	0.036	0.105	0.063	0.056	0.050	0.07
HR	0.026	0.018	0.036	--	0.080	0.037	0.039	0.028	0.046
HJ	0.091	0.083	0.105	0.080	--	0.120	0.099	0.092	0.114
HB	0.051	0.034	0.063	0.037	0.120	--	0.061	0.051	0.071
HC	0.043	0.035	0.056	0.039	0.099	0.061	--	0.046	0.063
HS	0.038	0.030	0.050	0.028	0.092	0.051	0.046	--	0.058
SZ	0.054	0.047	0.070	0.046	0.114	0.071	0.063	0.058	--

601

602 **Table A5.** Population genetic summary statistics for North American ragweed

Statistic	Value	Std.Dev.	c.i.2.5%	c.i.97.5%	Description
Num	2	0	2	2	Number of alleles
Eff_num	1.163	0.001	1.161	1.165	Effective number of alleles
Ho	0.108	0.001	0.107	0.11	Observed heterozygosity
Hs	0.141	0.001	0.14	0.142	Heterozygosity within populations
Ht	0.15	0.001	0.148	0.151	Total heterozygosity
H't	0.15	0.001	0.149	0.152	Corrected total heterozygosity
Gis	0.23	0.001	0.228	0.233	Inbreeding coefficient
Gst	0.061	0.001	0.06	0.062	Fixation index
G'st(Nei)	0.064	0.001	0.063	0.065	Nei, corrected fixation index
G'st(Hed)	0.071	0.001	0.07	0.072	Hedrick, standardised fixation index
G"st	0.075	0.001	0.073	0.076	Corrected standardised fixation index
D_est	0.011	0	0.011	0.011	Jost, differentiation

603

604 **Table A6.** Population genetic summary statistics for European ragweed

Statistic	Value	Std.Dev.	c.i.2.5%	c.i.97.5%	Description
Num	2	0	2	2	Number of alleles
Eff_num	1.224	0.001	1.221	1.226	Effective number of alleles
Ho	0.137	0.001	0.136	0.138	Observed heterozygosity
Hs	0.192	0.001	0.19	0.193	Heterozygosity within populations
Ht	0.202	0.001	0.2	0.204	Total heterozygosity
H't	0.203	0.001	0.201	0.205	Corrected total heterozygosity
Gis	0.285	0.002	0.281	0.289	Inbreeding coefficient
Gst	0.051	0.001	0.05	0.052	Fixation index
G'st(Nei)	0.057	0.001	0.055	0.059	Nei, corrected fixation index
G'st(Hed)	0.065	0.001	0.063	0.066	Hedrick, standardised fixation index
G"st	0.071	0.001	0.069	0.073	Corrected standardised fixation index
D_est	0.014	0	0.014	0.015	Jost, differentiation

605

606 **Table A7.** AMOVA output for North American and European ragweed

Source of Variation	Nested in	%var	F-stat	F-value	Std.Dev.	c.i.2.5%	c.i.97.5%	P-value	F'-value
Within Individual	--	0.795	F _{it}	0.205	0.001	0.204	0.207	--	--
Among Individual	Population	0.162	F _{is}	0.169	0.001	0.167	0.171	0.001	--
Among Population	Continent	0.042	F _{sc}	0.042	0	0.041	0.043	0.001	0.048
Among Continents	--	0.002	F _{ct}	0.002	0	0.001	0.002	0.017	0.002

607

Figure A1. Splitsree results for invasive and native ragweed populations

Figure A2. STRUCTURE results for all ragweed populations. Populations are ordered first by continent and then from lowest to highest latitudes within continents.