***Agrobacterium*-mediated transient transformation of sorghum leaves for accelerating functional genomics and genome editing studies.**

Rita Sharma^{1,2,3#}, Yan Liang^{1,2#}, Mi Yeon Lee^{1,2}, Venkataramana R. Pidatala^{1,2}, Jenny C. Mortimer^{1,2*} and Henrik V. Scheller^{1,2,4,*}

[#]Equal Contribution

^{*}Corresponding Authors

¹Joint BioEnergy Institute, Emeryville, CA 94608, USA; ²Environmental Genomics and Systems Biology Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720; ³Crop Genetics and Informatics Group, School of Computational & Integrative Sciences, Jawaharlal Nehru University, New Delhi 110067, India; ⁴Department of Plant and Microbial Biology, University of California Berkeley, Berkeley, CA 94720, USA.

ABSTRACT

Objectives

Sorghum is one of the most recalcitrant species for transformation. Considering the time and effort required for stable transformation in sorghum, establishing a transient system to screen the efficiency and full functionality of vector constructs is highly desirable.

Results

Here, we report an *Agrobacterium*-mediated transient transformation assay with intact sorghum leaves using green fluorescent protein as marker. It also provides a good monocot alternative to tobacco and protoplast assays with a direct, native and more reliable system for testing single guide RNA (sgRNA) expression construct efficiency. Given the simplicity and ease of transformation, high reproducibility, and ability to test large constructs, this method can be widely adopted to speed up functional genomic and genome editing studies.

Keywords: *Agrobacterium*, CRISPR, sgRNA, sorghum, transformation, transient

INTRODUCTION

Sorghum is a gluten-free C4 crop, important as both a human dietary staple and animal feed, but more recently also as a potential feedstock for biofuel production[1]. With high collinearity and synteny with other grass genomes, sorghum also provides an ideal template to serve as model for other grasses[2]. However, realizing the full potential of sorghum as feedstock requires bioengineering efforts aimed at tailoring sorghum biomass for biorefining applications [3, 4]. Indeed, while the sorghum genome sequence was completed a decade ago [2], only a handful of genes have been characterized using transgenic approaches.

A major factor in the lack of progress is the low efficiency and time-consuming nature of stable transformation. Indeed, sorghum is one of the most recalcitrant crops to transformation and regeneration. The first sorghum transgenic plants were generated using particle bombardment in 1993 with only 0.28% transformation rate[5]. Subsequently, Zhao and coworkers[6] reported 2.12% transformation rate using *Agrobacterium*-mediated transformation. Although with recent advancements in technology and optimization of regeneration protocols, several labs have been able to now transform a few limited sorghum cultivars with improved efficiency; reproducibility and consistency still remain major issues[7-9].

When developing engineered plants, due to the time and cost involved, it is highly desirable to test construct functionality in a transient assay. This is particularly true for sorghum. Transient assays in grasses mostly rely on protoplasts[12-14]. However, expression of a gene in protoplasts may not always mimic *in planta* native state and, also experience inconsistent efficiency due to variability in quality of protoplasts and size of vector transformed[15]. Here, we have established a simplified transient assay with *Agrobacterium*, also known as agroinfiltration, for transient transformation of sorghum and demonstrated its application by confirming gene editing in sorghum leaves using GFP as a marker. Using our

method, researchers can directly test the *in planta* efficacy of binary constructs that may subsequently be used for stable transformation.

MAIN TEXT

METHODS

Plasmids and bacterial strains

Binary vectors C282 and C283 were built based on pTKan-p35S-attR1-GW-attR2 backbone vector[16] using Gateway (Invitrogen, CA, U.S.A.) to introduce codons for GFP (C282) or frame-shifted (fs)GFP (C283) for expression under the CaMV 35S promoter. The fsGFP has a 23 bp positive target control (PTC) sequence inserted after the start codon (5'-gcgcttcaaggtgcacatggagg-3')[21]. C286 contains GFP driven by maize Ubiquitin 1 promoter, described elsewhere[17, 18]. Binary vectors C475 and C476 were built based on pTKan-pNOS-DsRed-pZmUBQ1-attR1-GW-attR2 backbone vector[17]. The C476 cassette (pTKan-pNOS-DsRed-tNOS-pZmUBQ1-CAS9p-pOsU3-PTC_gRNA-p35S-fsGFP) contains a sgRNA (5'-gcgcttcaaggtgcacatgg-3') targeting the PTC sequence in fsGFP. CAS9p is a plant codon optimized CAS9 from *Streptococcus pyogenes*[19]. The C475 cassette (pTKan-pNOS-DsRed-tNOS-pZmUBQ1-CAS9p-pOsU3-nongRNA-p35S-fsGFP) lacking a sgRNA targeting sequence was used as a negative control. Plasmids are available from the JBEI registry: <https://registry.jbei.org>

Binary vectors were transformed into *Agrobacterium tumefaciens* strain GV3101 using electroporation, and grown in Luria Bertani (LB) medium containing 100/30/50 µg/mL rifampicin/gentamicin/spectinomycin at 28°C. Similarly, *A. tumefaciens* strain C58C1 containing the P19 suppressor of gene-silencing protein was grown in LB media containing 100/5/50 µg/mL rifampicin/tetracycline/kanamycin.

86

87 **Leaf infiltration**

88 For agroinfiltration, *Agrobacterium* was grown in liquid culture (5 mL, 24 h, 30°C),
 89 and cells were pelleted (5000 xg, 5 min), and resuspended in infiltration medium containing
 90 50 mM MES, pH 5.6, 2 mM Na₃PO₄, 0.5% (w/v) dextrose, 200 µM acetosyringone and 0.01%
 91 Silwet L-77 with an OD₆₀₀ of 0.5. The P19 strain was mixed with each of the other strains to
 92 ¼ of the final volume. Prior to infiltration, the *Agrobacterium* suspension was incubated
 93 without shaking at 30°C for about 2 h. The *Nicotiana benthamiana* plants were grown in a
 94 growth chamber under 16/8 h and 26/24°C day/night cycle, and plants of ~4-weeks-old used
 95 for infiltration. *Sorghum bicolor* (L.) Moench inbred line Tx430 plants were grown in a plant
 96 growth room under 14/10 h 29/26°C day/night cycle. Plants at the three-leaf stage (3-4 weeks
 97 old), were used for co-infiltration (Fig. 1). The fully expanded sorghum leaves were
 98 mechanically wounded with a syringe needle (0.8 mm X 40 mm) several times to make the
 99 epidermis more conducive to infiltration. No injury was caused to tobacco leaves. The
 100 *Agrobacterium* strains suspended in infiltration medium were infiltrated into leaves using a 1
 101 mL syringe without needle. The boundaries of regions infiltrated with *Agrobacterium* were
 102 marked with a permanent marker for later visualization.

103

104 **Microscopy**

105 About 3-4 days after infiltration (DAI), tobacco and sorghum leaves were detached
 106 from the plant and observed under a Leica D4000B fluorescence microscope coupled with a
 107 Leica DC500 camera using appropriate filters for GFP and DsRed.

108

RESULTS

Expression of GFP in infiltrated leaves of tobacco and sorghum

We tested binary constructs C282 containing 35S_{pro}::GFP and the modified plasmid C283 with 35S_{pro}::fsGFP (frame-shifted GFP) by agroinfiltration in both tobacco and sorghum leaves. At 3DAI, the GFP signal was examined in detached leaves under a fluorescent microscope. Both sorghum and tobacco leaves infiltrated with C282 showed high and consistent expression of GFP (Fig. 2). However, those infiltrated with C283, containing fsGFP, exhibited no signal. It was noted that the area of detectable GFP expression was much smaller in sorghum as compared to tobacco. This is likely due to the limited infiltration of *Agrobacterium* suspension in sorghum leaves. The signal could be observed up to 7 DAI, after which the signal declined.

Ubiquitin promoter is more effective for sorghum

We compared infiltration of plasmid C282 (35S_{pro}::GFP) with C286 (Ubq_{pro}::GFP) in sorghum. While a higher intensity of GFP signal was observed in tobacco leaves with the 35S promoter compared to sorghum leaves (Fig. 2); GFP expression driven by the maize ubiquitin1 promoter exhibited higher intensity in sorghum leaves.

Demonstration of gene editing in sorghum leaves using GFP as target gene

To test whether we can use our transient *Agrobacterium*-mediated transformation method to determine sgRNA gene editing efficiency in sorghum, we used the binary vectors, C475 and C476 for agroinfiltration. Tobacco leaves were also infiltrated as a comparison. Both C475 and C476 contained constitutively expressed DsRed under NOS promoter, fsGFP driven by 35S promoter and pUbi-driven CAS9p for CRISPR-mediated genome editing. C476 contained a sgRNA targeting the PTC sequence in fsGFP. As C475 lacked the targeting

sgRNA, GFP expression was only expected with C476 vector and only when editing occurs to correct the GFP frame shift.

Following agroinfiltration, DsRed expression could be detected in both sorghum and tobacco leaves with both the constructs, confirming successful infiltration (Fig. 3). However, GFP expression was observed only in the leaves infiltrated with C476 demonstrating successful editing in the intact leaves of both tobacco and sorghum (Fig. 3).

DISCUSSION

Plant transformation is indispensable for elucidating gene function and engineering plant genomes for improved agronomic traits. Several biological, mechanical, chemical and electrical methods of DNA delivery have been developed to facilitate plant transformation over past several decades[22, 23]. Among biological methods, the soil-borne gram-negative bacterium *A. tumefaciens* is no doubt the most popular and widely used vehicle for DNA delivery in plant cells[24]. Although monocots are outside the host range of this bacterium, *Agrobacterium*-mediated transformation is now routinely used for transforming monocot genomes as well, though with lower efficiency[25, 26]. Agroinfiltration is also routinely used in several plant species due to rapidity, versatility and convenience[27-33]. However, success of this method in monocot species is very limited primarily due to extensive epidermal cuticular wax, high silica content, and low volume of intercellular space. These morphological features prevent the infiltration of bacterial cells into grasses *via* the application of simple pressure. Although microprojectile bombardment may be used to introduce expression constructs in cereals, the set-up cost for establishing microprojectile bombardment is high. Moreover, it only targets single cells limiting the scope of screening[34], and often leads to cell damage. Earlier, Andrieu and coworkers[35] reported *Agrobacterium*-mediated transient gene expression and silencing in rice leaves by mechanically wounding leaves followed by direct incubation in

Agrobacterium suspension. However, we made several attempts to transform sorghum leaves at different stages of development, using their methodology, but could not detect any expression of GFP (data not shown).

Virus-based vectors provide an alternative opportunity for elucidating monocot gene functions. However, instability of the recombinant vector, improper orientation of insert and inconsistency due to inadequate infectivity, inoculation methods, replication/movement of virus in the host, pose serious challenges[36]. Another recent study demonstrated application of nanoparticles in transformation of wheat leaves by combining wounding treatment with syringe infiltration of the nanoparticles[37]. However, the size of plasmid that can be loaded onto nanoparticles is a major constraint due to size exclusion limit of the plant cell wall (~20 nm).

To overcome these constraints, we attempted syringe infiltration with recombinant *Agrobacterium*, containing vectors for *in planta* GFP expression, at different stages of development in sorghum leaves. As expected, strength of signal in sorghum leaves was higher with the maize ubiquitin promoter as compared to cauliflower mosaic virus 35S promoter, which is reported to perform better in dicots[38]. In our system, although infiltration medium could enter the mature leaves, GFP expression was only detected in the infiltrated younger leaves of 3-4-week-old plants. The expression of GFP seemed to localize to where bacteria were initially infiltrated through mechanical pressure. We did not observe a spread of signal in the adjacent areas, unlike that reported by Andrieu and coworkers[35] for siRNAs in rice. This observation indicated that although bacteria could enter sorghum leaf cells through the wounded regions, they could not passively diffuse to other cells without mechanical pressure in sorghum leaves. We also tried dipping the leaf in *Agrobacterium* suspension after clipping the leaf from the top, as well as wounding by needle, however *Agrobacterium* could not detectably enter the sorghum leaves without applied mechanical pressure.

Further, we demonstrated the application of our method to test the efficiency of sgRNA in genome editing constructs. CRISPR-associated Cas9 is a powerful genome editing tool for engineering plants[39]. Although the design of sgRNAs and preparation of constructs is straightforward, the accuracy and efficiency of the method relies on the choice of sgRNAs[40]. Several *in silico* prediction tools are available to predict the efficiency of sgRNAs based on the sequence features. However, predicted sgRNAs often have vastly different editing efficiencies in planta[18]. Protoplasts have been commonly used to test sgRNA efficiency. However, obtaining high quality protoplasts for genome editing needs extensive standardization, especially for plants such as sorghum. Secondly, additional cloning steps have to be performed to obtain a smaller vector for protoplast transformation. Thirdly and most importantly, the efficiency predicted in protoplasts may not correlate with the efficiency observed in intact plant tissue[40]. Therefore, screening of sgRNAs to achieve high accuracy and efficiency remains a challenge. We adopted our *Agrobacterium*-mediated transient transformation strategy to test sgRNA-mediated editing efficiency in sorghum leaves. The editing was observed in the transformed tissue within three days after infiltration, thereby providing a reliable assay for testing sgRNAs under native conditions.

We used GFP as a reporter in our study as it allows direct visualization in living tissues without being invasive or destructive and does not need any substrate. Gao and workers[41] demonstrated successful use of GFP as marker for stable transformation in sorghum, avoiding use of antibiotics or herbicides. This strategy can be easily applied in our system to quickly assess the full functionality of the vector constructs. For sgRNAs targeting endogenous genes, efficacy can be tested using RT-PCR or sequencing.

Overall, our study demonstrated that *in planta Agrobacterium*-mediated transient expression of transgenes is achievable in sorghum leaves. High reproducibility, simplicity, rapidity and feasibility to transform large constructs, which can directly be used for stable

transformation, are the key advantages of our method. Though this method can be used for subcellular localization studies and physiological assays, the ability to test sgRNA targeting efficiency should be of particular interest.

LIMITATIONS

1. The efficiency of agroinfiltration is much less compared to that observed in tobacco plants and therefore infiltration of more plants may be necessary if significant amount of materials are required for downstream analysis.
2. Since we were targeting a transgene in our editing assays, editing of an endogenous sorghum gene and confirmation of successful editing by sequencing would be an important step to confirm wide applicability of this method.

DECLARATIONS

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and material

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests

Funding

R.S. acknowledges IUSSTF-DBT for GETin fellowship. This work was funded by DOE Joint BioEnergy Institute (<http://www.jbei.org>) supported by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research through Contract DEAC0205CH11231 between Lawrence Berkeley National Laboratory and the U.S. Department of Energy.

Author's Contributions

RS, JCM and HVS conceptualized the study and designed the experiments. RS, YL, VRP and MYL performed the experiments and analyzed the data. RS drafted the manuscript. All authors read and approved the final manuscript.

Acknowledgements

Not applicable

REFERENCES

- Mathur S, Umakanth AV, Tonapi VA, Sharma R, Sharma MK: **Sweet sorghum as biofuel feedstock: recent advances and available resources**. *Biotechnol Biofuels* 2017, **10**:146.
- Paterson AH, Bowers JE, Bruggmann R, Dubchak I, Grimwood J, Gundlach H, Haberer G, Hellsten U, Mitros T, Poliakov A *et al*: **The Sorghum bicolor genome and the diversification of grasses**. *Nature* 2009, **457**(7229):551-556.
- van der Weijde T, Alvim Kamei CL, Torres AF, Vermerris W, Dolstra O, Visser RG, Trindade LM: **The potential of C4 grasses for cellulosic biofuel production**. *Front Plant Sci* 2013, **4**:107.
- Bhatia R, Gallagher JA, Gomez LD, Bosch M: **Genetic engineering of grass cell wall polysaccharides for biorefining**. *Plant Biotechnol J* 2017, **15**(9):1071-1092.
- Casas AM, Kononowicz AK, Zehr UB, Tomes DT, Axtell JD, Butler LG, Bressan RA, Hasegawa PM: **Transgenic sorghum plants via microprojectile bombardment**. *Proc Natl Acad Sci U S A* 1993, **90**(23):11212-11216.
- Zhao ZY, Cai T, Tagliani L, Miller M, Wang N, Pang H, Rudert M, Schroeder S, Hondred D, Seltzer J *et al*: **Agrobacterium-mediated sorghum transformation**. *Plant Mol Biol* 2000, **44**(6):789-798.
- Ahmed RI, Ding A, Xie M, Kong Y: **Progress in Optimization of Agrobacterium-Mediated Transformation in Sorghum (Sorghum bicolor)**. *Int J Mol Sci* 2018, **19**(10).

- 265 8. Che P, Anand A, Wu E, Sander JD, Simon MK, Zhu W, Sigmund AL, Zastrow-Hayes
266 G, Miller M, Liu D *et al*: **Developing a flexible, high-efficiency Agrobacterium-**
267 **mediated sorghum transformation system with broad application.** *Plant Biotechnol*
268 *J* 2018, **16**(7):1388-1395.
- 269 9. Nelson-Vasilchik K, Hague J, Mookkan M, Zhang ZJ, Kausch A: **Transformation of**
270 **Recalcitrant Sorghum Varieties Facilitated by Baby Boom and Wuschel2.** *Curr*
271 *Protoc Plant Biol* 2018, **3**(4):e20076.
- 272 10. Girijashankar V, Swathisree V: **Genetic transformation of Sorghum bicolor.** *Physiol*
273 *Mol Biol Plants* 2009, **15**(4):287-302.
- 274 11. Jeoung JM, Krishnaveni S, Muthukrishnan S, Trick HN, Liang GH: **Optimization of**
275 **sorghum transformation parameters using genes for green fluorescent protein and**
276 **beta-glucuronidase as visual markers.** *Hereditas* 2002, **137**(1):20-28.
- 277 12. Burris KP, Dlugosz EM, Collins AG, Stewart CN, Jr., Lenaghan SC: **Development of**
278 **a rapid, low-cost protoplast transfection system for switchgrass (Panicum**
279 **virgatum L.).** *Plant Cell Rep* 2016, **35**(3):693-704.
- 280 13. Lin CS, Hsu CT, Yang LH, Lee LY, Fu JY, Cheng QW, Wu FH, Hsiao HC, Zhang Y,
281 Zhang R *et al*: **Application of protoplast technology to CRISPR/Cas9 mutagenesis:**
282 **from single-cell mutation detection to mutant plant regeneration.** *Plant Biotechnol*
283 *J* 2018, **16**(7):1295-1310.
- 284 14. Zhang Y, Su J, Duan S, Ao Y, Dai J, Liu J, Wang P, Li Y, Liu B, Feng D *et al*: **A highly**
285 **efficient rice green tissue protoplast system for transient gene expression and**
286 **studying light/chloroplast-related processes.** *Plant Methods* 2011, **7**(1):30.
- 287 15. Rehman L, Su X, Guo H, Qi X, Cheng H: **Protoplast transformation as a potential**
288 **platform for exploring gene function in Verticillium dahliae.** *BMC Biotechnol* 2016,
289 **16**(1):57.
- 290 16. Gonzalez TL, Liang Y, Nguyen BN, Staskawicz BJ, Loque D, Hammond MC: **Tight**
291 **regulation of plant immune responses by combining promoter and suicide exon**
292 **elements.** *Nucleic Acids Res* 2015, **43**(14):7152-7161.
- 293 17. Liang Y, Richardson S, Yan J, Benites VT, Cheng-Yue C, Tran T, Mortimer J,
294 Mukhopadhyay A, Keasling JD, Scheller HV *et al*: **Endoribonuclease-Based Two-**
295 **Component Repressor Systems for Tight Gene Expression Control in Plants.** *ACS*
296 *Synth Biol* 2017, **6**(5):806-816.
- 297 18. Liang Y, Eudes A, Yogiswara S, Jing B, Benites VT, Yamanaka R, Cheng-Yue C,
298 Baidoo EE, Mortimer JC, Scheller HV *et al*: **A screening method to identify efficient**
299 **sgRNAs in Arabidopsis, used in conjunction with cell-specific lignin reduction.**
300 *Biotechnol Biofuels* 2019, **12**:130.
- 301 19. Ma X, Zhang Q, Zhu Q, Liu W, Chen Y, Qiu R, Wang B, Yang Z, Li H, Lin Y *et al*: **A**
302 **robust CRISPR/Cas9 system for convenient, high-efficiency multiplex genome**
303 **editing in monocot and dicot Plants.** *Molecular plant* 2015, **8**(8):1274-1284.
- 304 20. Sparkes IA, Runions J, Kearns A, Hawes C: **Rapid, transient expression of**
305 **fluorescent fusion proteins in tobacco plants and generation of stably transformed**
306 **plants.** *Nat Protoc* 2006, **1**(4):2019-2025.
- 307 21. Jiang W, Zhou H, Bi H, Fromm M, Yang B, Weeks DP: **Demonstration of**
308 **CRISPR/Cas9/sgRNA-mediated targeted gene modification in Arabidopsis,**
309 **tobacco, sorghum and rice.** *Nucleic Acids Res* 2013.
- 310 22. Altpeter F, Springer NM, Bartley LE, Blechl AE, Brutnell TP, Citovsky V, Conrad LJ,
311 Gelvin SB, Jackson DP, Kausch AP *et al*: **Advancing Crop Transformation in the**
312 **Era of Genome Editing.** *Plant Cell* 2016, **28**(7):1510-1520.
- 313 23. Barampuram S, Zhang ZJ: **Recent advances in plant transformation.** *Methods Mol*
314 *Biol* 2011, **701**:1-35.

24. Gelvin SB: **Agrobacterium-mediated plant transformation: the biology behind the "gene-jockeying" tool.** *Microbiol Mol Biol Rev* 2003, **67**(1):16-37, table of contents.
25. Wu E, Zhao ZY: **Agrobacterium-Mediated Sorghum Transformation.** *Methods Mol Biol* 2017, **1669**:355-364.
26. Hiei Y, Ishida Y, Komari T: **Progress of cereal transformation technology mediated by Agrobacterium tumefaciens.** *Front Plant Sci* 2014, **5**:628.
27. Bhaskar PB, Venkateshwaran M, Wu L, Ane JM, Jiang J: **Agrobacterium-mediated transient gene expression and silencing: a rapid tool for functional gene assay in potato.** *PLoS One* 2009, **4**(6):e5812.
28. Circelli P, Donini M, Villani ME, Benvenuto E, Marusic C: **Efficient Agrobacterium-based transient expression system for the production of biopharmaceuticals in plants.** *Bioeng Bugs* 2010, **1**(3):221-224.
29. Figueiredo JF, Romer P, Lahaye T, Graham JH, White FF, Jones JB: **Agrobacterium-mediated transient expression in citrus leaves: a rapid tool for gene expression and functional gene assay.** *Plant Cell Rep* 2011, **30**(7):1339-1345.
30. Kim MJ, Baek K, Park CM: **Optimization of conditions for transient Agrobacterium-mediated gene expression assays in Arabidopsis.** *Plant Cell Rep* 2009, **28**(8):1159-1167.
31. Krenek P, Samajova O, Luptovciak I, Daskocilova A, Komis G, Samaj J: **Transient plant transformation mediated by Agrobacterium tumefaciens: Principles, methods and applications.** *Biotechnol Adv* 2015, **33**(6 Pt 2):1024-1042.
32. Li JF, Nebenfuhr A: **FAST technique for Agrobacterium-mediated transient gene expression in seedlings of Arabidopsis and other plant species.** *Cold Spring Harb Protoc* 2010, **2010**(5):pdb prot5428.
33. Zheng L, Liu G, Meng X, Li Y, Wang Y: **A versatile Agrobacterium-mediated transient gene expression system for herbaceous plants and trees.** *Biochem Genet* 2012, **50**(9-10):761-769.
34. Schweizer P, Pokorny J, Schulze-Lefert P, Dudler R: **Technical advance. Double-stranded RNA interferes with gene function at the single-cell level in cereals.** *Plant J* 2000, **24**(6):895-903.
35. Andrieu A, Breitler JC, Sire C, Meynard D, Gantet P, Guiderdoni E: **An in planta, Agrobacterium-mediated transient gene expression method for inducing gene silencing in rice (Oryza sativa L.) leaves.** *Rice (N Y)* 2012, **5**(1):23.
36. Kant R, Dasgupta I: **Gene silencing approaches through virus-based vectors: speeding up functional genomics in monocots.** *Plant Mol Biol* 2019.
37. Demirer GS, Zhang H, Matos JL, Goh NS, Cunningham FJ, Sung Y, Chang R, Aditham AJ, Chio L, Cho MJ *et al*: **High aspect ratio nanomaterials enable delivery of functional genetic material without DNA integration in mature plants.** *Nat Nanotechnol* 2019.
38. Somssich M: **A Short History of the CaMV 35S Promoter.** *PeerJ PrePr* 2018, **6**(e27096v2):1-16.
39. Khatodia S, Bhatotia K, Passricha N, Khurana SM, Tuteja N: **The CRISPR/Cas Genome-Editing Tool: Application in Improvement of Crops.** *Front Plant Sci* 2016, **7**:506.
40. Doench JG, Hartenian E, Graham DB, Tothova Z, Hegde M, Smith I, Sullender M, Ebert BL, Xavier RJ, Root DE: **Rational design of highly active sgRNAs for CRISPR-Cas9-mediated gene inactivation.** *Nat Biotechnol* 2014, **32**(12):1262-1267.
41. Gao Z, Jayaraj J, Muthukrishnan S, Claflin L, Liang GH: **Efficient genetic transformation of Sorghum using a visual screening marker.** *Genome* 2005, **48**(2):321-333.

Figure Legends

Figure 1. Image of sorghum seedling depicting the stage of sorghum plant required for efficient agroinfiltration. Leaves used for syringe-mediated infiltration on abaxial side are marked by white arrows.

Figure 2. Results of agroinfiltration with *Agrobacterium* suspension in sorghum and tobacco leaves. Column A shows bright field images and column B depicts GFP expression detected using fluorescence microscope. Scale bar: 100 μ m.

Figure 3. Successful editing of GFP in tobacco and sorghum leaves using agroinfiltration. Column A presents bright field images, whereas, columns B and C present expression of GFP and DsRed, respectively. The C476 vector construct contained sgRNA required for editing, while C475 lacked the sgRNA and serves as negative control. Expression of GFP in leaves transformed with C476 demonstrates successful editing. Scale bar: 100 μ m.

Sorghum

A

B

Tobacco

A

B

35S_{pro}::fsGFP

35S_{pro}::GFP

Ubi_{pro}::GFP

A**B****C****C475** **Tobacco****C476** **C475** **Sorghum****C476**