Bacteriological Monitoring and Evaluation of Cleaning-disinfection of Computer-related Equipment in an Obstetric and Gynecology Hospital

Meiling Li^{1*}, Tingyan Lu², Hongwei Zhang³, Shufang Chen^{1*}, Xueying Mao⁴, Li Shen², Yan Lu⁵ and Shufang Leng¹

¹ Department of hospital infection management, The International Peace Maternity & Child Health Hospital of China welfare institute attached to Shanghai Jiao Tong university school of medicine, Shanghai, China

² Department of Clinical laboratory, The International Peace Maternity & Child Health Hospital of China welfare institute attached to Shanghai Jiao Tong university school of medicine, Shanghai, China

³ Department of Epidemiology, Secondary Military Medical University, Shanghai, China.

⁴ Centre for Molecular Oncology, Barts Cancer Institute, Barts and the London School of Medicine and Dentistry, Queen Mary University of London, London, UK

⁵ Department of Internal medicine, The International Peace Maternity & Child Health Hospital of China welfare institute attached to Shanghai Jiao Tong university school of medicine, Shanghai, China

Corresponding author: Meiling Li^{1*}: E-mail address: limeilinghua@163.com; Shufang Chen^{1*}:E-mail address: 18017316228@163.com

Abstract

It is already known that computer keyboards and mice in hospitals are contaminated with different kinds of bacteria. However, the mouse pad has been neglected with regard to both research and regular cleaning and disinfection in hospitals. In our study, we monitored and evaluated the bacteriology degrees of 74 computers' keyboards, mice and mouse pads from six departments. The results showed that before cleaning-disinfection, the contamination rate of the mouse pad ranked second following the keyboards. *Enterococcus Faecium* was cultured from the mouse pads. The computer-related equipment in the wards and outpatient rooms were much more contaminated than that in the operating rooms. *Acinetobacter spp.* was only isolated from the doctor's computers. After cleaning-disinfection, 4 strains of MRSA were isolated from the keyboards and the mice, one and 3 were cultured at day 3 and day 5

after cleaning-disinfection, respectively. One strain of *Pseudomonas Aeruginosa* was isolated from the mice at day 3 after cleaning-disinfection. These demonstrated that the bacterial contamination of the mouse pads must be as emphasized as that of the keyboards and mice. Furthermore, It is better to clean and disinfect the computer-related equipment(keyboards, mice, mouse pads) at least once a day.

Keywords: Contamination; Computer-related equipment; Cleaning-disinfection;

Introduction

On a global scale, hospital-acquired infections (HAIs) have become one of the most important causes of morbidity and mortality in medical institutions[1-5] and also threaten the safety of health-care providers[6]. According to a survey from the World Health Organization, there are approximately 1.7 million and 4.5 million HAI patients in USA and Europe, respectively, accounting for 37,000 and 100,000 deaths each year. Many pathogens, such as MRSA, VRE, *Acinetobacter*, *Klebsiella*, *Listeria*, *Escherichia coli*, *Mycobacterium tuberculosis*, *Pseudomonas aeruginosa* and the Noe virus, can survive on a dry object surface for several months or even a year [7-9]. Therefore, cleaning and disinfecting the high-touch object surfaces is an important measure for controlling HAIs [10].

There have been many studies emphasizing the importance of cleaning and disinfecting the computer keyboard and mouse in healthcare settings, representing an important type of high-touch object surface. One study demonstrated that 95% of keyboards in a teaching hospital had growth of one or more microorganisms, and 5% were positive for pathogens known to be associated with HAI transmission, such as *Staphylococcus aureus* and *Enterococci* [11]. Some studies showed that the keyboard or mouse was one of the most likely bacterial vehicles in the ICU and that the degree of contamination cannot be neglected [12-14]. A survey of two acute district general hospitals indicated that MRSA had been identified on computer terminals (24%), and five of the MRSA-positive terminals were from hospital A, which had a significantly higher rate of MRSA transmission than hospital B [15]. However, the mouse pads have been neglected in both research and the regular cleaning-disinfection in hospitals.

In our research, we aimed to address four issues: 1) The bacteriological characteristics of computer-related equipment, especially the mouse pad; 2) The bacteriological

characteristics of computer-related equipment in different clinical departments; 3) The bacteriological characteristics of doctor's and nurse's computer-related equipment in the wards; 4) How often we should clean and disinfect computer-related equipment.

Materials and methods

Study Object Selection

chosen between October 2014 and December 2015 ([Supplementary Table 1](#)). In the wards, 1 nurse's station computer, 1 doctor's office computer and 1 doctor's mobile computer from each obstetric and gynecology ward were selected randomly for testing. Five samples were collected from every surface, including before cleaning-disinfection, immediately after cleaning-disinfection and day 1, day 3 and day 5 after cleaning-disinfection.

Sample Method

Samples were collected from the keyboard (including the Number keys, Character keys, Enter key, Shift key, and Space bar), the mouse (except the underside) and the mouse pad (area $\geq$ 100cm²). Sterile swabs dipped with sterile saline solution or neutralizing agent were smeared and rolled evenly back and forth five times on the surfaces. The hand-contacted part of the swabs was cut off, and the rest was put into a sampling tube containing 10 ml of sterile saline solution or neutralizing agent. All samples (947 samples) were sent to the clinical laboratory immediately.

Bacteriology Identification

In a biological safety cabinet, the sampling tubes were shaken for 30s on an oscillator, 100 μ l of each sample was transferred to blood-agar culture medium plates, and the plates were cultured for 48 hours at 35°C in an incubator. Colonies were counted and identified by Gram stain, catalase test, oxidase test, plasma coagulase test, biochemical tube test or using a VITEK-2 instrument for bacterial identification. Drug-sensitive testing was only used for detecting MRSA.

Statistical analysis

SPSS 17.0 software was used for statistical analysis and used the following parameters: $\alpha = 0.05$, which may be calibrated according to the specific statistical data:

$\alpha' = 2\alpha \div [R \times (R - 1)]$, in which R was the number of sample rates that had to be compared in pairs. The contamination rate (%) was the proportion of samples with bacterial colonies >10 cfu/cm².

Results

Bacteriological Analysis before Cleaning-disinfection

The bacterial contamination of keyboards, mice and mouse pads

As shown in Table 1, there were significant differences in the contamination rate between the keyboard group and the mouse group, as well as between the mouse group and mouse pad group, from high to low was keyboards, mouse pads and mice, respectively. The potentially pathogenic bacteria cultured from the computer-related equipment was as shown in Supplementary Table 2, One isolate of *Enterococcus Faecium* was cultured from the mouse pad. *Klebsiella. Pneumoniae*, *Pseudomonas*, and *Enterobacter cloacae* were isolated from the keyboard. *Acinetobacter lwoffii* were mainly cultured from the mouse pad and keyboard.

Table 1 The contamination rates of keyboard, mouse and mouse pad.

	N	Contamination Rate (%)	Median	IQR	P-value ¹
Keyboard	74	39.1	9.0	1.8-18.3	0.000 ²
Mouse	74	12.2	2.5	1.0-6.0	0.004 ³
Mouse pad	47	34.0	5.0	2.0-16.0	0.568 ⁴

¹ P-value was calculated for the contamination rate, $\alpha' = 0.017$

² P-value was calculated between the Keyboard group and Mouse group

³ P-value was calculated between the Mouse group and Mouse pad group

⁴ P-value was calculated between the Keyboard group and Mouse pad group

The bacterial contamination of the computer-related equipment in different departments

As shown in Table 2, the computer-related equipment in the wards and outpatient rooms were much more contaminated than that in the other departments. In total, 8 isolates of *Staphylococcus aureus* were cultured, 5, 1, 1, and 1 from the wards, outpatient rooms, neonatal dept., and delivery room, respectively. *Enterobacter cloacae*, and *Pseudomonas*

were cultured from the wards. *Enterococcus Faecium* was from the neonatal dept. *Klebsiella. Pneumoniae* was isolated from the operating rooms (Supplementary Table 3). **Table 2** The contamination rate of computer-related equipment in different departments before cleaning-disinfection

	N	¹ Contamination Rate (%)	Median	IQR	P-value
Wards	72	38.9	8.5	3.0-17.5	0.000
Outpatient Room	42	33.3	5.5	2.0-14.5	0.001
Delivery Room	14	28.6	4.0	1.0-15.3	0.036
Medical Dept.	10	30.0	3.0	2.0-15.0	0.051
Neonatal Dept.	20	15.0	0.0	0.0-5.8	0.325
Operating Room ²	39	5.1	1.0	0.0-4.0	

¹ Contamination rate is the subject of P-value calculation, $\alpha' = 0.005$

² Operating Room as the control group

The bacterial contamination of the doctor's and nurse's computer-related equipment in the obstetric and gynecology wards

There was no significant difference in the contamination rate between the doctor's office/mobile computer-related equipment and the nurse's computer-related equipment in the obstetric and gynecology wards (Table 3). The species of potentially pathogenic bacteria from the doctor's computers was more than that from the nurse's computers, *Acinetobacter lwoffii* and *Acinetobacter ursingii* were isolated from the doctor's computers. One strain of *Enterobacter cloacae* was from the nurse's computers in the gynecology wards, 2 isolated of *Pseudomonas* were cultured from the obstetric wards (Supplementary Table 4-5).

Table 3 The contamination rate of computer-related equipment in obstetric and gynecology wards before cleaning-disinfection

	N	Contamination Rate (%)	Median	IQR	P-value ¹
Obstetric wards	40	42.5	9.0	5.0-17.5	
Doctor's office computers	14	50.0	11.5	4.5-19.3	0.310
Doctor's mobile computers	13	46.2	10.0	5.5-50.0	0.420
Nurse's station computers ²	13	30.8	7.0	2.0-12.5	
Gynecology wards	27	40.7	6.0	2.0-22.0	

Doctor's office computers	10	70.0	20.5	5.5-32.3	0.070
Doctor's mobile computers	8	25.0	2.5	2.0-11.8	1.000
Nurse's station computers ²	9	22.2	5.0	2.0-12.0	

¹ calculated for the contamination rate, $\alpha = 0.05$

² control group

Bacteriological Analysis after Cleaning-disinfection

As shown in Table 4, at day 1 and day 3 after cleaning-disinfection, the contamination rates of the computer-related equipment gradually increased, and the contamination rate of mouse pads ranked the second following the keyboards. 4 strains of MRSA were isolated from the keyboards and the mice, one and 3 were cultured at day 3 and day 5 after cleaning-disinfection, respectively. Furthermore, the strain of *Staphylococcus aureus* gradually increased (7, 8 and 10 strains). 11, 7, and 7 were isolated from the keyboards, and mice and mouse pads, respectively. One strain of *Pseudomonas aeruginosa* was isolated from the mice at day 3 after cleaning-disinfection (Supplementary Table 6).

Table 4 The contamination rate of computer-related equipment after cleaning-disinfection.

	N	Cleaning-disinfection Rate	Contamination Rate (%) ¹ BCD	Contamination Rate (%) ² ACD Day 1	Contamination Rate (%) ² ACD Day 3	Contamination Rate (%) ² ACD Day 5
Keyboard	74	98.5	39.1	46.6	54.8	54.7
Mouse	74	97.9	12.2	17.8	20.6	18.8
Mouse pad	47	99.3	34.0	36.2	37.2	27.9

¹ BCD: Before Cleaning-disinfection

² ACD: After Cleaning-disinfection

Discussion

Cleaning and disinfecting object in the hospital is significantly important for controlling hospital associated infections [16-19]. In this study, we found that the contamination rate of mouse pads ranked second following the keyboards (34.0% vs 39.1%). The mouse pad is one of the high-touch objects so that it can be a “container” for pathogens. In another study was the contamination rate of the mouse pad researched, and

the results were as same as in our study [20]. The mouse pads have been relatively disregarded in the medical settings. Furthermore, the contamination rates of computer-related equipment in the wards and outpatient rooms were significantly higher than that in the operating rooms. In the gynecology wards, the contamination rate of the doctor's computer-related equipment was higher than that of nurse's computer-related equipment.

The most common bacteria cultured from the computer-related equipment was *Coagulase-negative staphylococcus*, This finding was similar to the results of William's study[21]. In total, 60 isolates of *Acinetobacter* were detected including 41 isolates of *Acinetobacter lwoffii*, 16 isolates of *Acinetobacter ursingii*, and 3 isolates of *Acinetobacter baumannii*. As an opportunistic pathogen, *A.baumannii* is one of the most clinically significant multidrug-resistant bacteria, which can cause of the nosocomial infections, especially in intensive care units [22-24]. It can persist and form biofilms on various abiotic materials in the hospital environment [24]. Contamination of ambient air with *Acinetobacter baumannii* was also a transmission way in Luis A study [25]. Despite *Acinetobacter spp.* (*A. lwoffii*, *A. ursingii*) other than *A. baumannii* were often considered relatively avirulent bacteria, they were able to be the opportunists in the presence of indwelling medical devices and caused invasive diseases [26]. A former research found that indwelling catheter-related with *A. lwoffii* bacteremia in immunocompromised hosts appeared to be associated with a low risk of mortality [27]. A bacteremia caused by *A. ursingii* in a patient with a pulmonary adenocarcinoma confirmed that it was an opportunistic human pathogen for the first time [28]. 33 strains of *Staphylococcus aureus* were detected, including 4 strains of MRSA. MRSA was previously detected from healthcare personnel computers [15, 29]. The other importantly isolated bacteria included *Enterococcus Faecalis*, *Enterococcus Faecium*, *Klebsiella. Pneumoniae*, *Enterobacter cloacae*, and *Pseudomonas Aeruginosa*.

The above isolated potentially pathogenic bacteria were also cultured from the samples of the HAI patients in our hospital, their detection rates were as shown in Table 5. the most common pathogens from HAI patients were *Enterococcus Faecalis*. this may be associated with the characteristic of maternity hospitals,with the large number of samples taken from the genital tract. *Staphylococcus aureus* and *Coagulase-negative staphylococcus* were also

the main pathogens from HAI patients. The majority of *Staphylococcus aureus* were cultured from surgical incisions, and 12 cases were MRSA positive. In the process of “patient-object-patient” pathogens transmission, hand carriage plays an important role.

Table 5 The strain and percentage of the associated pathogenic bacteria from HAI patients in 2014-2016

	Strain (%)
Enterococcus Faecalis	113 (14.9)
Staphylococcus aureus	38 (5.0)
MRSA	12 (1.6)
Coagulase-negative staphylococcus	31 (4.1)
Klebsiella. pneumoniae	26 (3.4)
Enterococcus Faecium	25 (3.3)
Enterobacter cloacae	9 (1.2)
Pseudomonas Aeruginosa	9 (1.2)
Acinetobacter baumannii	3 (0.4)
Acinetobacter lwoffii	1 (0.1)
Micrococcus	1 (0.1)

A limitation of the study was the absence of bacteria homology detection among different computer equipment or between the computer equipment and HAI patients who were infected with the same bacteria. It will be further explored in the future study.

Conclusions

In summary, we found that the cleaning and disinfection of mouse pads must be brought to attention in the hospitals. Furthermore, it was better to clean and disinfect the computer-related equipment at least 1 time/day. At the same time, health-care workers should stick to good hand hygiene.

Acknowledgements

Not applicable

References

1. Schumacher M, Allignol A, Beyersmann J, Binder N, Wolkewitz M. 2013. Hospital-acquired infections appropriate statistical treatment is urgently needed! *Int J Epidemiol* **42**:1502-1508.

- 217 2. **Allegranzi B, Bagheri Nejad S, Combescure C, et al.** 2011, Burden of endemic
218 health-care-associated infection in developing countries: systematic review and meta-analysis. *Lancet*
219 **377**:228–241.
- 220 3. **Klevens RM, Edwards JR, Richards CL, et al.** 2007. Estimating health care-associated infections
221 and deaths in U.S. hospitals, 2002. *Public Health Rep* **122**:160–166.
- 222 4. **ECDC.** Annual epidemiological report on communicable diseases in Europe, 2008: Report on the
223 State of Communicable Diseases in the EU and EEA/EFTA Countries. 2008.
- 224 5. **Yatin Mehta, Abhinav Gupta, Subhash Todi, et al.** 2014. Guidelines for prevention of hospital
225 acquired Infections. *Indian J Crit Care Med* **18**: 149-163.
- 226 6. **Flanagan ME, Welsh CA, Kiess C, et al.** 2011. A national collaborative for reducing health care
227 associated infections: current initiatives, challenges, and opportunities. *Am J Infect Control* **39**:685-689.
- 228 7. **Dancer SJ.** 2009. Mopping up hospital infection. *J Hosp Infect* **43**: 85-100.
- 229 8. **Dancer SJ.** 2008. Importance of the environment in meticillin resistant *Staphylococcus aureus*
230 acquisition: the case for hospital cleaning. *Lancet Infect Dis* **8**:101-113.
- 231 9. **Kramer A, Schwebke I, Kampf G.** 2006. How long do nosocomial pathogens persist on inanimate
232 surfaces? *BMC Infect Dis* **16**:130.
- 233 10. **Murni IK, Duke T, Kinney S, et al.** 2015. Reducing hospital-acquired infections and improving
234 the rational use of antibiotics in a developing country: an effectiveness study. *Arch Dis Child* **100**:
235 454-459.
- 236 11. **Schultz M, Gill J, Zubairi S, Huber R, Gordin F.** 2003. Bacterial Contamination of Computer
237 Keyboards in a Teaching Hospital. *Infect Control Hosp Epidemiol* **24**:302-303.
- 238 12. **Ghamdi AA, Shukri H, Yamani A, Hawsawi H, Bagatadah K, Gharawi L, Shukri N, AlEnazi**
239 **W.** 2011. Computer keyboards and mice contamination at intensive care unit in Western Region in
240 Kingdom of Saudi Arabia. *J Crit Care* **26**:38–39.
- 241 13. **Hartmann B, Benson M, Junger A, Quinzio L, Röhrig R, Fengler B, Färber UW, Wille**
242 **B, Hempelmann G.** 2004. Computer Keyboard and Mouse as a Reservoir of Pathogens in an Intensive
243 Care Unit. *J Clin Monit Comput* **18**:7-12.
- 244 14. **Moore G, Muzslay M, Wilson AP.** 2013. The Type, Level, and Distribution of Microorganisms
245 within the Ward Environment: A Zonal Analysis of an Intensive Care Unit and a Gastrointestinal
246 Surgical Ward. *Infect Control Hosp Epidemiol* **34**:500-506.

- 247 15. **CookeRP, WrightEP.**2001.Is methicillin-resistant Staphylococcus aureus (MRSA) contamination o
248 f ward-based computer terminals asurrogate marker for nosocomial MRSA transmission and handwashi
249 ng compliance? *J Hosp Infect* **48**:72-75.
- 250 16. **Dancer SJ.** 2009. The role of environmental cleaning in the control of hospital-acquired infection. *J*
251 *Hosp Infect* **73**:378-385.
- 252 17. **Pereira da Fonseca TA, Pessoa R, Felix AC, Sanabani SS.** 2016.
253 Diversity of Bacterial Communities on Four Frequently Used Surfaces ina Large Brazilian Teaching
254 Hospital. *Int J Environ Res Public Health* **13**:152.
- 255 18. **Donskey CJ.** 2013. Does improving surface cleaning and sterilization reduce health care-associated
256 infections? *Am J Infect Control* **41**:12-19.
- 257 19. **Otter JA, Yezli S, Salkeld JA, French GL.** 2013. Evidence that contaminated surfaces contribute
258 to the transmission of hospital pathogens and an overview of strategies to address contaminated surfaces
259 in hospital settings. *Am J Infect Control* **41**:6-11.
- 260 20. **Man GS, Olapoju M, Chadwick MV, Vuddamalay P, Hall AV, Edwards A, Kerr JR.** 2002.
261 Bacterial contamination of ward-based computer terminals. *J Hosp Infect* **52**:314-315.
- 262 21. **Rutala WA, White MS, Gergen MF, Weber DJ .** 2006. Bacterial Contamination of Keyboards:
263 Efficacy and Functional Impact of Disinfectants. *Infect Control Hosp Epidemiol* **27**:372-377.
- 264 22. **Munoz-Price LS, Weinstein RA.** 2008. Acinetobacter infection. *N Engl J Med* **358**:1271-1281.
- 265 23. **Atlanta, GA.** 2013. Antibiotic resistance threats in the United States. Centers for Disease Control
266 and Prevention 2013.
- 267 24. **Paweł K, Agnieszka C, Monika P, Dorota R, Jadwiga WM.** 2017. Acinetobacter baumannii isolated
268 from hospital-acquired infection: biofilm production and drug susceptibility. *APMIS* **125**: 1017 – 1026.
- 269 25. **Luis A. Shimose, Yohei Doi, Robert A. Bonomo, Dennise De Pascale, Roberto A. Viau,**
270 **Timothy Cleary, Nicholas Namias, Daniel H. Kett, L. Silvia Munoz-Price.** 2015. Contamination of
271 Ambient Air with Acinetobacter baumannii on Consecutive Inpatient Days. *J Clin Microbiol* **53**:
272 2346-2348.
- 273 26. **H. Seifert, A. Strate, A. Schulze, G. Pulverer.** 1994. Bacteremia due to Acinetobacter Species
274 Other than Acinetobacter baumannii. *Infection* **22**:379-385.
- 275 27. **Ku SC, Hsueh PR., Yang PC, Luh KT.** 2000. Clinical and Microbiological Characteristics of
276 Bacteremia Caused by Acinetobacter lwoffii. *Eur J Clin Microbiol Infect Dis* **19**:501–505.

28. Julien L, Liliana MA, Anne F, Etienne P, Gerard H, Patrick AD, Grimont AB. 2003. Bacteremia Caused by *Acinetobacter ursingii*. *J Clin Microbiol* **41**: 1337–1338.

29. Oguzkaya-Artan M, Baykan Z, Artan C, Avsarogullari L. 2015. Prevalence and risk factors for methicillin resistant *Staphylococcus aureus* carriage among emergency department workers and bacterial contamination on touch surfaces in Erciyes University Hospital, Kayseri, Turkey. *Afr Health Sci* **15**: 1289-1294.

Authors' contributions

Meiling Li designed the study, finished all experimental tests, collected and analyzed the data and wrote the manuscript. Shufang Chen made a contribution to design the study, analyze data and wrote the manuscript. Tingyan Lu guided the bacteria test and data analysis. Hongwei Zhang contributed to guide the data analysis and modify the manuscript. Xueying Mao done a great job to revise and polish the manuscript in English. Li Shen took part in doing the experimental test. Yan Lu took part in the design of this study. Shufang Leng took part in collect the data. All authors read and approved the final manuscript.

Additional information

Competing interests: The authors declared no conflict of interest in the manuscript.

Funding: This research was funded by the hospital funding (GFY5508)