1 Genomic epidemiology of a complex, multi-species plasmid-borne *bla*_{KPC}
2 carbapenemase outbreak in Enterobacterales in the UK, 2009-2014

3

4 Stoesser N^{*#1, 2}/Phan HTT^{*2,3}, Anna C. Seale^{1,4}, Zoie Aiken⁵, Stephanie Thomas⁵,
 5 Matthew Smith⁵, David Wyllie¹, Ryan George⁵, Robert Sebra⁶, Amy J Mathers^{7,8},
 6 Alison Vaughan¹, Tim EA Peto^{1,2,9}, Matthew J Ellington¹⁰, Katie L Hopkins¹⁰,
 7 Derrick W Crook^{1,2,9}, Alex Orlek^{1,2}, William Welfare¹¹, Julie Cawthorne⁵, Cheryl
 8 Lenney⁵, Andrew Dodgson^{5, 12}, Neil Woodford¹⁰, A Sarah Walker^{1,2,9}, and the
 9 TRACE Investigators' Group

10 * These authors contributed equally to this work

11 # Corresponding author

12

13 ¹Nuffield Department of Clinical Medicine, University of Oxford, Oxford, UK

14 ²NIHR Health Protection Research Unit in Healthcare Associated Infections and
 15 Antimicrobial Resistance, University of Oxford, Oxford, UK

16 ³Faculty of Medicine, University of Southampton, Southampton, UK

17 ⁴ Department of Infectious Disease Epidemiology, London School of Hygiene and
 18 Tropical Medicine, UK

19 ⁵ Manchester University NHS Foundation Trust, Manchester, UK

20 ⁶ Icahn Institute and Department of Genetics and Genomic Sciences, Icahn School of
 21 Medicine, Mount Sinai, New York, USA

22 ⁷ Division of Infectious Diseases and International Health, Department of Medicine,
 23 University of Virginia Health System, Charlottesville, Virginia, USA

24 ⁸ Clinical Microbiology Laboratory, Department of Pathology, University of Virginia
 25 Health System, Charlottesville, Virginia, USA

26 ⁹ NIHR Biomedical Research Centre, University of Oxford, Oxford, UK

27 ¹⁰ Antimicrobial Resistance and Healthcare Associated Infections (AMRHAI)

28 Reference Unit, National Infection Service, Public Health England, London, UK

29 ¹¹ Public Health England North West, Manchester, UK

30 ¹² Public Health Laboratory, Public Heath England, Manchester, UK

31

32 **Keywords:** carbapenemase, Enterobacterales, KPC, whole genome sequencing, short-

33 read sequencing, long-read sequencing, outbreak analysis

34

35 **Corresponding author:** Nicole Stoesser

36 Email: nicole.stoesser@ndm.ox.ac.uk

37 Tel: +44 (0) 1865 220856

38 ABSTRACT

39 Carbapenem resistance in Enterobacterales is a public health threat. *Klebsiella*
40 *pneumoniae* carbapenemase (encoded by alleles of the *bla*_{KPC} family) is one of the
41 commonest transmissible carbapenem resistance mechanisms worldwide. The
42 dissemination of *bla*_{KPC} has historically been associated with distinct *K. pneumoniae*
43 lineages (clonal group 258 [CG258]), a particular plasmid family (pKpQIL), and a
44 composite transposon (Tn4401). In the UK, *bla*_{KPC} has caused a large-scale, persistent
45 outbreak focused on hospitals in North-West England. This outbreak has evolved to
46 be polyclonal and poly-species, but the genetic mechanisms underpinning this
47 evolution have not been elucidated in detail; this study used short-read whole genome
48 sequencing of 604 *bla*_{KPC}-positive isolates (Illumina) and long-read assembly
49 (PacBio)/polishing (Illumina) of 21 isolates for characterisation. We observed the
50 dissemination of *bla*_{KPC} (predominantly *bla*_{KPC-2}; 573/604 [95%] isolates) across eight
51 species and more than 100 known sequence types. Although there was some variation
52 at the transposon level (mostly Tn4401a, 584/604 (97%) isolates; predominantly with
53 ATTGA-ATTGA target site duplications, 465/604 [77%] isolates), *bla*_{KPC} spread
54 appears to have been supported by highly fluid, modular exchange of larger genetic
55 segments amongst plasmid populations dominated by IncFIB (580/604 isolates),
56 IncFII (545/604 isolates) and IncR replicons (252/604 isolates). The subset of
57 reconstructed plasmid sequences also highlighted modular exchange amongst non-
58 *bla*_{KPC} and *bla*_{KPC} plasmids, and the common presence of multiple replicons within
59 *bla*_{KPC} plasmid structures (>60%). The substantial genomic plasticity observed has
60 important implications for our understanding of the epidemiology of transmissible
61 carbapenem resistance in Enterobacterales, for the implementation of adequate
62 surveillance approaches, and for control.

63 **IMPORTANCE**

64 Antimicrobial resistance is a major threat to the management of infections, and
65 resistance to carbapenems, one of the “last line” antibiotics available for managing
66 drug-resistant infections, is a significant problem. This study used large-scale whole
67 genome sequencing over a five-year period in the UK to highlight the complexity of
68 genetic structures facilitating the spread of an important carbapenem resistance gene
69 (*bla_{KPC}*) amongst a number of bacterial species that cause disease in humans. In
70 contrast to a recent pan-European study from 2012-2013(1), which demonstrated the
71 major role of spread of clonal *bla_{KPC}-Klebsiella pneumoniae* lineages in continental
72 Europe, our study highlights the substantial plasticity in genetic mechanisms
73 underpinning the dissemination of *bla_{KPC}*. This genetic flux has important
74 implications for: the surveillance of drug resistance (i.e. making surveillance more
75 difficult); detection of outbreaks and tracking hospital transmission; generalizability
76 of surveillance findings over time and for different regions; and for the
77 implementation and evaluation of control interventions.

78 INTRODUCTION

79 Antimicrobial resistance (AMR) in Enterobacterales is a critical public health threat.
80 Carbapenem resistance is of particular concern, and outbreaks involving multiple
81 species of carbapenemase-producing Enterobacterales (CPE) are increasingly
82 reported(2-5). Exchange of AMR genes, including carbapenem resistance genes,
83 happens at multiple genetic levels(6), and is often facilitated by their presence on
84 plasmids [circular DNA structures of variable size (2kb~>1Mb)], and/or other smaller
85 mobile genetic elements (MGEs) such as transposons and insertion sequences (IS),
86 that form part of the accessory genome.

87
88 Whole genome sequencing (WGS) has significantly improved our understanding of
89 infectious diseases epidemiology and is used in both community-associated and
90 nosocomial transmission analyses(7, 8). Although useful for delineating transmission
91 routes in clonal, strain-based outbreaks, standard phylogenetic approaches and
92 comparative analyses have been more difficult for outbreaks involving multiple
93 bacterial strains/species and transmissible resistance genes(6). Reconstruction of the
94 genetic structures of plasmids carrying relevant antimicrobial resistance genes using
95 long-read sequencing has improved our understanding of the genetic complexity of
96 these resistance gene outbreaks, but has been difficult to undertake on a large scale.

97
98 Although approximately 40 *Klebsiella pneumoniae* carbapenemase (KPC; encoded by
99 *bla_{KPC}*) variants have now been described (as per NCBI's AMR reference gene
100 catalogue, available at <https://www.ncbi.nlm.nih.gov/pathogens/isolates#/refgene/>),
101 only two have been most widely reported globally, namely KPC-2 and KPC-3
102 (H272Y with respect to KPC-2; single nucleotide difference in *bla_{KPC}* [C814T])(9,

103 10). In the UK, the first KPC isolate identified was a KPC-4-containing *Enterobacter*
104 sp. isolated in Scotland in 2003(11), with subsequent identification of KPC-3 in
105 isolates in the UK in 2007. From 2007, increasing numbers of suspected KPC isolates
106 were referred to Public Health England (PHE's) Antimicrobial Resistance and
107 Healthcare Associated Infections (AMRHAI) Reference Unit, with the majority of
108 confirmed KPC-producers (>95%) coming from an evolving KPC-2-associated
109 outbreak in hospitals in North-West England, first recognised in 2008(12). These
110 isolates were predominantly *bla*_{KPC}-positive Enterobacterales cultured from patients
111 in the Central Manchester University Hospitals NHS Foundation Trust (CMFT; now
112 part of Manchester University NHS Foundation Trust)(13). *bla*_{KPC} is thought to have
113 been introduced into the region via a pKpQIL-like plasmid(14, 15), a plasmid
114 backbone previously associated with the global dissemination of *bla*_{KPC} in *K.*
115 *pneumoniae* clonal group 258, and already observed in other *K. pneumoniae* sequence
116 types (STs) and species in an analysis of 44 UK KPC-Enterobacterales from 2008-
117 2010(15).

118
119 We used WGS to undertake a large-scale retrospective study of this multi-species,
120 polyclonal, *bla*_{KPC} outbreak in North-West England from 2009, generating complete
121 genome structures, including *bla*_{KPC} plasmids, for a subset of isolates. We
122 contextualised our analysis of regional outbreak strains using isolates from a national
123 *bla*_{KPC} surveillance programme, with the goal of understanding the genetic structures
124 associated with the regional emergence of *bla*_{KPC} in this setting.

125

126 RESULTS

127 Of 742 isolates identified for sequencing, 60 (8%) were not retrievable or cultivable
 128 from the laboratory archives. After de-duplicating by taking the first *bla*_{KPC}-positive
 129 Enterobacterales (KPC-E) per patient, and excluding sequencing failures, any
 130 sequences without *bla*_{KPC} (assumed lost in culture), and mixtures (identified from
 131 genomic data analysis, see Methods), 604 evaluable isolate sequences were included.
 132 These included: 327 archived isolates (54%) from inpatients in the early stages of the
 133 observed outbreak (2009-2011), of which 309 and 18 isolates were from CMFT and
 134 the University Hospital of South Manchester NHS Foundation Trust (UHSM; now
 135 part of Manchester University NHS Foundation Trust) respectively; 78 (13%) later
 136 isolates from CMFT/UHSM (2012-2014); 119 (20%) isolates from other hospitals
 137 (n=15 hospitals) in North-West England (2009-2014, excluding CMFT and UHSM,
 138 up to the first 25 consecutive KPC-E isolates per hospital); 72 (12%) isolates from
 139 UK and Republic of Ireland hospitals (n=72 locations [n=4 from the Republic of
 140 Ireland]) outside the North-West (2009-2014) (first KPC-E isolate per hospital); and 8
 141 (1%) isolates from English outpatient/primary care settings.

142
 143 Although three *bla*_{KPC} variants were observed in the 604 included isolates, *bla*_{KPC-2}
 144 dominated (n=573, 95%); *bla*_{KPC-3} [n=27, 4%] and *bla*_{KPC-4} [n=4, 1%] were also
 145 observed. Two isolates (0.3%; trace524, trace534) showed evidence of mixed
 146 populations of *bla*_{KPC-2} and *bla*_{KPC-3}. The median *bla*_{KPC} copy number estimate was 1.8
 147 (IQR: 1.6-2.1), with a maximum of 8.2. Across the three main species, *bla*_{KPC} copy
 148 numbers were higher in *K. pneumoniae* (n=525 [87%], median 1.8 [IQR: 1.6-2.1]),
 149 than *E. coli* (40 [7%]: 1.7 [1.5-1.9]) or *E. cloacae* (26 [4%], 1.6 [1.4-2.0]) (Kruskal-
 150 Wallis; p=0.0003; Fig.1A). Amongst common STs, copy number was highest in *K.*

151 *pneumoniae* ST258 (n=65 [11%], median 2.4 [IQR: 1.8-2.9]) versus other species/STs
 152 (n=531 [89%], median 1.8 [1.6, 2.0]) (Kruskal-Wallis; p=0.0001; Fig.1B, Fig.S1).
 153
 154 Other broad or extended-spectrum beta-lactamase genes were also commonly present
 155 across isolates, including: *bla*_{TEM} (n=452, all *bla*_{TEM-1}), *bla*_{OXA} (n=492; Δ *bla*_{OXA-9}
 156 [n=425], *bla*_{OXA-1} [n=138]), *bla*_{SHV} (n=497) and *bla*_{CTX-M} (n=89; *bla*_{CTX-M-15} [n=57],
 157 *bla*_{CTX-M-9} [n=28]). Aminoglycoside resistance genes were also widely prevalent: *aac*
 158 (n=243), *aph* (n=196), *ant* (n=93) and *aadA* (n=280). In terms of acquired quinolone
 159 resistance, 160 isolates contained *qnr* variants, and 137 isolates contained *aac(6')-Ib-*
 160 *cr*; no *qep* variants were seen.
 161
 162 Insertion sequences (ISs) have been shown to be key in the reshaping and
 163 streamlining of bacterial genomes, as well as exerting more subtle effects in the
 164 regulation of gene expression(16). The median number of different IS types in isolates
 165 was 15 (IQR: 13-16), with a maximum of 32. Four hundred distinct IS profiles were
 166 identified amongst the 604 isolates, with only five identical profiles shared amongst
 167 ≥ 10 isolates - these included a distinct profile seen only in national *K. pneumoniae*
 168 ST258 isolates (IS1F, IS1R, IS3000, IS6100, IS903B, ISEcl1, ISKpn1, ISKpn14,
 169 ISKpn18, ISKpn25, ISKpn26, ISKpn28, ISKpn31, ISKpn6, ISKpn7), and other unique
 170 profiles seen in small groups of *K. pneumoniae* ST588, ST11, ST321 and ST54. This
 171 highlights the significant flux of small mobile genetic elements within and between
 172 lineages in our dataset.
 173
 174 Tn4401 is a ~10kb transposon that has been the major transposable context for *bla*_{KPC}
 175 to date(17, 18). A predominant Tn4401 isoform was associated with both *bla*_{KPC-2} and

176 *bla*_{KPC-3} in this study, namely Tn4401a(17), which occurred in 584/604 (97%)
177 isolates. Other known variants included Tn4401b (n=7) and Tn4401d (n=3). Only
178 20/584 (3%) isolates demonstrated evidence of SNV-level variation in Tn4401a
179 (homozygous calls at 6 positions; heterozygous calls [i.e. mixed populations] at 3
180 positions). *bla*_{KPC-2}-Tn4401a (n=539 isolates) was predominantly flanked by a 5-bp
181 target site duplication (TSD) ATTGA, with 465/604 (77%; 465/539 [86%] of this
182 sub-type) isolates with this Tn4401/TSD combination (Fig.2A). In 74 other *bla*_{KPC-2}-
183 Tn4401a isolates, the Tn4401a was flanked by other target site sequence (TSS)
184 combinations, consistent with additional transposition events. Thirty-two of these
185 were TSDs (16 AATAT-AATAT, 16 AGTTG-AGTTG), which have been described
186 as more consistent with inter-plasmid transposition of Tn4401(19), and 35 were non-
187 duplicate TSS combinations (ATTGA with either ATATA, TGGTA, CTGCC,
188 AATAA, AGGAT), described as more consistent with intra-plasmid transposition.
189 Evidence of multiple TSSs around *bla*_{KPC-2}-Tn4401a within single isolates was seen
190 in 6 cases (i.e. multiple right and/or left Tn4401 TSSs); 1 case had a right TSS
191 present, but no left TSS identified.
192
193 The 604 isolates contained 91 unique combinations of plasmid Inc types (a crude
194 proxy of plasmid populations present); no isolate was replicon negative. However,
195 there were seven predominant combinations (Fig.2B) represented in 443/604 (73%)
196 isolates, and these included six major Inc types, namely IncF (FIB [found in n=580
197 isolates], FII [n=545]), FIA (n=103), IncR (n=252), ColRNAI (n=86), and IncX3
198 (n=60). For many of the plasmid families, several different reference replicon
199 sequences exist in the PlasmidFinder database, with a degree of homology amongst
200 sequences in the same family, making it difficult to establish exactly which exact sub-

201 type of replicon is present. However, restricting to 100% matches to reference
 202 replicon types for these common families, top matches included:
 203 IncFIB(pQil)_JN233705 (n=300) and IncFIB(K)_1_Kpn3_JN233704 (n=107);
 204 ColRNAI_1_DQ298019 (n=84); IncR_1_DQ449578 (n=70); and
 205 IncFII_1_pKP91_CP000966 (n=62; plasmid MLST IncFII_{K4}),
 206 IncFII(K)_1_CP000648 (n=51; plasmid MLST IncFII_{K1}) and IncFII_1_AY458016
 207 (n=19; plasmid MLST IncFII_{K2}).

208

209 Species and lineage diversity in the outbreak was substantial, with eight different
 210 species amongst sequenced isolates, and many different known STs, including: *K.*
 211 *pneumoniae* (n=525 isolates, 70 known STs), *E. coli* (n=40, 20 known STs),
 212 *Enterobacter cloacae* (n=26, 9 known STs), *Klebsiella oxytoca* (n=6, 3 known STs),
 213 *Raoultella ornithinolytica* (n=4), *Enterobacter aerogenes* (n=2), *Serratia marcescens*
 214 (n=1) and *Kluyvera ascorbata* (n=1). The most common STs were all *K. pneumoniae*,
 215 including ST258 (n=66), ST11 (n=35), ST491 (n=31), ST1162 (n=29) and ST54
 216 (n=27) (Fig.2C). Therefore, although some of the earliest sequenced isolates were
 217 KPC-*K. pneumoniae* ST258 and ST11 (both in 2009) [two major KPC strains from
 218 CG258 circulating globally and in China at the time(9, 20)] and although KPC-
 219 producing *K. pneumoniae* ST258 appears to have been one of the earliest strains
 220 observed in CMFT and UHSM, multiple diverse STs and species were subsequently
 221 rapidly recruited to the outbreak in 2010 and 2011. This was most likely by the
 222 widespread sharing of a *bla*_{KPC-2}-Tn4401a-ATTGA-ATTGA transposon within and
 223 between IncFIB, IncFII and IncR plasmid populations (Fig.2B, 2C).

224

225 *Long-read sequencing analyses*

226 In addition to short-read data, to resolve genetic structures fully we obtained long-
 227 read PacBio data for 23 isolates, chosen to maximise the *bla*_{KPC} plasmid diversity
 228 assayed and focussing on isolates collected from the two main Manchester hospitals
 229 (12 CMFT isolates, 5 UHSM; plus 2 from other hospitals in North-West England, 4
 230 from other UK locations). These included the two presumed earliest *bla*_{KPC} isolates
 231 from both CMFT and UHSM, as well as isolates sharing the same species/ST but with
 232 different plasmid replicon combinations or from North West regional versus national
 233 locations, same-species isolates with different STs, and isolates of different species.
 234 One PacBio sequencing dataset represented a clear isolate mixture (trace597 [UHSM]
 235 of *E. cloacae* ST133 and *K. pneumoniae* ST258), and for one isolate (trace457
 236 [CMFT]), there were discrepancies between the short-read and long-read sequencing
 237 datasets, suggesting a laboratory error (*E. cloacae* ST45 long-read, *E. coli* ST88
 238 short-read). These two assemblies were excluded, leaving 21 assemblies for further
 239 analysis (Table S1).

240
 241 Of the 153 contigs from these 21 assemblies, 30 were clearly chromosomal, 77
 242 plasmid, one chromosomal with an integrated plasmid, and 45 with unclear
 243 provenance (i.e. possibly phage, plasmid, or chromosomal). Overall 78/153 [51%]
 244 contigs were circularised, including 56/77 [73%] clear plasmid sequences. Thirty-one
 245 contigs (21 (68%) circularised) harboured *bla*_{KPC}, of which one (trace552, *K.*
 246 *pneumoniae* ST11) had *bla*_{KPC} integrated into the chromosome. Four isolates had
 247 multiple copies of *bla*_{KPC} (trace205 [*K. pneumoniae*, ST468; 2 copies, 1 contig],
 248 trace457 [*E. cloacae*, ST45; 5 copies [1 truncated], 2 contigs], trace75 [*K.*
 249 *pneumoniae*, ST252; 2 copies, 2 contigs], trace149 [9 copies, 9 contigs]). Trace149
 250 (*E. coli*, ST1642), which had nine copies of *bla*_{KPC}, had one copy each on: a long,

251 incomplete *bla*_{KPC} contig (203kb), seven nearly identical complete *bla*_{KPC}-containing
252 circularised sequences of size ~10kb (possibly representing circularised translocatable
253 intermediates(21)), and a short linear *bla*_{KPC} contig (~18kb).

254

255 We observed *bla*_{KPC} in multiple plasmid backgrounds (Fig.3), including a majority of
256 *bla*_{KPC} plasmids with multiple replicons (13/21 [60%] clear plasmid contigs, as
257 represented in Fig.3), particularly with IncFIB/IncFII and/or IncR, consistent with
258 replicon patterns in the isolates overall (Fig.2). For the IncFII group, for which we
259 had 17 plasmid sequences with an IncFII(K)_CP000648-like replicon (plasmidFinder
260 match; 5 *bla*_{KPC}-negative [i.e. not represented in Fig.3] and 12 *bla*_{KPC}-positive), there
261 was evidence of significant exchange and rearrangement of plasmid components
262 between both *bla*_{KPC}-positive and *bla*_{KPC}-negative plasmids, integration of IncFII_K and
263 IncR plasmids, and gene duplication events of Tn4401/*bla*_{KPC}, as well as sharing
264 between STs and species (Fig.4).

265

266 In addition to their plasticity, part of the success of these *bla*_{KPC} plasmids may also be
267 attributable to the presence of toxin-antitoxin plasmid addiction systems (*ccdA/ccdB*
268 *n*=4 *bla*_{KPC} plasmids; *higA* *n*=6; *vapB/vapC* *n*=11); anti-restriction mechanisms (*klcA*
269 *n*=16, previously shown to promote *bla*_{KPC} dissemination(22)); and heavy metal
270 resistance (*terB* [tellurite] *n*=3; *ars* operon [arsenicals] *n*=3; chromate resistance *n*=1;
271 *cop* operon/*pcoC/pcoE* [copper] *n*=7; *mer* operon [mercury] *n*=10).

272

273 *bla*_{KPC} plasmid typing

274 Attempts to identify complete plasmids (as opposed to plasmid replicon typing) from
275 short-read data by comparison to a reference plasmid database has been estimated as

276 being correct in only ~45%-85% of cases in previous studies(6, 23). However, 13/14
 277 (93%) of isolates for which we had hybrid assemblies with only one completely
 278 reconstructed *bla*_{KPC} plasmid had the correct top match using our *bla*_{KPC} plasmid
 279 typing method (Table S2). Noting that any complete plasmid typing approach from
 280 short-read data is sub-optimal, we compared all short-read sequences with our
 281 reference *bla*_{KPC} plasmid database (see Methods); matches to one or more reference
 282 *bla*_{KPC} plasmid sequences were identified in 554/604 (92%) isolates. Filtering the
 283 single match with the highest score at the predefined ≥ 0.80 threshold left a subset of
 284 428/554 (77%) for evaluation. These 428 isolates had matches to 12 *bla*_{KPC} plasmid
 285 clusters.

286

287 Based on classification by these top plasmid-cluster matches, *bla*_{KPC} plasmid clusters
 288 were shared across a median (IQR) of 3 (1-6) STs, with pKpQIL-like plasmids being
 289 most widespread across species/STs (7 species, 75 STs), and clearly playing a major
 290 role in the North-West England outbreak, as well as being spread regionally and
 291 nationally (Fig.5). Other plasmid types identified as top-matches across the entire
 292 dataset included those fully resolved by long-read sequencing performed within this
 293 study, some of which were seen in $\geq 5\%$ of study isolates (e.g. pKPC-trace75 [a non-
 294 typeable replicon]), and in non-North-West settings, likely reflecting recombination
 295 and generation of new *bla*_{KPC} plasmid variants in North-West England and their
 296 subsequent dissemination.

297

298 **DISCUSSION**

299 We present the largest WGS-based analysis of *bla*_{KPC}-positive isolates (n=604) to our
 300 knowledge, focused on assessing genetic diversity around the carbapenemase gene

301 itself rather than limiting the analysis based on species type, and incorporating a
302 sampling frame from UK regional and national collections, over five years. *bla*_{KPC}
303 remains one of the three most common carbapenemases observed in the UK,
304 accounting for ~15% of cases referred to the AMRHAI Reference Unit in 2017(24),
305 and presenting a significant challenge to hospitals in North-West England, including
306 Manchester, where it accounted for >97% of carbapenem resistance through
307 2015(25).
308
309 Our study provides an interesting context in which to consider the findings of a
310 recently published pan-European survey of carbapenem non-susceptible *K.*
311 *pneumoniae* (the EuSCAPE study; 6 months, 2013-2014; 244 hospitals, 32
312 countries)(1). In EuSCAPE, 684 carbapenemase-producing *Klebsiella* spp. isolates
313 were Illumina sequenced, and similar to our study, most cases were healthcare-
314 exposed (<2% from outpatients). EuSCAPE carbapenemase-producing isolates were
315 also predominantly *bla*_{KPC} (~45%, n=311 isolates), but mostly *bla*_{KPC-3} (232/311
316 [75%] versus 27/604 [5%] in our study), and ST258/ST512 (226/311 [73%] versus
317 107/525 (20%) of *K. pneumoniae* overall in our study). Based on identifying genetic
318 “nearest-neighbours” in their data, the EuSCAPE team found 51% of *bla*_{KPC}-*K.*
319 *pneumoniae* were most closely related to another isolate from the same hospital. The
320 authors concluded that there was strong geographic structuring of strains, and that the
321 expansion of a handful of clonal lineages was predominantly responsible for the
322 spread of carbapenemases in *K. pneumoniae* in Europe, with onward nosocomial
323 transmission. Like *bla*_{KPC-3} in EuSCAPE, *bla*_{KPC-2} has also been linked with the clonal
324 expansion of ST258 in Australia(26), where 48% of 176 *K. pneumoniae* isolates
325 sequenced were *bla*_{KPC-2}-containing ST258.

326

327 However, instead of clonal expansion, in our study we found rapid dissemination of
 328 mobile backgrounds supporting *bla*_{KPC-2}, similar to observations from sequencing of
 329 other polyclonal *bla*_{KPC} outbreaks reported elsewhere, including the US(6, 27).
 330 Tn4401a, associated with high levels of *bla*_{KPC} expression(28), has been previously
 331 predominantly seen in *K. pneumoniae*, and in isolates from the US, Israel and Italy,
 332 and similarly most commonly with an ATTGA-ATTGA TSD(10). Thus our findings
 333 are consistent with the importation of the predominant *bla*_{KPC-2}-Tn4401a-ATTGA-
 334 ATTGA motif into CMFT/North-West England and subsequent horizontal spread.
 335 Notably, as in EuSCAPE, 46/72 (64%) singleton isolates we sampled from UK
 336 hospitals were also CG258, but our detailed sampling within a region reflected a very
 337 different molecular epidemiology. Although the EuSCAPE study is large and
 338 impressive, its breadth may have been limiting in understanding regional diversity -
 339 for example, the subset of *bla*_{KPC}-*K. pneumoniae* from the UK that were analysed in
 340 EuSCAPE consisted of 11 isolates submitted from six centres
 341 (https://microreact.org/project/EuSCAPE_UK). The focus was also more on analysing
 342 species-specific clonal relationships, with no analysis of other species or MGEs.

343

344 Although in our study diversification occurred at all genetic levels (Tn4401+TSSs,
 345 plasmids, plasmid populations, strains, species), there was more limited variation
 346 observed within the transposon and its flanking regions, and the spread of *bla*_{KPC}
 347 appears to have been supported by highly plastic modular exchange of larger genetic
 348 segments within a distinct plasmid population, particularly IncFIB/IncFII (found in
 349 580 and 545 of the 604 isolates respectively) and IncR replicons (252/604 isolates). A
 350 previous study, in which 11 transformed *bla*_{KPC} plasmids from the UK (2008-2010)

351 were sequenced (Roche 454/assembly, PCR+sequencing based gap closure),
 352 identified a UK variant of the pKpQIL plasmid, designated pKpQIL-UK (IncFII_{K2} by
 353 plasmid MLST), that was highly similar to pKpQIL (maximum 32 SNVs diversity),
 354 and several other IncFII_{K2} pKpQIL-like plasmids, but with novel segmental genetic
 355 rearrangements (gains/losses; pKpQIL-D1, pKpQIL-D2)(15). Our data support the
 356 importance of IncFII_{K2}-like plasmids in this *bla*_{KPC} outbreak too, but also that other
 357 IncFII_K-like plasmids (e.g. IncFII_{-K1}, -K4, -K7, -K15) and replicons (IncFIB, IncR) have
 358 also been a significant feature. In addition to their plasticity, the plasmids identified
 359 frequently harboured AMR genes other than *bla*_{KPC} which might offer a selective
 360 advantage, alongside heavy metal resistance genes, and plasmid toxin-antitoxin
 361 addiction systems. The plasticity and association of IncFII_K plasmids with resistance
 362 genes and IncFIB replicons has been supported by findings of a recent analysis of
 363 IncFII_K plasmids(29).

364
 365 The problem of accurately classifying plasmid populations from short-read data was
 366 exemplified in this analysis, and highlighted by our smaller long-read/short-read
 367 hybrid assembly-based analysis, which demonstrated significant diversity within
 368 structures assigned as similar by short-read based typing approaches. With this caveat,
 369 it was interesting that even with relatively relaxed thresholds, 29% of isolates did not
 370 have a match to our reference *bla*_{KPC} plasmid database (based on clustering of all
 371 publicly available reference sequences, as in Methods), consistent with rapid
 372 diversification in the genetic background of Tn4401/*bla*_{KPC} elements in this setting.

373
 374 Our findings demonstrated that it is also important to consider plasmids without the
 375 resistance gene of interest in a population, as these may be relevant to a wider

understanding of the transmission and evolution of smaller mobile genetic elements harbouring resistance genes (Fig.4). This was also shown to be relevant in a previous analysis of a large KPC-*E. coli* outbreak in the same setting in 2015-2016, in which a circulating *bla*_{KPC}-negative plasmid, pCAD3 (IncFIB/FII), acquired Tn4401 from a IncHI2/HI2A *bla*_{KPC}-positive plasmid, and went on to dominate within a clonal *E. coli* lineage(25). Most studies in general however tend to focus on analysing AMR plasmids of interest. Fortunately, long-read sequencing is becoming increasingly low cost and high-throughput, and hybrid assembly is able to reconstruct plasmid sequences in Enterobacterales(30, 31). New developments in large-scale comparative genomics of complete genomes, including plasmid structures, are essential for future large-scale analysis of AMR gene outbreaks.

There are several limitations to our study. The reconstructed genomes generated using long-read PacBio data remained incomplete (49% of all contigs uncircularised). Improvements in long-read technology and assembly approaches will likely overcome this(30). Our short-read and long-read datasets were generated from the same frozen stocks of isolates, but from separate sub-cultures (because we used the short-read data to inform selection for long-read sequencing); ideally they would have been generated from the same DNA extract. PacBio sequencing library preparation incorporates size selection, and this may have led to short plasmid sequences (<15kb) being lost. Our interpretation of the evolution of backgrounds supporting *bla*_{KPC} was limited by the diversity present, and the inability to capture sequential evolutionary events, even with this large study. Lastly, very limited epidemiological data linked to the isolates were available, meaning that we were unable to ascertain any epidemiological drivers which might be contributing to the enormous heterogeneity of *bla*_{KPC} transmission

401 over apparently short timeframes; the latter finding also precluded the useful
 402 application of standard phylogenetic approaches based on identifying variants core to
 403 and within species. In addition, the collection of isolates by PHE as part of regional
 404 and national surveillance was dictated by referral patterns of isolates from the
 405 hospitals surveyed, and we do not have any denominator information on cultures
 406 (either *bla*_{KPC}-positive or *bla*_{KPC}-negative) to corroborate details on the robustness of
 407 this referral process.

408

409 In conclusion, our large analysis highlights the difficulty and complexity of these
 410 outbreaks once important AMR genes have “escaped” the genetic confines of
 411 particular mobile genetic elements and bacterial species/lineages, with important
 412 implications for surveillance. These include the need to consider multiple bacterial
 413 species and plasmids as potential hosts of *bla*_{KPC}, and invest resource in sequencing
 414 approaches to adequately reconstruct genetic structures and avoid misinterpreting the
 415 molecular epidemiology. It also demonstrates that regional differences in AMR gene
 416 epidemiology may be quite marked, which may affect the generalizability of control
 417 methods. Finally, it is important to consider the wider genetic background of host
 418 strains and plasmids in understanding the evolution and dissemination of important
 419 AMR genes, as AMR gene transfer between plasmid backgrounds within bacteria
 420 may occur over short timescales, and the interaction of several plasmids (i.e. not just
 421 those harbouring the AMR gene of interest at any given time) in a population may be
 422 highly relevant to the persistence and dissemination of the AMR gene itself.

423

424 MATERIAL AND METHODS

425 *Study isolates and setting*

We sequenced archived carbapenem-resistant Enterobacterales isolates from two hospital groups in Manchester (formerly known as CMFT and UHSM), aiming to include all inpatient isolates archived in the early stages of the observed outbreak, 2009-2011, and a subset of *bla*_{KPC}-positive Enterobacterales (KPC-E) isolates sequenced as part of regional and national surveillance undertaken by Public Health England (PHE, 2009-2014). The PHE set included: (i) up to the first 25 consecutive KPC-E isolates from any hospital in North-West England (2009-2014) and referred to the PHE reference laboratory (2009-2014); (ii) the first KPC-E isolate from any other hospital in the UK and the Republic of Ireland referred to PHE (2009-2014); and, (iii) any KPC-E isolates from outpatient/primary care settings in the UK referred to PHE (2009-2014).

Ethical approval was not required as only bacterial isolates were sequenced, and their collection was part of outbreak investigation and management.

DNA extraction and sequencing

For short-read Illumina sequencing (HiSeq 2500, 150bp PE reads), DNA was extracted using Quickgene (Fujifilm, Japan), with an additional mechanical lysis step following chemical lysis (FastPrep, MP Biomedicals, USA). Sequencing libraries were constructed using the NEBNext Ultra DNA Sample Prep Master Mix Kit (NEB) with minor modifications and a custom automated protocol on a Biomek FX (Beckman). Ligation of adapters was performed using Illumina Multiplex Adapters, and ligated libraries were size-selected using Ampure magnetic beads (Agencourt). Each library was PCR-enriched with custom primers (Index primer plus dual index PCR primer) (32). Enrichment and adapter extension of each preparation was

451 obtained using 9ul of size-selected library in a 50ul PCR reaction. Reactions were
452 then purified with Ampure beads (Agencourt/Beckman) on a Biomek NXp after 10
453 cycles of amplification (as per Illumina recommendations). Final size distributions of
454 libraries were determined using a Tapestation 1DK system (Agilent/Lab901), and
455 quantified by Qubit fluorometry (Thermofisher).

456

457 For long-read sequencing (PacBio [n=28]), DNA was extracted using the Qiagen
458 Genomic tip 100/G kit (Qiagen, Netherlands). DNA extracts were initially sheared to
459 an average length of 15kb using g-tubes, as specified by the manufacturer (Covaris).
460 Sheared DNA was used in SMRTbell library preparation, as recommended by the
461 manufacturer. Quantity and quality of the SMRTbell libraries were evaluated using
462 the High Sensitivity dsDNA kit and Qubit Fluorimeter (Thermo Fisher Scientific) and
463 DNA 12000 kit on the 2100 Bioanalyzer (Agilent). To obtain the longest possible
464 SMRTbell libraries for sequencing (as recommended by the manufacturer), a further
465 size selection step was performed using the PippinHT pulsed-field gel electrophoresis
466 system (Sage Science), enriching for the SMRTbell libraries >15kb for loading onto
467 the instrument. Sequencing primer and P6 polymerase were annealed and bound to
468 the SMRTbell libraries, and each library was sequenced using a single SMRT cell on
469 the PacBio RSII sequencing system.

470

471 Sequencing data have been deposited in the NCBI (BioProject Accession:
472 PRJNA564424).

473

474 *Sequence data processing and assembly*

475 We used Kraken(33) to assign species to sequenced isolates. SPAdes(34) v3.6 was
 476 used to *de novo* assemble reads (default options; subsequent removal of contigs
 477 shorter than 500bp and assembly coverage <2X). Isolates with sequence assemblies
 478 >6.5Mb were excluded to ensure that potentially mixed sequences were not included
 479 in the analyses. MLST was derived in silico by blasting *de novo* assemblies against
 480 publicly available MLST databases for *E. coli*
 481 (<http://mlst.warwick.ac.uk/mlst/dbs/Ecoli>), *K. pneumoniae*, *E. cloacae* and *K. oxytoca*
 482 (<https://pubmlst.org/>). Isolates with mixed MLST outputs were excluded.
 483 Antimicrobial resistance (AMR) genes, plasmid replicon (Inc) types and insertion
 484 sequences (IS) were identified using resistType
 485 (https://github.com/hangphan/resistType_docker; curated AMR gene database as
 486 in(35), plasmid replicon reference sequences from PlasmidFinder(36), ISs from the
 487 ISFinder platform(37); ≥80% identity used as a threshold). *bla*_{KPC} copy number for
 488 each isolate was estimated by dividing coverage of the contig containing *bla*_{KPC} by
 489 the average coverage for the assembly. Plasmid MLST for common family types
 490 identified in short read data and for resolved genomes (i.e. based on PacBio and Pilon
 491 assemblies - see below) was confirmed by 100% sequence matches to reference
 492 alleles for families catalogued in the plasmidMLST website
 493 (<https://pubmlst.org/plasmid/>; IncA/C, IncHI1/2, IncN).
 494
 495 PacBio sequencing data were assembled using the HGAP pipeline(38), and polished
 496 with the corresponding Illumina datasets using Pilon (version 1.18, default
 497 parameters)(39). Chromosomal sequences and plasmid sequences were then manually
 498 curated where possible to create complete, closed, circular structures by using
 499 BLASTn to identify overlaps at the end of assembled contigs. Those with overlapping

ends larger than 1000bp with sequence identity >99% were considered circularised/complete, and trimmed appropriately for resolution. Complete sequences were annotated using PROKKA (version 1.11)(40); annotations were used to determine genes known to encode toxin-antitoxin systems, heavy metal resistance, and anti-restriction mechanisms.

505

506 *Tn4401* typing

Tn4401 typing was performed using TETyper(10), using the *Tn4401*, SNP and structural profile reference files included with the package (<https://github.com/aesheppard/TETyper>; Aug 2019), and a flanking length of 5bp, representative of the known target site signature sequence indicative of *Tn4401* transposition(41).

512

513 *Plasmid database for bla_{KPC} plasmid typing*

A reference *bla_{KPC}* plasmid sequence database composed of *bla_{KPC}*-harbouring contigs/plasmids from long-read sequencing of isolates in this study and all complete *bla_{KPC}* plasmids from (42-44) (August 2018) was used for *bla_{KPC}* plasmid typing within this study. To construct this database, all 279/6018 evaluable plasmid sequences carrying *bla_{KPC}* were first compared using *dnadiff*(45) to obtain the pairwise similarity between any two plasmid sequences p_i and p_j . The similarity was defined as a function of their lengths l_i , l_j , and the aligned bases l_{ij} , l_{ji} as reported by:

$$(p_i, p_j) = \frac{1}{2} \left(\frac{l_{ij}}{l_i} + \frac{l_{ji}}{l_j} \right) \times \min \left(\frac{l_i}{l_j}, \frac{l_j}{l_i} \right)$$

521

The score was designed to penalise differences in length of the compared sequences, i.e. to make sequences of different lengths proportionately more different. The

523

524 resulting similarity matrix was used to perform clustering of plasmid sequences using
 525 the affinity propagation clustering technique, suitable for graph clustering problems
 526 with sparse similarity matrix and uneven cluster size and cluster number(46), and
 527 resulted in 34 clusters of 1-43 plasmids per cluster (Table S3). The largest cluster was
 528 the set of pKpQIL-like plasmids comprising 43 related sequences. Representative
 529 sequences of each *bla*_{KPC} plasmid cluster in this network were chosen randomly, to
 530 generate a set (*KPC-pDB*) of plasmids ranging from 7,995bp (NC_022345.1; plasmid
 531 pAP-2) to 447,095bp (NZ_CP029436.1; plasmid pKPC_CAV2013) in the final
 532 database used for *bla*_{KPC} plasmid typing in this study.

533

534 Subsequently, *bla*_{KPC} plasmid typing for each study isolate sequence was performed
 535 as follows: (1) assembled sequences for each isolate were BLASTed (BLASTn)
 536 against *KPC-pDB*; (2) any >1kb contig with >90% nucleotide identity and >80% total
 537 coverage match to sequences in *KPC-pDB* was retained; (3) for any sequence p_i in
 538 *KPC-pDB*, a score s_i was calculated by dividing the total matched bases of all contigs
 539 matched to p_i by p_i 's length; and (4) an isolate was assumed to plausibly carry p_i if s_i
 540 ≥ 0.80 . An isolate could have several *bla*_{KPC} plasmid matches; we restricted to the top
 541 match for each isolate in our analyses.

542

543 *Statistical analysis and data visualisation*

544 Statistical analysis (Kruskal-Wallis testing) was carried out in Stata 14.2. Plots for
 545 figures 1, 2, 5 and S1 were generated using ggplot2 in R (version 1.1.463). Figure 4
 546 was generated using the GenomeDiagram package(47) in Biopython(48).

547

Acknowledgements and funding: We are grateful to and acknowledge the sharing of isolates by microbiology and clinical teams from contributing UK hospitals, and from Martin Cormican and the contributing laboratories in Dublin, Republic of Ireland. We are also grateful to the microbiology laboratory staff and infection control teams at Manchester University NHS Foundation Trust (formerly CMFT and UHSM); the staff of the Manchester Medical Microbiology Partnership; and the research laboratory, informatics and project management teams working as part of the Modernising Medical Microbiology consortium (Oxford).

Contemporaneous investigation by CMFT, UHSM and PHE was undertaken as part of routine activity. The retrospective investigation was funded by the National Institute for Health Research Health Protection Research Unit (NIHR HPRU) in Healthcare Associated Infections and Antimicrobial Resistance at Oxford University in partnership with Public Health England (PHE) [grant HPRU-2012-10041] and supported by the NIHR Biomedical Research Centre, Oxford. The views expressed in this publication are those of the authors and not necessarily those of the NHS, the National Institute for Health Research, the Department of Health or Public Health England. NS is funded by a PHE/University of Oxford Academic Clinical Lectureship. TEAP, DWC and ASW are NIHR Senior Investigators.

The Transmission of Carbapenemase-producing Enterobacteriaceae (TRACE) study investigators are listed alphabetically, and include several of the authors also listed by name in the main author list: Zoie Aiken, Oluwafemi Akinremi, Aiysha Ali, Julie Cawthorne, Paul Cleary, Derrick W. Crook, Valerie Decraene, Andrew Dodgson, Michel Doumith, Matthew J. Ellington, Ryan George, John Grimshaw, Malcolm

573 Guiver, Robert Hill, Katie L. Hopkins, Rachel Jones, Cheryl Lenney, Amy J.
 574 Mathers, Ashley McEwan, Ginny Moore, Mark Neilson, Sarah Neilson, Tim E.A.
 575 Peto, Hang T.T. Phan, Mark Regan, Anna C. Seale, Nicole Stoesser, Jay Turner-
 576 Gardner, Vicky Watts, A. Sarah Walker, Jimmy Walker, David Wyllie, William
 577 Welfare and Neil Woodford.
 578
 579 None of the authors has any conflicts of interest to declare.

580 **References**

- 581 1. David S, Reuter S, Harris SR, Glasner C, Feltwell T, Argimon S, Abudahab
 582 K, Goater R, Giani T, Errico G, Aspbury M, Sjunnebo S, Eu SWG, Group ES,
 583 Feil EJ, Rossolini GM, Aanensen DM, Grundmann H. 2019. Epidemic of
 584 carbapenem-resistant *Klebsiella pneumoniae* in Europe is driven by
 585 nosocomial spread. *Nat Microbiol* doi:10.1038/s41564-019-0492-8.
- 586 2. Mathers AJ, Stoesser N, Sheppard AE, Pankhurst L, Giess A, Yeh AJ, Didelot
 587 X, Turner SD, Sebra R, Kasarskis A, Peto T, Crook D, Sifri CD. 2015.
 588 *Klebsiella pneumoniae* Carbapenemase (KPC)-Producing *K-pneumoniae* at a
 589 Single Institution: Insights into Endemicity from Whole-Genome Sequencing.
 590 *Antimicrobial Agents and Chemotherapy* 59:1661-1668.
- 591 3. Cerqueira GC, Earl AM, Ernst CM, Grad YH, Dekker JP, Feldgarden M,
 592 Chapman SB, Reis-Cunha JL, Shea TP, Young S, Zeng Q, Delaney ML, Kim
 593 D, Peterson EM, O'Brien TF, Ferraro MJ, Hooper DC, Huang SS, Kirby JE,
 594 Onderdonk AB, Birren BW, Hung DT, Cosimi LA, Wortman JR, Murphy CI,
 595 Hanage WP. 2017. Multi-institute analysis of carbapenem resistance reveals
 596 remarkable diversity, unexplained mechanisms, and limited clonal outbreaks.
 597 *Proc Natl Acad Sci U S A* 114:1135-1140.

- 598 4. Ruiz-Garbajosa P, Curiao T, Tato M, Gijon D, Pintado V, Valverde A,
599 Baquero F, Morosini MI, Coque TM, Canton R. 2013. Multiclonal dispersal of
600 KPC genes following the emergence of non-ST258 KPC-producing *Klebsiella*
601 *pneumoniae* clones in Madrid, Spain. *J Antimicrob Chemother* 68:2487-92.
- 602 5. Martin J, Phan H, Findlay J, Stoesser N, Pankhurst L, Navickaite I, De Maio
603 N, Eyre D, Toogood G, Orsi N, Kirby A, Young N, Turton J, Hill R, Hopkins
604 K, Woodford N, Peto T, Walker A, Crook D, Wilcox M. 2017. Covert
605 dissemination of carbapenemase-producing *Klebsiella pneumoniae* (KPC) in a
606 successfully controlled outbreak: long and short-read whole-genome
607 sequencing demonstrate multiple genetic modes of transmission. *Journal of*
608 *Antimicrobial Chemotherapy*.
- 609 6. Sheppard AE, Stoesser N, Wilson DJ, Sebra R, Kasarskis A, Anson LW,
610 Giess A, Pankhurst LJ, Vaughan A, Grim CJ, Cox HL, Yeh AJ, Modernising
611 Medical Microbiology Informatics G, Sifri CD, Walker AS, Peto TE, Crook
612 DW, Mathers AJ. 2016. Nested Russian Doll-Like Genetic Mobility Drives
613 Rapid Dissemination of the Carbapenem Resistance Gene blaKPC.
614 *Antimicrob Agents Chemother* 60:3767-78.
- 615 7. Quainoo S, Coolen JPM, van Hijum S, Huynen MA, Melchers WJG, van
616 Schaik W, Wertheim HFL. 2017. Whole-Genome Sequencing of Bacterial
617 Pathogens: the Future of Nosocomial Outbreak Analysis. *Clin Microbiol Rev*
618 30:1015-1063.
- 619 8. Walker TM, Monk P, Smith EG, Peto TE. 2013. Contact investigations for
620 outbreaks of *Mycobacterium tuberculosis*: advances through whole genome
621 sequencing. *Clin Microbiol Infect* 19:796-802.

- 622 9. Munoz-Price LS, Poirel L, Bonomo RA, Schwaber MJ, Daikos GL, Cormican
623 M, Cornaglia G, Garau J, Gniadkowski M, Hayden MK, Kumarasamy K,
624 Livermore DM, Maya JJ, Nordmann P, Patel JB, Paterson DL, Pitout J,
625 Villegas MV, Wang H, Woodford N, Quinn JP. 2013. Clinical epidemiology
626 of the global expansion of *Klebsiella pneumoniae* carbapenemases. *Lancet*
627 *Infect Dis* 13:785-96.
- 628 10. Sheppard AE, Stoesser N, German-Mesner I, Vegesana K, Sarah Walker A,
629 Crook DW, Mathers AJ. 2018. TETyper: a bioinformatic pipeline for
630 classifying variation and genetic contexts of transposable elements from short-
631 read whole-genome sequencing data. *Microb Genom* 4.
- 632 11. Palepou MFW, N.; Hope, R.; Colman, M.; Glover, J.; Kaufmann, M.; Lafong,
633 C.; Reynolds, R.; Livermore, D. M. 2005. Novel class A carbapenemase,
634 KPC-4, in an *Enterobacter* isolate from Scotland, abstr abstr. 1134_01_20.
635 Prog. Abstr. 15th Eur. Cong. Clin. Microbiol. Infect. Dis., Copenhagen,
636 Denmark.
- 637 12. Public Health England. 2011. Carbapenemase-producing Enterobacteriaceae:
638 laboratory confirmed cases, 2003 to 2013.
639 [https://www.gov.uk/government/publications/carbapenemase-](https://www.gov.uk/government/publications/carbapenemase-producing-enterobacteriaceae-laboratory-confirmed-cases/carbapenemase-producing-enterobacteriaceae-laboratory-confirmed-cases-2003-to-2013)
640 [producing-enterobacteriaceae-laboratory-confirmed-](https://www.gov.uk/government/publications/carbapenemase-producing-enterobacteriaceae-laboratory-confirmed-cases/carbapenemase-producing-enterobacteriaceae-laboratory-confirmed-cases-2003-to-2013)
641 [cases/carbapenemase-producing-enterobacteriaceae-laboratory-](https://www.gov.uk/government/publications/carbapenemase-producing-enterobacteriaceae-laboratory-confirmed-cases/carbapenemase-producing-enterobacteriaceae-laboratory-confirmed-cases-2003-to-2013)
642 [confirmed-cases-2003-to-2013](https://www.gov.uk/government/publications/carbapenemase-producing-enterobacteriaceae-laboratory-confirmed-cases/carbapenemase-producing-enterobacteriaceae-laboratory-confirmed-cases-2003-to-2013). Accessed 02/09/2016.
- 643 13. Donker T, Henderson KL, Hopkins KL, Dodgson AR, Thomas S, Crook DW,
644 Peto TEA, Johnson AP, Woodford N, Walker AS, Robotham JV. 2017. The
645 relative importance of large problems far away versus small problems closer

- 646 to home: insights into limiting the spread of antimicrobial resistance in
647 England. BMC Med 15:86.
- 648 14. Findlay J, Hopkins KL, Doumith M, Meunier D, Wiuff C, Hill R, Pike R, Loy
649 R, Mustafa N, Livermore DM, Woodford N. 2016. KPC enzymes in the UK:
650 an analysis of the first 160 cases outside the North-West region. J Antimicrob
651 Chemother 71:1199-206.
- 652 15. Doumith M, Findlay J, Hirani H, Hopkins KL, Livermore DM, Dodgson A,
653 Woodford N. 2017. Major role of pKpQIL-like plasmids in the early
654 dissemination of KPC-type carbapenemases in the UK. J Antimicrob
655 Chemother 72:2241-2248.
- 656 16. Siguier P, Gourbeyre E, Chandler M. 2014. Bacterial insertion sequences:
657 their genomic impact and diversity. FEMS Microbiol Rev 38:865-91.
- 658 17. Cuzon G, Naas T, Nordmann P. 2011. Functional characterization of Tn4401,
659 a Tn3-based transposon involved in blaKPC gene mobilization. Antimicrob
660 Agents Chemother 55:5370-3.
- 661 18. Cuzon G, Naas T, Truong H, Villegas MV, Wisell KT, Carmeli Y, Gales AC,
662 Venezia SN, Quinn JP, Nordmann P. 2010. Worldwide diversity of Klebsiella
663 pneumoniae that produce beta-lactamase blaKPC-2 gene. Emerg Infect Dis
664 16:1349-56.
- 665 19. He S, Hickman AB, Varani AM, Siguier P, Chandler M, Dekker JP, Dyda F.
666 2015. Insertion Sequence IS₂₆ Reorganizes Plasmids in Clinically
667 Isolated Multidrug-Resistant Bacteria by Replicative Transposition. mBio
668 6:e00762-15.

- 669 20. Qi Y, Wei Z, Ji S, Du X, Shen P, Yu Y. 2011. ST11, the dominant clone of
670 KPC-producing *Klebsiella pneumoniae* in China. *J Antimicrob Chemother*
671 66:307-12.
- 672 21. Harmer CJ, Hall RM. 2016. IS26-Mediated Formation of Transposons
673 Carrying Antibiotic Resistance Genes. *mSphere* 1.
- 674 22. Liang W, Xie Y, Xiong W, Tang Y, Li G, Jiang X, Lu Y. 2017. Anti-
675 Restriction Protein, KlcAHS, Promotes Dissemination of Carbapenem
676 Resistance. *Front Cell Infect Microbiol* 7:150.
- 677 23. Arredondo-Alonso S, Willems RJ, van Schaik W, Schurch AC. 2017. On the
678 (im)possibility of reconstructing plasmids from whole-genome short-read
679 sequencing data. *Microb Genom* 3:e000128.
- 680 24. Public Health England. 2018. English Surveillance Programme for
681 Antimicrobial Utilisation and Resistance (ESPAUR), Report 2018.
- 682 25. Decraene V, Phan HTT, George R, Wyllie DH, Akinremi O, Aiken Z, Cleary
683 P, Dodgson A, Pankhurst L, Crook DW, Lenney C, Walker AS, Woodford N,
684 Sebra R, Fath-Ordoubadi F, Mathers AJ, Seale AC, Guiver M, McEwan A,
685 Watts V, Welfare W, Stoesser N, Cawthorne J, Group TI. 2018. A Large,
686 Refractory Nosocomial Outbreak of *Klebsiella pneumoniae* Carbapenemase-
687 Producing *Escherichia coli* Demonstrates Carbapenemase Gene Outbreaks
688 Involving Sink Sites Require Novel Approaches to Infection Control.
689 *Antimicrob Agents Chemother* 62.
- 690 26. Sherry NL, Lane CR, Kwong JC, Schultz M, Sait M, Stevens K, Ballard S,
691 Goncalves da Silva A, Seemann T, Gorrie CL, Stinear TP, Williamson DA,
692 Brett J, van Diemen A, Easton M, Howden BP. 2019. Genomics for Molecular

693 Epidemiology and Detecting Transmission of Carbapenemase-Producing
694 Enterobacterales in Victoria, Australia, 2012 to 2016. *J Clin Microbiol* 57.

695 27. Weingarten RA, Johnson RC, Conlan S, Ramsburg AM, Dekker JP, Lau AF,
696 Khil P, Odom RT, Deming C, Park M, Thomas PJ, Program NCS, Henderson
697 DK, Palmore TN, Segre JA, Frank KM. 2018. Genomic Analysis of Hospital
698 Plumbing Reveals Diverse Reservoir of Bacterial Plasmids Conferring
699 Carbapenem Resistance. *MBio* 9.

700 28. Cheruvanky A, Stoesser N, Sheppard AE, Crook DW, Hoffman PS, Weddle
701 E, Carroll J, Sifri CD, Chai W, Barry K, Ramakrishnan G, Mathers AJ. 2017.
702 Enhanced *Klebsiella pneumoniae* Carbapenemase Expression from a Novel
703 Tn4401 Deletion. *Antimicrob Agents Chemother* 61.

704 29. Bi D, Zheng J, Li JJ, Sheng ZK, Zhu X, Ou HY, Li Q, Wei Q. 2018. In Silico
705 Typing and Comparative Genomic Analysis of IncFIIK Plasmids and Insights
706 into the Evolution of Replicons, Plasmid Backbones, and Resistance
707 Determinant Profiles. *Antimicrob Agents Chemother* 62.

708 30. De Maio N, Shaw LP, Hubbard A, George S, Sanderson ND, Swann J, Wick
709 R, AbuOun M, Stubberfield E, Hoosdally SJ, Crook DW, Peto TEA, Sheppard
710 AE, Bailey MJ, Read DS, Anjum MF, Walker AS, Stoesser N, On Behalf Of
711 The Rehab C. 2019. Comparison of long-read sequencing technologies in the
712 hybrid assembly of complex bacterial genomes. *Microb Genom*
713 doi:10.1099/mgen.0.000294.

714 31. Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Completing bacterial genome
715 assemblies with multiplex MinION sequencing. *Microb Genom* 3:e000132.

716 32. Lamble S, Batty E, Attar M, Buck D, Bowden R, Lunter G, Crook D, El-
717 Fahmawi B, Piazza P. 2013. Improved workflows for high throughput library

718 preparation using the transposome-based nextera system. BMC Biotechnology
719 13:104.

720 33. Wood DE, Salzberg SL. 2014. Kraken: ultrafast metagenomic sequence
721 classification using exact alignments. *Genome Biol* 15:R46.

722 34. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS,
723 Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV,
724 Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new
725 genome assembly algorithm and its applications to single-cell sequencing. *J*
726 *Comput Biol* 19:455-77.

727 35. Stoesser N, Batty EM, Eyre DW, Morgan M, Wyllie DH, Del Ojo Elias C,
728 Johnson JR, Walker AS, Peto TE, Crook DW. 2013. Predicting antimicrobial
729 susceptibilities for *Escherichia coli* and *Klebsiella pneumoniae* isolates using
730 whole genomic sequence data. *J Antimicrob Chemother* 68:2234-44.

731 36. Carattoli A, Zankari E, Garcia-Fernandez A, Voldby Larsen M, Lund O, Villa
732 L, Moller Aarestrup F, Hasman H. 2014. In silico detection and typing of
733 plasmids using PlasmidFinder and plasmid multilocus sequence typing.
734 *Antimicrob Agents Chemother* 58:3895-903.

735 37. Siguier P, Perochon J, Lestrade L, Mahillon J, Chandler M. 2006. ISfinder: the
736 reference centre for bacterial insertion sequences. *Nucleic Acids Res* 34:D32-
737 6.

738 38. Chin CS, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A,
739 Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013.
740 Nonhybrid, finished microbial genome assemblies from long-read SMRT
741 sequencing data. *Nat Methods* 10:563-9.

- 742 39. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo
743 CA, Zeng Q, Wortman J, Young SK, Earl AM. 2014. Pilon: an integrated tool
744 for comprehensive microbial variant detection and genome assembly
745 improvement. *PLoS One* 9:e112963.
- 746 40. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation.
747 *Bioinformatics* 30:2068-9.
- 748 41. Naas T, Cuzon G, Villegas MV, Lartigue MF, Quinn JP, Nordmann P. 2008.
749 Genetic structures at the origin of acquisition of the beta-lactamase bla KPC
750 gene. *Antimicrob Agents Chemother* 52:1257-63.
- 751 42. Villa L, Feudi C, Fortini D, Brisse S, Passet V, Bonura C, Endimiani A,
752 Mammina C, Ocampo AM, Jimenez JN, Doumith M, Woodford N, Hopkins
753 K, Carattoli A. 2017. Diversity, virulence, and antimicrobial resistance of the
754 KPC-producing *Klebsiella pneumoniae* ST307 clone. *Microbial Genomics* 3.
- 755 43. Orlek A, Phan H, Sheppard AE, Doumith M, Ellington M, Peto T, Crook D,
756 Walker AS, Woodford N, Anjum MF, Stoesser N. 2017. A curated dataset of
757 complete Enterobacteriaceae plasmids compiled from the NCBI nucleotide
758 database. *Data in Brief* 12:423-426.
- 759 44. Stoesser N, Sheppard AE, Peirano G, Anson LW, Pankhurst L, Sebra R, Phan
760 HTT, Kasarskis A, Mathers AJ, Peto TEA, Bradford P, Motyl MR, Walker
761 AS, Crook DW, Pitout JD. 2017. Genomic epidemiology of global *Klebsiella*
762 *pneumoniae* carbapenemase (KPC)-producing *Escherichia coli*. *Sci Rep*
763 7:5917.
- 764 45. Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C,
765 Salzberg SL. 2004. Versatile and open software for comparing large genomes.
766 *Genome Biol* 5:R12.

- 767 46. Frey BJ, Dueck D. 2007. Clustering by passing messages between data points.
768 Science 315:972-6.
- 769 47. Pritchard L, White JA, Birch PR, Toth IK. 2006. GenomeDiagram: a python
770 package for the visualization of large-scale genomic data. Bioinformatics
771 22:616-7.
- 772 48. Cock PJ, Antao T, Chang JT, Chapman BA, Cox CJ, Dalke A, Friedberg I,
773 Hamelryck T, Kauff F, Wilczynski B, de Hoon MJ. 2009. Biopython: freely
774 available Python tools for computational molecular biology and
775 bioinformatics. Bioinformatics 25:1422-3.
776

777 **Figure legends**

778 **Figure 1.** Estimated *bla*_{KPC} copy number distributions within major species (Fig.1A),
 779 and the top nineteen commonest species/ST combinations (Fig.1B) observed within
 780 the study (other ST/species combinations assigned as “Other” or “Other species/ST”
 781 respectively). Dots represent estimated copy number for single isolates; boxplots
 782 represent median estimated *bla*_{KPC} copy number $\pm 1.58 \cdot \text{IQR} / \sqrt{n}$. Boxplots are
 783 ordered by most common species and species/ST categories, left-to-right, except for
 784 the “Other”, “Other species/ST”, assigned to the right of the plots. For species
 785 assignments, “Kpne” = *Klebsiella pneumoniae*, “Ecol” = *Escherichia coli*, and “Eclo”
 786 = *Enterobacter cloacae*.

787
 788 **Figure 2.** Incidence plots for 604 isolates included in the analysis. Dots are coloured
 789 by location of isolate collection, as defined in Methods. (A) Incidence plot of
 790 Tn4401/target site sequence (TSS; categories including ≥ 10 isolates); Tn4401a-1 is
 791 *bla*_{KPC-2}/Tn4401a, Tn4401a-2 is *bla*_{KPC-3}/Tn4401a; Tn4401a-unknown comprises a set
 792 of Tn4401a (n=18) with mutational variation including C684T, G962A
 793 C3042Y|G4739R, C4121T, G5583A and C7187T. (B) Incidence plot of plasmid
 794 replicon combinations (categories including ≥ 10 isolates). (C) Incidence plot of
 795 species/ST (top 20 common categories as in Fig.1).

796
 797 **Figure 3.** Schematic of *bla*_{KPC} plasmid types and sizes identified from long-
 798 read/short-read hybrid sequencing approach by species/ST and year of collection (NB
 799 only 21 contigs clearly designated as plasmid are represented). Closed circles denote
 800 circularised contigs (i.e. complete plasmids); circle colours denote replicon types
 801 assigned to each plasmid sequence (i.e. multiple colours represent multi-replicon

802 plasmids). Plasmids from isolates from the wider UK collection are denoted with a
803 “*”.

804

805 **Figure 4.** Alignments of plasmid sequences harbouring an IncFII(K)_1_CP000648-
806 like replicon, includes *bla*_{KPC}-negative and *bla*_{KPC}-positive sequences. All sequences
807 were re-orientated to start at IncFII for the purposes of alignment visualization (this
808 also includes incomplete sequences, for which the exact structure and order may
809 therefore be a proxy only). Loci of interest have been coloured and annotated as
810 shown. Shading between sequences denotes regions of homology, with light pink
811 shading denoting areas $\geq 90\%$ nucleotide identity, dark pink areas $\geq 50\%$ nucleotide
812 identity, and light blue areas $\geq 90\%$ nucleotide identity in reverse orientation. The
813 order of sequences is adjusted to highlight genetic overlap between sequences, but not
814 to imply any specific direct exchange events.

815

816 **Figure 5.** Short-read *bla*_{KPC} plasmid typing results (top match) for isolates by species
817 and date. Dots are coloured by location of isolate collection, as defined in Methods.

818

819 **Supplementary Tables and Figures**

820 **Figure S1.** Estimated *bla*_{KPC} copy number distributions for species/ST combinations.
821 Dots represent estimated copy number for single isolates; boxplots represent median
822 estimated *bla*_{KPC} copy number $\pm 1.58 \cdot \text{IQR} / \sqrt{n}$. For species assignments, “Eclo”
823 = *Enterobacter cloacae*, “Ecol” = *Escherichia coli*, “Ente” = *Enterobacter* spp.,
824 “Kluy” = *Kluvera* spp., “Koxy” = *Klebsiella oxytoca*, “Kpne” = *Klebsiella*
825 *pneumoniae*, “Raou” = *Raoultella ornithinolytica*, “Serr” = *Serratia marcescens*.

826 **Table S1.** Details of isolates assembled using short-read (Illumina) and long-read
827 (PacBio) datasets.

828

829 **Table S2.** Plasmid typing matches for isolates with short-read (Illumina) and long-
830 read (PacBio) assemblies and reconstructed plasmid structures.

831

832 **Table S3.** Assignment of *bla*_{KPC} plasmids in study reference database to clusters for
833 plasmid typing.

Top Tn4401 types and target site sequences within isolates

Top plasmid populations within isolates defined by Inc types

Top species and multilocus sequence types

2009–2011 cmft/uhsm 2012–2014 cmft/uhsm 2009–2014 nw region 2009–2014 other uk hospital 2009–2014 outpatient