1 **Neural Correlates of Causal Confounding**

2 Mimi Liljeholm

3 Department of Cognitive Sciences, University of California, Irvine

4

5 Corresponding author:

6 Mimi Liljeholm

7 Department of Cognitive Sciences

8 2312 Social and Behavioral Sciences Gateway

9 University of California

10 Irvine, CA, 92697-5100

11 E-mail: m.liljeholm@uci.edu

12

13 **Keywords:** Causal induction, Uncertainty, confounding, prediction errors, fMRI

14 **Abstract**

15 As scientists, we are keenly aware that if putative causes perfectly co-vary, the
 16 independent influence of neither can be discerned – a “no confounding” constraint on
 17 inference, fundamental to philosophical and statistical perspectives on causation.
 18 Intriguingly, a substantial behavioral literature suggests that naïve human reasoners,
 19 adults and children, are tacitly sensitive to causal confounding. Here, a combination
 20 of fMRI and cognitive computational modeling was used to investigate neural
 21 substrates mediating such sensitivity. While being scanned, participants observed
 22 and judged the influences of various putative causes with confounded or non-
 23 confounded, deterministic or stochastic, influences. During judgments requiring
 24 generalization of causal knowledge from a feedback-based learning context to a
 25 transfer probe, activity in the dorsomedial prefrontal cortex (DMPFC) was better
 26 accounted for by a Bayesian causal model, sensitive to both confounding and
 27 stochasticity, than a purely error-driven algorithm, sensitive only to stochasticity.
 28 Implications for the detection and estimation of distinct forms of uncertainty, and for a
 29 neural mediation of domain general constraints on causal induction, are discussed.

30

31

32

33 Introduction

34 Consider two scenarios involving a target cause, *C*, an alternative cause, *A*, and the
 35 presence (+) or absence (-) of some effect. In the first scenario, the effect occurs
 36 when *C* and *A* are presented in combination, but not when *A* is presented alone [*A*-,
 37 *AC*+]]. In the second scenario, the *A*- trials are removed, so that both *C* and *A* occur
 38 only in combination with each other [*AC*+]]. How would your judgment about the
 39 influence of *C* differ across these two scenarios? Philosophers, scientists and
 40 statisticians alike recognize that perfect co-variation of *C* and *A* in the second scenario
 41 occludes the independent influences of each putative cause, rendering any judgment
 42 about their respective effects fraught with uncertainty. Importantly, naïve human
 43 reasoners also appear to, tacitly, apply this domain-general constraint on causal
 44 induction, as evidenced by behavioral data showing sensitivity to confounding in a
 45 wide range of causal and predictive judgments, by children as well as adults
 46 (Spellman, 1996; Kushnir and Gopnik, 2005; Meder et al., 2006; Schulz and Bonawitz,
 47 2007; Schulz et al., 2007; Liljeholm, 2015). Very little is known, however, about the
 48 neural substrates mediating sensitivity to, and uncertainty associated with, causal
 49 confounding. The current study aims to identify such neural computations.

50 A second source of uncertainty in causal and predictive inference, that has been
 51 thoroughly investigated both behaviorally and neutrally, is the stochasticity, or
 52 variance, of the outcome variable, which is greatest when the probability distribution
 53 over possible outcome states is uniform. As with causal confounding, behavioral
 54 sensitivity to outcome stochasticity is well established (e.g., Holt & Laury, 2002).
 55 Moreover, several neuroimaging studies have implicated the dorsomedial prefrontal

56 cortex (DMPFC), anterior insula, thalamus and dorsolateral prefrontal cortex (DLPFC)
 57 in the neural representation of outcome stochasticity (Volz et al., 2003; Huettel et al.,
 58 2005; Grinband et al., 2006; Abler et al., 2009). An important open question is
 59 whether, and which of, these neural regions also implement signals associated with
 60 causal confounding. In the current study, participants were scanned with functional
 61 MRI while observing and judging the influences of various putative causes with
 62 confounded or non-confounded, deterministic or stochastic, influences.

63 Note that, if a particular set of confounded causes always occur in the same
 64 configuration, and the only goal is to predict the state of the outcome based on that
 65 particular configuration, then confounding does not pose a problem. Indeed, most
 66 individual, non-confounded, causes can presumably themselves be broken into a set
 67 of always co-occurring, and thus confounded, elements. The critical question, thus, is
 68 whether it is necessary to tease apart the influences of individual elements, such as
 69 when a confounded cause suddenly occurs on its own, or in a novel combination with
 70 some other cause. In the current study, participants were required both to make
 71 predictions about the outcome given a particular, recurring, configuration of
 72 confounded causes and, in other instances, to make explicit judgments about the
 73 individual influences of those same causes. Only in the latter case would uncertainty
 74 due to confounding be warranted.

75 Formally, two dominant approaches to predictive uncertainty can be discerned in the
 76 psychological literature. First, in associative learning theory, an error-driven
 77 representation of uncertainty about a cue's predictive strength is captured by the
 78 "Pearce-Hall" (PH) algorithm, which relates the "associability" of a cue to a weighted

average of the absolute prediction error on previous trials involving that cue. Since the frequency and size of prediction errors increases with unpredictability, this quantity is proportional to the stochasticity of the outcome. Conversely, in Bayesian Causal Models, uncertainty about the predictive strength of a cause is reflected in the entropy of the posterior distribution over its possible strengths, which depends on the variance of the effect variable, but also on the independent occurrence of alternative causes. While both the associative and Bayesian causal model predicts sensitivity to stochasticity, only the causal model accounts for uncertainty due to causal confounding. Here, a combination of neuroimaging and cognitive modeling was used to dissociate neural signals scaling with error-driven and causal uncertainty. In particular, judgments requiring generalization of causal knowledge from a feedback-based learning context to a transfer probe were expected to elicit strong neural responses to both stochasticity and confounding.

Materials & Methods

Participants. 20 healthy normal volunteers (mean age = 20.9 ± 2.4 , range: 18-27, 12 females) participated in the study. A power analysis performed on data from a pilot study on error-driven uncertainty, described in detail below, indicated that a sample size of 16 would yield a power of 0.9 at a Gaussian random field theory corrected threshold of 0.05 in regions of interest. One participant was excluded prior to any analyses due to excessive head movement (> 6mm), leaving a sample size of 19. The volunteers were pre-assessed to exclude those with a history of neurological or

psychiatric illness. All subjects gave informed consent and the study was approved by the Institutional Review Board of the University of California, Irvine.

Task & procedure. Participants were scanned with functional MRI while performing a causal induction task in which they assumed the role of a research scientist assessing the influence of various allergy medicines on headache – a potential side effect. At the beginning of the study, participants were instructed that each medicine could either produce headache or have no influence on headache (i.e., there were no preventive causes) and, further, that the influence of a given medicine might be stochastic, so that even if that medicine was indeed a cause of headache, it may still not produce headache every time it was administered. Three target medicines, C, D and S, mnemonically labeled here to indicate “confounded”, “deterministic” and “stochastic” influences respectively, occurred only in combination with some alternative medicine during feedback-based learning. Specifically, with +, - and $\pm$ respectively indicating a 1.0, 0 and 0.5 probability of the effect, and with lower case letters indicating non-target causes, there were 5 different types of medicine treatments: e-, s' $\pm$, De+, Se $\pm$ and Cc'+. None of the allergy patients had headache in the absence of any medicine, which was explicitly stated in the initial instructions as well as apparent on several “no medicine” (n-) trials.

Note that medicines C and c' both occurred *only* in combination with one another, so that target medicine C was perfectly confounded with its compound counterpart. Medicine D also occurred only in combination with another medicine, e, however, “elemental” medicine e also occurred by itself, allowing for an estimation of the independent causal influence of medicine D across the constant presence of e (i.e.,

123 across e- and De+ trials). Medicine S was identical to medicine D, except that the
 124 probability of headache given the Se compound was 0.5 rather than 1.0; a
 125 comparison of medicines S and D, therefore, identifies a difference between
 126 deterministic and stochastic causation. Finally, medicine s' also produced headache
 127 with a probability of 0.5 but unlike medicine S, never occurred in combination with
 128 any other medicine, so that contrasting S with s' identifies differences due to
 129 compound presentation. From a modeling perspective, an algorithm that relies
 130 solely on outcome variance to track uncertainty would only respond to medicines S
 131 and s'. In contrast, a computation that treats *both* confounding and stochasticity as
 132 sources of uncertainty should generate an increased signal to medicine C as well as
 133 S and s': These divergent predictions are respectively instantiated by the error-
 134 driven and Bayesian causal model specified in the subsequent section.

Figure 1. Task illustration. A. Prediction and outcome screens from a feedback-based learning trial. Screens were separated by a 2 sec. pause during which the word "Wait" (not shown in figure) was displayed. B. Causal query regarding the individual influence of a particular medicine on headache.

135
 136 On each trial of feedback-based learning (see Figure 1A), participants were
 137 presented with an individual allergy patient and were told either that the patient had
 138 not received any medicine or that a particular medicine, or combination of

139 medicines, had been administered. Medicines were color coded and labeled
140 accordingly (i.e., Medicine “B” was blue) with color assignments counterbalanced
141 across participants. On the first screen, the allergy patient’s state was obscured by
142 a question mark and participants were asked to indicate whether or not the patient
143 had a headache, pressing the “Y” key for “yes” and the “N” key for “no”. After a
144 prediction was made, there was a brief (2000ms) pause during which the word “wait”
145 was displayed on the screen, followed by a screen revealing the allergy patient’s
146 state (i.e., with or without headache).

147 Each of 6 distinct trial types was presented 8 times in each of 4 sessions, for a total
148 of 192 trials, with sessions separated by 2-minute breaks, during which the scanner
149 was turned off. Each session was further divided into 4, non-delineated, blocks, with
150 each type of trial occurring once in random order in each block. All trials were
151 separated by a jittered 4-second ITI. In every other block, each trial (except for n-
152 trials) was followed by a query regarding the individual causal influence of the target
153 medicine present on that trial (i.e., C, D or S), or of medicine e or s’ on non-target
154 trials. Specifically, participants were asked to choose between the following three
155 options regarding the relevant medicine: “is a cause of headache”, “is not a cause
156 headache” and an ambivalent “may or may not be a cause of headache” (Figure
157 1B). Categorical response options, rather than the rating scales commonly
158 employed in the causal learning literature (Lu et al., 2008; Liljeholm & Cheng, 2009;
159 Liljeholm, 2015) were used for two main reasons: First, because the limited number
160 of response buttons in the scanner would have necessitated a scale slider, the initial

positioning of which might potentially bias ratings, and second, to reduce movement due to back-and-forth scale scrolling.

Computational models. Two formal accounts of predictive strength and uncertainty were implemented. First, an error-driven, associative learning algorithm updates predictive strength, on each trial, using the difference between the observed state of the outcome and the expected state based on all present cues:

$$V_{c,n+1} \leftarrow V_{c,n} + \alpha_{c,n} (\lambda - \sum_i V_i)_n \quad (1)$$

where V is the predictive strength of a particular cue, c , $\sum V_i$ is the summed strength of all cues present on trial n , and λ is the observed state of the outcome on that trial. The associability of cue c on trial n , $\alpha_{c,n}$, is defined as:

$$\alpha_{c,n} = \gamma \left| \lambda - \sum_i V_i \right|_{n-1} + (1 - \gamma) \alpha_{c,n-1} \quad (2)$$

where $n-1$ references to the previous trial involving cue c , and γ is a free parameter accounting for the weighting of that previous trial relative to preceding ones. In addition to scaling the influence of the prediction error on predictive strength, α is taken to reflect current levels of uncertainty about the cue's influence (e.g., Esberg & Haselgrove, 2011). The strengths and uncertainties associated with each cue were initialized to 0.5. On feedback-based trials with target causes, which always occurred in compound with an alternative cause, the α of the target cause (always greater than or equal to that of its compound counterpart) was used to indicate uncertainty, while

180 the summed strength, ΣV , was used to predict the outcome, and to compute the
181 prediction error.

182 A second formal account is provided by a Bayesian Causal Model, in which reasoners
183 make inferences over causal structures potentially responsible for the observed data
184 (Griffiths & Tenenbaum, 2005; Lu et al., 2008; Liljeholm, 2015). Here, as illustrated in
185 Figure 2, three possible graphs (*Graphs 0-2*) were defined for a candidate cause, C , an
186 alternative cause, A , and a constantly present background cause, B , where a causal
187 link to the effect exists for neither C nor A (G_0), C only (G_1), or both C and A (G_2).

Figure 2. Possible causal structures involving a background cause, B , a target cause, C , an alternative cause, A , and an effect, E . A link to the effect always exists for B , and may exist for neither C nor A ($Graph_0$), C only ($Graph_1$), or both C and A ($Graph_2$).

188
189 Each graph has a set of parameters θ , which are strengths w_i associated with causal
190 links in each graph (i.e., w_B , w_C and w_A for links associated with B , C and A respectively
191 in G_2). Sequential estimates of these parameters were modeled, for each candidate
192 cause, using the smallest possible *focal set* (Cheng & Novick, 1992) – a set of events
193 across which alternative causes can be assumed to occur with the same probability in
194 the presence and absence of the relevant cause. Thus, estimates of the strength of
195 the “no medicine” background cause was modeled using only n- trials in G_0 . Estimates
196 of elemental (non-target) candidate causes s' and e were respectively modeled based
197 on trials with those causes and the “no medicine” trials (i.e., n- and $s' \pm$ for candidate

s', and n- and e- for candidate e), using G_I . Finally, for each target candidate cause, which always occurred in compound with some alternative cause, estimation was modeled given G_2 , using the trials relevant for the particular target cause (i.e., n- and Cc' for target C; n-, e- and De+ for target D, and n-, e- and Se± for target S).

The likelihoods $P(d|\theta, G_i)$ were computed using a noisy-OR parameterization (Cheng, 1997). Specifically, summarizing data d by contingencies $N(e, c, a)$, the frequencies of each combination of the presence versus absence of the effect, target cause and alternative cause, the likelihood term for G_2 is:

$$P(d|w_B, w_C, w_A, G_2) = \left(\frac{N(c^-, a^-)}{N(e^+, c^-, a^-)} \right) \times \left(\frac{N(c^+, a^-)}{N(e^+, c^+, a^-)} \right) \times \left(\frac{N(c^-, a^+)}{N(e^+, c^-, a^+)} \right) \times \left(\frac{N(c^+, a^+)}{N(e^+, c^+, a^+)} \right) w_B^{N(e^+, c^-, a^-)} (1-w_B)^{N(e^-, c^-, a^-)} [w_B + w_C - w_B w_C]^{N(e^+, c^+, a^-)} [1-w_B - w_C + w_B w_C]^{N(e^-, c^+, a^-)} [w_B + w_A - w_B w_A]^{N(e^+, c^-, a^+)} [1-w_B - w_A + w_B w_A]^{N(e^-, c^-, a^+)} [w_B + w_C + w_A - w_B w_C - w_B w_A - w_C w_A + w_B w_C w_A]^{N(e^+, c^+, a^+)} [1-w_B - w_C - w_A + w_B w_C + w_B w_A + w_C w_A - w_B w_C w_A]^{N(e^-, c^+, a^+)}. \quad (3)$$

where $\binom{n}{k}$ denotes the number of ways of picking k unordered outcomes from n possibilities. $N(c^+)$ indicates the frequency of events in which the target cause is present, with analogous definitions for the other $N(.)$ terms. The likelihood terms for G_0 and G_I are similarly specified, where frequencies $N(.)$ are summed across the presence and absence of relevant events, and $w_i=0$ for any cause that does not have link to the effect in the relevant graph.

The marginal posterior distribution over strengths for a particular cause is obtained by applying Bayes' rule, and integrating out the parameters for other causes in the graph, such that, for G_2 ,

$$P(w_C | d, G_2) = \int_0^1 \int_0^1 \frac{P(d | w_B, w_C, w_A, G_2) P(w_B, w_C, w_A | G_2)}{P(d)} dw_B dw_A, \quad (4)$$

where $P(d | w_B, w_C, w_A, G_2)$ is the likelihood term, $P(w_B, w_C, w_A | G_2)$ refers to the prior probabilities of causal strength parameters, and $P(d)$ is the normalizing term, denoting the probability of the observed data.

Uncertainty about the strength of a particular cause, the focus of the study, was modeled as the Shannon entropy $H(w_C)$ of its marginal posterior distribution $P(w_C | d)$:

$$H(w_C) = - \int_0^1 P(w_C | d) \ln P(w_C | d) dw_C. \quad (5)$$

Point estimates of causal strength for each elemental and target cause were modeled as the mean of the relevant marginal posterior distribution (Lu et al., 2008; Liljeholm, 2015), given by:

$$\bar{w}_C = \int_0^1 w_C P(w_C | d) dw_C. \quad (6)$$

The model was sequentially implemented on a trial-by-trial basis, such that, for each cause, likelihoods were computed on each trial of feedback-based learning that yielded information relevant to that cause, using only the data point provided on that trial, with the posterior on the previous trial being used as the prior to obtain the posterior on the current trial. On the first trial in which data relevant to a particular candidate cause was presented, the priors were assigned independent uniform distributions. Finally, as an analog to the model-free prediction error, the Kullback-

Leibler (KL) divergence between the prior (Q) and posterior (P) on each trial, commonly referred to as “Bayesian surprise” (Itti & Baldi, 2006) was computed as:

$$KL(P\|Q) = \int_0^1 p(w_c | d) \log \frac{p(w_c | d)}{q(w_c | d)} dw_c. \quad (7)$$

As noted, target causes only occurred in compound with alternative causes during feedback-based learning. On such trials, the entropy and KL divergence of the marginal distribution for the target cause was used to indicate uncertainty and surprise respectively, while predictions regarding the occurrence of the effect were generated using a noisy-OR integration of causal strengths.

Both models assumed that, during feedback-based learning, participants selected “Yes”/“No” responses to questions about whether headache would occur using probabilities generated by a softmax distribution, in which a free noise parameter controls the influence of predictive strength (i.e., the strength with which the medicine(s) present on a given trial predicted headache; noisy-logical $P(e|w_i)$ and ΣV in causal and error-driven models respectively) on choice behavior. An additional noise parameter controlled the influence of uncertainty (i.e., α and $H(w_c)$ in error-driven and causal models respectively) on the proportion of ambivalent “May or may not be a cause” judgments during queries about the individual influence of each medicine. The first two instances of each trial type, and the first causal judgment for each medicine, were excluded from model fitting to eliminate transient noise due to task adjustment, and to minimize the influence of assumptions about priors. Free parameters were fit to behavioral data by minimizing the negative log likelihood of

255 observed responses for each individual and the Bayesian Information Criterion (BIC)
256 was used for model selection.

257 The model-derived probabilities of uncertain “May or may not be a cause” judgments
258 are plotted in Figure 3 for the causal and error-driven account of uncertainty
259 respectively. Note that, as was foreshadowed in the description of the study design,
260 the Bayesian causal model predicts high levels of uncertainty for both confounded, C,
261 and stochastic, S and s', causes, while the error-driven algorithm predicts that only
262 stochasticity will generate high levels of uncertainty.

Figure 3. Mean (normalized) probabilities of uncertain “May or may not be a cause” judgments derived from the Bayesian causal model and error-driven algorithm respectively. Confounded, deterministic and stochastic target causes (C, D & S) occurred only in compound with alternative causes during feedback-based learning, while non-target causes occurred only individually (s'), or individually as well as in compound (e). Error bars=SEM.

263

264 *Imaging procedure and analysis.* A 3 Tesla scanner (Phillips Achieva) was used to
 265 acquire structural T1-weighted images and T2*-weighted echoplanar images
 266 (repetition time = 2.65 s; echo time = 30 ms; flip angle = 90°; 45 transverse slices;
 267 matrix = 64 × 64; field of view = 192 mm; thickness = 3 mm; slice gap = 0 mm) with
 268 BOLD contrast. The first four volumes of images were discarded to avoid T1
 269 equilibrium effects. All remaining volumes were corrected for differences in the time of
 270 slice acquisition, realigned to the first volume, spatially normalized to the Montreal
 271 Neurological Institute (MNI) echoplanar imaging template, and spatially smoothed with
 272 a Gaussian kernel (8 mm, full width at half-maximum). High-pass filtering with a cutoff
 273 of 128 s was used. All effects were reported at a whole brain FWE cluster corrected
 274 threshold of $p < 0.05$, calculated using the Statistical nonParametric Mapping toolbox
 275 (SnPM13; <http://warwick.ac.uk/snpm>; Nichols & Holmes, 2002), with 5000
 276 permutations, no variance smoothing and an uncorrected height threshold of $p <$
 277 0.001 .

278 To address collinearity due to the shared prediction by causal and error-driven
 279 accounts that uncertainty increases with increased stochasticity, the two algorithms
 280 were analyzed using separate 1st level General Linear Models (GLM), and were
 281 directly contrasted at the group level using Bayesian Model Selection. Specifically, for
 282 each participant, two GLMs were specified, one for the causal and one for the error-
 283 driven computational model. In each such GLM, three regressors respectively
 284 modeled 1) the onset of the prediction screen on each trial during feedback-based
 285 learning, 2) the onset of the outcome screen on each trial during feedback-based
 286 learning, and 3) the onsets of the screens soliciting judgments about the influences of

individual medicines. In one GLM, prediction-screen onsets and causal judgment onsets were modulated by trial-by-trial error-driven estimates of uncertainty and strength, while outcome-screen onsets were modulated by trial-by-trial absolute values of the prediction errors. In the second GLM, prediction-screen and causal judgment onsets were modulated by trial-by-trial estimates of uncertainty and strength by the causal model while outcome-screen onsets were modulated by trial-by-trial estimates of Bayesian surprise. In both models, the first two instances of each trial type, and corollary causal judgments, were modeled by separate regressors, to eliminate transient noise due to task adjustment and reduce the influence of, non-modeled, prior beliefs. Moreover, in each model, two onset regressors respectively modeled the responses on feedback-based and causal query trials, three regressors indicated separate sessions, and six additional regressors accounted for the residual effects of head motion. Except for those indicating sessions and head motion, all regressors were convolved with a canonical hemodynamic response function. No orthogonalization was applied.

To directly contrast error-driven and causal accounts, BMS analyses was performed using a set of GLMs with the same onsets as those specified above, but with a single parametric modulator (e.g., error-driven uncertainty at prediction screen onsets) entered for each GLM. Thus, model comparisons isolated the relative contribution of a particular computational variable to neural activity. First-level Bayesian estimation procedure was used to compute a log-model evidence map for every subject and each GLM and inferences at the group level were modeled by applying a random effects approach (Rosa et al., 2010) at every voxel of the log evidence data. Group-

level exceedance probability maps, reflecting the probability that one model is more likely than the other, were thresholded to identify voxels in which the exceedance probability was greater than 0.95. Classical inferences were then assessed using BMS masks, within which a particular computational variable was more likely than its competing counterpart, at an exceedance probability greater than 0.95: For example, significant effects of error-driven uncertainty during the prediction period of feedback-based learning trials were reported only for those regions in which error-driven uncertainty during this event period was more likely than causal uncertainty during the same period, according to the BMS.

A pilot study on error-driven uncertainty. Previous research has implicated the anterior insula and dorsal medial frontal cortex in outcome stochasticity, using a range of decision-making tasks (Mohr et al., 2010; Volz et al., 2003; Huettel et al., 2005; Grinband et al., 2006). To ensure that such effects, used here as a benchmark against which to assess neural representations of uncertainty due to confounding, also emerge in our causal learning task, a pilot study (n=10) was conducted that was highly similar to that reported here, with the following exceptions: First, only stochasticity, not confounding, was manipulated across putative causes, second, both generative and preventive causal influences were included and, finally, judgments of individual causal influences were solicited at the end of, rather than throughout, each scanning session, and measured on a scale ranging from -100 (strongly removes headache) to +100 (strongly produces headache). The error-driven model of uncertainty was implemented as described above, fit to behavior during feedback-based learning, and regressed against the BOLD data. A single region of interest

(ROI) was constructed from the anterior insula and dorsal medial frontal cortex, using the “Willard” functional parcellation atlas (Richiardi et al., 2015). A power analysis performed with NeuroPower (<http://neuropowertools.org>; Durnez et al., 2015), using a screening threshold of $z=2.3$, revealed that a sample size of 16 would yield a power of 0.9 to detect effects of error-driven predictive uncertainty in this ROI at a Gaussian random field corrected alpha level of 0.05.

Results

Behavioral results

All statistical tests of behavioral data were planned comparisons, employing two-tailed t-tests, and were calculated using $n=19$. Confidence intervals (CI) and effect sizes (Cohen’s d) are reported for all comparisons. The distributions of causal judgments across response options are shown, with associated response times, for each putative cause in Table 1. Note that the proportion of ambivalent “May or may not be a cause” judgments (3rd row in Table 1) were greatest for the confounded cause (C), intermediate, although still substantial, for the two stochastic causes (S and s’), and virtually absent for deterministic causes (D and e), exactly as predicted by the causal, but not the error-driven, model (cfr Figure 3; see Liljeholm, 2015 for similar results). Note also that the distribution of “Is a cause” and “Is not a cause” judgments differed markedly across stochastic and confounded causes, such that participants almost never select the latter option for the confounded cause, consistent with the normative increase in the likelihood that target cause C is causal given Cc+ trials, but distributed

355 their responses fairly evenly across these two options for stochastic causes, reflecting,
356 perhaps, the outcome on trials immediately preceding each judgment.

357 Planned comparisons revealed that the mean proportion of ambivalent causal
358 judgments were significantly lower for the deterministic (D) target cause than for both
359 confounded (C), $t(18)=6.78$, $p<0.001$, $CI=[0.35\ 0.67]$, $d=1.79$, and stochastic (S),
360 $t(18)=4.09$, $p<0.001$, $CI=[0.14\ 0.44]$, $d=1.11$, target causes. The difference between
361 the confounded and stochastic cause was also significant, with the mean proportion of
362 ambivalent judgments being greater for C than for S $t(18)=2.65$, $p<0.05$, $CI=[0.05\ 0.40]$,
363 $d=0.68$. Recall that all target causes, including S, occurred only in compound
364 with alternative causes during feedback-based learning. The proportion of ambivalent
365 judgments for the two stochastic causes, one always occurring in isolation (s') and the
366 other always in compound (S), was marginally significant, $p=0.08$.

367 Judgment response times, averaged across response options for each cause, were
368 significantly faster for judgments about the deterministic target cause D than
369 stochastic target cause S, $t(18)=2.20$, $p<0.05$, $CI=[0.03\ 1.02]$, $d=0.35$, but did not differ
370 significantly between D and C, nor between C and S, $p>0.1$. Moreover, response
371 times did differ significantly across the two stochastic causes, being significantly
372 slower for target cause S, which never occurred in isolation during feedback-based
373 learning, than for cause s', $t(18)=3.42$, $p<0.005$, $CI=[0.29\ 1.21]$, $d=0.52$.

374 With respect to model fitting, the mean best fitting values of the free parameters for
375 the error driven model were 0.35 ± 0.27 for the weighting parameter, 3.25 ± 0.95 for the
376 prediction noise parameter, and 1.08 ± 0.88 for the judgment noise parameter. The

mean best fitting values of the free parameters for the causal model were 4.13 ± 1.44 for the prediction noise parameter and 1.03 ± 1.05 for the judgment noise parameter. The mean BIC was significantly lower, indicating superior performance, for the causal model (220.00 ± 44.15) than for the error-driven model (237.09 ± 35.58), $t(18) = 2.51$, $p < 0.05$, $CI = [2.81 \ 31.37]$, $d = 0.43$.

Neuroimaging results

All results reported below survived inclusive masking with voxels identified by Bayesian model selection analyses. A complete list of the non-masked significant effects of each model is provided in Tables 2 and 3. Bar graphs show effect sizes extracted from 8mm spheres centered on peak MNI coordinates, using rfxplot (Glascher, 2009).

Uncertainty signals during causal judgments. During causal queries, as participants made judgments about the causal influences of individual medicines, selective and significant effects of the causal model of uncertainty emerged in the medial and lateral superior frontal gyrus and the right dorsolateral prefrontal cortex (Figure 4). Note that the effect size of BOLD responses (bar graphs in Figure 4) across putative causes were greatest for the confounded cause (C), intermediate for stochastic causes (S and s'), and smallest for deterministic causes (D and e), exactly as predicted by the causal, but not the error-driven, model (cfr Figure 3). To further probe these effects, response times (RTs) were included as a parametric modulator of judgment uncertainty in 1st level, individual-subject, models, and as a between-subjects covariate in group-level analyses. Neither of these additional analyses revealed any

modulatory influence by RTs on the neural effects of the causal model of judgment
uncertainty. No effects of the error-driven measure of uncertainty during causal
judgments survived whole-brain correction.

Figure 4. Selective effects of the causal model of uncertainty during judgments about the causal influences of individual medicines, showing effects in the dorsomedial and dorsolateral prefrontal cortex (DMPFC and DLPFC), and the lateral Superior Frontal Gyrus (LSFG). Bar plots show effect sizes extracted from spheres centered on peak coordinates for each queried medicine. Error bars=SEM.

Uncertainty signals during prediction. During the prediction period of feedback-based learning trials, the error-driven, but not causal, measure of uncertainty was significantly positively correlated with activity throughout the inferior occipital and lingual gyri, as well as with activity in the supplementary motor area and right anterior

407 insula (Figure 5). Moreover, activity negatively correlated with error-driven predictive
408 uncertainty was observed in the posterior middle temporal gyrus.

Figure 5. Selective effects of the error-driven model of uncertainty during the prediction period of feedback-based learning trials, showing effects in the right anterior insula (aINS), posterior middle temporal gyrus (pMTG) and lingual gyrus (LG). Bar plots show effect sizes extracted from spheres centered on peak coordinates for each type of medicine trial. Error bars=SEM.

409

410 These effects of the error-driven account of uncertainty appear relatively robust, as
411 they closely mirror those from the pilot study, in which, at an uncorrected threshold of
412 0.005, extensive (cluster size > 100) effects of error-driven predictive uncertainty
413 emerged in the dorsal medial frontal cortex, bilateral anterior insula, and lingual gyrus

414 during feedback-based learning. In contrast to the widespread effects of error-driven
 415 predictive uncertainty during feedback-based learning, selective and significant effects
 416 of the causal measure of uncertainty emerged only in the right dorsal thalamus.

Figure 6. Strength predictions derived from the causal and error-driven model, together with effect sizes extracted from spheres centered on peak coordinates in the precentral gyrus, for each type of medicine trial during feedback-based learning. Error bars=SEM.

417
 418 *Strength & surprise signals.* While uncertainty in causal inferences, due to
 419 confounding or stochasticity, is of primary interest here, estimates of the strength with
 420 which a cause, or combination of causes, predicts the effect, and of the surprise at the
 421 outcome on each trial of feedback-based learning, were also modeled for
 422 completeness. During the prediction period of feedback-based learning trials, the
 423 causal measure of predictive strength was selectively and significantly positively
 424 correlated with activity in the calcarine and superior occipital gyrus, as well as

negatively correlated with activity in the left lateral superior frontal and precentral gyri (Figure 6). In contrast, a selective and significant negative correlation with the error-driven measure of predictive strength emerged in the posterior MTG. No significant effects of either the causal or error-driven account of predictive strength were found during causal judgments.

Figure 7. Outcome surprise signals derived from the causal and error-driven model (top), together with effect sizes (bottom) extracted from spheres centered on peak coordinates in the anterior caudate nucleus (aCN) and anterior insula (aINS), for each type of medicine trial during feedback-based learning. Error bars=SEM.

For measures of surprise during the outcome period of feedback-based learning trials, the absolute prediction error generated by the error-driven account was selectively

and significantly positively correlated with activity in the bilateral anterior insula, the medial and lateral dorsal prefrontal cortex and the inferior parietal gyrus. In addition, significant negative correlations with this measure were identified in the middle occipital gyrus, fusiform gyrus, the superior parietal lobule, precentral gyrus and calcarine sulcus. In contrast, a selective and significant negative correlation with the causal measure of (Bayesian) surprise was observed in the anterior caudate.

Discussion

Sensitivity to the “no confounding” constraint – a requirement that putative causes do not co-vary, fundamental to philosophical and statistical perspectives on causation – has been reliably demonstrated in a range of behavioral judgments by children as well as adults (Spellman, 1996; Kushnir and Gopnik, 2005; Meder et al., 2006; Schulz and Bonawitz, 2007; Schulz et al., 2007; Liljeholm, 2015). The current study used fMRI and cognitive computational modeling to investigate neural substrates mediating uncertainty associated with causal confounding. While being scanned, participants studied and judged the influences of various putative causes with confounded or non-confounded, deterministic or stochastic, influences. During judgments requiring generalization of causal knowledge from a feedback-based learning context to a transfer probe, activity in the dorsomedial prefrontal cortex (DMPFC) was better accounted for by a Bayesian causal model, sensitive to both confounding and stochasticity, than a purely error-driven algorithm, sensitive only to stochasticity.

The DMPFC has been implicated in encoding uncertainty across a variety of tasks (Volz et al., 2003; Grinband et al., 2006; Xue et al., 2008; Michael et al., 2015). For example, scanning participants with fMRI as they classified stimuli according to a

variable category boundary, Grinband et al. (2006) found that activity in the DMPFC scaled with the proximity of a stimulus to a category boundary. An important aspect of this and other demonstrations of the involvement of the DMPFC in uncertainty is that the level of uncertainty is directly related to the degree of experienced errors: that is, the greater the uncertainty on a given trial, the more likely it is that the prediction on that trial will be incorrect, as indicated by explicit feedback. In contrast, during feedback-based learning in the current task, the outcome on confounded trials was deterministic. Consequently, the increased activity in the DMPFC, as participants generated judgments about confounded cause, cannot be attributed to a history of errors. Instead, these results suggest that the DMPFC encodes a more abstract representation of uncertainty, divorced from the immediate consequences of choice, and irrespective of whether the specific source of uncertainty is confounding or stochasticity.

Several neuroeconomic studies (Hsu et al., 2005; Krain et al., 2006; Huettel et al., 2006; Bach et al., 2009; Levy et al., 2010; Pushkarskaya et al., 2015) have contrasted gambles involving stochasticity with “ambiguous” gambles, for which outcome probabilities are completely unknown, with some finding greater activity in the DMPFC in response to ambiguity than stochasticity (e.g., Hsu et al., 2005). One might argue that, in the current study, the influence of the confounded medicine is unknown, just as the outcome probabilities of ambiguous gambles in previous studies were unknown. But what does it mean to “know” something? Critically, all target medicines occurred only in compound with alternative medicines during feedback-based learning: consequently, when presented individually during causal queries, each was

479 equally novel and, presumably, equally unknown. In other words, differences in
 480 ambiguity between confounded and other target causes cannot be attributed to a
 481 generalization decrement based on changes in stimulus features. Moreover, ambiguity
 482 due to small samples and uncertainty due to confounding call for very different plans
 483 of exploratory action: while in former case, repeated observation of the same stimulus
 484 can resolve the uncertainty, the latter case requires an intervention that unconfounds
 485 the stimulus configuration. Further work is needed to assess the overlap between
 486 neural representations of confounding and ambiguity.

487 In addition to stochasticity and ambiguity, the DMPFC has been heavily implicated in
 488 cognitive control (Ridderinkhof et al., 2004; Fellows & Farah, 2005; Egner, 2009;
 489 Taren et al., 2011; Shenhav et al., 2013), raising the question of whether the currently
 490 observed effects in this region might reflect similar processes. In particular,
 491 judgments about individual causal influences may be associated with efforts to
 492 retrieve instances of the queried cause occurring in isolation from episodic memory –
 493 a search that is futile, and thus presumably more effortful, for confounded causes.
 494 However, such retrieval-induced control demands are unlikely to account for the
 495 current effects in the DMPFC for several reasons: First, effort associated with episodic
 496 memory retrieval does not address responses to stochasticity – in contrast, the causal
 497 uncertainty account explains why DMPFC responses are greater for *both* confounded
 498 and stochastic causes, relative to non-confounded and deterministic causes. It may
 499 be argued, of course, that stochasticity elicits additional control processes mediated
 500 by the DMPFC, such as, for example, response conflict (Botnivick et al., 2001;
 501 Ridderinkhof et al., 2004; Kerns et al., 2004). Recall, however that *none* of the target

causes (C, S and D) were presented in isolation during feedback-based learning and that, consequently, *all* target causes should involve some level of searching for non-presented information during causal judgments. If DMPFC activity reflected this search process as well as response conflict due to stochasticity, one would expect a difference between neural responses to causes S and s', both of which are stochastic, but only one of which, s', was presented in isolation during feedback-based learning, eliminating the need to search for an un-experienced memory. Notably, while RTs (Table 1), commonly used as a measure of cognitive load, are indeed consistent with the notion of an increased control demand for cause S over s', the DMPFC responses (Figure 4) are not. Finally, and perhaps most conclusively, the inclusion of RTs as a parametric modulator of judgment uncertainty in 1st level models, and as a covariate in group analyses, did not have any impact on the neural effects of model-based judgment uncertainty. Taken together, these aspects of the results make causal uncertainty the more parsimonious explanation of the DMPFC effects observed here.

As with the DMPFC and DLPFC, the anterior insula has been implicated in outcome stochasticity, or "risk", in several previous neuroimaging studies (e.g., Volz et al., 2003; Huettel et al., 2005; d'Acremont et al., 2013; Preuschoff et al., 2008; Critchley et al., 2001). Here, the dorsal anterior right insula was selectively recruited by the error-driven model of uncertainty during the prediction period of feedback-based learning trials. Plots of effect sizes for each type of trial (in Figure 5) suggest that these responses were indeed driven by stochasticity-based uncertainty about the state of the outcome on any given trial. Moreover, the fact that, unlike the DMPFC, activity in the anterior insula did not scale with either an error-driven or causal measure of

uncertainty during judgments about individual causal influences indicates that representations in this region pertain primarily to the immediate consequences of choice, rather than generalizable knowledge. Consistent with this interpretation, the bilateral anterior insula also responded selectively to stochastic conditions during the outcome period of feedback-based learning trials, suggesting that this region is recruited by both the anticipation and detection of errors. While, as noted, it is difficult to discern based on the current results whether the unsigned error signals observed in the anterior insula are Bayesian or model-free, they are consistent with a substantial literature implicating this region in expectancy violations across a wide range of tasks (Klein et al., 2007; Metereau et al., 2012; Allen et al., 2016; Bastin et al., 2016).

Selective effects of the error-driven account of predictive uncertainty were also identified throughout large portions of the posterior lingual gyrus. In a recent neuroimaging study, Causse et al., (2013) assessed decision making under uncertainty using an aviation task in which participants had to either land a plane or abort landing, given certain (100% or 0%) vs. uncertain (50%) information regarding landing conditions. Consistent with the current results, they found greater activity in the posterior lingual gyrus during high than during low uncertainty conditions. Our results are also consistent with those of Payzan-LeNestour et al. (2016), who found that the lingual gyrus scaled specifically with uncertainty due to outcome stochasticity, as opposed to uncertainty due to changes in the statistical structure of the environment or due to ambiguity. Payzan-LeNestour et al. argued that the lingual gyrus might not typically emerge in studies assessing outcome stochasticity because those studies did not dissociate stochasticity from other uncertainty components. A

similar case can be made here, that the inclusion of an alternative, generative, model that captures additional aspects of uncertainty may have increased sensitivity to detect stochasticity-specific signals in the lingual gyrus.

An important aspect of the current study is the direct comparison of a Bayesian inference model with an error-driven, model-free, algorithm. While not uncommon (Courville et al, 2006; Hampton et al., 2006; Payzan-LeNestour & Bossaerts, 2011; Prevost et al., 2013), the contrasting of these quite different computational frameworks somewhat complicates interpretation. In particular, it raises the question of whether selective effects of the causal account of uncertainty indeed reflect sensitivity to confounding, or instead some extraneous, perhaps incidental, feature of the different modeling frameworks. Here, a significant clue is provided by neural and behavioral responses to each queried putative cause, shown in Figure 4 & Table 1 respectively, which clearly correspond to the increased uncertainty associated with confounded and stochastic causes predicted by the causal model (cfr Figure 3). It is worth noting that, far from incidental, the failure of the error-driven account to predict sensitivity to confounding is intrinsic to the model-free approach, due to its lack of a representation of the independence of causal influences. Critically, a configural solution (e.g., Pearce, 1987; 2002), such that individually queried causes are treated as non-overlapping with the compounds in which they occur during feedback-based learning, would fail to account for both behavioral and neural differences between confounded and deterministic target causes, both of which occurred only in compound with alternative causes during feedback-based learning.

570 While the effects of causal uncertainty in the DMPFC likely reflect sensitivity to causal
 571 confounding, other neuroimaging results are clearly unrelated and, in some cases,
 572 largely incidental to the specific model implementation. In particular, as can be seen
 573 in Figures 6 and 7, the ordinal mean values derived from causal and error-driven
 574 accounts across trial types during feedback-based learning are identical for both
 575 strength and surprise measures, suggesting that model-selection may reflect more
 576 granular differences. For example, because of the Σ term in Equations 1 and 2 of the
 577 error-driven account, the variance in the occurrence of the outcome across e-, De+
 578 and Se± trials results in a constantly fluctuating strength, and persistent prediction
 579 errors, even on deterministic e- and De+, trials. In contrast, in the implementation of
 580 the causal model, such non-conditional variance is eliminated by the use of a focal set
 581 (e.g., Cheng & Novick, 1992; Spellman, 1996). Behaviorally, the focal set assumption
 582 is strongly supported by the lack of variability in judgments about non-target cause e.
 583 This cause, which is paired with the outcome on half of trials in which it occurs, but
 584 never when occurring in isolation, is almost exclusively judged to be non-causal,
 585 suggesting that participants conditioned their inferences about its influence on trials
 586 across which alternative causes could be assumed to be held constant. Nonetheless,
 587 at the neural level, several regions may instead have been tracking the unconditional
 588 outcome variance computed by the error-driven model.

589 As the mantra “covariation does not equal causation” implies, requirements for causal
 590 induction extend far beyond mere statistical regularity. Countless studies have
 591 demonstrated neural and behavioral effects of error-driven uncertainty, whether due to
 592 stochasticity, insufficient samples or changes in the statistical structure of the

environment. The current study reveals that, given violation of a basic boundary condition on causal inference, and in spite of the same constraints on perceptual generalization, differences in neural representations of uncertainty can emerge across equally deterministic, static and well-sampled contingencies. These results contribute to a growing literature on the role of generative models in neural representations of uncertainty, and shed light on a possible neural implementation of domain general, a priori, constraints on causal induction.

501

502 **Acknowledgements**

503 This work was supported by a start-up fund from the University of California Irvine to
504 M.L.

505

506

507

508

References

- 509 Abler, B., Herrnberger, B., Grön, G., & Spitzer, M. (2009). From uncertainty to reward:
510 BOLD characteristics differentiate signaling pathways. *BMC neuroscience*, *10*(1),
511 154.
- 512 Allen, M., Fardo, F., Dietz, M. J., Hillebrandt, H., Friston, K. J., Rees, G., & Roepstorff,
513 A. (2016). Anterior insula coordinates hierarchical processing of tactile mismatch
514 responses. *Neuroimage*, *127*, 34-43.
- 515 Bach, D. R., Seymour, B., & Dolan, R. J. (2009). Neural activity associated with the
516 passive prediction of ambiguity and risk for aversive events. *Journal of*
517 *Neuroscience*, *29*(6), 1648-1656.
- 518 Baetens, K., Ma, N., Steen, J., & Van Overwalle, F. (2013). Involvement of the
519 mentalizing network in social and non-social high construal. *Social cognitive and*
520 *affective neuroscience*, *9*(6), 817-824.
- 521 Baetens, K. L., Ma, N., & Van Overwalle, F. (2017). The dorsal medial prefrontal
522 cortex is recruited by high construal of non-social stimuli. *Frontiers in behavioral*
523 *neuroscience*, *11*, 44.
- 524 Bastin, J., Deman, P., David, O., Gueguen, M., Benis, D., Minotti, L., ... & Kahane, P.
525 (2016). Direct recordings from human anterior insula reveal its leading role within
526 the error-monitoring network. *Cerebral Cortex*, bhv352.
- 527 Causse, M., Péran, P., Dehais, F., Caravasso, C. F., Zeffiro, T., Sabatini, U., & Pastor,
528 J. (2013). Affective decision making under uncertainty during a plausible aviation
529 task: An fMRI study. *NeuroImage*, *71*, 19-29.

530 Cheng P (1997) From covariation to causation: A causal power theory. *Psychological*
531 *Review*:367-405.

532 Cheng, P. W., & Novick, L. R. (1992). Covariation in natural causal induction.
533 *Psychological Review*, 99(2), 365.

534 Courville, A. C., Daw, N. D., & Touretzky, D. S. (2006). Bayesian theories of
535 conditioning in a changing world. *Trends in cognitive sciences*, 10(7), 294-300.

536 Critchley, H. D., Mathias, C. J., & Dolan, R. J. (2001). Neural activity in the human
537 brain relating to uncertainty and arousal during anticipation. *Neuron*, 29(2), 537-
538 545.

539 d'Acremont, M., Fornari, E., & Bossaerts, P. (2013). Activity in inferior parietal and
540 medial prefrontal cortex signals the accumulation of evidence in a probability
541 learning task. *PLoS computational biology*, 9(1), e1002895.

542 Durnez, J., Degryse, J., Seurinck, R., Moerkerke, B., & Nichols, T. E. (2015).
543 Prospective power estimation for peak inference with the toolbox neuropower.
544 In *Second Belgian Neuroinformatics Congress* (Vol. 9). Frontiers Media SA.

545 Esber, G. R., & Haselgrove, M. (2011). Reconciling the influence of predictiveness
546 and uncertainty on stimulus salience: a model of attention in associative
547 learning. *Proceedings of the Royal Society of London B: Biological*
548 *Sciences*, 278(1718), 2553-2561.

549 Egner, T. (2009). Prefrontal cortex and cognitive control: motivating functional
550 hierarchies. *Nature neuroscience*, 12(7), 821.

551 Fellows, L. K., & Farah, M. J. (2005). Is anterior cingulate cortex necessary for
552 cognitive control?. *Brain*, 128(4), 788-796.

553 Forman, S. D., Cohen, J. D., Fitzgerald, M., Eddy, W. F., Mintun, M. A., & Noll, D. C.
554 (1995). Improved assessment of significant activation in functional magnetic
555 resonance imaging (fMRI): use of a cluster-size threshold. *Magnetic Resonance in*
556 *medicine*, 33(5), 636-647.

557 Glymour C (1998) Psychological and normative theories of causal power and the
558 probabilities of causes. In: *Uncertainty in artificial intelligence* (F. CG, S. M, eds):
559 Kaufmann.

560 Gläscher, J. (2009) Visualization of group inference data in functional neuroimaging.
561 *Neuroinformatics*, 7(1), 73-82.

562 Griffiths, T. L., & Tenenbaum, J. B. (2005). Structure and strength in causal
563 induction. *Cognitive psychology*, 51(4), 334-384.

564 Grinband, J., Hirsch, J., & Ferrera, V. P. (2006). A neural representation of
565 categorization uncertainty in the human brain. *Neuron*, 49(5), 757-763.

566 Holt, C. A., & Laury, S. K. (2002). Risk aversion and incentive effects. *American*
567 *economic review*, 92(5), 1644-1655.

568 Hampton, A. N., Bossaerts, P., & O'doherty, J. P. (2006). The role of the ventromedial
569 prefrontal cortex in abstract state-based inference during decision making in
570 humans. *Journal of Neuroscience*, 26(32), 8360-8367.

571 Huettel, S. A., Stowe, C. J., Gordon, E. M., Warner, B. T., & Platt, M. L. (2006).
572 Neural signatures of economic preferences for risk and ambiguity. *Neuron*, 49(5),
573 765-775.

574 Huettel SA, Song AW, McCarthy G. (2005) Decisions under uncertainty: probabilistic
575 context influences activity of prefrontal and parietal cortices. *J Neurosci.* 25:3304–
576 3311

577 Hsu, M., Bhatt, M., Adolphs, R., Tranel, D., & Camerer, C. F. (2005). Neural systems
578 responding to degrees of uncertainty in human decision-
579 making. *Science*, 310(5754), 1680-1683.

580 Itti, Laurent, and Pierre F. Baldi. "Bayesian surprise attracts human
581 attention." *Advances in neural information processing systems*. 2006.

582 Kerns, J. G., Cohen, J. D., MacDonald, A. W., Cho, R. Y., Stenger, V. A., & Carter, C.
583 S. (2004). Anterior cingulate conflict monitoring and adjustments in
584 control. *Science*, 303(5660), 1023-1026.

585 Klein, T. A., Endrass, T., Kathmann, N., Neumann, J., von Cramon, D. Y., &
586 Ullsperger, M. (2007). Neural correlates of error awareness. *Neuroimage*, 34(4),
587 1774-1781.

588 Krain, A. L., Wilson, A. M., Arbuckle, R., Castellanos, F. X., & Milham, M. P. (2006).
589 Distinct neural mechanisms of risk and ambiguity: a meta-analysis of decision-
590 making. *Neuroimage*, 32(1), 477-484.

591 Kushnir T, Gopnik A (2005) Young children infer causal strength from probabilities and
592 interventions. *Psychological science* 16:678-683.

593 Levy, I., Snell, J., Nelson, A. J., Rustichini, A., & Glimcher, P. W. (2010). Neural
594 representation of subjective value under risk and ambiguity. *Journal of*
595 *neurophysiology*, 103(2), 1036-1047.

596 Liljeholm, M., & Cheng, P. W. (2009). The influence of virtual sample size on
597 confidence and causal-strength judgments. *Journal of Experimental Psychology:*
598 *Learning, Memory, and Cognition*, 35(1), 157.

599 Liljeholm, M. (2015). How multiple causes combine: independence constraints on
700 causal inference. *Frontiers in psychology*, 6.

701 Lu, H., Yuille, A. L., Liljeholm, M., Cheng, P. W., & Holyoak, K. J. (2008). Bayesian
702 generic priors for causal learning. *Psychological review*, 115(4), 955.

703 Botvinick, M. M., Braver, T. S., Barch, D. M., Carter, C. S., & Cohen, J. D. (2001).
704 Conflict monitoring and cognitive control. *Psychological review*, 108(3), 624.

705 Meder B, Hagmayer Y, Waldmann M (2006) Understanding the Causal Logic of
706 Confounds. In: Twenty-Eighth Annual Conference of the Cognitive Science
707 Society, pp 579-584: Erlbaum.

708 Metereau, E., & Dreher, J. C. (2012). Cerebral correlates of salient prediction error for
709 different rewards and punishments. *Cerebral Cortex*, 23(2), 477-487.

710 Mohr, P. N., Biele, G., & Heekeren, H. R. (2010). Neural processing of risk. *Journal of*
711 *Neuroscience*, 30(19), 6613-6619.

712 Nichols, T. E., & Holmes, A. P. (2002). Nonparametric permutation tests for functional
713 neuroimaging: a primer with examples. *Human brain mapping*, 15(1), 1-25.

714 Payzan-LeNestour, E., Dunne, S., Bossaerts, P., & O'Doherty, J. P. (2013). The
715 neural representation of unexpected uncertainty during value-based decision
716 making. *Neuron*, 79(1), 191-201.

717 Payzan-LeNestour, E., & Bossaerts, P. (2011). Risk, unexpected uncertainty, and
718 estimation uncertainty: Bayesian learning in unstable settings. *PLoS computational*
719 *biology*, 7(1), e1001048.

720 Pearce, J. M. (1987). A model for stimulus generalization in Pavlovian
721 conditioning. *Psychological review*, 94(1), 61.

722 Pearce, J. M. (2002). Evaluation and development of a connectionist theory of
723 configural learning. *Animal Learning & Behavior*, 30(2), 73-95.

724 Preuschoff, K., Quartz, S. R., & Bossaerts, P. (2008). Human insula activation reflects
725 risk prediction errors as well as risk. *Journal of Neuroscience*, 28(11), 2745-2752.

726 Prévost, C., McNamee, D., Jessup, R. K., Bossaerts, P., & O'Doherty, J. P. (2013).
727 Evidence for model-based computations in the human amygdala during Pavlovian
728 conditioning. *PLoS computational biology*, 9(2), e1002918.

729 Pushkarskaya, H., Smithson, M., Joseph, J. E., Corbly, C., & Levy, I. (2015). Neural
730 correlates of decision-making under ambiguity and conflict. *Frontiers in behavioral*
731 *neuroscience*, 9, 325.

732 Richiardi, J., Altmann, A., Milazzo, A. C., Chang, C., Chakravarty, M. M.,
733 Banaschewski, T., ... & Conrod, P. (2015). Correlated gene expression supports
734 synchronous activity in brain networks. *Science*, 348(6240), 1241-1244.

735 Ridderinkhof, K. R., Ullsperger, M., Crone, E. A., & Nieuwenhuis, S. (2004). The role
736 of the medial frontal cortex in cognitive control. *science*, 306(5695), 443-447.

737 Rosa, M. J., Bestmann, S., Harrison, L., & Penny, W. (2010). Bayesian model
738 selection maps for group studies. *Neuroimage*, 49(1), 217-224.

739 Schulz L, Bonawitz E (2007) Serious fun: preschoolers engage in more exploratory
740 play when evidence is confounded. *Developmental psychology* 43:1045-1050.

741 Schulz L, Gopnik A, Glymour C (2007) Preschool children learn about causal structure
742 from conditional interventions. *Dev Sci* 10:322-332.

743 Shenhav, A., Botvinick, M. M., & Cohen, J. D. (2013). The expected value of control:
744 an integrative theory of anterior cingulate cortex function. *Neuron*, 79(2), 217-240.

745 Spellman B (1996) Acting as Intuitive Scientists: Contingency Judgments Are Made
746 While Controlling for Alternative Potential Causes. *Psychological science* 7:337-
747 342.

748 Taren, A. A., Venkatraman, V., & Huettel, S. A. (2011). A parallel functional
749 topography between medial and lateral prefrontal cortex: evidence and
750 implications for cognitive control. *Journal of Neuroscience*, 31(13), 5026-5031.

751 Volz KG, Schubotz RI, von Cramon DY. (2003) Predicting events of varying
752 probability: uncertainty investigated by fMRI. *Neuroimage*. ;19:271–280.

753 Xue, G., Lu, Z., Levin, I. P., Weller, J. A., Li, X., & Bechara, A. (2008). Functional
754 dissociations of risk and reward processing in the medial prefrontal
755 cortex. *Cerebral Cortex*, 19(5), 1019-1027.

756
757
758
759

Figure Captions

Figure 1. Task illustration. A. Prediction and outcome screens from a feedback-based learning trial. Screens were separated by a 2 sec. pause during which the word “Wait” (not shown in figure) was displayed. B. Causal query regarding the individual influence of a particular medicine on headache.

Figure 2. Possible causal structures involving a background cause, B, a target cause, C, an alternative cause, A, and an effect, E. A link to the effect always exists for B, and may exist for neither C nor A (Graph₀), C only (Graph₁), or both C and A (Graph₂).

Figure 3. Mean (normalized) probabilities of uncertain “May or may not be a cause” judgments derived from the Bayesian causal model and error-driven algorithm respectively. Confounded, deterministic and stochastic target causes (C, D & S) occurred only in compound with alternative causes during feedback-based learning, while non-target causes occurred only individually (s’), or individually as well as in compound (e). Error bars=SEM.

Figure 4. Selective effects of the causal model of uncertainty during judgments about the causal influences of individual medicines, showing effects in the dorsomedial and dorsolateral prefrontal cortex (DMPFC and DLPFC), and the lateral Superior Frontal Gyrus (LSFG). Bar plots show effect sizes extracted from spheres centered on peak coordinates for each queried medicine. Error bars=SEM.

Figure 5. Selective effects of the error-driven model of uncertainty during the prediction period of feedback-based learning trials, showing effects in the right anterior

781 insula (aINS), posterior middle temporal gyrus (pMTG) and lingual gyrus (LG). Bar
782 plots show effect sizes extracted from spheres centered on peak coordinates for each
783 type of medicine trial. Error bars=SEM.

784 *Figure 6.* Strength predictions derived from the causal and error-driven model (top),
785 together with effect sizes (bottom) extracted from spheres centered on peak
786 coordinates in the precentral gyrus, for each type of medicine trial during feedback-
787 based learning. Error bars=SEM.

788 *Figure 7.* Outcome surprise signals derived from the causal and error-driven model
789 (top), together with effect sizes (bottom) extracted from spheres centered on peak
790 coordinates in the anterior caudate nucleus (aCN) and anterior insula (aINS), for each
791 type of medicine trial during feedback-based learning. Error bars=SEM.

792

Table 1. Mean proportions of “Is a cause”, “Is not a cause” and “May or may not be a cause” judgments, and associated response times, with standard deviations, for target causes, C, D and S, and non-target causes, s’ and e’.

	C	D	S	s’	e
<i>Proportions</i>					
Is a cause	0.36±0.33	0.84±0.25	0.38±0.25	0.40±0.27	0.02±0.03
Is not a cause	0.03±0.04	0.06±0.09	0.23±0.11	0.30±0.17	0.96±0.05
May be a cause	0.61±0.35	0.10±0.21	0.39±0.31	0.30±0.28	0.02±0.04
<i>Response Times</i>					
Is a cause	2.92±1.88	2.52±1.23	4.03±3.62	2.82±2.59	2.86±2.11
Is not a cause	4.26±2.69	3.94±3.36	4.19±3.64	2.72±1.23	2.44±1.15
May be a cause	3.79±2.68	3.76±1.46	2.99±1.94	2.59±1.32	4.08±4.78

Table 2: Non-masked significant, whole-brain corrected, neural effects of the causal model of uncertainty, strength and surprise, during causal judgments and feedback-based learning, with peak MNI coordinates and cluster sizes at $p < 0.001$. R/L = Right/Left, (-) = negative correlation.

Contrast	Region	Peak MNI	Cluster Size
Judgment Uncertainty	DMPFC	-8, 28, 46	1900
	R. IPG	50, -52, 40	292
	Dorsolateral PFC	42, 34, 30	201
Prediction Uncertainty	Fusiform	28, -72, -12	7195
	DMPFC	-10, 16, 52	1387
	R. Insula	-30, 18, 4	726
	L. Insula	34, 30, -4	470
	Thalamus	6, -22, 2	867
Prediction Strength	Calcarine/Occipital	18, -96, 4	236
Prediction Strength (-)	Lingual Gyrus	14, -62, -2	2232
	Superior Frontal Gyrus	-26, -8, 58	458
	Superior Parietal Lobule	-28, -50, 62	304
	Inferior Parietal Gyrus	-46, -26, 36	336
	Supp. Motor Area	8, 2, 52	297
Surprise	Right Insula	36, 18, -2	652
	Left Insula	-28, 22, -6	482
	Dorsomedial PFC	4, 46, 34	1146
	Left Dorsolateral PFC	-46, 4, 42	192
	Right Dorsolateral PFC	44, 24, 30	325
Surprise (-)	Supramarginal Gyrus	-50, -26, 38	1858
	Caudate	18, 32, 0	505
	Mid Cingulum	-10, -30, 42	472
	Postcentral Gyrus	50, -20, 48	2399
	Parahippocampal	34, -32, -12	227
	Precuneus	16, -52, 14	221

Table 3: Non-masked significant, whole-brain corrected, neural effects of the error-driven model of uncertainty, strength and surprise during feedback-based learning, with peak MNI coordinates and cluster sizes at $p < 0.001$. R/L = Right/Left, (-) = negative correlation.

Contrast	Region	Peak MNI	Cluster Size
Prediction Uncertainty	Calcarine/Lingual Gyrus	14, -88, 0	3116
	Supp. Motor Area	10, 20, 48	153
	Left Insula	-24, 22, 2	65
	Right Insula	34, 28, -2	109
Prediction Uncertainty (-)	Middle Temporal Gyrus	-46, -62, 20	341
	Middle Temporal Gyrus	-56, -22, -10	84
	Ventromedial PFC	-2, 30, -14	157
	Middle Temporal Gyrus	46, -58, 22	64
Prediction Strength	Lingual Gyrus	-20, -78, -10	2801
Prediction Strength (-)	Superior Parietal Lobule	-18, -52, 64	407
	Supramarginal Gyrus	-60, -28, 30	390
	Precentral Gyrus	26, -12, 50	240
	Supramarginal Gyrus	58, -38, 28	728
	Middle Temporal Gyrus	-54, -60, 2	603
	Precentral Gyrus	-22, -14, 64	383
	Thalamus	-12, -10, -4	214
Prediction Error	Right Insula	38, 18, -4	912
	Left Insula	-40, 20, 4	945
	Precentral Gyrus	-42, -2, 40	713
	Anterior Cingulate	4, 38, 30	2021
	Inferior Parietal Gyrus	-28, -52, 42	246
	Dorsolateral PFC	44, 28, 26	885
Prediction Error (-)	Middle Occipital Gyrus	-42, -74, 26	735
	Fusiform Gyrus	-34, -42, -12	253
	Postcentral Gyrus	54, -26, 52	906